

Semidefinite Programming and Integer Programming

Monique Laurent * and Franz Rendl **

* CWI, Kruislaan 413, 1098 SJ Amsterdam, The Netherlands

monique@cwi.nl

** Universität Klagenfurt, Institut für Mathematik

Universitätstrasse 65-67, A-9020 Klagenfurt, Austria

franz.rendl@uni-klu.ac.at

December 17, 2002

Contents

1	Introduction	3
2	Semidefinite Programming: Duality, Algorithms, Complexity, and Geometry	3
2.1	Duality	3
2.2	Algorithms	7
2.3	Complexity	9
2.4	Geometry	11
3	Semidefinite Programming and Integer 0/1 Programming	15
3.1	A general paradigm	15
3.2	Introduction on cutting planes and lift-and-project methods	17
3.3	A canonical lifting construction	19
3.4	The Balas-Ceria-Cornuéjols, Lovász-Schrijver, Serali-Adams, and Lasserre methods .	20
3.5	Application to the stable set problem	24
3.6	Application to the max-cut problem	25
3.7	Further results	27
3.8	Extensions to polynomial programming	29
4	Semidefinite Relaxation for the Maximum Stable Set Problem	32
4.1	The basic linear relaxation	32
4.2	The theta function $\vartheta(G)$ and the basic semidefinite relaxation $\text{TH}(G)$	33
4.3	Colouring and finding maximum stable sets in perfect graphs	36
4.4	Sharpening the theta function	37
5	Semidefinite Relaxation for the Max-Cut Problem	39
5.1	The basic linear relaxation	39
5.2	The basic semidefinite relaxation	40
5.3	The Goemans-Williamson randomized approximation algorithm for max-cut	42
5.4	How to improve the Goemans-Williamson algorithm?	44

6	Applications of Semidefinite Programming and the Rounding Hyperplane Technique to Other Combinatorial Optimization Problems	47
6.1	Approximating quadratic programming	47
6.2	Approximating the maximum bisection problem	49
6.3	Approximating the max k -cut problem	52
6.4	Approximating graph colouring	54
6.5	Approximating the maximum stable set and vertex cover problems	60
6.6	Approximating MAX SAT	63
6.7	Approximating the maximum directed cut problem	69
7	Further Topics	72
7.1	Approximating polynomial programming using semidefinite programming	72
7.2	Approximating combinatorial problems using copositive programming	75
8	Semidefinite Programming and the Quadratic Assignment Problem	78
9	Epilogue: Semidefinite Programming and Algebraic Connectivity	82
10	Appendix: Surveys, Books and Software	85

1 Introduction

Linear optimization is a relatively young area of applied mathematics. Even though the world is nonlinear, as physicists never stop to point out, it seems that in many practical situations a linearized model describes key features of a problem quite accurately.

The success of linear optimization in many real-world applications has led to the study of integer linear programming, which permits to model optimal decision making under finitely many alternatives. A natural way to approach these types of problems consists in using again linear theory, in this case polyhedral combinatorics, to solve them. Mathematically, one tries to find (at least) a (partial) linear description of the convex hull of all integral solutions. While this approach was successful for many combinatorial optimization problems, it turned out that some graph optimization problems, such as Max-Cut or Max-Clique, can not be approximated tightly by purely linear methods.

Stronger relaxation methods have therefore attracted the focus of recent research. The extension of linear optimization to semidefinite optimization has turned out to be particularly interesting for the following reasons. First, algorithmic ideas can be extended quite naturally from linear to semidefinite optimization. Secondly, there is theoretical evidence that semidefinite models are sometimes significantly stronger than purely linear ones, justifying the computational overhead to solve them.

It is the purpose of this chapter to explain in detail how semidefinite programming is used to solve integer programming problems. Specifically, we start out in the next section with explaining the relevant mathematical background underlying semidefinite programming by summarizing the necessary duality theory, explaining algorithmic ideas and recalling computational complexity results related to semidefinite programming. In Section 3 we show how semidefinite relaxations arise from integer 0/1 programming by lifting the problem formulated in \mathbb{R}^n to a problem in the space of symmetric matrices.

A detailed study of two prominent special graph optimization problems follows in Section 4, dealing with the stable set problem, and Section 5, devoted to Max-Cut. For both these problems the extension of polyhedral to semidefinite relaxations has led to a significant improvement in the approximation of the original problem. Section 5 also introduces the hyperplane rounding idea of Goemans and Williamson, which opened the way to many other approximation approaches, many of which are discussed in Section 6.

Section 7 discusses possible alternatives to the use of semidefinite models to get stronger relaxations of integer programs.

Finally, we summarize in Section 8 some recent semidefinite and other nonlinear relaxations applied to the Quadratic Assignment Problem, which have led to a computational break-through in Branch and Bound computations for this problem.

2 Semidefinite Programming: Duality, Algorithms, Complexity, and Geometry

2.1 Duality

To develop a duality theory for semidefinite programming problems, we take a more general point of view, and look at *Linear Programs over Cones*.

Suppose K is a closed convex cone in \mathbb{R}^n , $c \in \mathbb{R}^n$, $b \in \mathbb{R}^m$ and A is an $m \times n$ matrix. The problem

$$p^* := \sup\{c^T x : Ax = b, x \in K\} \tag{1}$$

is called **Cone-LP**, because we optimize a linear function subject to linear equations, and we have the condition that the decision variable x lies in the cone K .

The dual cone K^* is defined as follows:

$$K^* := \{y \in \mathbb{R}^n : y^T x \geq 0 \forall x \in K\}.$$

It is a well known fact, not hard to verify, that K^* is also a closed convex cone.

We will derive the dual of (1) by introducing Lagrange multipliers for the equality constraints and using the Minimax Inequality. Let $y \in \mathbb{R}^m$ denote the Lagrange multipliers for $Ax = b$. Using the Lagrangian $L(x, y) := c^T x + y^T (b - Ax)$ we get

$$\inf_y L(x, y) = \begin{cases} c^T x & \text{if } Ax = b \\ -\infty & \text{otherwise.} \end{cases}$$

Therefore,

$$p^* = \sup_{x \in K} \inf_y L(x, y) \leq \inf_y \sup_{x \in K} L(x, y).$$

The inequality is usually called ‘Minimax inequality’, and holds for any real-valued function $L(x, y)$ where x and y are from some ground sets X and Y , respectively.

We can rewrite L as $L = b^T y - x^T (A^T y - c)$. The definition of K^* implies the following. If $A^T y - c \notin K^*$ then there exists $x \in K$ such that $x^T (A^T y - c) < 0$. Therefore we conclude

$$\sup_{x \in K} L(x, y) = \begin{cases} b^T y & \text{if } A^T y - c \in K^* \\ \infty & \text{otherwise.} \end{cases}$$

This translates into

$$p^* \leq \inf \{b^T y : y \in \mathbb{R}^m, A^T y - c \in K^*\} =: d^*. \quad (2)$$

The problem on the right side of the inequality sign is again a Cone-LP, but this time over the cone K^* . We call this problem the dual to (1). By construction, a pair of dual cone-LP satisfies weak duality.

Lemma 1. (Weak Duality) *Let $x \in K, y \in \mathbb{R}^m$ be given with $Ax = b, A^T y - c \in K^*$. Then, $c^T x \leq b^T y$.*

One crucial issue in duality theory consists in identifying sufficient conditions that insure equality in (2), also called *Strong Duality*. The following condition insures strong duality. We say that the cone-LP (1) satisfies the *Slater constraint qualification* if there exists $x \in \text{int}(K)$ such that $Ax = b$. (A similar definition holds for the dual problem.) Duffin [70] shows the following result.

Theorem 2. *If (1) satisfies the Slater constraint qualification and p^* is finite, then $p^* = d^*$, and the dual infimum is attained.*

Returning to semidefinite programs, we consider the vector space \mathcal{S}_n of symmetric $n \times n$ matrices as the ground set for the primal problem. It is equipped with the usual inner product $\langle X, Y \rangle = \text{Tr}(XY)$ for $X, Y \in \mathcal{S}_n$. The Frobenius norm of a matrix $X \in \mathcal{S}_n$ is defined by $\|X\|_F := \sqrt{\text{Tr}(X^T X)}$. A linear operator A , mapping symmetric matrices into \mathbb{R}^m , is most conveniently represented by

$A(X)_i := \text{Tr}(A_i X)$ for given symmetric matrices $A_i, i = 1, \dots, m$. The adjoint in this case has the representation $A^T(y) = \sum y_i A_i$. From Fejer's Theorem, which states that

$$A \succeq 0 \text{ if and only if } \text{Tr}(AB) \geq 0 \ \forall B \succeq 0,$$

we see that the cone of positive semidefinite matrices is selfdual. Hence we arrive at the following primal-dual pair of semidefinite programs:

$$\max\{\text{Tr}(CX) : A(X) = b, X \succeq 0\}, \quad (3)$$

$$\min\{b^T y : A^T(y) - C \succeq 0\}. \quad (4)$$

In our combinatorial applications, we usually have the property that both the primal and the dual problem satisfy the Slater constraint qualification, hence we have strong duality and both optima are attained.

Stronger duals for semidefinite programs have been introduced having the property that there is no duality gap, in particular, by Borwein and Wolkowicz [45], Ramana [201] (see [203] for a comparison). In Section 2.3, we will come back briefly to the implications for the complexity of semidefinite programming.

The semidefiniteness of a matrix X can equivalently be expressed as X having only nonnegative eigenvalues. Thus there is some close connection between semidefinite programs and spectral theory of matrices. The following simple examples of semidefinite programs throw some more light onto this connection. Throughout, I denotes the identity matrix and I_k the identity matrix of order k .

Example 3. *Let C be a symmetric matrix. Consider*

$$\max \text{Tr}(CX) \text{ such that } \text{Tr}(X) = 1, X \succeq 0.$$

The dual is

$$\min y \text{ such that } yI - C \succeq 0.$$

Both problems clearly satisfy the Slater constraint qualification. In fact, dual feasibility implies that $y \geq \lambda_{\max}(C)$, hence at the optimum $y = \lambda_{\max}(C)$. It is, in fact, well known that the primal semidefinite program is equivalent to

$$\max x^T C x \text{ such that } x^T x = 1,$$

by taking $X = xx^T$.

Example 4. *More generally, the sum $\lambda_1 + \dots + \lambda_k$ of the k largest eigenvalues of $C \in \mathcal{S}_n$ can be expressed as the optimum value of the following semidefinite program:*

$$\max \text{Tr}(CX) \text{ such that } I \succeq X \succeq 0, \text{Tr}(X) = k \quad (5)$$

which is equivalent to

$$\max \text{Tr}(CYY^T) \text{ such that } Y \text{ is an } n \times k \text{ matrix with } Y^T Y = I_k. \quad (6)$$

The fact that $\lambda_1 + \dots + \lambda_k$ is equal to the optimum value of (6) is known as Fan's theorem; see [185] for discussion.

Let us sketch the proof. The fact that the optimum values of the two programs (5) and (6) are equal follows from a nice geometric property of the feasible set of (5) (namely, that its extreme points correspond to the feasible solutions of (6); cf. Lemma 7 below). Let y_1, \dots, y_k be a set of orthonormal eigenvectors of C for its k largest eigenvalues and let Y be the matrix with columns y_1, \dots, y_k . Then Y is feasible for (6) and $\text{Tr}(CYY^T) = \sum_{i=1}^k \text{Tr}(y_i^T C y_i) = \sum_{i=1}^k \lambda_i$, which shows that $\sum_{i=1}^k \lambda_i$ is less than or equal to the maximum of (6). Conversely, let Y be an $n \times k$ matrix such that $Y^T Y = I_k$; we show that $\text{Tr}(CYY^T) \leq \sum_{i=1}^k \lambda_i$. For this, let $C = Q^T D Q$ where $Q \in \mathcal{S}_n$ with $Q^T Q = I_n$ and $D := \text{diag}(\lambda_1, \dots, \lambda_n)$. Set $Z := QY$ and $X := ZZ^T$. As Z is an $n \times k$ matrix with $Z^T Z = I_k$, it follows that the only nonzero eigenvalue of X is 1 with multiplicity k and thus X is feasible for (5). Hence, $\text{Tr}(CYY^T) = \text{Tr}(DX) = \sum_{i=1}^n \lambda_i x_{ii} \leq \sum_{i=1}^k \lambda_i$ since $0 \leq x_{ii} \leq 1$ for all i .

By taking the dual of the semidefinite program (5), we obtain the following alternative formulation for the sum of the k largest eigenvalues of C :

$$\lambda_1 + \dots + \lambda_k = \min kz + \text{Tr}(Z) \quad \text{such that } zI + Z \succeq C, \quad Z \succeq 0. \quad (7)$$

This latter formulation permits to derive the following semidefinite programming characterization for minimizing the sum of the k largest eigenvalues of a symmetric matrix satisfying linear constraints (cf. [8]):

$$\begin{aligned} & \min \lambda_1(X) + \dots + \lambda_k(X) \quad \text{such that } X \in \mathcal{S}_n, \quad \text{Tr}(A_j X) = b_j \quad (j = 1, \dots, m) \\ & = \min kz + \text{Tr}(Z) \quad \text{such that } zI + Z - X \succeq 0, \quad Z \succeq 0, \quad \text{Tr}(A_j X) = b_j \quad (j = 1, \dots, m). \end{aligned}$$

More recently, Anstreicher and Wolkowicz showed a strong connection between a theorem of Hoffman and Wielandt and semidefinite programming.

Theorem 5. (Hoffman and Wielandt [118]) *Let A and B be symmetric matrices of order n with spectral decomposition $A = PDP^T$, $B = QEQ^T$. We assume that the diagonal matrix D contains the eigenvalues of A in nondecreasing order, and E contains the eigenvalues of B in nonincreasing order. Furthermore, $PP^T = QQ^T = I$. Then*

$$\min\{\text{Tr}(AXBX^T) : X^T X = I\} = \text{Tr}(DE). \quad (8)$$

Moreover, the minimum is attained for $X = PQ^T$.

A proof of this theorem can be found for instance in [118], the result can be traced back to the work of John von Neumann [183]. Anstreicher and Wolkowicz [18] have recently shown that the nonconvex quadratic minimization problem (8) over the set of orthogonal matrices can equivalently be expressed through semidefinite programming. This connection will be a useful tool to bound the Quadratic Assignment Problem, so we recall how this connection can be established. We have:

$$\text{Tr}DE = \min\{\text{Tr}AYBY^T : YY^T = I\} = \min\{\text{Tr}DXEX^T : XX^T = I\}.$$

The second equation follows because the mapping $X = P^T Y Q$ is a bijection on the set of orthogonal matrices. We next introduce Lagrange multipliers S and T for the equations $XX^T = I, X^T X = I$, and get

$$\text{Tr}DE = \min_X \max_{S, T} \text{Tr}(DXEX^T + S(I - XX^T) + T(I - X^T X))$$

$$\geq \max_{S,T} \min_{x=\text{vec}(X)} \text{Tr}S + \text{Tr}T + x^T(E \otimes D - I \otimes S - T \otimes I)x.$$

If $X = (x_1, \dots, x_n)$ is a matrix with columns x_i , we define $x = \text{vec}(X) = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}$ to be the vector obtained from stacking the columns of X . The vec -operator leads to the following identity, see [119]:

$$\text{vec}(AXB) = (B^T \otimes A)\text{vec}(X). \quad (9)$$

$A \otimes B$ denotes the Kronecker product of A and B . Formally,

$$A \otimes B = (a_{ij}B).$$

The inner minimization is bounded only if $E \otimes D - I \otimes S - T \otimes I \succeq 0$. Since D and E are diagonal, we may restrict also S and T to be diagonal, $S = \text{diag}(s), T = \text{diag}(t)$. (If s is a vector, $\text{diag}(s)$ denotes the diagonal matrix with s on the main diagonal.) This leads to

$$\text{Tr}DE \geq \max\{\sum s_i + \sum t_i : d_i e_j - s_i - t_j \geq 0 \forall i, j\}.$$

The last problem is the dual of the assignment problem. Therefore we get

$$\text{Tr}DE \geq \min\{\sum_{ij} d_i e_j z_{ij} : Z = (z_{ij}) \text{ doubly stochastic}\} = \text{Tr}DE.$$

The first term equals the last, so there must be equality throughout. We summarize this as follows.

Theorem 6. [18] *Let A and B be symmetric matrices. Then,*

$$\min\{\text{Tr}AXBX^T : XX^T = I\} = \max\{\text{Tr}S + \text{Tr}T : B \otimes A - I \otimes S - T \otimes I \succeq 0\}.$$

2.2 Algorithms

Semidefinite programs are convex minimization problems, hence they can be solved in polynomial time to any fixed prescribed precision using for instance the ellipsoid method [100]. More recently, interior point methods have turned out to be the method of choice to solve SDP, since they give faster algorithms than the ellipsoid method whose running time is prohibitively high in practice; see for instance the monograph [232].

We will now review the main ideas underlying the interior point approach for SDP. The basic assumption is that both the primal (3) and the dual (4) problem satisfy the Slater constraint qualification, which means we assume that there exists a triple (X, y, Z) such that

$$X \succ 0, Z \succ 0, A(X) = b, Z = A^T(y) - C.$$

To avoid trivialities, it is usually also assumed that the linear equations $A(X) = b$ are linearly independent. In view of Theorem 2, we get the following necessary and sufficient optimality conditions.

A triple (X, y, Z) solves (3) and (4) if and only if

$$A(X) = b, X \succeq 0 \quad (\text{primal feasibility}) \quad (10)$$

$$A^T(y) - Z = C, Z \succeq 0 \quad (\text{dual feasibility}) \quad (11)$$

$$ZX = 0 \quad (\text{complementarity}) \quad (12)$$

To see how (12) follows from Theorem 2, we note that both the primal and the dual optimum are attained, and the duality gap is 0. If (X, y, Z) is optimal, we get

$$0 = b^T y - \text{Tr}CX = y^T(A(X)) - \text{Tr}CX = \text{Tr}(A^T(y) - C)X = \text{Tr}ZX.$$

Since $X \succeq 0$, $Z \succeq 0$, we also have $X = UU^T$, $Z = VV^T$, for U and V of appropriate size. Thus

$$0 = \text{Tr}ZX = \text{Tr}VV^TUU^T = \|V^T U\|_F^2,$$

hence $V^T U = 0$, so that $ZX = VV^TUU^T = 0$.

In the interior point approach, the condition $ZX = 0$ is replaced by $ZX = \mu I$, leading to a parameterized system of equations:

$$F_\mu(X, y, Z) := \begin{pmatrix} A(X) - b \\ Z - A^T(y) + C \\ ZX - \mu I \end{pmatrix} = 0. \quad (13)$$

Under our assumptions, there exists a unique solution (X, y, Z) for every $\mu > 0$, see for instance [232] (Chapter 10). (To get this result, one interprets (13) as the KKT system of a convex problem with strictly convex cost function.) Denoting this solution by (X_μ, y_μ, Z_μ) , it is not too hard to show that the set

$$\{(X_\mu, y_\mu, Z_\mu) : \mu > 0\}$$

defines a smooth curve parameterized by μ , which is usually called the ‘central path’.

The interior point approach, more precisely the ‘primal-dual interior-point path-following method’, consists in applying Newton’s method to follow this curve until $\mu \rightarrow 0$. This sounds straightforward, and it is, except for the following aspect. The equation (13) has $2\binom{n+1}{2} + m$ variables, but $\binom{n+1}{2} + n^2 + m$ equations. The difference arises from $ZX - \mu I$, which need not be symmetric, even if X and Z are. Therefore, some sort of symmetrization of the last equation in (13) is necessary to overcome this problem.

The first papers exploiting this approach [115, 138] use some ad-hoc ideas to symmetrize the last equation. Later, Monteiro [176] and Zhang [237] introduced a rather general scheme to deal with the equation $ZX = \mu I$. Let P be invertible. Zhang considers the mapping $H_P(M) := \frac{1}{2} [PMP^{-1} + (PMP^{-1})^T]$ and shows that, for $X \succ 0$, $Z \succ 0$,

$$H_P(ZX) = \mu I \text{ if and only if } ZX = \mu I.$$

Of course, different choices for P produce different search directions after replacing $ZX = \mu I$ by $H_P(ZX) = \mu I$. Various choices for P have been proposed and investigated with respect to theoretical properties and behaviour in practice.

Todd [225] reviews about 20 different variants for the choice of P and investigates some basic theoretical properties of the resulting search directions. The main message seems to be at present

that there is no clear champion among these choices in the sense that it would dominate both with respect to theoretical convergence properties and practical efficiency.

The following variant was introduced by [115], and independently by [138]. It is simple, and yet computationally quite efficient. To simplify the presentation, we assume that there is some starting triple (X, y, Z) which satisfies $A(X) = b$, $A^T(y) - Z = C$ and $X \succ 0$, $Z \succ 0$. If this triple would lie on the central path, its ‘path parameter’ μ would be $\mu = \frac{1}{n} \text{Tr} ZX$. We do not assume that it lies on the central path, but would like to move from this triple towards the central path, and follow it until $\mu \approx 0$. Therefore we head for a point on the central path, given by the path parameter

$$\mu = \frac{1}{2n} \text{Tr} ZX.$$

Applying a Newton step to $F_\mu(X, y, Z) = 0$ at (X, y, Z) , with μ as above, leads to

$$A(\Delta X) = 0 \tag{14}$$

$$\Delta Z = A^T(\Delta y) \tag{15}$$

$$Z(\Delta X) + (\Delta Z)X = \mu I - ZX. \tag{16}$$

The second equation can be used to eliminate ΔZ , the last to eliminate ΔX :

$$\Delta X = \mu Z^{-1} - X - Z^{-1} A^T(\Delta y) X.$$

Substituting this into the first equation gives the following linear system for Δy :

$$A(Z^{-1} A^T(\Delta y) X) = \mu A(Z^{-1}) - b.$$

This system is positive definite, see [115], and can therefore be solved quite efficiently by standard methods, yielding Δy . Backsubstitution gives ΔZ , which is symmetric, and ΔX , which needs not be. Taking the symmetric part of ΔX gives the following new point (X^+, y^+, Z^+) :

$$\begin{aligned} X^+ &= X + t \frac{1}{2} (\Delta X + \Delta X^T) \\ y^+ &= y + t \Delta y \\ Z^+ &= Z + t \Delta Z. \end{aligned}$$

The stepsize $t > 0$ is chosen so that $X^+ \succ 0$, $Z^+ \succ 0$. In practice one starts with $t = 1$ (full Newton step), and backtracks by multiplying the current t with a factor smaller than 1, such as 0.8, until positive definiteness of X^+ and Z^+ holds.

A theoretical convergence analysis shows the following. Let a small scalar $\epsilon > 0$ be given. If the path parameter μ to start a new iteration is chosen properly, then the full step ($t = 1$ above) is feasible in each iteration, and a primal feasible solution X and a dual feasible solution y , whose duality gap $b^T y - \text{Tr}(CX)$ is less than ϵ , can be found after $O(\sqrt{n} |\log \epsilon|)$ iterations; see the handbook [232], chapter 10.

2.3 Complexity

We consider here complexity issues for semidefinite programming. We saw above that for semidefinite programs satisfying the Slater constraint qualification, the primal problem (3) and its dual (4) can be solved in polynomial time to any fixed prescribed precision using interior point methods.

However, even if all input data A_1, \dots, A_m, C, b are rational valued, no polynomial bound has been established for the bitlengths of the intermediate numbers occurring in interior point algorithms. Therefore, interior point algorithms for semidefinite programming are shown to be polynomial in the real number model only, not in the bit number model of computation.

As a matter of fact, there are semidefinite programs with no rational optimum solution. For instance, the matrix $\begin{pmatrix} 1 & x \\ x & 2 \end{pmatrix} \oplus \begin{pmatrix} 2x & 2 \\ 2 & x \end{pmatrix}$ is positive semidefinite if and only if $x = \sqrt{2}$. (Given two matrices A, B , $A \oplus B$ denotes the matrix $\begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}$.) This contrasts with the situation of linear programming, where every rational linear program has a rational optimal solution whose bitlength is polynomially bounded in terms of the bit lengths of the input data (see [208]).

Another ‘pathological’ situation which may occur in semidefinite programming is that all feasible solutions are doubly exponential. Consider, for instance, the matrix (taken from [201]): $Q(x) := Q_1(x) \oplus \dots \oplus Q_n(x)$, where $Q_1(x) := (x_1 - 2)$ and $Q_i(x) := \begin{pmatrix} 1 & x_{i-1} \\ x_{i-1} & x_i \end{pmatrix}$ for $i = 2, \dots, n$. Then, $Q(x) \succeq 0$ if and only if $Q_i(x) \succeq 0$ for all $i = 1, \dots, n$ which implies that $x_i \geq 2^{2^{i-1}}$ for $i = 1, \dots, n$. Therefore, every rational feasible solution has exponential bitlength.

Semidefinite programs can be solved in polynomial time to an arbitrary prescribed precision in the bit model using the ellipsoid method (see [100]). More precisely, let K denote the set of feasible solutions to (3) and, given $\epsilon > 0$, set $S(K, \epsilon) := \{Y \mid \exists X \in K \text{ with } \|X - Y\| < \epsilon\}$ (‘the points that are in the ϵ -neighborhood of K ’) and $S(K, -\epsilon) := \{X \in K \mid \|X - Y\| > \epsilon \text{ for all } Y \notin K\}$ (‘the points in K that are at distance at least ϵ from the border of K ’). Let L denote the maximum bit size of the entries of the matrices A_1, \dots, A_m and the vector b and assume that there is a constant $R > 0$ such that $\exists X \in K$ with $\|X\| \leq R$ if $K \neq \emptyset$. Then, the ellipsoid based algorithm, given $\epsilon > 0$, either finds $X \in S(K, \epsilon)$ for which $\text{Tr}(CY) \leq \text{Tr}(CX) + \epsilon$ for all $Y \in S(K, -\epsilon)$, or asserts that $S(K, -\epsilon) = \emptyset$. Its running time is polynomial in n, m, L , and $\log \epsilon$.

One of the fundamental open problems in semidefinite programming is the complexity of the following *semidefinite programming feasibility problem*¹ (F):

Given integral $n \times n$ symmetric matrices Q_0, Q_1, \dots, Q_m , decide whether there exist real numbers x_1, \dots, x_m such that $Q_0 + x_1 Q_1 + \dots + x_m Q_m \succeq 0$.

This problem belongs obviously to NP in the real number model (since one can test whether a matrix is positive semidefinite in polynomial time using Gaussian elimination), but it is not known whether it belongs to NP in the bit model of computation. Ramana [201] shows that problem (F) belongs to co-NP in the real number model, and that (F) belongs to NP if and only if it belongs to co-NP in the bit model. These two results are based on an extended exact duality theory for semidefinite programming. Namely, given a semidefinite program (P), Ramana [201] defines another semidefinite program (D) whose number of variables and coefficients bitlengths are polynomial in terms of the size of data in (P) and with the property that (P) is feasible if and only if (D) is infeasible.

Porkolab and Khachiyan [196] show that problem (F) can be solved in polynomial time (in the bit model) for *fixed* n or m . (More precisely, problem (F) can be solved in $O(mn^4) + n^{O(\min(m, n^2))}$ arithmetic operations over $Ln^{O(\min(m, n^2))}$ -bit numbers, where L is the maximum bitlength of the entries

¹The following is an equivalent form for the feasibility region of a semidefinite program (3). Indeed, a matrix X is of the form $Q_0 + \sum_{i=1}^m x_i Q_i$ if and only if it satisfies the system: $\text{Tr}(A_j X) = b_j$ ($j = 1, \dots, p$), where A_1, \dots, A_p span the orthogonal complement of the subspace of \mathcal{S}_n generated by Q_1, \dots, Q_m and $b_j = \text{Tr}(A_j Q_0)$ for $j = 1, \dots, p$.

of matrices Q_0, \dots, Q_m .) Moreover, for any *fixed* m , one can decide in polynomial time (in the bit model) whether there exist *rational* numbers x_1, \dots, x_m such that $Q_0 + x_1 Q_1 + \dots + x_m Q_m \succeq 0$ ([129]); this extends Lenstra's result ([156]) about polynomial time solvability of integer linear programming in fixed dimension to semidefinite programming. More generally, given a convex semi-algebraic set $K \subseteq \mathbb{R}^n$, one can find in polynomial time an integral point in K (if some exists) for any fixed dimension n [130]. When all the polynomials defining K are quadratic, this result still holds without the convexity assumption [31]. Further results have been recently given in [98].

A special instance of the semidefinite programming feasibility problem is the *semidefinite matrix completion problem* (MC), which consists of deciding whether a partially specified matrix can be completed to a positive semidefinite matrix. The complexity of problem (MC) is not known in general, even not for the class of partial matrices whose entries are specified on the main diagonal and on the positions corresponding to the edge set of a circuit. However, for circuits (and, more generally, for graphs with no K_4 -minor), problem (MC) is known to be polynomial-time solvable in the real number model [147]. In the bit model, problem (MC) is known to be polynomial time solvable when the graph corresponding to the positions of the specified entries is chordal or can be made chordal by adding a *fixed* number of edges [147]. A crucial tool is a result of [99] asserting that a partial matrix A whose entries are specified on the edge set of a chordal graph can be completed to a positive semidefinite matrix if and only if every fully specified principal submatrix of A is positive semidefinite.

As mentioned above, one of the difficulties in the complexity analysis of semidefinite programming is the possible nonexistence of rational solutions. However, in the special case of the matrix completion problem, no example is known of a rational partial matrix having only irrational positive semidefinite completions. (Obviously, a rational completion exists if a positive *definite* completion exists.)

Further conditions are known for existence of positive semidefinite matrix completions, involving cut and metric polyhedra (see [144]); see the surveys [121], [146] for more information. In practice, positive semidefinite matrix completions can be computed using, e.g., the interior point algorithm of Johnson et al. [122]. This algorithm solves the problem:

$$\min f(X) \quad \text{subject to } X \succeq 0,$$

where $f(X) := \sum_{i,j=1}^n (h_{ij})^2 (x_{ij} - a_{ij})^2$. Here H is a given nonnegative symmetric matrix with a positive diagonal and A is a given symmetric matrix corresponding to the partial matrix to be completed; the condition $h_{ij} = 0$ means that entry x_{ij} is free while $h_{ij} > 0$ puts a weight on forcing entry x_{ij} to be as close as possible to a_{ij} . The optimum value of the above program is equal to 0 precisely when there is a positive semidefinite matrix completion of A , where the entries of A corresponding to $h_{ij} = 0$ are unspecified.

2.4 Geometry

We discuss here some geometric properties of semidefinite programming. We refer to Chapter 3 in [232] for a detailed treatment. Let

$$K := \{X \in \text{PSD}_n \mid \text{Tr}(A_i X) = b_i \text{ for } i = 1, \dots, m\}$$

denote the feasible region of a semidefinite program, where $A_1, \dots, A_m \in \mathcal{S}_n$ and $b \in \mathbb{R}^m$. The set K is a convex set (called a *spectrahedron* in [202]) which inherits several of the geometric properties of the positive semidefinite cone PSD_n , in particular, concerning the structure of its faces. Recall that a set $F \subseteq K$ is a *face* of K if $X, Y \in F$ and $Z := \alpha X + (1 - \alpha)Y \in K$ for some $0 < \alpha < 1$ implies

that $Z \in F$. Given $A \in K$, $F_K(A)$ denotes the smallest face of K containing A . A point $A \in K$ is an *extreme point* if $F_K(A) = \{A\}$. It is well known (see [117]) that, given a matrix $A \in \text{PSD}_n$, the smallest face $F_{\text{PSD}}(A)$ of PSD_n that contains A is given by

$$F_{\text{PSD}}(A) = \{X \in \text{PSD}_n \mid \ker A \subseteq \ker X\}.$$

(For a matrix X , $\ker X := \{x \in \mathbb{R}^n \mid Xx = 0\}$.) Hence, if A has rank r , then $F_{\text{PSD}}(A)$ is isomorphic to the cone PSD_r and thus has dimension $\binom{r+1}{2}$. As K is the intersection of PSD_n with the affine space

$$\mathcal{A} := \{X \in \mathcal{S}_n \mid \text{Tr}(A_i X) = b_i \text{ for } i = 1, \dots, m\},$$

the face $F_K(A)$ for $A \in K$ is given by

$$F_K(A) = F_{\text{PSD}}(A) \cap \mathcal{A} = \{X \in K \mid \ker A \subseteq \ker X\}.$$

One can compute the dimension of faces of K in the following manner (see Chapter 31.5 in [68]).

Let r denote the rank of A and let $A = QQ^T$, where Q is a $n \times r$ matrix of rank r . A matrix $B \in \mathcal{S}_n$ is called a *perturbation* of A if $A \pm tB \in K$ for some small $t > 0$. One can verify that B is a perturbation of A if and only if $B = QRQ^T$ for some matrix $R \in \mathcal{S}_r$ satisfying $\text{Tr}(RQ^T A_i Q) = 0$ for all $i = 1, \dots, m$. Then the dimension of $F_K(A)$ is equal to the rank of the set of perturbations of A and, therefore,

$$\dim F_K(A) = \binom{r+1}{2} - \text{rank}\{Q^T A_i Q \mid i = 1, \dots, m\}.$$

This implies:

$$A \text{ is an extreme point of } K \iff \binom{r+1}{2} = \text{rank}\{Q^T A_i Q \mid i = 1, \dots, m\}. \quad (17)$$

We will use semidefinite programs as relaxations for 0/1 polytopes associated to combinatorial optimization problems; often the rank one matrices in the feasible region K correspond to the integer solutions of the combinatorial problem at hand. With this in mind, it is desirable to find a matrix $A \in K$ optimizing a given linear objective function over K having smallest possible rank. The smallest possible ranks are obviously achieved at extremal matrices of K . Some results have been obtained along these lines which we now mention.

As an application of (17), we have that if $K \neq \emptyset$ and $\text{rank}\{A_i \mid i = 1, \dots, m\} < \binom{r+2}{2}$, then there exists a matrix $X \in K$ with $\text{rank } X \leq r$ ([32], [190]). In fact, every extremal matrix X of K has this property; we will see below how to construct extremal matrices.

Barvinok [33] shows the following refinement. Suppose that K is a nonempty bounded set and that $\text{rank}\{A_i \mid i = 1, \dots, m\} = \binom{r+2}{2}$ for some $1 \leq r \leq n - 2$, then there exists a matrix $X \in K$ with $\text{rank } X \leq r$. Barvinok's proof is nonconstructive and it is an open question how to find efficiently such X .

Barvinok [32] suggests the following approach for finding an extremal matrix in K . Let $C \in \mathcal{S}_n$ be a positive definite matrix and let $A \in K$ minimize $\text{Tr}(CX)$ over K . Barvinok shows that if C is sufficiently generic then A is an extremal point of K .

The following algorithm for constructing an extreme point of K has been suggested by several authors (see [6], [190]). Suppose we want to minimize the objective function $\text{Tr}(CX)$ over K and

assume that the minimum is finite. Given $A \in K$, the algorithm will construct an extremal matrix $A' \in K$ with objective value $\text{Tr}(CA') \leq \text{Tr}(CA)$. Using (17), one can verify whether A is an extreme point of K . If yes, then stop and return $A' = A$. Otherwise, one can find a nonzero matrix R belonging to the orthogonal complement in \mathcal{S}_r of the space spanned by $Q^T A_i Q$ ($i = 1, \dots, m$); then $B := QRQ^T$ is a perturbation of A . If $\text{Tr}(CB) > 0$ then replace B by $-B$. Let t be the largest possible scalar for which $A + tB \succeq 0$. Then, $A + tB$ belongs to the boundary of the face $F_K(A)$ and thus the face $F_K(A + tB)$ is strictly contained in $F_K(A)$. We iterate with $A + tB$ in place of A . In at most n iterations the algorithm returns an extreme point of K .

We conclude with some examples.

The Max-Cut spectrahedron. The following spectrahedron

$$\mathcal{E}_n := \{X \in \text{PSD}_n \mid X_{ii} = 1 \ \forall i = 1, \dots, n\}$$

underlies the semidefinite relaxation for Max-Cut and will be treated in detail in Section 5. Its geometric properties have been investigated in [153, 154]. In particular, it is shown there that the only vertices (that is, the extreme points having a full dimensional normal cone) of \mathcal{E}_n are its rank one matrices (corresponding to the cuts, i.e., the combinatorial objects in which we are interested). The spectrum of possible dimensions for the faces of \mathcal{E}_n is shown to be equal to

$$\left[0, \binom{k_n}{2}\right] \cup \bigcup_{r=k_n+1}^n \left[\binom{r+1}{2} - n, \binom{r}{2}\right],$$

where $k_n := \lfloor \frac{n}{2} \rfloor + 1$. Moreover it is shown that the possible dimensions for the polyhedral faces of \mathcal{E}_n are all integers k satisfying $\binom{k+1}{2} \leq n$. Geometric properties of other tighter spectrahedra for max-cut are studied in [15], [151].

Sum of largest eigenvalues. We introduced in Example 4 two programs (5) and (6) permitting to express the sum of the k largest eigenvalues of a symmetric matrix. Let K and \mathcal{Y} denote their respective feasible regions; that is,

$$K := \{X \in \mathcal{S}_n \mid I \succeq X \succeq 0, \text{Tr}(X) = k\},$$

$$\mathcal{Y} := \{YY^T \mid Y \in \mathbb{R}^{n \times k} \text{ with } Y^T Y = I_k\}.$$

Lemma 7. *The extreme points of the set K are the matrices of \mathcal{Y} . Therefore, K is equal to the convex hull of the set \mathcal{Y} .*

PROOF. Let X be an extreme point of K . Then all its eigenvalues belong to the segment $[0, 1]$. As $\text{Tr}(X) = k$, it follows that X has at least k nonzero eigenvalues and thus $\text{rank}(X) \geq k$. In fact, $\text{rank}(X) = k$ since X is an extreme point of K . Now this implies that the only nonzero eigenvalue of X is 1 with multiplicity k and thus $X \in \mathcal{Y}$. Conversely, every matrix of \mathcal{Y} is obviously an extreme point of K . ■

Note the resemblance of the above result to the Birkhoff-König theorem asserting that the set of stochastic matrices is equal to the convex hull of the set of permutation matrices.

Euclidean distance matrix completions. Let $G = (V, E; d)$ be a weighted graph with $V = \{1, \dots, n\}$ and nonnegative edge weights $d \in \mathbb{Q}_+^E$. Given an integer r , we say that G is r -realizable if there exist points $v_1, \dots, v_n \in \mathbb{R}^r$ such that $d_{ij} = \|v_i - v_j\|$ for all edges $ij \in E$; G is said to be *realizable* if it is r -realizable for some r . The problem of testing existence of a realization is known as the *Euclidean distance matrix completion problem* (EDM) (cf. [146] and Chapter 18 in [232] for surveys). It has important applications, e.g., to molecular conformation problems in chemistry and distance geometry (see [60]). As is well known, problem (EDM) can be formulated as a semidefinite programming problem. Namely, G is realizable if and only if the system:

$$X \succeq 0, \quad X_{ii} + X_{jj} - 2X_{ij} = (d_{ij})^2 \quad \text{for } ij \in E \quad (18)$$

is feasible; moreover G is r -realizable if and only if the system (18) has a solution X with $\text{rank } X \leq r$. It follows from the above mentioned results about ranks of extremal points that if G is realizable, then G is r -realizable for some r satisfying $\binom{r+1}{2} \leq |E|$. Such a realization can be found using the above mentioned algorithm for finding extremal points (see [6], [32]).

It is also well known that the Euclidean distance matrix completion problem can be recast in terms of the positive semidefinite matrix completion problem (MC) treated earlier in Section 2.3 (see [145] for details). As a consequence, the complexity results mentioned earlier for problem (MC) also hold for problem (EDM). Namely, problem (EDM) can be solved in polynomial time in the bit number model when G can be made chordal by adding a fixed number of edges, and (EDM) can be solved in polynomial time in the real number model when G has no K_4 -minor [147].

An interior point algorithm is proposed in [7] for computing graph realizations. Alfakih [4, 5] studies rigidity properties of graph realizations in terms of geometric properties of certain associated spectrahedra.

When the graph G is not realizable, one can look for the smallest distortion needed to be applied to the edge weights in order to ensure existence of a realization. Namely, define this smallest distortion as the smallest scalar C for which there exist points $v_1, \dots, v_n \in \mathbb{R}^n$ satisfying

$$\frac{1}{C}d_{ij} \leq \|v_i - v_j\| \leq d_{ij}$$

for all $ij \in E$. The smallest distortion can be computed using semidefinite programming. Bourgain [46] has shown that $C = O(\log n)$ if $G = K_n$ and d satisfies the triangle inequalities: $d_{ij} \leq d_{ik} + d_{jk}$ for all $i, j, k \in V$ (see also Chapter 10 in [68]). Since then research has been done for evaluating the minimum distortion for several classes of metric spaces including graph metrics (that is, when d is the path metric of the graph G); see in particular [158], [159], [160].

3 Semidefinite Programming and Integer 0/1 Programming

3.1 A general paradigm

Suppose we want to solve a 0/1 linear programming problem:

$$\max c^T x \text{ subject to } Ax \leq b, x \in \{0, 1\}^n. \quad (19)$$

The classic polyhedral approach to this problem consists of formulating (19) as a linear programming problem:

$$\max c^T x \text{ subject to } x \in P$$

over the polytope

$$P := \text{conv}(\{x \in \{0, 1\}^n \mid Ax \leq b\})$$

and of applying linear programming techniques to it. For this one has to find the linear description of P or, at least, good linear relaxations of P . An initial linear relaxation of P is

$$K := \{x \in \mathbb{R}_+^n \mid Ax \leq b\}$$

and, if $K \neq P$, one has to find ‘cutting planes’ permitting to strengthen the relaxation K by cutting off its fractional vertices. Extensive research has been done for finding (partial) linear descriptions for many polyhedra arising from specific combinatorial optimization problems by exploiting the combinatorial structure of the problem at hand. Next to that, research has also focused on developing general purpose methods applying to arbitrary 0/1 problems (or, more generally, integer programming problems).

An early such method, developed in the sixties by Gomory and based on integer rounding, permits to generate the so-called Chvátal-Gomory cuts. This class of cutting planes was later extended, in particular, by Balas [23] who introduced the disjunctive cuts. In the nineties several authors investigated lift-and-project methods for constructing cutting planes, the basic idea being of trying to represent a 0/1 polytope as the projection of a polytope lying in higher dimension. These methods aim at constructing good *linear* relaxations of a given 0/1 polytope, all with the exception of the lift-and-project method of Lovász and Schrijver which permits, moreover, to construct *semidefinite* relaxations. Further constructions for semidefinite relaxations have been recently investigated, based on algebraic results about representations of nonnegative polynomials as sums of squares of polynomials.

This idea of constructing semidefinite relaxations for a combinatorial problem goes back to the seminal work of Lovász [164] who introduced the semidefinite bound $\vartheta(G)$ for the stability number of a graph G , obtained by optimizing over a semidefinite relaxation $\text{TH}(G)$ (see (44)) of the stable set polytope. An important application is the polynomial time solvability of the maximum stable set problem in perfect graphs. This idea was later again used successfully by Goemans and Williamson [95] who, using a semidefinite relaxation of the cut polytope, could prove an approximation algorithm with a good performance guarantee for the max-cut problem. Since then semidefinite programming has been widely used for approximating a variety of combinatorial optimization problems. This will be discussed in detail in further sections of this chapter.

For now we want to go back to the basic question of how to embed the 0/1 linear problem (19) in a semidefinite framework. A natural way of involving positive semidefiniteness is to introduce the matrix variable

$$Y = \begin{pmatrix} 1 \\ x \end{pmatrix} (1 \ x^T).$$

Then Y can be constrained to satisfy

$$(i) Y \succeq 0, \quad (ii) Y_{ii} = Y_{0i} \quad \forall i = 1, \dots, n.$$

Condition (ii) expresses the fact that $x_i^2 = x_i$ as $x_i \in \{0, 1\}$. One can write (i), (ii) equivalently as

$$Y = \begin{pmatrix} 1 & x^T \\ x & X \end{pmatrix} \succeq 0 \quad \text{where } x := \text{diag}(X). \quad (20)$$

The objective function $c^T x$ can be modeled as $\langle \text{diag}(c), X \rangle$. There are several possibilities for modeling a linear constraint $a^T x \leq \beta$ from the system $Ax \leq b$. The simplest way is to use the *diagonal representation*:

$$\langle \text{diag}(a), X \rangle \leq \beta. \quad (21)$$

One can also replace $a^T x \leq \beta$ by its square $(\beta - a^T x)^2 \geq 0$, giving the inequality $(\beta - a^T)Y \begin{pmatrix} \beta \\ -a \end{pmatrix} \geq 0$ which is however redundant under the assumption $Y \succeq 0$. Instead, when $a, \beta \geq 0$, one can use the *squared representation*: $(a^T x)^2 \leq \beta^2$; that is,

$$\langle aa^T, X \rangle \leq \beta^2 \quad (22)$$

or the *extended square representation*: $(a^T x)^2 \leq \beta (a^T x)$; that is,

$$\langle aa^T - \beta \text{diag}(a), X \rangle \leq 0. \quad (23)$$

Another possibility is to exploit the fact that the variable x_i satisfies $0 \leq x_i \leq 1$ and to multiply $a^T x \leq \beta$ by x_i and $1 - x_i$, which yields the system:

$$\sum_{j=1}^n a_j X_{ij} \leq \beta X_{ii} \quad (i = 1, \dots, n), \quad \sum_{j=1}^n a_j (X_{jj} - X_{ij}) \leq \beta(1 - X_{ii}) \quad (i = 1, \dots, n). \quad (24)$$

One can easily compare the strengths of these various representations of the inequality $a^T x \leq \beta$ and verify that, if (20) holds, then

$$(24) \implies (23) \implies (22) \implies (21).$$

Therefore, the constraints (24) define the strongest relaxation; they are, in fact, at the core of the lift-and-project methods by Lovász and Schrijver and by Sherali and Adams as we will see in Section 3.4. From an algorithmic point of view they are however the most expensive ones, as they involve $2n$ inequalities as opposed to one for the other relaxations. Helmberg et al. [116] made an experimental comparison of the various relaxations which seems to indicate that the best trade off between running time and quality is obtained when working with the squared representation.

Instead of treating each inequality of the system $Ax \leq b$ separately, one can also consider pairwise products of inequalities: $(\beta_i - a_i^T x) \cdot (\beta_j - a_j^T x) \geq 0$, yielding the inequalities: $(\beta_i - a_i^T)Y \begin{pmatrix} \beta_j \\ -a_j \end{pmatrix} \geq 0$. This operation is also central to the lift-and-project methods as we will see later in this section.

3.2 Introduction on cutting planes and lift-and-project methods

Given a set $F \subseteq \{0, 1\}^n$, we are interested in finding the linear description of the polytope $P := \text{conv}(F)$. A first (easy) step is to find a *linear programming formulation* for P ; that is, to find a linear system $Ax \leq b$ for which the polytope $K := \{x \in \mathbb{R}^n \mid Ax \leq b\}$ satisfies $K \cap \{0, 1\}^n = F$. If all vertices of K are integral, then $P = K$ and we are done. Otherwise we have to find cutting planes permitting to tighten the relaxation K and possibly find P after a finite number of iterations.

One of the first methods, which applies to general integral polyhedra, is the method of Gomory for constructing cutting planes. Given a linear inequality $\sum_i a_i x_i \leq \alpha$ valid for K where all the coefficients a_i are integers, the inequality $\sum_i a_i x_i \leq \lfloor \alpha \rfloor$ (known as a *Gomory-Chvátal cut*) is still valid for P but may eliminate some part of K . The *Chvátal closure* K' of K is defined as the solution set of all Chvátal-Gomory cuts; that is,

$$K' := \{x \in \mathbb{R}^n \mid u^T Ax \leq \lfloor u^T b \rfloor \text{ for all } u \geq 0 \text{ such that } u^T A \text{ integral}\}.$$

Then,

$$P \subseteq K' \subseteq K. \tag{25}$$

Set $K^{(1)} := K'$ and define recursively $K^{(t+1)} := (K^{(t)})'$ for $t \geq 1$. Chvátal [52] proved that K' is a polytope and that $K^{(t)} = \text{conv}(K)$ for some t ; the smallest t for which this is true is the *Chvátal rank* of the polytope K . The Chvátal rank may be very large as it depends not only on the dimension n but also on the coefficients of the inequalities involved. However, when K is assumed to be contained in the cube $[0, 1]^n$, its Chvátal rank is bounded by $O(n^2 \log n)$; if, moreover, $K \cap \{0, 1\}^n = \emptyset$, then the Chvátal rank is at most n [42, 74]. Even if we can optimize a linear objective function over K in polynomial time, optimizing a linear objective function over the first Chvátal closure K' is a co-NP-hard problem in general [73].

Further classes of cutting planes have been investigated; in particular, the class of split cuts [56] (they are a special case of the disjunctive cuts studied in [23]). An inequality $a^T x \leq \alpha$ is a *split cut* for K if it is valid for the polytope $\text{conv}\left(\left(K \cap \{x \mid c^T x \leq c_0\}\right) \cup \left(K \cap \{x \mid c^T x \geq c_0 + 1\}\right)\right)$ for some integral $c \in \mathbb{Z}^n$, $c_0 \in \mathbb{Z}$. Split cuts are known to be equivalent to Gomory's mixed integer cuts (see, e.g., [57]). The *split closure* K' of K , defined as the solution set to all split cuts, is a polytope which satisfies again (25) [56]. One can iterate this operation of taking the split closure and it follows from results in [23] that P is found after n steps. However, optimizing over the first split closure is again a hard problem [47]. (An alternative proof for NP-hardness of the membership problem in the split closure and in the Chvátal closure, based on a reduction from the closest lattice vector problem, is given in [59].) If we consider only the split cuts obtained from the disjunctions $x_j \leq 0$ and $x_j \geq 1$, then we obtain a tractable relaxation of K which coincides with the relaxation obtained in one iteration of the Balas-Ceria-Cornuéjols lift-and-project method (which will be described later in Section 3.4).

Another popular approach is to try to represent P as the projection of another polytope Q lying in a higher (but preferably still polynomial) dimensional space, the idea behind being that the projection of a polytope Q may have more facets than Q itself. Hence it could be that even if P has an exponential number of facets, such Q exists having only a polynomial number of facets and lying in a space whose dimension is polynomial in the original dimension of P (such Q is then called a *compact representation* of P). If this is the case then we have a proof that any linear optimization problem over P can be solved in polynomial time. At this point let us stress that it is not difficult to find a lift Q of P with a simple structure and lying in a space of *exponential* dimension; indeed, as pointed out in Section 3.3,

any n -dimensional 0/1 polytope can be realized as projection of a canonical simplex lying in the $(2^n - 1)$ -space.

This idea of finding compact representations has been investigated for several polyhedra arising from combinatorial optimization problems; for instance, Barahona [26], Barahona and Mahjoub [29, 30], Ball, Liu and Pulleyblank [25], Maculan [169], Liu [162] have provided such representations for certain polyhedra related to Steiner trees, stable sets, metrics, etc. On the negative side, Yannakakis [233] proved that the matching polytope cannot have a compact representation satisfying a certain symmetry assumption.

Several general purpose methods have been developed for constructing projection representations for general 0/1 polyhedra; in particular, by Balas, Ceria and Cornuéjols [24] (the BCC method), by Sherali and Adams [212] (the SA method), by Lovász and Schrijver [167] (the LS method) and, recently, by Lasserre [142]. [These methods are also known under the following names: lift-and-project for BCC, Reformulation-Linearization Technique (RLT) for SA, and matrix-cuts for LS.] A common feature of these methods is the construction of a hierarchy

$$K \supseteq K_1 \supseteq K_2 \supseteq \dots \supseteq K_n \supseteq P$$

of linear or semidefinite relaxations of P which finds the exact convex hull in n steps; that is, $K_n = P$. The methods also share the following important algorithmic property: If one can optimize a linear objective function over the initial relaxation K in polynomial time, then the same holds for the next relaxations K_t for any *fixed* t , when applying the BCC, SA or LS constructions; for the Lasserre construction, this is true under the more restrictive assumption that the matrix A has a polynomial number of rows.

The first three methods (BCC, SA and LS) provide three hierarchies of *linear* relaxations of P satisfying the following inclusions: the Sherali-Adams relaxation is contained in the Lovász-Schrijver relaxation which in turn is contained in the Balas-Ceria-Cornuéjols relaxation. All three can be described following a common recipe: Multiply each inequality of the system $Ax \leq b$ by certain products of the bound inequalities $x_i \geq 0$ and $1 - x_i \geq 0$, replace each square x_i^2 by x_i , and linearize the products $x_i x_j$ ($i \neq j$) by introducing a new variable $y_{ij} = x_i x_j$. In this way, we obtain polyhedra in a higher dimensional space whose projection on the subspace \mathbb{R}^n of the original x variable contains P and is contained in K . The three methods differ in the way of choosing the variables employed as multipliers and of iterating the basic step. The Lovász-Schrijver method can be strengthened by requiring positive semidefiniteness of the matrix (y_{ij}) , which leads then to a hierarchy of positive semidefinite relaxations of P .

The construction of Lasserre produces a hierarchy of *semidefinite* relaxations of P which refines each of the above three hierarchies (BCC, SA and LS, even its positive semidefinite version). It was originally motivated by results about moment sequences and the dual theory of representation of nonnegative polynomials as sums of squares. It is however closely related to the SA method as both can be described in terms of requiring positive semidefiniteness of certain principal submatrices of the moment matrices of the problem.

We present in Section 3.3 some preliminary results which permit to show the convergence of the Lasserre and SA methods and to prove that every 0/1 polytope can be represented as the projection of a simplex in the $(2^n - 1)$ -space. Then we describe in Section 3.4 the four lift-and-project methods and Sections 3.5, 3.6 and 3.7 contain applications of these methods to the stable set polytope, the cut polytope and some related polytopes. Section 3.8 presents extensions to (in general non convex) polynomial programming problems.

It will sometimes be convenient to view a polytope in \mathbb{R}^n as being embedded in the hyperplane $x_0 = 1$ of \mathbb{R}^{n+1} . The following notation will be used throughout these sections. For a polytope P in \mathbb{R}^n , its *homogenization*

$$\tilde{P} := \left\{ \lambda \begin{pmatrix} 1 \\ x \end{pmatrix} \mid x \in P, \lambda \geq 0 \right\}$$

is a cone in \mathbb{R}^{n+1} such that $P = \{x \in \mathbb{R}^n \mid \begin{pmatrix} 1 \\ x \end{pmatrix} \in \tilde{P}\}$. For a cone C in \mathbb{R}^n ,

$$C^* := \{y \in \mathbb{R}^n \mid x^T y \geq 0 \ \forall x \in C\}$$

denotes its *dual cone*.

3.3 A canonical lifting construction

Let $\mathcal{P}(V) := 2^V$ denote the collection of all subsets of $V = \{1, \dots, n\}$ and let Z be the square 0/1 matrix indexed by $\mathcal{P}(V)$ with entries

$$Z(I, J) = 1 \text{ if and only if } I \subseteq J. \quad (26)$$

As Z is upper triangular with ones on its main diagonal, it is nonsingular and its inverse Z^{-1} has entries

$$Z^{-1}(I, J) = (-1)^{|J \setminus I|} \text{ if } I \subseteq J, \quad Z^{-1}(I, J) = 0 \text{ otherwise.}$$

For $J \subseteq V$, let Z^J denote the J -th column of Z . [The matrix Z is known as the *Zeta matrix* of the lattice $\mathcal{P}(V)$ and the matrix Z^{-1} as its *Möbius matrix*.]

Given a subset $\mathcal{J} \subseteq \mathcal{P}(V)$, let $\mathcal{C}_{\mathcal{J}}$ denote the cone in $\mathbb{R}^{\mathcal{P}(V)}$ generated by the columns Z^J ($J \in \mathcal{J}$) of Z and let $P_{\mathcal{J}}$ be the 0/1 polytope in \mathbb{R}^n defined as the convex hull of the incidence vectors of the sets in \mathcal{J} . Then $\mathcal{C}_{\mathcal{J}}$ is a simplicial cone,

$$\mathcal{C}_{\mathcal{J}} = \{y \in \mathbb{R}^{\mathcal{P}(V)} \mid Z^{-1}y \geq 0, (Z^{-1}y)_J = 0 \text{ for } J \in \mathcal{P}(V) \setminus \mathcal{J}\},$$

and $P_{\mathcal{J}}$ is the projection on \mathbb{R}^n of the simplex $\mathcal{C}_{\mathcal{J}} \cap \{y \mid y_{\emptyset} = 1\}$. This shows therefore that any 0/1 polytope in \mathbb{R}^n is the projection of a simplex lying in $\mathbb{R}^{2^n - 1}$.

Given $y \in \mathbb{R}^{\mathcal{P}(V)}$, let $M_V(y)$ be the square matrix indexed by $\mathcal{P}(V)$ with entries

$$M_V(y)(I, J) := y(I \cup J) \quad (27)$$

for $I, J \subseteq V$; $M_V(y)$ is known as the *moment matrix* of the sequence y . (See Section 7.1 for motivation and further information.) As noted in [167], we have:

$$M_V(y) = Z \text{diag}(Z^{-1}y) Z^T.$$

Therefore, the cone $\mathcal{C}_{\mathcal{P}(V)}$ can be alternatively characterized by any of the following linear and positive semidefinite conditions:

$$y \in \mathcal{C}_{\mathcal{P}(V)} \iff Z^{-1}y \geq 0 \iff M_V(y) \succeq 0. \quad (28)$$

Suppose that \mathcal{J} corresponds to the set of 0/1 solutions of a semi-algebraic system

$$g_{\ell}(x) \geq 0 \text{ for } \ell = 1, \dots, m$$

where the g_ℓ 's are polynomials in x . One can assume without loss of generality that each g_ℓ has degree at most one in every variable x_i and then one can identify g_ℓ with its sequence of coefficients indexed by $\mathcal{P}(V)$. Given $g, y \in \mathbb{R}^{\mathcal{P}(V)}$, define $g * y \in \mathbb{R}^{\mathcal{P}(V)}$ by

$$g * y := M(y)g; \text{ that is, } g * y(J) := \sum_I g_I y_{I \cup J} \text{ for } J \subseteq V. \quad (29)$$

It is noted in [150] that the cone $\mathcal{C}_{\mathcal{J}}$ can be alternatively characterized by the following positive semidefinite conditions:

$$y \in \mathcal{C}_{\mathcal{J}} \iff M_V(y) \succeq 0 \text{ and } M_V(g_\ell * y) \succeq 0 \text{ for } \ell = 1, \dots, m. \quad (30)$$

This holds, in particular, when \mathcal{J} corresponds to the set of 0/1 solutions of a linear system $Ax \leq b$, i.e., in the case when each polynomial g_ℓ has degree 1.

3.4 The Balas-Ceria-Cornu ejols, Lov asz-Schrijver, Sherali-Adams, and Lasserre methods

Consider the polytope $K = \{x \in [0, 1]^n \mid Ax \leq b\}$ and let $P = \text{conv}(K \cap \{0, 1\}^n)$ be the 0/1 polytope whose linear description is to be found. It is convenient to assume that the bound constraints $0 \leq x_i \leq 1$ ($i \in \{1, \dots, n\}$) are explicitly present in the linear description of K ; let us rewrite the two systems $Ax \leq b$ and $0 \leq x_i \leq 1$ ($i \in \{1, \dots, n\}$) as $\tilde{A}x \leq \tilde{b}$ and let m denote the number of rows of A .

The Balas-Ceria-Cornu ejols construction. Fix an index $j \in \{1, \dots, n\}$. Multiply the system $\tilde{A}x \leq \tilde{b}$ by x_j and $1 - x_j$ to obtain the nonlinear system: $x_j(\tilde{A}x - \tilde{b}) \leq 0$, $(1 - x_j)(\tilde{A}x - \tilde{b}) \leq 0$. Replace x_j^2 by x_j and linearize by introducing new variables $y_i = x_i x_j$ ($i = 1, \dots, n$); thus $y_j = x_j$. This defines a polytope in the (x, y) -space defined by $2(m + 2n)$ inequalities: $\tilde{A}y - \tilde{b}x_j \leq 0$, $\tilde{A}(x - y) - \tilde{b}(1 - x_j) \leq 0$. Its projection $P_j(K)$ on the subspace \mathbb{R}^n indexed by the original x -variable satisfies

$$P \subseteq P_j(K) \subseteq K.$$

Iterate by defining $P_{j_1 \dots j_t}(K) := P_{j_t}(P_{j_{t-1}} \dots (P_{j_1}(K)) \dots)$. It is shown in [24] that

$$P_{j_1 \dots j_t}(K) = \text{conv}(K \cap \{x \mid x_{j_1}, \dots, x_{j_t} \in \{0, 1\}\}). \quad (31)$$

Therefore,

$$P = P_{j_1 \dots j_n}(K) \subseteq P_{j_1 \dots j_{n-1}}(K) \subseteq \dots \subseteq P_{j_1}(K) \subseteq K.$$

The Sherali-Adams construction. The first step is analogue to the first step of the BCC method except that we now multiply the system $\tilde{A}x \leq \tilde{b}$ by x_j and $1 - x_j$ for *all* indices $j \in \{1, \dots, n\}$. More generally, for $t = 1, \dots, n$, the t -th step goes as follows. Multiply the system $\tilde{A}x \leq \tilde{b}$ by each product $f_t(J_1, J_2) := \prod_{j \in J_1} x_j \cdot \prod_{j \in J_2} (1 - x_j)$ where J_1 and J_2 are disjoint subsets of V with $|J_1 \cup J_2| = t$. Replace each square x_i^2 by x_i and linearize each product $\prod_{i \in I} x_i$ by a new variable y_I . This defines a polytope $R_t(K)$ in the space of dimension $n + \binom{n}{2} + \dots + \binom{n}{T}$ where $T := \min(t + 1, n)$ (defined by $2^t \binom{n}{t} (m + 2n)$ inequalities) whose projection $S_t(K)$ on the subspace \mathbb{R}^n of the original x -variable satisfies

$$P \subseteq S_n(K) \subseteq \dots \subseteq S_{t+1}(K) \subseteq S_t(K) \subseteq \dots \subseteq S_1(K) \subseteq K$$

and $P = S_n(K)$. The latter equality follows from facts in Section 3.3 as we now see.

Write the linear system $\tilde{A}x \leq \tilde{b}$ as $g_\ell^T \begin{pmatrix} 1 \\ x \end{pmatrix} \geq 0$ ($\ell = 1, \dots, m + 2n$) where $g_\ell \in \mathbb{R}^{n+1}$. Extend g_ℓ to a vector in $\mathbb{R}^{\mathcal{P}(V)}$ by adding zero coordinates. The linearization of the inequality $g_\ell^T \begin{pmatrix} 1 \\ x \end{pmatrix} \cdot f_t(I, J) \geq 0$ reads:

$$\sum_{I \subseteq H \subseteq I \cup J} (-1)^{|H \setminus I|} g_\ell * y(H) \geq 0.$$

Using relation (28), one can verify that the set $R_t(K)$ can be alternatively described by the positive semidefinite conditions:

$$\begin{aligned} M_U(g_\ell * y) \succeq 0 & \text{ for } \ell = 1, \dots, m \text{ and } U \subseteq V \text{ with } |U| = t, \\ M_U(y) \succeq 0 & \text{ for } U \subseteq V \text{ with } |U| = t + 1 \end{aligned} \quad (32)$$

(where g_1, \dots, g_m correspond to the system $Ax \leq b$). It then follows from (30) that the projection $S_n(K)$ of $R_n(K)$ is equal to P .

The Lovász-Schrijver construction. Let U be another linear relaxation of P which is also contained in the cube $Q := [0, 1]^n$; write U as $\{x \in \mathbb{R}^n \mid u_r^T \begin{pmatrix} 1 \\ x \end{pmatrix} \geq 0 \ \forall r = 1, \dots, s\}$. Multiply each inequality $g_\ell^T \begin{pmatrix} 1 \\ x \end{pmatrix} \geq 0$ by each inequality $u_r^T \begin{pmatrix} 1 \\ x \end{pmatrix} \geq 0$ to obtain the nonlinear system: $u_r^T \begin{pmatrix} 1 \\ x \end{pmatrix} \cdot g_\ell^T \begin{pmatrix} 1 \\ x \end{pmatrix} \geq 0$ for all $\ell = 1, \dots, m + 2n$, $r = 1, \dots, s$. Replace each x_i^2 by x_i and linearize by introducing a new matrix variable $Y = \begin{pmatrix} 1 \\ x \end{pmatrix} (1 \ x^T)$. This defines the set $M(K, U)$ consisting of the symmetric matrices $Y = (y_{ij})_{i,j=0}^n$ satisfying

$$y_{jj} = y_{0j} \text{ for } j = 1, \dots, n, \quad (33)$$

$$u_r^T Y g_\ell \geq 0 \text{ for all } r = 1, \dots, s, \ \ell = 1, \dots, m + 2n \text{ [equivalently, } Y \tilde{U}^* \subseteq \tilde{K}]. \quad (34)$$

The first LS relaxation of P is defined as

$$N(K, U) := \{x \in \mathbb{R}^n \mid \begin{pmatrix} 1 \\ x \end{pmatrix} = Y e_0 \text{ for some } Y \in M(K, U)\}.$$

Then, $P \subseteq N(K, U) \subseteq N(K, Q) \subseteq K$ and $N(K, K) \subseteq N(K, U)$ if $K \subseteq U$. One can obtain stronger relaxations by adding positive semidefiniteness. Let $M_+(K, U)$ denote the set of positive semidefinite matrices in $M(K, U)$ and $N_+(K, U) := \{x \in \mathbb{R}^n \mid \begin{pmatrix} 1 \\ x \end{pmatrix} = Y e_0 \text{ for some } Y \in M_+(K, U)\}$. Then,

$$P \subseteq N_+(K, U) \subseteq N(K, U) \subseteq K.$$

The most extensively studied choice for U is $U := Q$, leading to the N operator. Set $N(K) := N(K, Q)$ and, for $t \geq 2$, $N^t(K) := N(N^{t-1}(K)) = N(N^{t-1}(K), Q)$. It follows from condition (34) that $N(K) \subseteq \text{conv}(K \cap \{x \mid x_j = 0, 1\}) = P_j(K)$, the first BCC relaxation, and thus

$$N(K) \subseteq N_0(K) := \bigcap_{j=1}^n P_j(K). \quad (35)$$

[One can verify that $N_0(K)$ consists of the vectors $x \in \mathbb{R}^n$ for which $\begin{pmatrix} 1 \\ x \end{pmatrix} = Y e_0$ for some matrix Y (not necessarily symmetric) satisfying (33) and (34) (with $U = Q$).] More generally, $N^t(K) \subseteq P_{j_1 \dots j_t}(K)$ and, therefore, $P = N^n(K)$.

The choice $U := K$ leads to the stronger operator N' , where we define $N'(K) := N(K, K)$ and, for $t \geq 2$,

$$(N')^t(K) := N((N')^{t-1}(K), K). \quad (36)$$

This operator is considered in [149] when applied to the cut polytope.

When using the relaxation $U = Q$, the first steps in the SA and LS constructions are identical; that is, $S_1(K) = N(K)$. The next steps are however distinct. A main difference between the two methods is that the LS procedure constructs the successive relaxations by a succession of t lift-and-project steps, each lifting taking place in a space of dimension $O(n^2)$, whereas the SA procedure carries out only one lifting step, occurring now in a space of dimension $O(n^{t+1})$; moreover, the projection step is not mandatory in the SA construction.

The Lasserre construction. We saw in relation (32) that the SA method can be interpreted as requiring positive semidefiniteness of certain principal submatrices of the moment matrices $M_V(y)$ and $M_V(g_\ell * y)$. The Lasserre method consists of requiring positive semidefiniteness of certain other principal matrices of those moment matrices. Namely, given an integer $t = 0, \dots, n$, let $P_t(K)$ be defined by the conditions

$$M_{t+1}(y) \succeq 0, \quad M_t(g_\ell * y) \succeq 0 \text{ for } \ell = 1, \dots, m \quad (37)$$

and let $Q_t(K)$ denote the projection of $P_t(K)$ on \mathbb{R}^n . (For a vector $z \in \mathbb{R}^{\mathcal{P}(V)}$, $M_t(z)$ denotes the principal submatrix of $M_V(z)$ indexed by all sets $I \subseteq V$ with $|I| \leq t$.) Then,

$$P \subseteq Q_n(K) \subseteq Q_{n-1}(K) \subseteq \dots \subseteq Q_1(K) \subseteq Q_0(K) \subseteq K$$

and it follows from (30) that $P = Q_n(K)$.

The construction of Lasserre [140, 142] was originally presented in terms of moment matrices indexed by integer sequences (rather than subsets of V) and his proof of convergence used results about moment theory and the representation of nonnegative polynomials as sums of squares. The presentation and the proof of convergence given here are taken from [150].

How do the four hierarchies of relaxations relate? The following inclusions hold among the relaxations $P_{j_1 \dots j_t}(K)$ (BCC), $S_t(K)$ (SA), $N^t(K)$ and $N_+^t(K)$ (LS), and $Q_t(K)$ (Lasserre):

- (i) $Q_1(K) \subseteq N_+(K) \subseteq Q_0(K)$
- (ii) [167] For $t \geq 1$, $S_t(K) \subseteq N^t(K) \subseteq P_{j_1 \dots j_t}(K)$
- (iii) [150] For $t \geq 1$, $S_t(K) \subseteq N(S_{t-1}(K))$, $Q_t(K) \subseteq N_+(Q_{t-1}(K))$,
and thus $Q_t(K) \subseteq S_t(K) \cap N_+^t(K)$.

Summarizing, the Lasserre relaxation is the strongest among all four types of relaxations.

Algorithmic aspects. Efficient approximations to linear optimization problems over the 0/1 polytope P can be obtained by optimizing over its initial relaxation K or any of the stronger relaxations constructed using the BCC, LS, SA and Lasserre methods. Indeed, if one can optimize in polynomial time any linear objective function over K [equivalently (by the results in [100]), one can solve the separation problem for K in polynomial time], then, for any *fixed* t , the same holds for each of the relaxations $P_{j_1 \dots j_t}(K)$, $S_t(K)$, $N^t(K)$, $N_+^t(K)$ in the BCC, SA, and LS hierarchies. This holds for the Lasserre relaxation $Q_t(K)$ under the more restrictive assumption that the linear system defining K has polynomial number of rows. Better approximations are obtained for higher values of t , at an increasing cost however. Computational experiments have been carried out using the various methods;

see, in particular, [24], [49], [50] for results using the BCC method, [214] (and further references there) for results using the SA method, and to [62] for a computational study of the N_+ operator.

Worst case examples where n iterations are needed for finding P . Let us define the *rank* of K with respect to a certain lift-and-project method as the smallest number of iterations needed for finding P . Specifically, the N -rank of K is the smallest integer t for which $P = N^t(K)$; define similarly the N_+ , N_0 , BCC, SA and Lasserre ranks. We saw above that n is a common upper bound for any such rank. We give below two examples of polytopes K whose rank is *equal* to n with respect to all procedures (except maybe with respect to the procedure of Lasserre, since the exact value of the Lasserre rank of these polytopes is not known).

As we will see in Section 3.5, the relaxation of the stable set polytope obtained with the Lovász-Schrijver N operator is much weaker than that obtained with the N_+ -operator. For example, the fractionnal stable set polytope of K_n (defined by nonnegativity and the edge constraints) has N -rank $n - 2$ while its N_+ -rank is equal to $1!$ However, in the case of max-cut, no graph is known for which a similar result holds. Thus it is not clear in which situations the N_+ -operator is significantly better, especially when applied iteratively. Some geometric results about the comparative strengths of the N , N_+ and N_0 operators are given in [93]. As a matter of fact, there exist polytopes K having N_+ -rank equal to n (thus, for them, adding positive semidefiniteness does not help!).

As a first example, let

$$K := \left\{ x \in [0, 1]^n \mid \sum_{i=1}^n x_i \geq \frac{1}{2} \right\}; \quad (38)$$

then $P = \{x \in [0, 1]^n \mid \sum_{i=1}^n x_i \geq 1\}$ and the Chvátal rank of K is therefore equal to 1. The N_+ -rank of K is equal to n [55, 62] and its SA-rank as well [150]. As a second example, let

$$K := \left\{ x \in [0, 1]^n \mid \sum_{i \in I} x_i + \sum_{i \notin I} (1 - x_i) \geq \frac{1}{2} \quad \forall I \subseteq \{1, \dots, n\} \right\}; \quad (39)$$

then $K \cap \{0, 1\}^n = \emptyset$ and thus $P = \emptyset$. Then the N_+ -rank of K is equal to n [55, 93] as well as its SA-rank [150]. In fact, the Chvátal rank of K is also equal to n [54]. The rank of K remains equal to n for the iterated operator N^* defined by $N^*(K) := N_+(K) \cap K'$, combining the Chvátal closure and the N_+ -operator [55, 62]. The rank is also equal to n if in the definition of N^* we replace the Chvátal closure by the split closure [58].

General setting in which the four methods apply. We have described above how the various lift-and-project methods apply to 0/1 linear programs, i.e., to the case when K is a polytope and $P = \text{conv}(K \cap \{0, 1\}^n)$. In fact, they apply in a more general context, still retaining the property that P is found after n steps. Namely, the Lovász-Schrijver method applies to the case when K and U are arbitrary convex sets, the condition (34) reading then $Y\tilde{U}^* \subseteq \tilde{K}$. The BCC and SA methods apply to mixed 0/1 linear programs [24, 213]. Finally, the Lasserre and Sherali-Adams methods apply to the case when K is a semi-algebraic set, i.e., when K is the solution set of a system of polynomial inequalities (since relation (30) holds in this context).

Moreover, various strengthenings of the basic SA method have been proposed involving, in particular, products of other inequalities than the bounds $0 \leq x_i \leq 1$ (cf., e.g., [49], [214], [215], [216]). A comparison between the Lasserre and SA methods for polynomial programming from the algebraic point of view of representations of positive polynomials is made in [143].

3.5 Application to the stable set problem

Given a graph $G = (V, E)$, a set $I \subseteq V$ is *stable* if no two nodes of I form an edge and the *stable set polytope* $\text{STAB}(G)$ is the convex hull of the incidence vectors χ^S of all stable sets S of G , where $\chi_i^S = 1$ if $i \in S$ and $\chi_i^S = 0$ if $i \in V \setminus S$. As linear programming formulation for $\text{STAB}(G)$, we consider the *fractional stable set polytope* $\text{FRAC}(G)$ which is defined by the nonnegativity constraints: $x \geq 0$ and the *edge inequalities*:

$$x_i + x_j \leq 1 \quad \text{for } ij \in E. \quad (40)$$

Let us indicate how the various lift-and-project methods apply to the pair $P := \text{STAB}(G)$, $K := \text{FRAC}(G)$.

The LS relaxations $N(\text{FRAC}(G))$ and $N_+(\text{FRAC}(G))$ are studied in detail in [167] where the following results are shown. The polytope $N(\text{FRAC}(G))$ is completely described by nonnegativity, the edge constraints (40) and the *odd hole inequalities*:

$$\sum_{i \in V(C)} x_i \leq \frac{|C| - 1}{2} \quad \text{for } C \text{ odd circuit in } G. \quad (41)$$

Moreover, $N(\text{FRAC}(G)) = N_0(\text{FRAC}(G))$. Therefore, this gives a compact representation for the stable set polytope of t -perfect graphs (they are the graphs whose stable set polytope is completely determined by nonnegativity together with edge and odd hole constraints).

Other valid inequalities for $\text{STAB}(G)$ include the *clique inequalities*:

$$\sum_{i \in Q} x_i \leq 1 \quad \text{for } Q \text{ clique in } G. \quad (42)$$

The smallest integer t for which (42) is valid for $N^t(\text{FRAC}(G))$ is $t = |Q| - 2$ while (42) is valid for $N_+(\text{FRAC}(G))$. Hence the N_+ operator yields a stronger relaxation of $\text{STAB}(G)$ and equality $N_+(\text{FRAC}(G)) = \text{STAB}(G)$ holds for perfect graphs (they are the graphs for which $\text{STAB}(G)$ is completely determined by nonnegativity and the clique inequalities; cf. Theorem 9). Odd antihole and odd wheel inequalities are also valid for $N_+(\text{FRAC}(G))$.

Given a graph G on n nodes with stability number $\alpha(G)$ (i.e., the maximum size of a stable set in G), the following bounds hold for the N -rank t of $\text{FRAC}(G)$ and its N_+ -rank t_+ :

$$\frac{n}{\alpha(G)} - 2 \leq t \leq n - \alpha(G) - 1, \quad t_+ \leq \alpha(G).$$

See Lipták and Tunçel [161] for a detailed study of further properties of the N and N_+ operators applied to $\text{FRAC}(G)$; in particular, they show the bound $t_+ \leq \frac{n}{3}$ for the N_+ -rank of $\text{FRAC}(G)$.

The Sherali-Adams method does not seem to give a significant improvement, since the quantity $\frac{n}{\alpha(G)} - 2$ remains a lower bound for the SA-rank [150].

The Lasserre hierarchy refines the sequence $N_+^t(\text{FRAC}(G))$. Indeed, it is shown in [150] that, for $t \geq 1$, the set $Q_t(\text{FRAC}(G))$ can be alternatively described as the projection of the set

$$M_{t+1}(y) \succeq 0, \quad y_{ij} = 0 \quad \text{for all edges } ij \in E, \quad y_\emptyset = 1. \quad (43)$$

This implies that $Q_{\alpha(G)-1}(\text{FRAC}(G)) = \text{STAB}(G)$; that is, the Lasserre rank of $\text{FRAC}(G)$ is at most $\alpha(G) - 1$. The inclusion $Q_{\alpha(G)-1}(\text{FRAC}(G)) \subseteq N_+^{\alpha(G)-1}(\text{FRAC}(G))$ is strict, for instance, when G is the line graph of K_n (n odd) since the N_+ -rank of $\text{FRAC}(G)$ is then equal to $\alpha(G)$ ([223]).

Let us mention a comparison with the basic semidefinite relaxation of $\text{STAB}(G)$ by the theta body $\text{TH}(G)$, which is defined by

$$\text{TH}(G) := \{x \in \mathbb{R}^n \mid \begin{pmatrix} 1 \\ x \end{pmatrix} = Y e_0 \text{ for some } Y \succeq 0 \text{ s.t. } Y_{ii} = Y_{0i} \ (i \in V), Y_{ij} = 0 \ (ij \in E)\}. \quad (44)$$

When maximizing $\sum_i x_i$ over $\text{TH}(G)$, we obtain the theta number $\vartheta(G)$. Comparing with (43), we see that $Q_t(\text{FRAC}(G))$ ($t \geq 1$) is a natural generalization of the SDP relaxation $\text{TH}(G)$ satisfying the following chain of inclusions:

$$Q_t(\text{FRAC}(G)) \subseteq Q_1(\text{FRAC}(G)) \subseteq N_+(\text{FRAC}(G)) \subseteq \text{TH}(G) \subseteq Q_0(\text{FRAC}(G)).$$

Section 4.2 below contains a detailed treatment of the relaxation $\text{TH}(G)$.

Feige and Krauthgamer [80] study the behaviour of the N_+ operator applied to the fractional stable set polytope of $G_{n,1/2}$, a random graph on n nodes in which two nodes are joined by an edge with probability $1/2$. It is known that the independence number of $G_{n,1/2}$ is equal, almost surely, to roughly $2 \log_2 n$ and that its theta number is, almost surely, $\Theta(\sqrt{n})$. Feige and Krauthgamer [80] show that the maximum value of $\sum_i x_i$ over $N_+(\text{FRAC}(G_{n,1/2}))$ is, almost surely, roughly $\sqrt{\frac{n}{2^r}}$ when $r = o(\log n)$. This value can be computed efficiently if $r = O(1)$. Therefore, in that case, the typical value of these relaxations is smaller than that of the theta number by no more than a constant factor. Moreover, it is shown in [80] that the N_+ -rank of a random graph $G_{n,1/2}$ is almost surely $\Theta(\log n)$.

3.6 Application to the max-cut problem

We consider here how the various lift-and-project methods can be used for constructing relaxations of the cut polytope. Section 5 will focus on the most basic SDP relaxation of the cut polytope and, in particular, on how it can be used for designing good approximation algorithms for the max-cut problem. As is well known (cf. (75)), the max-cut problem can be formulated as an unconstrained quadratic ± 1 problem:

$$\max x^T A x \text{ subject to } x \in \{\pm 1\}^n \quad (45)$$

for some (suitably defined) symmetric matrix A .

As we are now working with ± 1 variables instead of $0/1$ variables, one should appropriately modify some of the definitions given earlier in this section. For instance, the condition (33) in the definition of the LS matrix operator M now reads $y_{ii} = y_{00}$ for all $i \in \{1, \dots, n\}$ (in place of $y_{ii} = y_{0i}$) and the (I, J) -th entry of the moment matrix $M_V(y)$ is now $y(I \Delta J)$ (instead of $y(I \cup J)$ as in (27)).

There are two possible strategies for constructing relaxations of the max-cut problem (45). The first possible strategy is to linearize the quadratic objective function, to formulate (45) as a linear problem

$$\max \langle A, X \rangle \text{ subject to } X \in \text{CUT}_n$$

over the *cut polytope*

$$\text{CUT}_n := \text{conv}(xx^T \mid x \in \{\pm 1\}^n),$$

and to apply the various lift-and-project methods to some linear relaxation of CUT_n . As linear programming formulation for CUT_n , one can take the *metric polytope* MET_n which is defined as the set of symmetric matrices X with diagonal entries 1 satisfying the triangle inequalities:

$$X_{ij} + X_{ik} + X_{jk} \geq -1, \quad X_{ij} - X_{ik} - X_{jk} \geq -1$$

for all distinct $i, j, k \in \{1, \dots, n\}$.

Given a graph $G = (V, E)$ ($V = \{1, \dots, n\}$), $\text{CUT}(G)$ and $\text{MET}(G)$ denote, respectively, the projections of CUT_n and MET_n on the subspace \mathbb{R}^E indexed by the edge set of G . Barahona and Mahjoub [29] show that $\text{CUT}(G) \subseteq \text{MET}(G)$ with equality if and only if G has no K_5 -minor. Laurent [149] studies how the Lovász-Schrijver construction applies to the pair $P := \text{CUT}(G)$ and $K := \text{MET}(G)$. The following results are shown there: Equality $N_0^t(\text{MET}(G)) = \text{CUT}(G)$ holds if G has a set of t edges whose contraction produces a graph with no K_5 -minor (recall the definition of N_0 from (35)). In particular, $N^{n-\alpha(G)-3}(\text{MET}(G)) = \text{CUT}(G)$ if G has a maximum stable set whose deletion leaves at most three connected components and $N^{n-\alpha(G)-3}(G) = \text{CUT}(G)$. Here, $N^t(G)$ denotes the projection on the subspace indexed by the edge set of G of the set $N^t(\text{MET}(K_n))$. The inclusion $N^t(G) \subseteq N^t(\text{MET}(G))$ holds obviously. Therefore, the N -rank of $\text{MET}(K_n)$ is at most $n - 4$, with equality for $n \leq 7$ (equality is conjectured for any n). A stronger relaxation is obtained when using the N' operator (recall the definition of N' from (36)). Indeed, $N'(\text{MET}(K_6)) = \text{CUT}(K_6)$ is strictly contained in $N(\text{MET}(K_6))$ and the N' -rank of $\text{MET}(K_n)$ is at most $n - 5$ for $n \geq 6$.

Another possible strategy is to apply the lift-and-project constructions to the set $K := [-1, 1]^n$ and to project on the subspace indexed by the set E_n of all pairs ij of points of V (instead of projecting on the space \mathbb{R}^n indexed by the singletons of V). The SA and Lasserre methods converge now in $n - 1$ steps (as there is no additional linear constraint beside the constraints expressing membership in the cube).

The t -th relaxation in the SA hierarchy is determined by all the inequalities valid for $\text{CUT}(K_n)$ that are induced by at most $t + 1$ points. Thus, the relaxation of order $t = 1$ is the cube $[-1, 1]^E$ while the relaxation of order $t = 2$ is the metric polytope $\text{MET}(K_n)$.

The t -th relaxation in the Lasserre hierarchy, denoted as $Q_t(G)$, is the projection on the subspace \mathbb{R}^E indexed by the edge set of G of the set of vectors y satisfying

$$M_{t+1}(y) = (y_{I\Delta J})_{\substack{I, J \subseteq V \\ |I|, |J| \leq t+1}} \succeq 0, \quad y_\emptyset = 1. \quad (46)$$

Equivalently, one can replace in (46) the matrix $M_{t+1}(y)$ by its principal submatrix indexed by the subsets whose cardinality has the same parity as $t + 1$. Therefore, for $t = 0$, $Q_0(K_n)$ corresponds to the basic semidefinite relaxation

$$\{X = (X_{ij})_{i,j=1}^n \mid X \succeq 0, X_{ii} = 1 \forall i \in \{1, \dots, n\}\}$$

of the cut polytope. For $t = 1$, $Q_1(K_n)$ consists of the vectors $x \in \mathbb{R}^{E_n}$ for which $\begin{pmatrix} 1 \\ x \end{pmatrix} = Y e_0$ for some matrix $Y \succeq 0$ indexed by $\{\emptyset\} \cup E_n$ satisfying

$$Y_{ij,ik} = Y_{\emptyset,jk}, \quad (47)$$

$$Y_{ij,hk} = Y_{ih,jk} = Y_{ik,jh} \quad (48)$$

for all distinct $i, j, h, k \in \{1, \dots, n\}$.

Applying Lagrangian duality to some extended formulation of the max-cut problem, Anjos and Wolkowicz [14] obtained a relaxation F_n of $\text{CUT}(K_n)$, which can be defined as the set of all $x \in \mathbb{R}^{E_n}$ for which $\begin{pmatrix} 1 \\ x \end{pmatrix} = Y e_0$ for some $Y \succeq 0$ indexed by $\{\emptyset\} \cup E_n$ satisfying (47). Thus

$$Q_1(K_n) \subseteq F_n$$

(with strict inclusion if $n \geq 5$). It is interesting to note that the relaxation F_n is stronger than the basic linear relaxation by the metric polytope [14]; that is,

$$F_n \subseteq \text{MET}(K_n).$$

Indeed, let $x \in F_n$ with $\begin{pmatrix} 1 \\ x \end{pmatrix} = Y e_0$ for some $Y \succeq 0$ satisfying (47). The principal submatrix X of Y

indexed by $\{\emptyset, 12, 13, 23\}$ has the form
$$\begin{matrix} & \emptyset & 12 & 13 & 23 \\ \emptyset & \begin{pmatrix} 1 & x_{12} & x_{13} & x_{23} \\ x_{12} & 1 & x_{23} & x_{13} \\ x_{13} & x_{23} & 1 & x_{12} \\ x_{23} & x_{13} & x_{12} & 1 \end{pmatrix} \end{matrix}$$
. Now $e^T X e = 4(1 + x_{12} + x_{13} + x_{23}) \geq 0$

implies one of the triangle inequalities for the triple (1,2,3); the other triangle inequalities follow by suitably flipping signs in X .

Laurent [151] shows that

$$Q_t(G) \subseteq N_+^{t-1}(G)$$

for any $t \geq 1$. Therefore, the second strategy seems to be the most attractive one. Indeed, the relaxation $Q_t(G)$ is at least as tight as $N_+^{t-1}(G)$ and, moreover, it has a simpler explicit description (given by (46)) while the set $N_+^{t-1}(G)$ has only a recursive definition. We refer to [151] for a detailed study of geometric properties of the set of (moment) matrices of the form (46). Laurent [152] shows that the smallest integer t for which $Q_t(K_n) = \text{CUT}(K_n)$ satisfies $t \geq \lceil \frac{n}{2} \rceil - 1$; equality holds for $n \leq 7$ and is conjectured to hold for any n .

Anjos [13] considers higher order semidefinite relaxations for the satisfiability problem involving similar types of constraints as the above relaxations for the cut polytope.

3.7 Further results

Lift-and-project relaxations for the matching and related polytopes. Let $G = (V, E)$ be a graph. A *matching* in G is a set of edges whose incidence vector x satisfies the inequalities

$$x(\delta(v)) = \sum_{e \in \delta(v)} x_e \leq 1 \quad \text{for all } v \in V. \quad (49)$$

(As usual, $\delta(v)$ denotes the set of edges adjacent to v .) Hence, the polytope K consisting of the vectors $x \in [0, 1]^E$ satisfying the inequalities (49) is a linear relaxation of the *matching polytope*² of G , defined as the convex hull of the incidence vectors of all matchings in G . If, in relation (49), we replace the inequality sign ' \leq ' by the equality sign '=' (resp., by the reverse inequality sign ' \geq '), then we obtain the notion of *perfect matching* (resp., of *edge cover*) and the corresponding polytope K is a linear relaxation of the *perfect matching polytope* (resp., of the *edge cover polytope*). Thus, depending on the inequality sign in (49), we obtain three different classes of polytopes.

We now let G be the complete graph on $2n + 1$ nodes. Stephen and Tunçel [223] show that n steps are needed for finding the matching polytope when using the N_+ operator applied to the linear relaxation K . Aguilera, Bianchi and Nasini [3] study the rank of the Balas-Ceria-Cornuéjols procedure and of the N and N_+ operators applied to the linear relaxation K for the three (matching, perfect matching, and edge cover) problems. They show the following results, summarized in Figure 8.

²Of course, the matching polytope of G coincides with the stable set polytope of the line graph L_G of G ; the linear relaxation K considered here is stronger than the linear relaxation $\text{FRAC}(L_G)$ considered in Section 3.5. This implies, e.g., that $N(K) \subseteq N(\text{FRAC}(L_G))$ and analogously for the other lift-and-project methods.

- (i) The BCC rank is equal to n^2 for the three problems.
- (ii) For the perfect matching problem, the rank is equal to n for both the N and N_+ operators.
- (iii) The rank is greater than n for the N operator applied to the matching problem, and for the N and N_+ operators applied to the edge cover problem.

	BCC	N	N_+
matching polytope	n^2	$> n$	n
perfect matching polytope	n^2	n	n
edge cover polytope	n^2	$> n$	$> n$

Figure 8

About the rank of the BCC procedure. Given a graph $G = (V, E)$, the polytope $\text{QSTAB}(G)$, consisting of the vectors $x \in \mathbb{R}_+^V$ satisfying the clique inequalities (42), is a linear relaxation of the stable set polytope $\text{STAB}(G)$, stronger than the fractional stable set polytope $\text{FRAC}(G)$ considered earlier in Section 3.5. Aguilera, Escalante and Nasini [2] show that the rank of the polytope $\text{QSTAB}(G)$ with respect to the Balas-Ceria-Cornuéjols procedure is equal to the rank of $\text{QSTAB}(\overline{G})$, where \overline{G} is the complementary graph of G .

Aguilera, Escalante and Nasini [1] define an extension of the Balas-Ceria-Cornuéjols procedure for up-monotone polyhedra K . Namely, given a subset $F \subseteq \{1, \dots, n\}$, they define the operator $\overline{P}_F(K)$ by

$$\overline{P}_F(K) = P_F(K \cap [0, 1]^n) + \mathbb{R}_+^n,$$

where $P_F(\cdot)$ is the usual BCC operator defined as in (31). Then, the *BCC rank* of K is defined as the smallest $|F|$ for which $\overline{P}_F(K)$ is equal to the convex hull of the integer points in K . It is shown in [1] that, for a clutter \mathcal{C} and its blocker $bl(\mathcal{C})$, the two polyhedra $P_{\mathcal{C}} = \{x \in \mathbb{R}_+^n \mid x(C) \geq 1 \ \forall C \in \mathcal{C}\}$ and $P_{bl(\mathcal{C})} = \{x \in \mathbb{R}_+^n \mid x(D) \geq 1 \ \forall D \in bl(\mathcal{C})\}$ have the *same* rank with respect to the extended BCC procedure.

An extension of lift operators to subset algebras. As we have seen earlier, the lift-and-project methods are based on the idea of lifting a vector $x \in \{0, 1\}^n$ to a higher dimensional vector $y \in \{0, 1\}^N$ (where $N > n$) such that $y_i = x_i$ for all $i = 1, \dots, n$. More precisely, let L denote the lattice of all subsets of $V = \{1, \dots, n\}$ with the set inclusion as order relation, and let Z_L be its Zeta matrix, defined by (26). Then, the lift of $x \in \{0, 1\}^n$ is the vector $y \in \{0, 1\}^L$ with components $y_I = \prod_{i \in I} x_i$ for $I \in L$; in other words, y is the column of Z_L indexed by x (after identifying a set with its incidence vector).

Bienstock and Zuckerberg [38] push this idea further and introduce a lifting to a lattice Σ , larger than L . Namely, let Σ denote the lattice of all subsets of $\{0, 1\}^n$, with the reverse set inclusion as order relation; that is, $\alpha \leq \beta$ in Σ if $\beta \subseteq \alpha$. Let Z_Σ denote the Zeta matrix of Σ , with (α, β) -entry 1 if $\alpha \leq \beta$ and 0 otherwise. Then, any vector $x \in \{0, 1\}^n$ can be lifted to the vector $z \in \{0, 1\}^\Sigma$ with components $z_\alpha = 1$ if and only if $x \in \alpha$ (for $\alpha \in \Sigma$); that is, z is the column of Z_Σ indexed by $\{x\}$.

Note that the lattice L is isomorphic to a sublattice of Σ . Indeed, if we set $H_I = \{x \in \{0, 1\}^n \mid x_i = 1 \ \forall i \in I\}$ for $I \subseteq V$, then $I \subseteq J \iff H_I \supseteq H_J \iff H_I \leq H_J$ (in Σ) and, thus, the mapping $I \mapsto H_I$ maps L to a sublattice of Σ . Therefore, given $x \in \{0, 1\}^n$ and, as above, y (resp., z) the column of Z_L (resp., of Z_Σ) indexed by x , then $z_{H_I} = y_I$ for all $I \in L$ and $z_{H_i} = x_i$ for all $i \in V$.

Let $F \subseteq \{0, 1\}^n$ be the set of 0–1 points whose convex hull $P := \text{conv}(F)$ has to be found, and let F_L (resp., F_Σ) be the corresponding set of columns of Z_L (resp., of Z_Σ). Then, a vector $x \in \mathbb{R}^n$ belongs to $\text{conv}(F)$ if and only if there exists $y \in \text{conv}(F_L)$ such that $y_i = x_i$ ($i \in V$) or, equivalently, if there exists $z \in \text{conv}(F_\Sigma)$ such that $z_{H_i} = x_i$ ($i \in V$). The SA, LS and Lasserre methods consist of requiring certain conditions on the lifted vector y (or projections of it); Bienstock and Zuckerberg [38] present analogous conditions for the vector z .

Bienstock and Zuckerberg work, in fact, with a lifted vector \tilde{z} indexed by a small subset ν of Σ ; this set ν is constructed on the fly, depending on the structure of F . Consider, for instance, the set covering problem, where F is the set of 0/1 solutions of a system: $x(A_1) \geq 1, \dots, x(A_m) \geq 1$ (with $A_1, \dots, A_m \subseteq \{1, \dots, n\}$). Then, the most basic lifting procedure presented in [38] produces a polyhedron $R^{(2)}$ (whose projection is a linear relaxation of P) in the variable $\tilde{z} \in \mathbb{R}^\nu$, where $\nu \subseteq \Sigma$ consists of F , $Y_i := \{x \in F \mid x_i = 1\}$, $N_i := F \setminus Y_i$ ($i = 1, \dots, n$), and $\cap_{i \in C} N_i$, $Y_{i_0} \cap \cap_{i \in C \setminus i_0} N_i$ ($i_0 \in C$), and $\cup_{S \subseteq C, |S| \geq 2} \cap_{i \in S} Y_i \cap \cap_{i \in C \setminus S} N_i$, for each of the distinct intersections $C = A_h \cap A_\ell$ ($h \neq \ell = 1, \dots, m$) with size ≥ 2 . The linear relaxation $R^{(2)}$ has $O(m^4 n^2)$ variables and constraints; hence, one can optimize over $R^{(2)}$ in polynomial time. Moreover, any inequality $a^T x \geq a_0$, valid for P with coefficients in $\{0, 1, 2\}$, is valid for (the projection of) $R^{(2)}$. Note that there exist set covering polytopes having exponentially many facets with coefficients in $\{0, 1, 2\}$. The new lifting procedure is more powerful in some cases. For instance, $R^{(2)} = P$ holds for the polytope K from (38), while the N_+ -rank of K is equal to n . As another example, consider the circulant set covering polytope:

$$P = \text{conv}(\{x \in \{0, 1\}^n \mid \sum_{i \neq j} x_i \geq 1 \ \forall j = 1, \dots, n\});$$

then the inequality $\sum_{i=1}^n x_i \geq 2$ is valid for P , it is not valid neither for $S_{n-3}(K)$ nor for $N_{n-3}^+(K)$, while it is valid for the relaxation $R^{(2)}$ [38].

A more sophisticated lifting procedure is proposed in [38] yielding stronger relaxations $R^{(k)}$ of P , with the following properties. For fixed $k \geq 2$, one can optimize in polynomial time over $R^{(k)}$; any inequality $a^T x \geq a_0$, valid for P with³ coefficients in $\{0, 1, \dots, k\}$, is valid for $R^{(k)}$. For instance, $R^{(3)} = \emptyset$ holds for the polytope K from (39), while n steps of the classic lift-and-project procedures are needed for proving that $P = \emptyset$.

Complexity of cutting plane proofs. Results about the complexity of cutting plane proofs using cuts produced by the various lift-and-project methods can be found, e.g., in [62], [63], [97].

3.8 Extensions to polynomial programming

Quadratic programming. Suppose we want to solve the program

$$p^* := \min g_0(x) \text{ subject to } g_\ell(x) \geq 0 \ (\ell = 1, \dots, m) \quad (50)$$

where g_0, g_1, \dots, g_m are quadratic functions of the form: $g_\ell(x) = x^T Q_\ell x + 2q_\ell^T x + \gamma_\ell$ (Q_ℓ symmetric $n \times n$ matrix, $q_\ell \in \mathbb{R}^n$, $\gamma_\ell \in \mathbb{R}$). For any ℓ , define the matrix $P_\ell := \begin{pmatrix} \gamma_\ell & q_\ell^T \\ q_\ell & Q_\ell \end{pmatrix}$. Then, $g_\ell(x) = \langle P_\ell, \begin{pmatrix} 1 & x^T \\ x & xx^T \end{pmatrix} \rangle$. This suggests the following natural positive semidefinite relaxation of (50):

$$\min \langle P_0, Y \rangle \text{ subject to } Y \succeq 0, Y_{00} = 1, \langle P_\ell, Y \rangle \geq 0 \ (\ell = 1, \dots, m). \quad (51)$$

³Validity holds, more generally, for any inequality $a^T x \geq a_0$ with $\text{pitch} \leq k$. If we order the indices in such a way that $0 < a_1 \leq a_2 \leq \dots \leq a_J$, $a_{J+1} = \dots = a_n = 0$, then the *pitch* is the smallest t for which $\sum_{j=1}^t a_j \geq a_0$.

Let $F := \{x \in \mathbb{R}^n \mid g_\ell(x) \geq 0 \ (\ell = 1, \dots, m)\}$ denote the feasible set of (50) and

$$\hat{F} := \{x \in \mathbb{R}^n \mid \begin{pmatrix} 1 \\ x \end{pmatrix} = Y e_0 \text{ for some } Y \succeq 0 \text{ satisfying } \langle P_\ell, Y \rangle \geq 0 \text{ for all } \ell = 1, \dots, m\} \quad (52)$$

its natural semidefinite relaxation. It is shown in [88, 139] that \hat{F} can be alternatively described by the following quadratic system:

$$\hat{F} = \{x \in \mathbb{R}^n \mid \sum_{\ell=1}^m t_\ell g_\ell(x) \geq 0 \text{ for all } t_\ell \geq 0 \text{ for which } \sum_{\ell=1}^m t_\ell Q_\ell \preceq 0\}. \quad (53)$$

If, in (52), one omits the condition $Y \succeq 0$ and, in (53), the condition $\sum_{\ell} t_\ell Q_\ell \preceq 0$ is replaced by $\sum_{\ell} t_\ell Q_\ell = 0$, then one obtains a linear relaxation \hat{F}_L of F such that $\text{conv}(F) \subseteq \hat{F} \subseteq \hat{F}_L$.

Using this construction of linear/semidefinite relaxations, Kojima and Tunçel [139] construct a hierarchy of successive relaxations of F that converges asymptotically to $\text{conv}(F)$. Lasserre [141] also constructs such a hierarchy which applies, more generally, to polynomial programs; we expose it below.

Polynomial programming. Consider now the program (50) where all the g_ℓ 's are polynomials in $x = (x_1, \dots, x_n)$. Let w_ℓ be the degree of g_ℓ , $v_\ell := \lceil \frac{w_\ell}{2} \rceil$ and $v := \max_{\ell=1, \dots, m} v_\ell$. We need some definitions.

Given a sequence $y = (y_\alpha)_{\alpha \in \mathbb{Z}_+^n}$ indexed by \mathbb{Z}_+^n , its *moment matrix* is

$$M^{\mathbb{Z}}(y) := (y_{\alpha+\beta})_{\alpha, \beta \in \mathbb{Z}_+^n} \quad (54)$$

and, given an integer $t \geq 0$, $M_t^{\mathbb{Z}}(y)$ is the principal submatrix of $M^{\mathbb{Z}}(y)$ indexed by the sequences $\alpha \in \mathbb{Z}_+^n$ with $|\alpha| := \sum_i \alpha_i \leq t$. [Note that the moment matrix $M_V(y)$ defined earlier in (27) corresponds to the principal submatrix of $M^{\mathbb{Z}}(y)$ indexed by the sequences $\alpha \in \{0, 1\}^n$, after replacing y_α by $y_{\alpha'}$ where $\alpha'_i := \min(\alpha_i, 1)$ for all i .] The operation from (29) extends to sequences indexed by \mathbb{Z}_+^n in the following way:

$$g, y \in \mathbb{R}^{\mathbb{Z}_+^n} \rightsquigarrow g * y := \left(\sum_{\beta} g_\beta y_{\alpha+\beta} \right)_{\alpha \in \mathbb{Z}_+^n}. \quad (55)$$

Given $x \in \mathbb{R}^n$, define the sequence $y \in \mathbb{R}^{\mathbb{Z}_+^n}$ with α -th entry $y_\alpha := \prod_{i=1}^n x_i^{\alpha_i}$ for $\alpha \in \mathbb{Z}_+^n$. Then, $M_t^{\mathbb{Z}}(y) = yy^T \succeq 0$ (where we use the same symbol y for denoting the truncated vector $(y_\alpha)_{|\alpha| \leq t}$) and $M_t^{\mathbb{Z}}(g_\ell * y) = g_\ell(x) \cdot M_t^{\mathbb{Z}}(y) \succeq 0$ if $g_\ell(x) \geq 0$. This observation leads naturally to the following relaxations of the set F , introduced by Lasserre [141].

For $t \geq v - 1$, let $\mathcal{Q}_t(F)$ be the convex set defined as the projection of the solution set to the system

$$M_{t+1}^{\mathbb{Z}}(y) \succeq 0, M_{t-v_\ell+1}^{\mathbb{Z}}(g_\ell * y) \succeq 0 \text{ for } \ell = 1, \dots, m, y_0 = 1 \quad (56)$$

on the subspace \mathbb{R}^n indexed by the variables y_α for $\alpha = (1, 0, \dots, 0), \dots, (0, \dots, 0, 1)$ (identified with x_1, \dots, x_n). Then,

$$\text{conv}(F) \subseteq \mathcal{Q}_{t+1}(F) \subseteq \mathcal{Q}_t(F).$$

Lasserre [141] shows that

$$\bigcap_{t \geq v-1} \mathcal{Q}_t(F) = \text{conv}(F);$$

that is, the hierarchy $(\mathcal{Q}_t(F))_t$ converges asymptotically to $\text{conv}(F)$. This equality holds under some technical assumption on F which holds, for instance, when F is the set of 0/1 solutions of a polynomial system and the constraints $x_i(1-x_i) = 0$ ($i \in \{1, \dots, n\}$) are present in the description of F , or when the set $\{x \mid g_\ell(x) \geq 0\}$ is compact for at least one of the constraints defining F . Lasserre's result relies

on a result about representations of positive polynomials as sums of squares, to which we will come back in Section 7.1.

In the quadratic case, when all g_ℓ are quadratic polynomials, one can verify that the first Lasserre relaxation $\mathcal{Q}_0(F)$ coincides with the basic SDP relaxation \hat{F} defined in (52); that is,

$$\mathcal{Q}_0(F) = \hat{F}.$$

Consider now the 0/1 case when F is the set of 0/1 solutions of a polynomial system; write F as

$$F = \{x \in \mathbb{R}^n \mid g_\ell(x) \geq 0 \ (\ell = 1, \dots, m), \ h_i(x) := x_i - x_i^2 = 0 \ (i = 1, \dots, n)\}.$$

One can assume without loss of generality that each g_ℓ has degree at most 1 in every variable. The set

$$K := \{x \in [0, 1]^n \mid g_\ell(x) \geq 0 \ (\ell = 1, \dots, m)\}$$

is a natural relaxation of F . We have constructed in Section 3.4 the successive relaxations $Q_t(K)$ of $\text{conv}(F)$ satisfying $\text{conv}(F) = Q_{n+v-1}(K)$; their construction used moment matrices indexed by the subsets of V while the definition of $Q_t(F)$ involves moment matrices indexed by integer sequences. However, the condition $M_t^{\mathbb{Z}}(h_i * y) = 0$ (present in the definition of $Q_t(F)$) permits to show that the two definitions are equivalent; that is,

$$Q_t(K) = Q_t(F) \ \text{for } t \geq v - 1.$$

See [150] for details.

In the quadratic 0/1 case, we find therefore that

$$\hat{F} = \mathcal{Q}_0(F) = Q_0(K).$$

As an example, given a graph $G = (V = \{1, \dots, n\}, E)$, consider the set

$$F := \{x \in \{0, 1\}^n \mid x_i x_j = 0 \ \text{for all } ij \in E\};$$

then $\text{conv}(F)$ is equal to the stable set polytope of G . It follows from the definitions that \hat{F} coincides with the basic SDP relaxation $\text{TH}(G)$ (defined in (44)). Therefore, $\mathcal{Q}_0(F) = \text{TH}(G)$ while the inclusion $\text{TH}(G) \subseteq Q_0(\text{FRAC}(G))$ is strict in general. Hence one obtains stronger relaxations for the stable set polytope $\text{STAB}(G)$ when starting from the above quadratic representation F for stable sets rather than from the linear relaxation $\text{FRAC}(G)$. Applying the equivalent definition (53) for \hat{F} , one finds that

$$\text{TH}(G) = \left\{ x \in \mathbb{R}^n \mid x^T M x - \sum_{i=1}^n M_{ii} x_i \leq 0 \ \text{for } M \succeq 0 \ \text{with } M_{ij} = 0 \ (i \neq j \in V, ij \notin E) \right\}. \quad (57)$$

(This formulation of $\text{TH}(G)$ also follows using the duality between the cone of completable partial positive semidefinite matrices and the cone of positive semidefinite matrices having zeros at the positions of unspecified entries; cf. [148].) See Section 4.2 for further information about the semidefinite relaxation $\text{TH}(G)$.

4 Semidefinite Relaxation for the Maximum Stable Set Problem

Given a graph $G = (V, E)$, its *stability number* $\alpha(G)$ is the maximum cardinality of a stable set in G , and its *clique number* $\omega(G)$ is the maximum cardinality of a clique in G . Given an integer $k \geq 1$, a k -*colouring* of G is an assignment of numbers from $\{1, \dots, k\}$ (*colours*) to the nodes of G in such a way that adjacent nodes receive distinct colours; in other words, a k -colouring is a partition of V into k stable sets. The *colouring number* (or *chromatic number*) $\chi(G)$ is the smallest integer k for which G has a k -colouring. With $\overline{G} = (V, \overline{E})$ denoting the complementary graph of G , the following holds trivially:

$$\alpha(\overline{G}) = \omega(G) \leq \chi(G).$$

The inequality $\omega(G) \leq \chi(G)$ is strict, for instance, for odd circuits of length ≥ 5 and their complements. Berge [36] defined a graph G to be *perfect* if $\omega(G') = \chi(G')$ for every induced subgraph G' of G and he conjectured that a graph is perfect if and only if it does not contain a circuit of length ≥ 5 or its complement as an induced subgraph; this is the well known *strong perfect graph conjecture*. M. Chudnovsky and P.D. Seymour announced in May 2002 that they could prove the strong perfect graph conjecture. Lovász [163] proved that the complement of a perfect graph is again perfect, solving another conjecture of Berge. As we will see later in this section, perfect graphs can also be characterized in terms of integrality of certain associated polyhedra.

Computing the stability number or the chromatic number of a graph are hard problems; more precisely, given an integer k , it is an NP-complete problem to decide whether $\alpha(G) \geq k$ or $\chi(G) \leq k$ [128]. Deciding whether a graph is 2-colourable can be done in polynomial time (as this happens if and only if the graph is bipartite). On the other hand, while every planar graph is 4-colourable (by the celebrated four colour theorem), it is NP-complete to decide whether a planar graph is 3-colourable [90]. When restricted to the class of perfect graphs, the maximum stable set problem and the colouring problem can be solved in polynomial time. This result relies on the use of the Lovász theta function $\vartheta(G)$ which can be computed (with an arbitrary precision) in polynomial time (as the optimum of a semidefinite program) and satisfies the ‘sandwich’ inequalities:

$$\alpha(G) \leq \vartheta(G) \leq \chi(\overline{G}).$$

The polynomial time solvability of the maximum stable set problem for perfect graphs is one of the first beautiful applications of semidefinite programming to combinatorial optimization and, up to today, no other purely combinatorial method is known for proving this.

4.1 The basic linear relaxation

As before, the *stable set polytope* $\text{STAB}(G)$ is the polytope in \mathbb{R}^V defined as the convex hull of the incidence vectors of the stable sets of G , $\text{FRAC}(G)$ is its linear relaxation defined by nonnegativity and the edge inequalities (40), and $\text{QSTAB}(G)$ denotes the linear relaxation of $\text{STAB}(G)$ defined by nonnegativity and the clique inequalities (42). Therefore,

$$\text{STAB}(G) \subseteq \text{QSTAB}(G) \subseteq \text{FRAC}(G)$$

and

$$\alpha(G) = \max(e^T x \mid x \in \text{STAB}(G))$$

setting $e := (1, \dots, 1)^T$. One can easily see that equality $\text{STAB}(G) = \text{FRAC}(G)$ holds if and only if G is a bipartite graph with no isolated nodes; thus the maximum stable set problem for bipartite

graphs can be solved in polynomial time as a linear programming problem over $\text{FRAC}(G)$. Fulkerson [89] and Chvátal [53] show:

Theorem 9. *A graph G is perfect if and only if $\text{STAB}(G) = \text{QSTAB}(G)$.*

This result does not (yet) help for computing efficiently $\alpha(G)$ for perfect graphs. Indeed, optimizing over the linear relaxation $\text{QSTAB}(G)$ is, unfortunately, a hard problem in general (as hard as the original problem, since the membership problem for $\text{QSTAB}(G)$ is nothing but a maximum weight clique problem in G .) Proving polynomiality requires the use of the semidefinite relaxation $\text{TH}(G)$ as we see later in this section.

4.2 The theta function $\vartheta(G)$ and the basic semidefinite relaxation $\text{TH}(G)$

Lovász [164] introduced the following parameter $\vartheta(G)$, known as the *theta number*:

$$\begin{aligned} \vartheta(G) := \max \quad & e^T X e \\ \text{s.t.} \quad & \text{Tr}(X) = 1 \\ & X_{ij} = 0 \quad (i \neq j, ij \in E) \\ & X \succeq 0. \end{aligned} \tag{58}$$

The theta number has two important properties: it can be computed with an arbitrary precision in polynomial time (as the optimum value of a semidefinite program) and it provides bounds for the stability and chromatic numbers. Namely,

$$\alpha(G) \leq \vartheta(G) \leq \chi(\overline{G}). \tag{59}$$

To see that $\alpha(G) \leq \vartheta(G)$, consider a maximum stable set S ; then the matrix $X := \frac{1}{|S|} \chi^S (\chi^S)^T$ is feasible for the program (58) and $\alpha(G) = e^T X e$. To see that $\vartheta(G) \leq \chi(\overline{G})$, consider a matrix X feasible for (58) and a partition $V = Q_1 \cup \dots \cup Q_k$ into $k := \chi(\overline{G})$ cliques. Then,

$$0 \leq \sum_{h=1}^k (k \chi^{Q_h} - e)^T X (k \chi^{Q_h} - e) = k^2 \text{Tr}(X) - k e^T X e = k^2 - k e^T X e,$$

which implies $e^T X e \leq k$ and thus $\vartheta(G) \leq \chi(\overline{G})$.

Several equivalent definitions are known for $\vartheta(G)$ that we recall below. (See [100] or [137] for a detailed treatment, and [102] for an algorithmic comparison.) The dual semidefinite program of (58) reads:

$$\min(t \mid tI + \sum_{ij \in E} \lambda_{ij} E_{ij} - J \succeq 0), \tag{60}$$

where $J := ee^T$ is the all ones matrix and E_{ij} is the elementary matrix with all zero entries except 1 at positions (i, j) and (j, i) . As the program (58) has a strictly feasible solution (e.g., $X = \frac{1}{n}I$), there is no duality gap and the optimum value of (60) is equal to the theta number $\vartheta(G)$. Setting $Y := J - \sum_{ij \in E} \lambda_{ij} E_{ij}$, $Z := tI - Y$ and $U := \frac{1}{t-1}Z$ in (60), we obtain the following reformulations for

$\vartheta(G)$:

$$\begin{aligned} \vartheta(G) = \min \quad & \lambda_{\max}(Y) \\ \text{s.t.} \quad & Y_{ij} = 1 \quad (i = j \text{ or } ij \in \overline{E}) \\ & Y \text{ symmetric matrix,} \end{aligned} \tag{61}$$

$$\begin{aligned} \vartheta(G) = \min t & & = \min t \\ \text{s.t. } Z_{ii} = t - 1 & \quad (i \in V) & \text{s.t. } U_{ii} = 1 & \quad (i \in V) \\ Z_{ij} = -1 & \quad (ij \in \overline{E}) & U_{ij} = -\frac{1}{t-1} & \quad (ij \in \overline{E}) \\ Z \succeq 0 & & U \succeq 0, t \geq 2. & \end{aligned} \quad (62)$$

The formulation (62) will be used later in Section 6 for the colouring and max k -cut problems. One can also express $\vartheta(G)$ as the optimum value of the linear objective function $e^T x$ maximized over a convex set forming a relaxation of $\text{STAB}(G)$. Namely, let \mathcal{M}_G denote the set of positive semidefinite matrices Y indexed by the set $V \cup \{0\}$ satisfying $y_{ii} = y_{0i}$ for $i \in V$ and $y_{ij} = 0$ for $i \neq j \in V$ adjacent in G , and set

$$\text{TH}(G) := \{x \in \mathbb{R}^V \mid \begin{pmatrix} 1 \\ x \end{pmatrix} = Y e_0 \text{ for some } Y \in \mathcal{M}_G\}, \quad (63)$$

where $e_0 := (1, 0, \dots, 0)^T \in \mathbb{R}^{n+1}$. (Same definition as (44).)

Lemma 10. *For any graph G , $\text{STAB}(G) \subseteq \text{TH}(G) \subseteq \text{QSTAB}(G)$.*

PROOF. If S is a stable set in G and $x := \chi^S$, then $Y := \begin{pmatrix} 1 \\ x \end{pmatrix} \begin{pmatrix} 1 & x^T \end{pmatrix} \in \mathcal{M}_G$ and $\begin{pmatrix} 1 \\ x \end{pmatrix} = Y e_0$; from this follows that $\text{STAB}(G) \subseteq \text{TH}(G)$. Let $x \in \text{TH}(G)$, $Y \in \mathcal{M}_G$ such that $\begin{pmatrix} 1 \\ x \end{pmatrix} = Y e_0$, and let Q be a clique in G . The principal submatrix Y_Q of Y whose rows and columns are indexed by the set $\{0\} \cup Q$ has the form $\begin{pmatrix} 1 & x^T \\ x & \text{diag}(x) \end{pmatrix}$. As $Y \succeq 0$, we have $Y_Q \succeq 0$, i.e., $\text{diag}(x) - x x^T \succeq 0$ (taking a Schur complement), which implies that $e^T (\text{diag}(x) - x x^T) e = e^T x (1 - e^T x) \geq 0$ and thus $e^T x = \sum_{i \in Q} x_i \leq 1$. This shows the inclusion $\text{TH}(G) \subseteq \text{QSTAB}(G)$. \blacksquare

Theorem 11. $\vartheta(G) = \max(e^T x \mid x \in \text{TH}(G))$.

PROOF. We use the formulation of $\vartheta(G)$ from (58). Let μ_G denote the maximum of $e^T x$ over $\text{TH}(G)$. We first show that $\vartheta(G) \leq \mu_G$. For this, let X be an optimum solution to the program (58). Let $v_1, \dots, v_n \in \mathbb{R}^n$ such that $x_{ij} = v_i^T v_j$ for all $i, j \in V$; thus $\vartheta(G) = \|\sum_{i=1}^n v_i\|^2$, $\sum_{i=1}^n (v_i)^2 = \text{Tr}(X) = 1$, and $v_i^T v_j = 0$ if i, j are adjacent in G . Set $P := \{i \in V \mid v_i \neq 0\}$, $u_0 := \frac{1}{\sqrt{\vartheta(G)}} \sum_{i=1}^n v_i$, $u_i := \frac{v_i}{\|v_i\|}$ for $i \in P$, and let u_i ($i \in V \setminus P$) be an orthonormal basis of the orthogonal complement of the space spanned by $\{v_i \mid i \in P\}$. Let D denote the diagonal matrix indexed by $\{0\} \cup V$ with diagonal entries $u_0^T u_i$ ($i = 0, 1, \dots, n$), let Z denote the Gram matrix of u_0, u_1, \dots, u_n and set $Y := D Z D$, with entries $y_{ij} = (u_i^T u_j)(u_0^T u_i)(u_0^T u_j)$ ($i, j = 0, 1, \dots, n$). Then, $Y \in \mathcal{M}_G$ with $y_{00} = 1$. It remains to verify that $\vartheta(G) \leq \sum_{i=1}^n y_{0i}$. By the definition of u_0 , we find

$$\vartheta(G) = \left(\sum_{i=1}^n u_0^T v_i \right)^2 = \left(\sum_{i \in P} u_0^T v_i \right)^2 = \left(\sum_{i \in P} u_0^T u_i \|v_i\| \right)^2 \leq \left(\sum_{i \in P} \|v_i\|^2 \right) \left(\sum_{i \in P} (u_0^T u_i)^2 \right) = \sum_{i=1}^n y_{0i},$$

where the inequality follows using the Cauchy-Schwartz inequality. We now show the converse inequality $\mu_G \leq \vartheta(G)$. For this, let $x \in \text{TH}(G)$ be optimum for the program defining μ_G , let $Y \in \mathcal{M}_G$ such that $\begin{pmatrix} 1 \\ x \end{pmatrix} = Y e_0$, and $v_0, v_1, \dots, v_n \in \mathbb{R}^{n+1}$ such that $y_{ij} = v_i^T v_j$ for all $i, j = 0, 1, \dots, n$. It suffices to construct X feasible for (58) satisfying $\sum_{i,j=1}^n x_{ij} \geq \mu_G$. Define the $n \times n$ matrix X with entries $x_{ij} := \frac{1}{\mu_G} v_i^T v_j$ ($i, j = 1, \dots, n$); then X is feasible for (58). Moreover, $\mu_G = \sum_{i=1}^n y_{0i} = \sum_{i=1}^n v_0^T v_i = v_0^T (\sum_{i=1}^n v_i)$ is less than or equal to $\|\sum_{i=1}^n v_i\|$ (by the Cauchy-Schwartz inequality, since $\|v_0\| = 1$).

As $\sum_{i,j=1}^n x_{ij} = \frac{1}{\mu_G} (\sum_{i=1}^n v_i)^2$, we find that $\mu_G \leq \sum_{i,j=1}^n x_{ij}$. ▀

An *orthonormal representation* of G is a set of unit vectors $u_1, \dots, u_n \in \mathbb{R}^N$ ($N \geq 1$) satisfying $u_i^T u_j = 0$ for all $ij \in \bar{E}$.

Theorem 12. $\vartheta(G) = \max_{d, v_i} \sum_{i \in V} (d^T v_i)^2$, where the maximum is taken over all unit vectors $d \in \mathbb{R}^N$ and all orthonormal representations $v_1, \dots, v_n \in \mathbb{R}^N$ of \bar{G} .

PROOF. Let $\vartheta(G) = e^T X e$, where X is an optimum solution to the program (58) and let b_1, \dots, b_n be vectors such that $X_{ij} = b_i^T b_j$ for $i, j \in V$. Set $d := \frac{\sum_{i \in V} b_i}{\|\sum_{i \in V} b_i\|}$, $P := \{i \in V \mid b_i \neq 0\}$ and $v_i := \frac{b_i}{\|b_i\|}$ for $i \in P$. Let v_i ($i \in V \setminus P$) be an orthonormal basis of the orthogonal complement of the space spanned by v_i ($i \in P$). Then, v_1, \dots, v_n is an orthonormal representation of \bar{G} . We have:

$$\sqrt{\vartheta(G)} = \left\| \sum_{i \in P} b_i \right\| = d^T \left(\sum_{i \in P} b_i \right) = \sum_{i \in P} \|b_i\| v_i^T d \leq \sqrt{\sum_{i \in P} \|b_i\|^2} \cdot \sqrt{\sum_{i \in P} (v_i^T d)^2} \leq \sqrt{\sum_{i \in V} (v_i^T d)^2}$$

(using the Cauchy-Schwartz inequality and $\text{Tr}(X) = 1$). This implies that $\vartheta(G) \leq \sum_{i \in V} (d^T v_i)^2$.

Conversely, let d be a unit vector and let v_1, \dots, v_n be an orthonormal representation of \bar{G} . Let Y denote the Gram matrix of the vectors $d, (d^T v_1)v_1, \dots, (d^T v_n)v_n$. Then, $Y \in \mathcal{M}_G$. Therefore, $((d^T v_1)^2, \dots, (d^T v_n)^2)^T \in \text{TH}(G)$ which implies that $\sum_{i \in V} (d^T v_i)^2 \leq \vartheta(G)$. ▀

Let A_G denote the convex hull of all vectors $((d^T v_1)^2, \dots, (d^T v_n)^2)^T$ where d is a unit vector and v_1, \dots, v_n is an orthonormal representation of \bar{G} , let B_G denote the set of $x \in \mathbb{R}_+^V$ satisfying the *orthonormal representation constraints*:

$$\sum_{i \in V} (c^T u_i)^2 x_i \leq 1 \tag{64}$$

for all unit vectors c and all orthonormal representations u_1, \dots, u_n of G , and let C_G denote the set of $x \in \mathbb{R}_+^V$ satisfying

$$\sum_{i \in V} x_i \leq \min_{c, u_i} \max_{i \in V} \frac{1}{(c^T u_i)^2}$$

where the minimum is taken over all unit vectors c and all orthonormal representations u_1, \dots, u_n of G .

Lemma 13. $A_G \subseteq \text{TH}(G) \subseteq B_G \subseteq C_G$.

PROOF. The inclusion $A_G \subseteq \text{TH}(G)$ follows from the second part of the proof of Theorem 12 and the inclusion $B_G \subseteq C_G$ is easy to verify. Let $x \in \text{TH}(G)$ and let $z := ((c^T u_1)^2, \dots, (c^T u_n)^2)^T$ where c is a unit vector and u_1, \dots, u_n is an orthonormal representation of G ; we show that $x^T z \leq 1$. By the above, $z \in A_{\bar{G}} \subseteq \text{TH}(\bar{G})$. Let $Y \in \mathcal{M}_G$ and $Z \in \mathcal{M}_{\bar{G}}$ such that $\begin{pmatrix} 1 \\ x \end{pmatrix} = Y e_0$ and $\begin{pmatrix} 1 \\ z \end{pmatrix} = Z e_0$. Denote by Y' the matrix obtained from Y by changing the signs on its first row and column. Then, $\langle Y', Z \rangle = 1 - 2 \sum_{i \in V} y_{0i} z_{0i} + \sum_{i \in V} y_{ii} z_{ii} = 1 - \sum_{i \in V} x_i z_i \geq 0$ (since $Y', Z \succeq 0$) and thus $x^T z \leq 1$. This shows the inclusion $\text{TH}(G) \subseteq B_G$. ▀

Theorem 14. $\vartheta(G) = \min_{c, u_i} \max_{i \in V} \frac{1}{(c^T u_i)^2}$, where the minimum is taken over all unit vectors c and all orthonormal representations u_1, \dots, u_n of G .

PROOF. The inequality $\vartheta(G) \leq \min \dots$ follows from the inclusion $\text{TH}(G) \subseteq C_G$ and Theorem 11. For the reverse inequality, we use the definition of $\vartheta(G)$ from (61). Let Y be a symmetric matrix with $Y_{ii} = 1$ ($i \in V$) and $Y_{ij} = 1$ ($ij \in \overline{E}$) and $\vartheta(G) = \lambda_{\max}(Y)$. As $\vartheta(G)I - Y \succeq 0$, there exist vectors b_1, \dots, b_n such that $b_i^2 = \vartheta(G) - 1$ ($i \in V$) and $b_i^T b_j = -1$ ($ij \in \overline{E}$). Let c be a unit vector orthogonal to all b_i (which exists since $\vartheta(G)I - Y$ is singular) and set $u_i := \frac{c + b_i}{\sqrt{\vartheta(G)}}$ ($i \in V$). Then, u_1, \dots, u_n is an orthonormal representation of G and $\vartheta(G) = \frac{1}{(c^T u_i)^2}$ for all i . \blacksquare

Theorems 12 and 14 and Lemma 13 show that one obtains the same optimum value when optimizing the linear objective function $e^T x$ over $\text{TH}(G)$ or over any of the sets A_G, B_G or C_G . In fact, the same remains true for an arbitrary linear objective function $w^T x$ where $w \in \mathbb{R}_+^V$, as the above extends easily to the weighted case. Therefore,

$$\text{TH}(G) = A_G = B_G = C_G.$$

Moreover, $\text{TH}(\overline{G})$ is the *antiblocker* of $\text{TH}(G)$; that is, $\text{TH}(\overline{G}) = \{z \in \mathbb{R}_+^V \mid x^T z \leq 1 \ \forall x \in \text{TH}(G)\}$. One can show that the only orthonormal representation inequalities (64) defining facets of $\text{TH}(G)$ are the clique inequalities. From this follows:

$$\text{TH}(G) \text{ is a polytope} \iff G \text{ is perfect} \iff \text{TH}(G) = \text{QSTAB}(G) \iff \text{TH}(G) = \text{STAB}(G).$$

We refer to ([204], chapter 12) for a detailed exposition on the theta body $\text{TH}(G)$.

4.3 Colouring and finding maximum stable sets in perfect graphs

The stability number $\alpha(G)$ and the chromatic number $\chi(G)$ of a perfect graph G can be computed in polynomial time. (Indeed, it suffices to compute an approximated value of $\vartheta(G)$ with precision $< 1/2$ in order to determine $\alpha(G) = \chi(\overline{G}) = \vartheta(G)$.) We now mention how to find in polynomial time a stable set of size $\alpha(G)$ and a $\chi(G)$ -colouring in a perfect graph. The weighted versions of these problems can also be solved in polynomial time (cf. [100] for details).

Finding a maximum cardinality stable set in a perfect graph. Let $G = (V, E)$ be a perfect graph and let v_1, \dots, v_n be an ordering of its nodes. We construct a sequence of graphs $G_0 := G \supseteq G_1 \supset \dots \supseteq G_i \supseteq G_{i+1} \supseteq \dots \supseteq G_n$ in the following manner: For each $i \geq 1$, compute $\alpha(G_{i-1} \setminus v_i)$; if $\alpha(G_{i-1} \setminus v_i) = \alpha(G)$, then set $G_i := G_{i-1} \setminus v_i$, otherwise set $G_i := G_{i-1}$. Then, $\alpha(G_i) = \alpha(G)$ for all i and G_n is a stable set, thus providing a maximum stable set in G . Therefore, a maximum stable set in a perfect graph G can be found by applying n times an algorithm for computing the theta function.

Finding a minimum colouring in a perfect graph. We follow the presentation of Schrijver [210]. Let $G = (V, E)$ be a perfect graph. A crucial observation is that it suffices to find a stable set S which intersects all the maximum cardinality cliques of G . Indeed, if such S is found, then one can recursively colour $G \setminus S$ with $\omega(G \setminus S) = \omega(S) - 1$ colours and thus G with $\omega(G) = \chi(G)$ colours. For $t \geq 1$, we grow iteratively a list Q_1, \dots, Q_t of maximum cardinality cliques. Suppose Q_1, \dots, Q_t have been found. We begin with finding a stable set S meeting each of Q_1, \dots, Q_t . For this, setting $w := \sum_{i=1}^t \chi^{Q_i}$, it suffices to find a maximum weight stable set S . (This can be done by applying the above maximum cardinality stable set algorithm to the graph G' obtained from G by replacing every

node i by a set W_i of w_i nonadjacent nodes, making two nodes $u \in W_i, v \in W_j$ adjacent in G' if the nodes i, j are adjacent in G .) Then S has weight t which means that S meets each of Q_1, \dots, Q_t . Now, if $\omega(G \setminus S) < \omega(G)$, then S meets *all* the maximum cardinality cliques in G and we are done. Otherwise, we find a clique Q_{t+1} in $G \setminus S$ of size $\omega(G)$ and add it to our list.

The algorithm has a polynomial running time since the number of iterations is bounded by $|V|$. To see it, consider the affine space $L_t := \{x \in \mathbb{R}^V \mid x(Q_i) = 1 \ \forall i = 1, \dots, t\}$. Then, $L_1 \supseteq L_2 \supseteq \dots \supseteq L_t \supseteq L_{t+1} \supseteq \dots$. The dimension of the spaces L_t decreases at each step since $\chi^S \in L_t \setminus L_{t+1}$, where S is the stable set constructed at the t -th iteration as above.

4.4 Sharpening the theta function

The number $\vartheta'(G)$. McEliece, Rodemich, Rumsey [173] and Schrijver [207] introduce the parameter $\vartheta'(G)$ as

$$\begin{aligned} \vartheta'(G) := \max \quad & e^T X e \\ \text{s.t.} \quad & \text{Tr}(X) = 1 \\ & X_{ij} = 0 \ (i \neq j, ij \in E) \\ & X \succeq 0, X \geq 0. \end{aligned} \tag{65}$$

Comparing with (58), it follows that

$$\alpha(G) \leq \vartheta'(G) \leq \vartheta(G).$$

As was done for $\vartheta(G)$ one can prove the following equivalent formulations for $\vartheta'(G)$:

$$\begin{aligned} \vartheta'(G) = \min \quad & \lambda_{\max}(Y) \\ \text{s.t.} \quad & Y_{ij} \geq 1 \ (i = j \text{ or } ij \in \overline{E}) \\ & Y \text{ symmetric matrix;} \end{aligned} \tag{66}$$

$$\begin{aligned} \vartheta'(G) = \min \quad & t & = \min \quad & t \\ \text{s.t.} \quad & Z_{ii} = t - 1 \quad (i \in V) & \text{s.t.} \quad & U_{ii} = 1 \quad (i \in V) \\ & Z_{ij} \leq -1 \quad (ij \in \overline{E}) & & U_{ij} \leq -\frac{1}{t-1} \quad (ij \in \overline{E}) \\ & Z \succeq 0 & & U \succeq 0, t \geq 2; \end{aligned} \tag{67}$$

and $\vartheta'(G) = \max(e^T x \mid \begin{pmatrix} 1 \\ x \end{pmatrix} = Y e_0$ for some nonnegative matrix $Y \in \mathcal{M}_G$). The inequality $\vartheta'(G) \leq \vartheta(G)$ is strict, for instance, for the graph with node set $\{0, 1\}^6$ where two nodes are adjacent if their Hamming distance (i.e., the number of positions where their coordinates are distinct) is at most 3 (then, $\vartheta(G) = \frac{16}{3}$ and $\vartheta'(G) = \alpha(G) = 4$).

The number $\vartheta^+(G)$. In a similar vein, Szegedy [224] introduced the following parameter $\vartheta^+(G)$ which provides a sharper lower bound for the chromatic number of \overline{G} :

$$\begin{aligned} \vartheta^+(G) := \max \quad & e^T X e \\ \text{s.t.} \quad & \text{Tr}(X) = 1 \\ & X_{ij} \leq 0 \ (i \neq j, ij \in E) \\ & X \succeq 0. \end{aligned} \tag{68}$$

We have $\vartheta(G) \leq \vartheta^+(G) \leq \chi(\overline{G})$. The first inequality is obvious and the second one can be proved in the same way as the inequality $\vartheta(G) \leq \chi(\overline{G})$ in Section 4.2. Therefore, the following chain of inequalities holds:

$$\alpha(G) \leq \vartheta'(G) \leq \vartheta(G) \leq \vartheta^+(G) \leq \chi(\overline{G}). \tag{69}$$

The parameters $\vartheta'(G)$, $\vartheta(G)$, and $\vartheta^+(G)$ are known, respectively, as the *vector chromatic number*, the *strict vector chromatic number*, and the *strong vector chromatic number* of \overline{G} ; see Section 6.4. As was done for $\vartheta(G)$, one can prove the following equivalent formulations for $\vartheta^+(G)$:

$$\begin{aligned} \vartheta^+(G) = \min \quad & \lambda_{\max}(Y) \\ \text{s.t.} \quad & Y_{ij} = 1 \quad (i = j \text{ or } ij \in \overline{E}) \\ & Y_{ij} \leq 1 \quad (ij \in E) \\ & Y \text{ symmetric matrix;} \end{aligned} \tag{70}$$

$$\begin{aligned} \vartheta^+(G) = \min \quad & t & = \min \quad & t \\ \text{s.t.} \quad & Z_{ii} = t - 1 \quad (i \in V) & \text{s.t.} \quad & U_{ii} = 1 \quad (i \in V) \\ & Z_{ij} = -1 \quad (ij \in \overline{E}) & & U_{ij} = -\frac{1}{t-1} \quad (ij \in \overline{E}) \\ & Z_{ij} \geq -1 \quad (ij \in E) & & U_{ij} \geq -\frac{1}{t-1} \quad (ij \in E) \\ & Z \succeq 0 & & U \succeq 0, \quad t \geq 2. \end{aligned} \tag{71}$$

The parameter $\vartheta^+(G)$ (in the formulation (71)) was introduced independently by Meurdesoif [174] who gives a graph G for which inequality $\vartheta(G) \leq \vartheta^+(G)$ is strict. See [224] for more about this parameter.

Bounding the Shannon capacity. The theta number $\vartheta(G)$ was introduced by Lovász [164] in connection with a problem of Shannon in coding theory. The strong product $G \cdot H$ of two graphs G and H has node set $V(G) \times V(H)$ with two distinct nodes (u, v) and (u', v') being adjacent if u, u' are equal or adjacent in G and v, v' are equal or adjacent in H . Then G^k is the strong product of k copies of G . The Shannon capacity of G is defined by

$$\Theta(G) := \sup_{k \geq 1} \sqrt[k]{\alpha(G^k)}.$$

As $\alpha(G^k) \geq (\alpha(G))^k$ and $\vartheta(G^k) \leq (\vartheta(G))^k$, one finds

$$\alpha(G) \leq \Theta(G) \leq \vartheta(G).$$

Using these inequalities, Lovász [164] could show that the Shannon capacity of C_5 is $\sqrt{5}$ (as $\alpha(C_5^2) = 5$ and $\theta(C_5) = \sqrt{5}$). For $n \geq 7$ odd, $\vartheta(C_n) = \frac{n \cos(\frac{\pi}{n})}{1 + \cos(\frac{\pi}{n})}$, but the value of $\Theta(C_n)$ is not known.

The theta number versus Delsarte's bound. Let G be a graph whose adjacency matrix can be written as $\sum_{i \in M} A_i$, where $M \subseteq \{1, \dots, N\}$ and A_0, A_1, \dots, A_N are 0/1 symmetric matrices forming an *association scheme*; that is, $A_0 = I$, $\sum_{i=0}^N A_i = J$, there exist scalars p_{ij}^k ($i, j, k = 1, \dots, N$) such that $A_i A_j = A_j A_i = \sum_{k=0}^N p_{ij}^k A_k$. As the matrices A_0, \dots, A_N commute, they have a common basis of eigenvectors and therefore positive semidefiniteness of a matrix $X := \sum_{i=0}^N x_i A_i$ can be expressed by a linear system of inequalities in x_1, \dots, x_N . Therefore, one finds that the theta numbers $\vartheta(G)$, $\vartheta'(G)$ can be computed by solving a linear programming problem. Based on this, Schrijver [207] shows that $\vartheta'(G)$ coincides with a linear programming bound introduced earlier by Delsarte [67].

These ideas have been extended to general semidefinite programs by Goemans and Rendl [92].

5 Semidefinite Relaxation for the Max-Cut Problem

We present here results dealing with the basic semidefinite relaxation of the cut polytope and its application to designing good approximation algorithms for the max-cut problem.

Given a graph $G = (V, E)$, the *cut* $\delta(S)$ induced by a vertex set $S \subseteq V$ is the set of edges with exactly one endpoint in S . Given edge weights $w \in \mathbb{Q}^E$, the *max-cut problem* consists of finding a cut $\delta(S)$ whose weight $w(\delta(S)) := \sum_{ij \in \delta(S)} w_{ij}$ is maximum. Let $\text{mc}(G, w)$ denote the maximum weight of a cut in G . A comprehensive survey about the max-cut problem can be found in [195]. The max-cut problem is one of the basic NP-hard problems studied by Karp [128]. Moreover, it cannot be approximated with an arbitrary precision; namely, Håstad [114] shows that for $\rho > \frac{16}{17} = 0.94117$ there is no ρ -approximation algorithm for max-cut if $P \neq NP$. [A ρ -approximation algorithm is an algorithm that returns in polynomial time a cut whose weight is at least ρ times the maximum weight of a cut; ρ being called the *performance ratio* or *guarantee*.] On the other hand, Goemans and Williamson [95] prove a 0.878-approximation algorithm for max-cut that will be presented in Section 5.3 below.

5.1 The basic linear relaxation

As before, the *cut polytope* $\text{CUT}(G)$ is the polytope in \mathbb{R}^E defined as the convex hull of the vectors $z^S \in \{\pm 1\}^E$ for $S \subseteq V$, where $z_{ij}^S = -1$ if and only if $|S \cap \{i, j\}| = 1$. The weight of the cut $\delta(S)$ can be expressed as $\frac{1}{2} \sum_{ij \in E} w_{ij}(1 - z_{ij}^S)$. Hence the max-cut problem is the problem of optimizing the linear objective function

$$\frac{1}{2} \sum_{ij \in E} w_{ij}(1 - z_{ij}) \quad (72)$$

over $\text{CUT}(G)$. The *circuit inequalities*:

$$\sum_{ij \in F} x_{ij} - \sum_{ij \in E(C) \setminus F} x_{ij} \geq 2 - |C|, \quad (73)$$

where C is a circuit in G and F is a subset of $E(C)$ with an odd cardinality, are valid for $\text{CUT}(G)$ as they express the fact that a cut and a circuit must have an even intersection. Together with the bounds $-1 \leq x_{ij} \leq 1$ ($ij \in E$) they define the *metric polytope* $\text{MET}(G)$. Thus $\text{CUT}(G) \subseteq \text{MET}(G)$; moreover, the only ± 1 vectors in $\text{MET}(G)$ are the cut vectors z^S ($S \subseteq V$). An inequality (73) defines a facet of $\text{CUT}(G)$ if and only if C is a chordless circuit in G while an inequality $\pm x_{ij} \leq 1$ is facet defining if and only if ij does not belong to a triangle [29]. Hence the metric polytope $\text{MET}(K_n)$ is defined by the $4\binom{n}{3}$ *triangle inequalities*:

$$x_{ij} + x_{ik} + x_{jk} \geq -1, \quad x_{ij} - x_{ik} - x_{jk} \geq -1 \quad (74)$$

for all triples $i, j, k \in \{1, \dots, n\}$. Therefore, one can optimize any linear objective function over $\text{MET}(K_n)$ in polynomial time. The same holds for $\text{MET}(G)$, since $\text{MET}(G)$ is equal to the projection of $\text{MET}(K_n)$ on the subspace \mathbb{R}^E indexed by the edge set of G [26]. The inclusion $\text{CUT}(G) \subseteq \text{MET}(G)$ holds at equality if and only if G has no K_5 -minor [29]. Therefore, the max-cut problem can be solved in polynomial time for the graphs with no K_5 -minor (including the planar graphs).

The polytope

$$Q(G) := \{x \in [-1, 1]^E \mid \sum_{ij \in E(C)} x_{ij} \geq 2 - |C| \text{ for all odd circuits } C \text{ in } G\}$$

contains the metric polytope $\text{MET}(G)$ and its ± 1 -vectors correspond to the bipartite subgraphs of G . Therefore, the max-cut problem for nonnegative weights can be reformulated as the problem of maximizing (72) over the ± 1 -vectors in $Q(G)$. A graph G is said to be *weakly bipartite* when all the vertices of $Q(G)$ are ± 1 -valued. It is shown in [101] that one can optimize in polynomial time a linear objective function over $Q(G)$. Therefore, the max-cut problem can be solved in polynomial time for weakly bipartite graphs with nonnegative edge weights. Guenin [103] characterized the weakly bipartite graphs as those graphs *containing no odd K_5 -minor* (they include the graphs with no K_5 -minor, the graphs having two nodes covering all odd circuits, etc.), settling a conjecture posed by Seymour [211]. (See Schrijver [209] for a shorter proof.) Poljak [192] shows that, for nonnegative edge weights, one obtains in fact the same optimum value when optimizing (72) over $\text{MET}(G)$ or over $Q(G)$.

Let $\text{met}(G, w)$ denote the optimum value of (72) maximized over $x \in \text{MET}(G)$. When all edge weights are equal to 1, we also use the notation $\text{met}(G)$ in place of $\text{met}(G, w)$ (and analogously $\text{mc}(G)$ in place of $\text{mc}(G, w)$). How well does the polyhedral bound $\text{met}(G, w)$ approximate the max-cut value $\text{mc}(G, w)$? In order to compare the two bounds, we assume that all edge weights are nonnegative. Then,

$$\text{met}(G, w) \leq w(E) = \sum_{ij \in E} w_{ij} \text{ and } \text{mc}(G, w) \geq \frac{1}{2}w(E).$$

(To see the latter inequality, consider an optimum cut $\delta(S)$ and the associated partition $(S, V \setminus S)$. Then, for every node $i \in V$, the sum of the weights of the edges connecting i to the opposite class of the partition is greater than or equal to the sum of the weights of the edges connecting i to nodes in the same class, since otherwise moving i to the other class would produce a heavier cut.) Therefore,

$$\frac{\text{mc}(G, w)}{\text{met}(G, w)} \geq \frac{1}{2}.$$

In fact, the ratio $\frac{\text{mc}(G, w)}{\text{met}(G, w)}$ tends to $\frac{1}{2}$ for certain classes of graphs (cf. [192], [194]) which shows that in the worst case the metric polytope does not provide a better approximation than the trivial relaxation of $\text{CUT}(G)$ by the cube $[-1, 1]^E$.

5.2 The basic semidefinite relaxation

The max-cut problem can be reformulated as the following integer quadratic program:

$$\begin{aligned} \text{mc}(G, w) = \max \quad & \frac{1}{2} \sum_{ij \in E} w_{ij}(1 - x_i x_j) \\ \text{s.t.} \quad & x_1, \dots, x_n \in \{\pm 1\}. \end{aligned} \tag{75}$$

For $x \in \{\pm 1\}^n$, the matrix $X := xx^T$ is positive semidefinite with all diagonal elements equal to one. Thus relaxing the rank one condition on X , we obtain the following *semidefinite relaxation* for max-cut:

$$\begin{aligned} \text{sdp}(G, w) := \max \quad & \frac{1}{2} \sum_{ij \in E} w_{ij}(1 - x_{ij}) \\ \text{s.t.} \quad & x_{ii} = 1 \quad \forall i \in \{1, \dots, n\} \\ & X = (x_{ij}) \succeq 0. \end{aligned} \tag{76}$$

The set

$$\mathcal{E}_n := \{X = (x_{ij})_{i,j=1}^n \mid X \succeq 0 \text{ and } x_{ii} = 1 \quad \forall i \in \{1, \dots, n\}\} \tag{77}$$

is the basic semidefinite relaxation of the cut polytope $\text{CUT}(K_n)$. More precisely,

$$x \in \text{CUT}(K_n) \implies \text{mat}(x) \in \mathcal{E}_n \quad (78)$$

where $\text{mat}(x)$ is the $n \times n$ symmetric matrix with ones on its main diagonal and x_{ij} as off-diagonal entries.

The quantity $\text{sdp}(G, w)$ can be computed in polynomial time (with an arbitrary precision). The objective function in (76) is equal to $\frac{1}{4}\langle L_w, X \rangle$, where $L_w = (l_{ij})$ is the *Laplacian* matrix defined by $l_{ii} := w(\delta(i))$ and $l_{ij} := -w_{ij}$ for $i \neq j$ (assigning weight 0 to non edges). Hence, the dual of the semidefinite program (76) is

$$\frac{1}{4} \min \left\{ \sum_{i=1}^n y_i \mid \text{diag}(y) - L_w \succeq 0 \right\} \quad (79)$$

and there is no duality gap (since I is a strictly feasible solution to (76)). Set $s = \frac{1}{n}y^T e$ and $u = se - y$; then $u^T e = 0$ and $\text{diag}(y) - L_w = sI - \text{diag}(u) - L_w \succeq 0$ if and only if $\lambda_{\max}(L_w + \text{diag}(u)) \leq s$. Therefore, (79) can be rewritten as the following eigenvalue optimization problem:

$$\frac{n}{4} \min \left\{ \lambda_{\max}(L_w + \text{diag}(u)) \mid \sum_{i=1}^n u_i = 0 \right\}; \quad (80)$$

this eigenvalue upper bound for max-cut had been introduced and studied earlier by Delorme and Poljak [64, 65]. One can also verify directly that (80) is an upper bound for max-cut. Indeed, for $x \in \{\pm 1\}^n$ and $u \in \mathbb{R}^n$ with $\sum_i u_i = 0$, one has:

$$w(\delta(S)) = \frac{1}{4}x^T L_w x = \frac{1}{4}x^T (L_w + \text{diag}(u))x = \frac{n}{4} \frac{x^T (L_w + \text{diag}(u))x}{x^T x}$$

which is less than or equal to $\frac{n}{4}\lambda_{\max}(L_w + \text{diag}(u))$ by the Rayleigh principle. The program (80) can be shown to have a unique minimizer u (when $w \neq 0$); this minimizer u is equal to the null vector, for instance, when G is vertex transitive, in which case the computation of the semidefinite bound amounts to an eigenvalue computation [64]. Based on this, one can compute the semidefinite bound for unweighted circuits. Namely, $\text{mc}(C_{2k}) = \text{sdp}(C_{2k}) = 2k$ and $\text{mc}(C_{2k+1}) = 2k$ while $\text{sdp}(C_{2k+1}) = \frac{2k+1}{4}(2 + 2\cos(\frac{\pi}{2k+1}))$. Hence, $\frac{\text{mc}(C_5)}{\text{sdp}(C_5)} = \frac{32}{25+5\sqrt{5}} \sim 0.88445$; the same ratio is obtained for some other circulant graphs [175].

Much research has been done for evaluating the *integrality ratio* $\frac{\text{mc}(G, w)}{\text{sdp}(G, w)}$ and comparing the polyhedral and semidefinite bounds. Poljak [192] proved the following inequality relating the two bounds:

$$\frac{\text{met}(G, w)}{\text{sdp}(G, w)} \geq \frac{32}{25 + 5\sqrt{5}} \quad \text{for any graph } G \text{ and } w \geq 0. \quad (81)$$

Therefore, the inequality

$$\frac{\text{mc}(G, w)}{\text{sdp}(G, w)} \geq \frac{32}{25 + 5\sqrt{5}} \quad (82)$$

holds for any weakly bipartite graph (G, w) with $w \geq 0$. The bound (82) remains valid for unweighted line graphs and the better bound $\frac{8}{9}$ was proved for the complete graph K_n with edge weights $w_{ij} := b_i b_j$ (given $b_1, \dots, b_n \in \mathbb{R}_+$) or for Paley graphs [64]. Moreover, the integrality ratio is asymptotically equal to 1 for the random graphs $G_{n,p}$ (p denoting the edge probability) [64].

Goemans and Williamson [95] proved the following bound for the integrality ratio:

$$\frac{\text{mc}(G, w)}{\text{sdp}(G, w)} \geq \alpha_0 \quad \text{for any graph } G \text{ and } w \geq 0, \quad (83)$$

where $0.87856 < \alpha_0 < 0.87857$ and α_0 is defined by

$$\alpha_0 := \min_{0 < \theta \leq \pi} \frac{2}{\pi} \frac{\theta}{1 - \cos \theta}. \quad (84)$$

Moreover, they present a randomized algorithm producing a cut whose expected weight is at least $\alpha_0 \cdot \text{sdp}(G, w)$; their result will be described in the next subsection.

Until recently, no example was known of a graph having a worst integrality ratio than C_5 and it had been conjectured by Delorme and Poljak [64] that $\frac{32}{25+5\sqrt{5}}$ is the worst possible value for the integrality ratio. Feige and Schechtman [82, 83] disproved this conjecture and proved that the worst case value for the integrality ratio $\frac{\text{mc}(G, w)}{\text{sdp}(G, w)}$ is *equal* to the Goemans-Williamson quantity α_0 ; we will come back to this result later in this section.

5.3 The Goemans-Williamson randomized approximation algorithm for max-cut

The randomized approximation algorithm of Goemans and Williamson [95] for max-cut goes as follows; its analysis will need the assumption that the edge weights are nonnegative.

1. *The semidefinite optimization phase:* Solve the semidefinite program (76). Let $X = (x_{ij})$ be an optimum solution and let $v_1, \dots, v_n \in \mathbb{R}^d$ (for some $d \leq n$) such that $x_{ij} = v_i^T v_j$ for all $i, j \in \{1, \dots, n\}$.
2. *The random hyperplane rounding phase:* Generate a random unit vector r and set $S := \{i \mid v_i^T r \geq 0\}$. Then, $\delta(S)$ is the randomized cut returned by the algorithm.

The hyperplane H_r with normal r cuts the space into two half-spaces and an edge ij belongs to the cut $\delta(S)$ if and only if the vectors v_i and v_j do not belong to the same half-space. Hence the probability that an edge ij belongs to $\delta(S)$ is equal to $\frac{\arccos(v_i^T v_j)}{\pi}$ and the expected weight $E(w(S))$ of the cut $\delta(S)$ is equal to

$$E(w(S)) = \sum_{ij \in E} w_{ij} \frac{\arccos(v_i^T v_j)}{\pi} = \sum_{ij \in E} w_{ij} \frac{1 - v_i^T v_j}{2} \cdot \frac{2}{\pi} \frac{\arccos(v_i^T v_j)}{1 - v_i^T v_j} \geq \alpha_0 \cdot \text{sdp}(G, w).$$

The last inequality holds if we assume that $w \geq 0$. As $E(w(S)) \leq \text{mc}(G, w)$, we find

$$\frac{\text{mc}(G, w)}{\text{sdp}(G, w)} \geq \frac{E(w(S))}{\text{sdp}(G, w)} \geq \alpha_0 > 0.87856. \quad (85)$$

As a byproduct of the analysis, we obtain the following trigonometric reformulation for max-cut with $w \geq 0$:

$$\begin{aligned} \text{mc}(G, w) = \max & \sum_{ij \in E} w_{ij} \frac{\arccos(v_i^T v_j)}{\pi} \\ \text{s.t.} & v_1, \dots, v_n \text{ unit vectors in } \mathbb{R}^n. \end{aligned} \quad (86)$$

Mahajan and Ramesh [170] have shown that the above randomized algorithm can be derandomized, therefore giving a deterministic α_0 -approximation algorithm for max-cut. Let us stress that until then the best known approximation algorithm was the simple random partition algorithm (which assigns a node to either side of the partition independently with probability $\frac{1}{2}$) with a performance ratio of $\frac{1}{2}$.

As mentioned above, the integrality ratio $\frac{mc(G,w)}{sdp(G,w)}$ is equal to α_0 in the worst case. More precisely, Feige and Schechtman [82, 83] show that for every $\epsilon > 0$ there exists a graph G (unweighted) for which the ratio is at most $\alpha_0 + \epsilon$. The basic idea of their construction is as follows. Let θ_0 denote the angle where the minimum in the definition of $\alpha_0 = \min_{0 < \theta \leq \pi} \frac{2}{\pi} \frac{\theta}{1 - \cos \theta}$ is attained; $\theta_0 \sim 2.331122$ is the nonzero root of $\cos \theta + \theta \sin \theta = 1$. Let $[\theta_1, \theta_2]$ be the largest interval containing θ_0 satisfying $\theta \in [\theta_1, \theta_2] \implies \frac{2}{\pi} \frac{\theta}{1 - \cos \theta} \leq \alpha_0 + \epsilon$. Distribute n points v_1, \dots, v_n uniformly on the unit sphere S^{d-1} in \mathbb{R}^d and let G be the graph on n nodes where there is an edge ij if and only if the angle between v_i and v_j belongs to $[\theta_1, \theta_2]$. Applying the random hyperplane rounding phase to the vectors v_1, \dots, v_n , the above analysis shows that the expected weight of the returned cut satisfies

$$\frac{E(w(S))}{sdp(G)} \leq \alpha_0 + \epsilon.$$

The crucial part of the proof consists then of showing that for some suitable choice of the dimension d and of the distribution of the n points on the sphere S^{d-1} the expected weight $E(w(S))$ is not far from the max-cut value $mc(G)$.

Nesterov [180] shows the weaker bound:

$$\frac{E(w(S))}{sdp(G, w)} \geq \frac{2}{\pi} \sim 0.63661 \tag{87}$$

for the larger class of weight functions w satisfying $L_w \geq 0$. (Note indeed that $L_w \geq 0$ if $w \geq 0$.) Hence, the GW rounding technique applies to a larger class of instances at the cost of obtaining a weaker performance ratio. Cf. Section 6.1 for more details.

The above analysis of the GW algorithm shows that its performance guarantee is at least α_0 . Karloff [126] shows that it is, in fact, equal to α_0 . For this, he constructs a class of graphs G (edge weights are equal to 1) for which the ratio $\frac{E(w(S))}{sdp(G,w)}$ can be made arbitrarily close to α_0 . (The graphs constructed by Feige and Schechtman [82] display the same behaviour; the construction of Karloff has however a simpler proof.) These graphs are the Johnson graphs $J(m, \frac{m}{2}, b)$ for m even, $b \leq \frac{m}{12}$ having the collection of subsets of $\{1, \dots, m\}$ of cardinality $\frac{m}{2}$ as node set and two nodes being adjacent if their intersection has cardinality b . An additional feature of these graphs is that $mc(G, w) = sdp(G, w)$. Hence, one of the problems that the Karloff's example emphasizes is that although the semidefinite program already solves the max-cut problem at optimality, the GW approximation algorithm is not able to recognize this fact and to take advantage of it for producing a better cut. As a matter of fact, recognizing whether $sdp(G, w) = mc(G, w)$ for given weights w is an NP-complete problem [65, 153].

Goemans and Williamson [95] show that their algorithm behaves, in fact, better for graphs having $\frac{sdp(G,w)}{w(E)} \geq \frac{85}{100}$ (and thus for graphs having very large cuts). To express their result, set $h(t) := \frac{1}{\pi} \arccos(1 - 2t)$, $t_0 := \frac{1 - \cos \theta_0}{2} \sim 0.84458$, where $\theta_0 \sim 2.331122$ is the angle at which the minimum in the definition of $\alpha_0 = \min_{0 < \theta \leq \pi} \frac{2}{\pi} \frac{\theta}{1 - \cos \theta}$ is attained. Then, $\frac{h(t_0)}{t_0} = \alpha_0$ and it follows from the definition of α_0 that $h(t) \geq \alpha_0 t$ for $t \in [0, 1]$. Further, set

$$\alpha_{GW}(t) := \frac{h(t)}{t} \text{ if } t \in [t_0, 1] \text{ and } \alpha_{GW}(t) := \alpha_0 \text{ if } t \in [0, t_0].$$

One can verify that the function $\tilde{h}(t) := \alpha_{\text{GW}}(t)t$ is convex on $[0, 1]$ and $\tilde{h} \leq h$. From this follows that

$$\frac{E(w(S))}{\text{sdp}(G, w)} \geq \alpha_{\text{GW}}(A), \text{ where } A := \frac{\text{sdp}(G, w)}{w(E)}. \quad (88)$$

Indeed, setting $y_{ij} := \frac{1-v_i^T v_j}{2}$, we have:

$$\frac{E(w(S))}{w(E)} = \sum_{ij \in E} \frac{w_{ij}}{w(E)} h(y_{ij}) \geq \sum_{ij \in E} \frac{w_{ij}}{w(E)} \tilde{h}(y_{ij}) \geq \tilde{h} \left(\sum_{ij \in E} \frac{w_{ij}}{w(E)} y_{ij} \right) = \tilde{h}(A) = \alpha_{\text{GW}}(A) \cdot A$$

which implies (88). Therefore, the performance guarantee of the GW algorithm is at least $\alpha_{\text{GW}}(A)$ which is greater than α_0 when $A > t_0$ and tends to 1 as A tends to 1. Extending Karloff's result, Alon and Sudakov [10] construct (unweighted) graphs G for which $\text{mc}(G, w) = \text{sdp}(G, w)$ and $\frac{E(w(S))}{\text{sdp}(G, w)} = \alpha_{\text{GW}}(A)$ for any $A = \frac{\text{sdp}(G, w)}{w(E)} \geq t_0$, which shows that the performance guarantee of the GW algorithm is equal to $\alpha_{\text{GW}}(A)$. For the remaining values of A , $\frac{1}{2} \leq A < t_0$, Alon, Sudakov and Zwick [11] construct graphs satisfying $\text{mc}(G, w) = \text{sdp}(G, w)$ and $\frac{E(w(S))}{\text{sdp}(G, w)} = \alpha_0$ which shows that the analysis of Goemans and Williamson is also tight in this case.

5.4 How to improve the Goemans-Williamson algorithm?

There are several ways in which one can try to modify the basic algorithm of Goemans and Williamson in order to obtain an approximation algorithm with a better performance ratio.

Adding valid inequalities. Perhaps the most natural idea is to strengthen the basic semidefinite relaxation by adding inequalities valid for the cut polytope. For instance, one can add all triangle inequalities; denote by $\text{sdp}'(G, w)$ the optimum value of the semidefinite program obtained by adding the triangle inequalities to (76). The new integrality ratio $\frac{\text{mc}(G, w)}{\text{sdp}'(G, w)}$ is equal to 1 for graphs with no K_5 -minor (thus for C_5). For K_5 (with edge weights 1) it is equal to $\frac{24}{25} = 0.96$. However this is not the worst case; Feige and Schechtman [82] construct graphs for which the new integrality ratio is no better than roughly 0.891.

On the other hand, the example of Karloff shows that the GW randomized approximation algorithm applied to the tighter semidefinite relaxation does not have a better performance guarantee. The same remains true if we would add to the semidefinite relaxation *all* inequalities valid for the cut polytope (because the Karloff's graphs satisfy $\frac{E(w(S))}{\text{sdp}(G, w)} \sim \alpha_0$ while $\text{mc}(G, w) = \text{sdp}(G, w)$!). Therefore, in order to improve the performance guarantee, beside adding some valid inequalities, a new rounding technique will be needed. We now present two ideas along these lines: the first from [78] uses triangle inequalities and adds a 'local search' phase to the GW algorithm, the second from [240] can be seen as a mixing of the hyperplane rounding technique and the basic random algorithm.

Adding valid inequalities and a local search phase. Feige, Karpinski and Langberg [78] have presented an approximation algorithm for max-cut with a better performance guarantee for graphs with a bounded maximum degree Δ (edge weights are assumed to be equal to one). Their algorithm has two new features: triangle inequalities are added to the basic semidefinite relaxation (also some triangle equalities in the case $\Delta = 3$) and an additional 'greedy' phase is added after the GW hyperplane rounding phase.

Given a partition $(S, V \setminus S)$, a vertex v belonging, say, to S , is called *misplaced* if it has more neighbours in S than in $V \setminus S$; then the cut $\delta(S \setminus \{v\})$ has more edges than the cut $\delta(S)$. One of the basic

ideas underlying the FKL algorithm is that, if $(S, V \setminus S)$ is the partition produced by the hyperplane rounding phase and if all angles $\arccos(v_i^T v_j)$ are equal to θ_0 (which implies $E(w(S)) = \alpha_0 \cdot \text{sdp}(G, w)$), then there is a positive probability (depending on Δ alone) of finding a misplaced vertex in the partition and, therefore, one can improve the cut.

In the case $\Delta = 3$ the FKL algorithm goes as follows. In the first step one solves the semidefinite program (76) to which have been added all triangle inequalities as well as the triangle equalities $x_{ij} + x_{ik} + x_{jk} = -1$ for all triples (i, j, k) for which $ij, ik \in E$ (such equality is indeed valid for a maximum cut for, if not, the vertex i would be misplaced). Then the hyperplane rounding phase is applied to the optimum matrix X , producing a partition $(S, V \setminus S)$. After that comes an additional *greedy phase*: If the partition $(S, V \setminus S)$ has a misplaced vertex v , move it to the other side of the partition and repeat until no misplaced vertex can be found. If at some step there are several misplaced vertices, we move the misplaced vertex v for which the ratio between the number of edges gained in the cut by moving v and the number of triples (i, j, k) with $ij, ik \in E$ and i misplaced destroyed by this action, is maximal.

It is shown in [78] that the expected weight of the final partition returned by the FKL algorithm satisfies

$$E(w(S)) \geq 0.919 \cdot \text{sdp}(G, w). \quad (89)$$

For regular graphs of degree 3, one can show an approximation ratio of 0.924 and, for graphs with maximum degree Δ , a ratio of $\alpha_0 + \frac{1}{233\Delta^4}$. Note that, when $\Delta \geq 4$, one cannot incorporate the triangle equality $x_{ij} + x_{ik} + x_{jk} = -1$ (with $ij, ik \in E$) as it is no longer valid for maximum cuts.

Recently, Halperin, Livnat and Zwick [109] gave an improved approximation algorithm for max-cut in graphs of maximum degree 3 with performance guarantee 0.9326. Their algorithm has an additional preprocessing phase (which converts the input graph into a cubic graph satisfying some additional property) and performs the greedy phase in a more global manner; moreover, it applies to a more general problem than max-cut.

Mixing the random hyperplane and the basic random rounding techniques. We saw above that the performance guarantee of the GW algorithm is greater than α_0 for graphs with large cuts (with weight at least 85% of the total weight of edges). Zwick [240] presents a modification of the GW algorithm which, on the other hand, has a better performance guarantee for graphs having *no* large cuts.

Note that the simple randomized algorithm, which constructs a partition $(S, V \setminus S)$ by assigning a vertex with probability $\frac{1}{2}$ to either side of the partition, produces a cut with expected weight $\frac{w(E)}{2}$ and thus its performance ratio is

$$\alpha_{\text{rand}}(A) := \frac{1}{2A} \text{ where } A = \frac{\text{sdp}(G, w)}{w(E)}.$$

Note, moreover, that this algorithm is equivalent to applying the hyperplane rounding technique to the standard unit vectors e_1, \dots, e_n , with the identity matrix as Gram matrix. As $\alpha_{\text{rand}}(A) \geq \alpha_{\text{GW}}(A)$ when $\frac{1}{2} \leq A \leq \frac{1}{2\alpha_0} \sim 0.569113$, Zwick's idea is to make a 'mix' of the hyperplane rounding and basic random algorithms. For this, if X is the optimum matrix obtained when solving the basic semidefinite program (76), set

$$X' := (\cos^2 \gamma_A)X + (\sin^2 \gamma_A)I$$

where $\gamma_A \in [0, \pi]$ is suitably chosen. Namely, if $A \geq t_0$ then $\gamma_A := 0$ and if $\frac{1}{2}A \leq t_0$, then solve the

following equations for c and t :

$$\frac{\arccos(c(1-2t)) - \arccos c}{t} = \frac{2c}{\sqrt{1-c^2(1-2t)^2}}, \quad \frac{1 - \frac{t}{A}}{\sqrt{1-c^2}} = \frac{1-2t}{\sqrt{1-c^2(1-2t)^2}}$$

(there is a unique solution c_A, t_A such that $0 \leq c_A \leq 1$ and $\frac{3}{4} \leq t_A \leq t_0$) and set $\gamma_A := \arccos(\sqrt{c_A})$. Note that γ_A tends to $\frac{\pi}{2}$ as A tends to $\frac{1}{2}$. Then a randomized cut $\delta(S)$ is produced by applying the hyperplane rounding phase to the modified matrix X' . Zwick shows that

$$\frac{E(w(S))}{\text{sdp}(G, w)} \geq \alpha_{\text{rot}}(A) \text{ for any graph } G \text{ and } w \geq 0 \quad (90)$$

where $\alpha_{\text{rot}}(A) := \alpha_{\text{GW}}(A)$ for $A \geq t_0$ and, setting $h_c(t) := \frac{\arccos(c(1-2t))}{\pi}$,

$$\alpha_{\text{rot}}(A) := \left(\frac{1}{A} - \frac{1}{t_A}\right)h_{c_A}(0) + \frac{1}{t_A}h_{c_A}(t_A)$$

for $\frac{1}{2} \leq A \leq t_0$. The new performance guarantee is at least $\alpha_{\text{rot}}(A)$, which is greater than $\alpha_{\text{rand}}(A)$ and $\alpha_{\text{GW}}(A)$ when $A < t_0$. For instance, $\alpha_{\text{rot}}(A) \geq 0.88$ if $A \leq 0.75$, $\alpha_{\text{rot}}(A) \geq 0.91$ if $A \leq 0.6$. Alon, Sudakov and Zwick [11] show that the analysis is tight; for this they construct graphs having $\text{mc}(G, w) = \text{sdp}(G, w)$ and $\frac{E(w(S))}{\text{sdp}(G, w)} = \alpha_{\text{rot}}(A)$ for any $\frac{1}{2} \leq A \leq t_0$.

Inapproximability results. Summarizing, the best performance guarantee of an approximation algorithm for max-cut (with nonnegative weights) known so far is $\alpha_0 \sim 0.87856$. In fact, $\frac{16}{17} \sim 0.94117$ is the best performance guarantee that one can hope for. Indeed, Håstad [114] shows that, for any $\epsilon > 0$, there is no $(\frac{16}{17} + \epsilon)$ -approximation algorithm for max-cut if $P \neq NP$. Berman and Karpinski [37] show that it is NP-hard to approximate max-cut in cubic graphs beyond the ratio of 0.997 (while there is an 0.932-approximation algorithm as we saw above).

On the positive side, Arora, Karger and Karpinski [20] show that the max-cut problem has a polynomial time approximation scheme (that is, an $(1-\epsilon)$ -approximation algorithm for any $\epsilon > 0$) when restricted to dense graphs, that is, graphs with $O(n^2)$ edges. De la Vega [230] described independently a randomized approximation scheme for max-cut in graphs with minimum degree cn for some constant $c > 0$.

We have seen in Section 3.6 several techniques permitting to construct semidefinite relaxations of the cut polytope refining the basic one. Thus a natural and very interesting question is whether some of them can be used for proving a better integrality ratio (better than the Goemans-Williamson bound α_0) and for designing an approximation algorithm for max-cut with an improved performance ratio. The most natural candidate to consider might be the Lasserre relaxation $Q_1(K_n)$ (defined using (47) and (48)) or its subset, the Anjos-Wolkowicz relaxation F_n (defined using (47)).

6 Applications of Semidefinite Programming and the Rounding Hyperplane Technique to Other Combinatorial Optimization Problems

The method developed by Goemans and Williamson for approximating the max-cut problem has been applied and generalized to a large number of combinatorial optimization problems. Summarizing, their method consists of the following two phases:

1. *The semidefinite optimization phase*, which finds a set of vectors v_1, \dots, v_n providing a Cholesky factorization of an optimum solution to the SDP program relaxing the original combinatorial problem.
2. *The random hyperplane rounding phase*, which constructs a solution to the original combinatorial problem by looking at the positions of the vectors v_i with respect to some random hyperplane.

The basic method of Goemans and Williamson may have to be modified in order to be applied to some other combinatorial problems. In the first phase, one has to choose an appropriate SDP relaxation of the problem at hand and, in the second phase, one may have to adapt the rounding procedure. For instance, if one wants to approximate graph colouring and max k -cut problems, one should consider more general partitions of the space using more than one random hyperplane. One may also have to add an additional phase permitting to modify the returned solution; for instance, to turn the returned cut into a bisection if one wants to approximate the bisection problem. It turns out that the analysis of the extended approximation algorithms is often more complicated than that of the basic GW algorithm; it sometimes needs the evaluation of certain integral formulas that are hard to evaluate numerically.

In this section we present approximation algorithms based on these ideas for the following problems: general quadratic programming problems, maximum bisection and k -cut problems, colouring, stable sets, MAX SAT and maximum directed cut problems.

Of course, the above is not an exhaustive list of the problems for which semidefinite programming combined with randomized rounding permits to obtain good approximations. There are other interesting problems, that we could not cover here, to which these techniques apply; this is the case, e.g., for scheduling (see [220]).

6.1 Approximating quadratic programming

We consider here the boolean quadratic programming problem:

$$\begin{aligned} m^*(A) := \max & \quad x^T A x \\ \text{s.t.} & \quad x \in \{\pm 1\}^n \end{aligned} \tag{91}$$

where A is a symmetric matrix of order n , and its natural SDP relaxation:

$$\begin{aligned} s^*(A) := \max & \quad \langle A, X \rangle \\ \text{s.t.} & \quad X_{ii} = 1 \quad (i = 1, \dots, n) \\ & \quad X \succeq 0. \end{aligned} \tag{92}$$

Obviously, $m^*(A) \leq s^*(A)$. How well does the semidefinite bound $s^*(A)$ approximate $m^*(A)$? Obviously $m^*(A) = s^*(A)$ when all off-diagonal entries of A are nonnegative. We saw in Section 5.3 that

$\frac{m^*(A)}{s^*(A)} \geq \alpha_0$ (the GW ratio from (84)) in the special case when A is the Laplacian matrix of a graph; that is, when $Ae = 0$ and $A_{ij} \leq 0$ for all $i \neq j$. (Note that these conditions imply that $A \succeq 0$.) Nesterov [180] studies the quality of the SDP relaxation for general A . When $A \succeq 0$ he shows the lower bound $\frac{2}{\pi}$ for the ratio $\frac{m^*(A)}{s^*(A)}$ and, based on this, he gives upper bounds for the relative accuracy $s^*(A) - m^*(A)$ for indefinite A . The basic step consists in giving a trigonometric reformulation of the problem (91), analogous to the trigonometric reformulation (86) for max-cut.

Proposition 15. *Given a symmetric matrix A ,*

$$\begin{aligned} m^*(A) = \max \quad & \frac{2}{\pi} \langle A, \arcsin(X) \rangle \\ \text{s.t.} \quad & X_{ii} = 1 \quad (i = 1, \dots, n) \\ & X \succeq 0 \end{aligned} \quad (93)$$

setting $\arcsin(X) := (\arcsin(x_{ij}))_{i,j=1}^n$. Moreover, $m^*(A) \geq \frac{2}{\pi} s^*(A)$ if $A \succeq 0$.

PROOF. Denote by μ the maximum of the program (93). Let x be an optimum solution to the program (91) and set $X := xx^T$. Then X is feasible for (93) with objective value $\frac{2}{\pi} \langle A, \arcsin(X) \rangle = \langle A, xx^T \rangle = m^*(A)$, which shows that $m^*(A) \leq \mu$. Conversely, let X be an optimum solution to (93) and let v_1, \dots, v_n be vectors such that $X_{ij} = v_i^T v_j$ for all i, j . Let r be a random unit vector. Then the expected value of $\text{sign}(r^T v_i) \text{sign}(r^T v_j)$ is equal to

$$1 - 2 \text{prob}(\text{sign}(r^T v_i) \neq \text{sign}(r^T v_j)) = 1 - 2 \frac{\arccos(v_i^T v_j)}{\pi} = \frac{2}{\pi} \arcsin(v_i^T v_j).$$

Therefore, the expected value E_A of $\sum_{i,j} a_{ij} \text{sign}(r^T v_i) \text{sign}(r^T v_j)$ is equal to $\frac{2}{\pi} \sum_{i,j} a_{ij} \arcsin(v_i^T v_j) = \frac{2}{\pi} \langle A, \arcsin(X) \rangle = \mu$. On the other hand, $\sum_{i,j} a_{ij} \text{sign}(r^T v_i) \text{sign}(r^T v_j) \leq m^*(A)$, since the vector $(\text{sign}(r^T v_i))_{i=1}^n$ is feasible for (91) for any unit vector r . This implies that $E_A \leq m^*(A)$ and thus $\mu \leq m^*(A)$. Assume $A \succeq 0$. Then, $\langle A, \arcsin(X) \rangle = \langle A, \arcsin(X) - X \rangle + \langle A, X \rangle \geq \langle A, X \rangle$, using the fact that $\arcsin(X) - X \succeq 0$ if $X \succeq 0$. Hence, $m^*(A) \geq \frac{2}{\pi} s^*(A)$ if $A \succeq 0$. \blacksquare

Let $m_*(A)$ (resp. $s_*(A)$) denote the optimum value of the program (91) (resp. (92)) where we replace maximization by minimization. Applying the duality theorem for semidefinite programming, we obtain:

$$s^*(A) = \min(e^T y \mid \text{diag}(y) - A \succeq 0), \quad (94)$$

$$s_*(A) = \max(e^T z \mid A - \text{diag}(z) \succeq 0). \quad (95)$$

For $0 \leq \alpha \leq 1$, set $s_\alpha := \alpha s^*(A) + (1 - \alpha) s_*(A)$.

Lemma 16. *For $\alpha := \frac{2}{\pi}$, $s_*(A) \leq m_*(A) \leq s_{1-\alpha} \leq s_\alpha \leq m^*(A) \leq s^*(A)$.*

PROOF. We show the inequality $m_*(A) \leq s_{1-\alpha}(A)$, that is, $s^*(A) - m_*(A) \geq \frac{2}{\pi}(s^*(A) - s_*(A))$. Let y (resp. z) be an optimum solution to (94) (resp. (95)). Then,

$$s^*(A) - m_*(A) = e^T y + m^*(-A) = m^*(\text{diag}(y) - A) \geq \frac{2}{\pi} s^*(\text{diag}(y) - A)$$

by Proposition 15, since $\text{diag}(y) - A \succeq 0$. To conclude, note that $s^*(\text{diag}(y) - A) = e^T y + s^*(-A) = e^T y - s_*(A) = s^*(A) - s_*(A)$. The inequality $s_\alpha(A) \leq m^*(A)$ can be shown similarly. \blacksquare

The above lemma can be used for proving the following bounds on the relative accuracy $m^*(A) - s_\alpha$.

Theorem 17. Set $\alpha := \frac{2}{\pi}$ and $\beta := \frac{\alpha^2 + 2\alpha - 1}{3\alpha - 1}$. Then,

$$\frac{m^*(A) - s_\alpha}{m^*(A) - m_*(A)} \leq \frac{\pi}{2} - 1 < \frac{4}{7} \quad \text{and} \quad \frac{|m^*(A) - s_\beta(A)|}{m^*(A) - m_*(A)} \leq \frac{\pi - 2}{6 - \pi} < \frac{2}{5}.$$

The above results can be extended to quadratic problems of the form:

$$\max x^T A x \text{ subject to } [x]^2 \in F$$

where F is a closed convex set in \mathbb{R}^n and $[x]^2 := (x_1^2, \dots, x_n^2)$. See [228], chapter 13 in [232], [235], [238] for further results. Inapproximability results are given in [35].

6.2 Approximating the maximum bisection problem

The *maximum weight bisection problem* is a variant of the max-cut problem where one wants to find a cut $\delta(S)$ such that $|S| = \frac{n}{2}$ (a *bisection* or *equicut*) (n being assumed even) having maximum weight. This is an NP-hard problem, for which no approximation algorithm with a performance ratio $> \frac{16}{17}$ exists unless $P = NP$ [114]. Polynomial time approximation schemes are known to exist for this problem over dense graphs [20] and over planar graphs [120].

Extending the Goemans-Williamson approach to max-cut, Frieze and Jerrum [87] gave a randomized 0.651-approximation algorithm for the maximum weight bisection problem. Ye [236] improved the performance ratio to 0.6993 by combining the Frieze-Jerrum approach with some rotation argument applied to the optimum solution of the semidefinite relaxation. Halperin and Zwick [111] further improved the approximation ratio to 0.7016 by strengthening the SDP relaxation with the triangle inequalities. Details are given below.

Given a graph $G = (V, E)$ ($V = \{1, \dots, n\}$) and edge weights $w \in \mathbb{R}_+^E$, the maximum weight bisection problem reads:

$$\begin{aligned} \max \quad & \frac{1}{2} \sum_{ij \in E} w_{ij} (1 - x_i x_j) \\ \text{s.t.} \quad & \sum_{i=1}^n x_i = 0 \\ & x_1, \dots, x_n \in \{\pm 1\}. \end{aligned} \tag{96}$$

A natural semidefinite relaxation is:

$$\begin{aligned} W^* := \max \quad & \frac{1}{2} \sum_{ij \in E} w_{ij} (1 - X_{ij}) \\ \text{s.t.} \quad & X_{ii} = 1 \quad (i \in V) \\ & \langle J, X \rangle = 0 \\ & X \succeq 0 \end{aligned} \tag{97}$$

The Frieze-Jerrum approximation algorithm:

1. *The SDP optimization phase:* Solve the SDP (97), let X be an optimum solution and let v_1, \dots, v_n be vectors such that $X_{ij} = v_i^T v_j$ for all i, j .
2. *The random hyperplane rounding phase:* Choose a random unit vector r and define the associated cut $\delta(S)$ where $S := \{i \in V \mid r^T v_i \geq 0\}$.

3. *Constructing a bisection:* Without loss of generality, assume that $|S| \geq \frac{n}{2}$. For $i \in S$, set $W(i) := \sum_{j \notin S} w_{ij}$. Order the elements of S as $i_1, \dots, i_{|S|}$ in such a way that $W(i_1) \geq \dots \geq W(i_{|S|})$ and define $\tilde{S} := \{i_1, \dots, i_{\frac{n}{2}}\}$.

Then $\delta(\tilde{S})$ is a bisection whose weight satisfies

$$w(\delta(\tilde{S})) \geq \frac{n}{2|S|} w(\delta(S)). \quad (98)$$

Consider the random variables $W := w(\delta(S))$ and $C := |S|(n - |S|)$; W is the weight of the cut $\delta(S)$ in G while C is the number of pairs $(i, j) \in V^2$ that are cut by the partition $(S, V \setminus S)$ (that is, the cardinality of the cut $\delta(S)$ viewed as cut in the complete graph K_n). The analysis of the GW algorithm from Section 5.3 shows the following lower bounds for the expected values $E(W)$ and $E(C)$:

$$E(W) \geq \alpha_0 W^*, \quad (99)$$

$$E(C) \geq \alpha_0 C^* \quad (100)$$

where $C^* := \frac{n^2}{4}$. Define the random variable

$$Z := \frac{W}{W^*} + \frac{C}{C^*}. \quad (101)$$

Then, $Z \leq 2$ and $E(Z) \geq 2\alpha_0$.

Lemma 18. *If $Z \geq 2\alpha_0$ then $w(\delta(\tilde{S})) \geq 2(\sqrt{2\alpha_0} - 1)W^*$.*

PROOF. Set $w(\delta(S)) = \lambda W^*$ and $|S| = \sigma n$. Then, $Z = \lambda + 4\sigma(1 - \sigma) \geq 2\alpha_0$, implying $\lambda \geq 2\alpha_0 - 4\sigma(1 - \sigma)$. Using (98), we obtain that

$$w(\delta(\tilde{S})) \geq \frac{n}{2|S|} w(\delta(S)) = \frac{\lambda W^*}{2\sigma} \geq W^* \frac{2\alpha_0 - 4\sigma(1 - \sigma)}{2\sigma} \geq 2(\sqrt{2\alpha_0} - 1)W^*.$$

(The last inequality being a simple verification.) ■

As $E(Z) \geq 2\alpha_0$, the strategy employed by Frieze and Jerrum in order to find a bisection satisfying the conclusion of Lemma 18 is to repeat the above steps 2 and 3 of the algorithm N times, where N depends on some small $\epsilon > 0$ ($N = \lceil \frac{1}{\epsilon} \ln \frac{1}{\epsilon} \rceil$) and to choose as output bisection the heaviest among the N bisections produced throughout the N runs. Then, with high probability, the largest among the variables Z produced throughout the N runs will be greater than or equal to $2\alpha_0$. Therefore, it follows from Lemma 18 that the weight of the output bisection is at least $(2(\sqrt{2\alpha_0} - 1) - \epsilon)W^*$. For ϵ small enough, this shows a performance ratio of 0.651.

Ye [236] shows an improved approximation ratio of 0.6993. For this, he modifies the Jerrum-Frieze algorithm in the following way. Instead of applying the random hyperplane rounding phase to the optimum solution X of (97), he applies it to the modified matrix $\rho X + (1 - \rho)I$, where ρ is a parameter to be determined. This operation is analogue to the ‘outward rotation’ used by Zwick [240] for the max-cut problem and mentioned in Section 5.4.

The starting point is to replace relations (99) and (100) by

$$E(W) \geq \alpha W^* \text{ and } E(C) \geq \gamma C^* \quad (102)$$

where $\alpha = \alpha(\rho)$ and $\gamma = \gamma(\rho)$ are lower bounds to be determined on the ratios $\frac{E(W)}{W^*}$ and $\frac{E(C)}{C^*}$, respectively. In fact, the following choices can be made for α, γ :

$$\alpha(\rho) := \min_{-1 \leq x < 1} \frac{2 \arccos(\rho x)}{\pi (1-x)}, \quad (103)$$

$$\gamma(\rho) := \min_{-1 \leq x < 1} \frac{2 \arccos(\rho x) - x \arccos \rho}{\pi (1-x)}. \quad (104)$$

Indeed, $E(W) = \frac{1}{2} \sum_{ij \in E} w_{ij} \frac{2}{\pi} \arccos(\rho X_{ij}) \geq \alpha(\rho) W^*$. By the definition of $\gamma(\rho)$, $\frac{2}{\pi} \arccos(\rho x) \geq (1-x)\gamma(\rho) + \frac{2}{\pi} x \arccos \rho$ for $x \in [-1, 1]$. Therefore,

$$E(C) = \frac{1}{4} \sum_{i \neq j \in \{1, \dots, n\}} \frac{2}{\pi} \arccos(\rho X_{ij}) \geq \frac{1}{4} \gamma(\rho) \sum_{i \neq j} (1 - X_{ij}) + \frac{1}{2\pi} \arccos \rho \sum_{i \neq j} X_{ij} = \frac{n^2}{4} \gamma(\rho) - \frac{\arccos \rho}{2\pi} n.$$

For n large enough, the linear term can be ignored and the result follows.

Modify the definition of Z from (101) as

$$Z := \frac{W}{W^*} + \tau \frac{C}{C^*}$$

where $\tau := \frac{\alpha}{2\beta} \left(\frac{1}{\sqrt{1-\beta}} - 1 \right)$. The proof of Lemma 18 can be adapted to show that, if $Z \geq \alpha + \tau\gamma$, then $E(w(\tilde{S})) \geq \frac{\alpha}{1+\sqrt{1-\beta}} W^*$. For $\rho = 0.89$, one can compute that $\alpha(\rho) \geq 0.8355$, $\gamma(\rho) \geq 0.9621$, and $\frac{\alpha}{1+\sqrt{1-\beta}} > 0.6993$. Therefore, this shows that Ye's algorithm is a 0.6993-approximation algorithm.

Halperin and Zwick [111] can improve the performance ratio to 0.7016. They achieve this by adding one more ingredient to Ye's algorithm; namely, they strengthen the SDP relaxation (97) by adding the triangle inequalities:

$$X_{ij} + X_{ik} + X_{jk} \geq -1, \quad X_{ij} - X_{ik} - X_{jk} \geq -1 \quad \text{for distinct } i, j, k \in \{1, \dots, n\}.$$

Although triangle inequalities had already been used earlier by some authors to obtain better approximations (e.g., in [78] for the max-cut problem in bounded degree graphs as mentioned in Section 5.4), they were always analyzed from a local point of view (e.g., in the above mentioned example, in a local search phase, searching for misplaced vertices). In contrast, Halperin and Zwick are able to make a global analysis of the contribution of triangle inequalities. Namely, they show that the function $\gamma(\rho)$ from (104) can be replaced by

$$\gamma'(\rho) := \min_{-1 \leq x \leq -\frac{1}{3}} \frac{1}{\pi} \left(\arccos(\rho x) + \frac{3(x+1)}{4} \arccos \left(-\frac{\rho}{3} \right) + \frac{1-3x}{4} \arccos \rho \right),$$

which enables them to demonstrate a better performance ratio (using appropriate values for the parameters ρ and τ). (Note that $\gamma'(\rho) > \gamma(\rho)$ for $0 < \rho < 1$.)

Let us give a flavor of how the function $\gamma'(\rho)$ comes up. The goal is to find a lower bound for the ratio $\frac{E(C)}{C^*} = \frac{4}{\pi n^2} \sum_{1 \leq i < j \leq n} \arccos(\rho X_{ij})$. Let A (resp. B, C) denote the set of pairs ij for which $X_{ij} < -\frac{1}{3}$ (resp. $-\frac{1}{3} \leq X_{ij} \leq 0, 0 \leq X_{ij} \leq 1$). By the triangle inequalities, the graph on $\{1, \dots, n\}$ with edge set A is triangle free, which implies that $|A| \leq \frac{n^2}{4}$. Thus the optimum value of the following nonlinear program is a lower bound for $\frac{E(C)}{C^*}$:

$$\begin{aligned} \min \quad & \frac{4}{\pi n^2} \sum_{i < j} \arccos(\rho z_{ij}) \\ \text{s.t.} \quad & \sum_{i < j} z_{ij} = -\frac{n}{2} \\ & -1 \leq z_{ij} \leq 1 \quad (i < j) \\ & |\{ij \mid z_{ij} < -\frac{1}{3}\}| \leq \frac{n^2}{4}. \end{aligned}$$

Halperin and Zwick show then that the above minimum can be expressed in closed form as $\gamma'(\rho)$.

Feige, Karpinski and Langberg [79] design a 0.795-approximation algorithm for the maximum bisection problem restricted to regular graphs. One of their key results is the following: Given a cut $\delta(S)$ in a regular graph G , one can efficiently construct a bisection $\delta(S')$ whose weight is at least $0.9027 w(\delta(S))$. Hence, if we start with the cut $\delta(S)$ given as output of the Goemans-Williamson algorithm, then this gives an approximation algorithm with performance ratio $0.9027 \cdot 0.878 \sim 0.793$; a further improvement is demonstrated in [79].

Extensions to variations of the bisection problem. The following variations of the bisection problem have been studied in the literature: (i) the *maximum $\frac{n}{2}$ -vertex cover problem*, (ii) the *maximum $\frac{n}{2}$ -dense subgraph problem*, (iii) the *maximum $\frac{n}{2}$ -uncut problem*, which ask for a subset $S \subseteq V$ of size $\frac{n}{2}$ maximizing the total weight of the edges incident to S , contained in S , contained in S or its complement, respectively. Halperin and Zwick [111] treat these three problems (together with the maximum bisection problem as well as some directed analogues) in a unified framework and they can show the best approximation ratios known up to today, namely, 0.8452 for problem (i), 0.6221 for problem (ii), and 0.6436 for problem (iii).

6.3 Approximating the max k -cut problem

Given a graph $G = (V, E)$, edge weights $w \in \mathbb{R}_+^E$ and an integer $k \geq 2$, the *max k -cut problem* asks for a partition $\mathcal{P} = (S_1, \dots, S_k)$ of V whose weight $w(\mathcal{P}) := \sum_{1 \leq h < h' \leq k} \sum_{ij \in E \mid i \in S_h, j \in S_{h'}} w_{ij}$ is maximum. The set of edges whose end nodes belong to distinct classes of the partition is a *k -cut*, denoted as $\delta(S_1, \dots, S_k)$. For $k = 2$, we find the max-cut problem. For any $k \geq 2$, the max k -cut problem is NP-hard; moreover, there can be no polynomial time approximation algorithm for it with performance ratio $1 - \frac{1}{239k}$, unless $P=NP$ [124].

A simple heuristic for max k -cut is to partition V randomly into k sets. As the probability that two nodes fall in the same class is $\frac{1}{k}$, the expected weight of the k -cut produced in this way is $\sum_{ij \in E} w_{ij} (1 - \frac{1}{k}) \geq w(E)(1 - \frac{1}{k})$ and, therefore, the simple random partition heuristic has a performance guarantee of $1 - \frac{1}{k}$.

Frieze and Jerrum [87] present an approximation algorithm for max k -cut with performance guarantee α_k satisfying

(i) $\alpha_k > 1 - \frac{1}{k}$ and $\lim_{k \rightarrow \infty} \frac{\alpha_k - (1 - \frac{1}{k})}{2k^{-2} \ln k} = 1$,

(ii) $\alpha_2 = \alpha_0 \geq 0.878567$ (recall (84)), $\alpha_3 \geq 0.832718$, $\alpha_4 \geq 0.850304$, $\alpha_5 \geq 0.874243$, $\alpha_{10} \geq 0.926642$, $\alpha_{100} \geq 0.990625$.

In particular, the Frieze-Jerrum algorithm has a better performance guarantee than the simple random heuristic.

One can model the max k -cut problem on a graph $G = (V, E)$ ($V = \{1, \dots, n\}$) by having n variables x_1, \dots, x_n taking one of k possible values. For $k = 2$ the 2 possible values are ± 1 and for $k \geq 2$ one can choose as possible values a set of k unit vectors $a_1, \dots, a_k \in \mathbb{R}^{k-1}$ satisfying

$$a_i^T a_j = -\frac{1}{k-1} \text{ for } 1 \leq i \neq j \leq k.$$

(Such vectors exist since the matrix $\frac{k}{k-1}I_k - \frac{1}{k-1}J_k$ is positive semidefinite.) Hence the max k -cut problem can be formulated as

$$\begin{aligned} \text{mc}_k(G, w) := \max & \quad \frac{k-1}{k} \sum_{ij \in E} w_{ij} (1 - x_i^T x_j) \\ \text{s.t.} & \quad x_1, \dots, x_n \in \{a_1, \dots, a_k\} \end{aligned} \quad (105)$$

and the following is a semidefinite relaxation of (105):

$$\begin{aligned} \text{sdp}_k(G, w) := \max & \quad \frac{k-1}{k} \sum_{ij \in E} w_{ij} (1 - X_{ij}) \\ \text{s.t.} & \quad X_{ii} = 1 & (i \in V) \\ & \quad X_{ij} \geq -\frac{1}{k-1} & (i \neq j \in V) \\ & \quad X \succeq 0. \end{aligned} \quad (106)$$

The Frieze-Jerrum approximation algorithm for max k -cut:

1. Solve (106) to obtain unit vectors v_1, \dots, v_n satisfying $v_i^T v_j \geq -\frac{1}{k-1}$ ($i, j \in V$) and $\text{sdp}_k(G, w) = \frac{k-1}{k} \sum_{ij \in E} w_{ij} (1 - v_i^T v_j)$.
2. Choose k independent random vectors $r_1, \dots, r_k \in \mathbb{R}^n$. (This can be done by choosing their kn components as independent random variables from the standard normal distribution with mean 0 and variance 1.)
3. Partition V into S_1, \dots, S_k where S_h consists of the nodes $i \in V$ for which $v_i^T r_h = \max_{h'=1, \dots, k} v_i^T r_{h'}$. (Break ties arbitrarily as they occur with probability 0.)

When $k = 2$ the algorithm reduces to the Goemans-Williamson algorithm for max-cut. Given two unit vectors $u, v \in \mathbb{R}^n$, the probability that $\max_{1 \leq h \leq k} u^T r_h$ and $\max_{1 \leq h \leq k} v^T r_h$ are both attained by the same vector within r_1, \dots, r_k depends only on the angle between u and v , i.e., on $\rho := u^T v$, and it is equal to $k \cdot \text{prob}(u^T r_1 = \max_{1 \leq h \leq k} u^T r_h \text{ and } v^T r_1 = \max_{1 \leq h \leq k} v^T r_h)$; denote this probability as $kI(\rho)$. Then the expected weight of the k -cut $\delta(S_1, \dots, S_k)$ produced by the Frieze-Jerrum algorithm is equal to

$$\begin{aligned} & \sum_{ij \in E} w_{ij} \text{prob}(ij \in \delta(S_1, \dots, S_k)) = \sum_{ij \in E} w_{ij} (1 - kI(v_i^T v_j)) \\ & = \sum_{ij \in E} w_{ij} \left(\frac{k}{k-1} \frac{1 - kI(v_i^T v_j)}{1 - v_i^T v_j} \right) \left(\frac{k-1}{k} (1 - v_i^T v_j) \right) \geq \alpha_k \text{sdp}_k(G, w), \end{aligned}$$

setting

$$\alpha_k := \min_{-\frac{1}{k-1} \leq \rho < 1} \frac{k}{k-1} \frac{1 - kI(\rho)}{1 - \rho}. \quad (107)$$

For $k = 2$, $\alpha_2 = \alpha_0$ can be computed exactly. For $k \geq 3$, the evaluation of α_k is more complicated and relies on the computation of the function $I(\rho)$ which can be expressed as a multiple integral. Using a

Taylor series expansion for $I(\rho)$, Frieze and Jerrum could show the lower bounds for α_k mentioned at the beginning of this subsection.

For $k = 3$, de Klerk and Pasechnik [135] give a closed form expression for $I(\rho)$ which enables them to show that

$$\alpha_3 = \frac{7}{12} + \frac{3}{4\pi^2} \arccos^2(-1/4).$$

Thus $\alpha_3 > 0.836008$ (instead of the lower bound 0.832718 of Frieze and Jerrum). Goemans and Williamson [96] find the same expression for α_3 using another formulation for max 3-cut based on complex semidefinite programming.

De Klerk, Pasechnik and Warners [134] prove a better lower bound for α_k for small $k \geq 3$. For instance, they show that $\alpha_4 \geq 0.857487$ (instead of 0.850304). For this they present another approximation algorithm for max k -cut (equivalent to the Frieze-Jerrum algorithm for the graphs G with $\vartheta(\overline{G}) \leq k$) which enables them to reformulate the function $I(\rho)$ in terms of the volume of a spherical simplex and do more precise computations.

The minimum k -cut problem is also studied in the literature, in particular, because of its application to frequency assignment (see [71, 72]). Whereas good approximation algorithms exist for the maximum k -cut problem, the minimum k -cut problem cannot be approximated within a ratio of $O(|E|)$ unless P=NP. Semidefinite relaxations are nevertheless used in practice for deriving good lower bounds for the problem (see [71, 72]).

6.4 Approximating graph colouring

Determining the chromatic number of a graph is a hard problem. Lund and Yannakakis [168] show that there is a constant $\epsilon > 0$ for which there exists no polynomial algorithm which can colour any graph G using at most $n^\epsilon \chi(G)$ colours unless P=NP. Khanna, Linial and Safra [131] show that it is not possible to colour a 3-colourable graph with 4 colours in polynomial time unless P=NP.

On the positive side, Wigderson [231] shows that it is possible to colour in polynomial time a 3-colourable graph with $3\lceil\sqrt{n}\rceil$ colours and, more generally, a k -colourable graph with $2kn^{1-\frac{1}{k-1}}$ colours; we will come back to this result later in this section. Later Blum [39] gives a polynomial time algorithm colouring a 3-colourable graph with $O(n^{\frac{3}{8}} \log^{\frac{8}{5}} n)$. Using semidefinite programming and randomized rounding, Karger, Motwani and Sudan [125] present a randomized polynomial time algorithm which colours a 3-colourable graph with maximum degree Δ with $O(\Delta^{\frac{1}{3}} \sqrt{\log \Delta \log n})$ or $O(n^{\frac{1}{4}} \sqrt{\log n})$ colours and, more generally, a k -colourable graph with $O(\Delta^{1-\frac{2}{k}} \sqrt{\log \Delta \log n})$ or $O(n^{1-\frac{3}{k+1}} \sqrt{\log n})$ colours. This result was later refined by Halperin, Nathaniel and Zwick [110], who proved that a k -colourable graph with maximum degree Δ can be coloured in randomized polynomial time with $O(\Delta^{1-\frac{2}{k}} (\log \Delta)^{\frac{1}{k}} \log n)$. Further colouring results can be found in [40], [105], [110].

In what follows we present some of these results. We first prove a weaker version of the Karger-Motwani-Sudan result, namely, how to find a $O(n^{0.387})$ colouring for a 3-colourable graph. This enables us to introduce the basic tools used in [125]: vector k -colouring, k -semicolouring, hyperplane rounding, and a result of Wigderson [231]. Then we describe the Halperin-Nathaniel-Zwick algorithm for finding a $O(\Delta^{\frac{1}{3}} (\log \Delta)^{\frac{1}{3}} \log n)$ -colouring of a 3-colourable graph with maximum degree Δ . (For simplicity in the exposition we only treat the case $k = 3$.) This result is based on a new randomized rounding technique introduced in [125], using the standard n -dimensional normal distribution (instead of the distribution on the unit sphere) and vector projections. We finally describe the $O(n^{1-\frac{3}{k+1}} \sqrt{\log n})$ -colouring algorithm for k -colourable graphs of Karger, Motwani and Sudan.

Vector colouring. The first step in the Karger-Motwani-Sudan algorithm consists in solving a semidefinite relaxation for the colouring problem. We saw in Sections 4.2 and 4.4 that the theta number $\vartheta(\overline{G})$ and its variations $\vartheta'(\overline{G})$ and $\vartheta^+(\overline{G})$ constitute lower bounds for the chromatic number of G . Karger, Motwani and Sudan consider the SDP program (67) defining $\vartheta'(\overline{G})$ as a SDP relaxation for the colouring problem and they introduce the notion of vector colouring. A *vector k -colouring* of G is an assignment of vectors v_1, \dots, v_n to the nodes of G such that $v_i^T v_j \leq -\frac{1}{k-1}$ for every edge $ij \in E$. Then the *vector chromatic number* $\chi_v(G)$ is defined as the smallest $k \geq 2$ for which there exists a vector k -colouring. By the discussion above, $\chi_v(G) = \vartheta'(\overline{G})$. If in the definition of vector colouring one requires that the inequalities $v_i^T v_j \leq -\frac{1}{k-1}$ hold at equality for all edges, then we obtain the *strict vector chromatic number* which coincides with $\vartheta(\overline{G})$. More strongly, one can consider the *strong vector chromatic number* $\vartheta^+(\overline{G})$ which is defined by requiring $v_i^T v_j = -\frac{1}{k-1}$ for all edges and $v_i^T v_j \geq -\frac{1}{k-1}$ for all non edges. Therefore, the vector chromatic number is less than or equal to the strict vector chromatic number, which in turn is less than or equal to the strong vector chromatic number, which is a lower bound for the chromatic number (recall (69)).

Let us point out that the gap between the chromatic number and all these vector chromatic numbers can be arbitrarily large. Karger, Motwani and Sudan [125] construct a class of graphs having $\chi_v(G) = 3$ while $\chi(G) \geq n^{0.0113}$. Feige [75] shows that for all $\epsilon > 0$ there exist families of graphs with $\chi(G) \geq \vartheta(\overline{G})n^{1-\epsilon}$ and Charikar [51] proves an analogous result for the strong vector chromatic number.

Semicolouring. The hard part in the Karger-Motwani-Sudan algorithm consists of constructing a good proper colouring from a vector k -colouring. There are two steps: first construct a semicolouring and then from it a proper colouring. A *k -semicolouring* of a graph on n nodes is an assignment of k colours to at least half of the nodes in such a way that no two adjacent nodes receive the same colour. This is a useful notion, as an algorithm for semicolouring yields an algorithm for proper colouring.

Lemma 19. *Let $f : \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$ be a monotone increasing function. If there is a randomized polynomial time algorithm which $f(i)$ -semicolours every i -vertex subgraph of graph G , then this algorithm can colour G with $O(f(n) \log n)$ colours. Moreover, if there exists some $\epsilon > 0$ such that $f(i) = O(i^\epsilon)$ for all i , then the algorithm can colour G with $f(n)$ colours.*

PROOF. We show how to colour any p -vertex subgraph H of G . By assumption one can semicolour H with $f(p)$ colours. Let S denote the set of nodes of H that have not been coloured; then $|S| \leq \frac{p}{2}$. One can recursively colour the subgraph of H induced by S using a new set of colours.

Let $c(p)$ denote the maximum number of colours that the above algorithm needs for colouring an arbitrary p -vertex subgraph of G . Then,

$$c(p) \leq c\left(\frac{p}{2}\right) + f(p).$$

This recurrence relation implies that $c(p) = O(f(p) \log p)$. Moreover, if $f(p) = p^\epsilon$, one can easily verify that $c(p) = O(f(p))$. ▀

In view of Lemma 19, we are now left with the task of transforming a vector k -colouring into a good semicolouring.

Colouring a 3-colourable graph with $O(n^{0.387})$ -colours.

Theorem 20. *Every vector 3-colourable graph G with maximum degree Δ has a $O(\Delta^{\log_3 2})$ -semicolouring which can be constructed in polynomial time with high probability.*

PROOF. Let $v_1, \dots, v_n \in \mathbb{R}^n$ be unit vectors forming a vector 3-colouring of G , i.e., $v_i^T v_j \leq -\frac{1}{2}$ for all edges $ij \in E$; this means that the angle between v_i and v_j is at least $\frac{2\pi}{3}$ for all edges $ij \in E$. Choose independently N random hyperplanes. This induces a partition of the space \mathbb{R}^n into 2^N regions and one colours the nodes of G with 2^N colours depending in which region their associated vectors v_i are located. Then the probability that an edge is monochromatic is at most 3^{-N} and thus the expected number of monochromatic edges is at most $|E|3^{-N} \leq \frac{1}{2}n\Delta 3^{-N}$. By Markov's inequality, the probability that the number of monochromatic edges is more than twice the expected number is at most $\frac{1}{2}$. After repeating the process t times, we find with probability $\geq 1 - \frac{1}{2^t}$ a colouring of G for which the number of monochromatic edges is at most $n\Delta 3^{-N}$. Setting $N := 2 + \lceil \log_3 \Delta \rceil$, we have $n\Delta 3^{-N} \leq \frac{n}{4}$. As the number of nodes that are incident to a monochromatic edge is $\leq \frac{n}{2}$, we have found a semicolouring using $2^N \leq 8\Delta^{\log_3 2}$ colours. \blacksquare

As $\log_3 2 < 0.631$, Theorem 20 and Lemma 19 imply a colouring with $O(n^{0.631})$ colours. This is yet weaker than Wigderson's $O(\sqrt{n})$ -colouring algorithm. In fact, the result can be improved using the following idea of Wigderson.

Theorem 21. *There is a polynomial time algorithm which, given a 3-colourable graph G and a constant $\delta \leq n$, finds an induced subgraph H of G with maximum degree $\Delta_H < \delta$ and a $\frac{2n}{\delta}$ -colouring of $G \setminus H$.*

PROOF. If G has a node v of degree $\geq \delta$, colour the subgraph induced by $N(v)$ with two colours and delete $\{v\} \cup N(v)$ from G . We repeat this process using two new colours at each deleted neighborhood and stop when we arrive at a graph H whose maximum degree is less than δ . \blacksquare

Applying Theorem 21 with $\delta = \sqrt{n}$ and the fact that a graph with maximum degree Δ has a $(\Delta + 1)$ -colouring, one finds Wigderson's polynomial algorithm for colouring a 3-colourable graph with $3\lceil \sqrt{n} \rceil$ colours. More strongly, one can prove:

Theorem 22. *A 3-colourable graph can be coloured with $O(n^{0.387})$ colours by a polynomial time randomized algorithm.*

PROOF. Let G be a 3-colourable graph. Applying Theorem 21 with $\delta := n^{0.613}$, we find an induced subgraph H of maximum degree $\Delta_H < \delta$ and a colouring of $G \setminus H$ using $\frac{2n}{\delta} = O(n^{0.387})$ colours. By Theorem 20 and Lemma 19, H can be coloured with $O(\delta^{\log_3 2}) = O(n^{0.387})$ colours. This shows the result. \blacksquare

Improved colouring algorithm using 'rounding via vector projections'. In order to achieve the better $O(\Delta^{\frac{1}{3}}(\log \Delta)^{\frac{1}{3}} \log n)$ -colouring algorithm for a 3-colourable graph, one has to improve Theorem 20 and to show how to construct in randomized polynomial time a $O(\Delta^{\frac{1}{3}}(\log \Delta)^{\frac{1}{3}})$ -semicolouring.

(Indeed, the desired colouring follows then as a direct application of Lemma 19.) For this, Karger, Motwani and Sudan introduced another randomized technique for constructing a semicolouring from a vector colouring whose analysis has been refined by Halperin, Nathaniel and Zwick [110] and is presented below. The main step consists of proving the following result.

Theorem 23. *Let G be a vector 3-colourable graph on n nodes with maximum degree Δ . Then an independent set of size $\Omega\left(\frac{n}{\Delta^{\frac{1}{3}}(\log \Delta)^{\frac{1}{3}}}\right)$ can be found in randomized polynomial time.*

Indeed if Theorem 23 holds, then one can easily construct a $O(\Delta^{\frac{1}{3}}(\log \Delta)^{\frac{1}{3}})$ -semicolouring. For this, assign one colour to the nodes of the independent set found in Theorem 23 and recurse on the remaining nodes. One can verify that after $O(\Delta^{\frac{1}{3}}(\log \Delta)^{\frac{1}{3}})$ recursive steps, one has properly coloured at least half of the nodes; that is, one has constructed a $O(\Delta^{\frac{1}{3}}(\log \Delta)^{\frac{1}{3}})$ -semicolouring.

We now turn to the proof of Theorem 23. Let v_1, \dots, v_n be unit vectors forming a vector 3-colouring of G (i.e., $v_i^T v_j \leq -\frac{1}{2}$ for all edges ij) and set $c := \sqrt{\frac{2}{3} \ln \Delta - \frac{1}{3} \ln \ln \Delta}$. Choose a random vector r according to the standard n -dimensional normal distribution; this means that the components r_1, \dots, r_n of r are independent random variables, each being distributed according to the standard normal distribution.

Set $I := \{i \in \{1, \dots, n\} \mid r^T v_i \geq c\}$, $n' := |I|$, and let m (resp., m') denote the number of edges of G (resp. the number of edges of G contained in I). Then an independent set $J \subseteq I$ can be obtained by removing one vertex from each edge contained in I ; thus $|J| \geq n' - m'$. Intuitively there cannot be too many edges within I . Indeed the vectors assigned to the endpoints of an edge are rather far apart since their angle is at least $\frac{2\pi}{3}$, while the vectors assigned to the vertices in I should all be close to r since they have a large inner product with r . The proof consists of showing that the expected value of $n' - m'$ is $\Omega\left(\frac{n}{\Delta^{\frac{1}{3}}(\log \Delta)^{\frac{1}{3}}}\right)$.

The expected size of I is

$$E(n') = \sum_{i=1}^n \text{prob}(v_i^T r \geq c) = n \cdot \text{prob}(v_1^T r \geq c)$$

and the expected number of edges contained in I is

$$E(m') = \sum_{ij \in E} \text{prob}(v_i^T r \geq c \text{ and } v_j^T r \geq c) = m \cdot \text{prob}(v_1^T r \geq c \text{ and } v_2^T r \geq c)$$

where v_1 and v_2 denote two unit vectors satisfying $v_1^T v_2 \leq -\frac{1}{2}$. The following properties of the standard n -dimensional normal distribution will be used (see [125]).

Lemma 24. *Let u_1 and u_2 be unit vectors and let r be a random vector chosen according to the standard n -dimensional normal distribution. Let $N(x) = \int_x^\infty \phi(y) dy$ denote the tail of the standard normal distribution, where $\phi(x) = \frac{1}{\sqrt{2\pi}} \exp(-\frac{x^2}{2})$ is its density function.*

(i) *The inner product $r^T u_1$ is distributed according to the standard normal distribution. Therefore, $\text{prob}(u_1^T r \geq c) = N(c)$.*

(ii) *If u_1 and u_2 are orthogonal, then $u_1^T r$ and $u_2^T r$ are independent random variables.*

(iii) $(\frac{1}{x} - \frac{1}{x^3})\phi(x) \leq N(x) \leq \frac{1}{x}\phi(x)$ for $x > 0$.

It follows from Lemma 24 (i) that $E(n') = n \cdot N(c)$. We now evaluate $E(m')$. As before, v_1 and v_2 are two unit vectors such that $v_1^T v_2 \leq -\frac{1}{2}$. Since the probability $P_{12} := \text{prob}(v_1^T r \geq c \text{ and } v_2^T r \geq c)$ is a monotone increasing function of $v_1^T v_2$, it attains its maximum value when $v_1^T v_2 = -\frac{1}{2}$. We can therefore assume that $v_1^T v_2 = -\frac{1}{2}$. Karger et al. [125] show the upper bound $N(2c)$ for the probability P_{12} and, using a refinement of their method, Halperin et al. [110] prove the sharper bound $N(\sqrt{2}c)^2$.




Figure 25

Lemma 26. *If v_1 and v_2 are unit vectors such that $v_1^T v_2 = -\frac{1}{2}$, then $\text{prob}(v_1^T r \geq c \text{ and } v_2^T r \geq c) \leq N(\sqrt{2}c)^2$.*

PROOF. Let r' denote the orthogonal projection of r on the plane spanned by v_1 and v_2 . Then r' follows the standard 2-dimensional normal distribution and $v_i^T r' = v_i^T r$ for $i = 1, 2$. Hence we can work in the plane; Figure 25 will help visualize the argument. Write r' as $r' = \alpha \cdot cv_1 + \beta \cdot c(v_1 + 2v_2)$ for some scalars α, β . As v_1 is orthogonal to $v_1 + 2v_2$, we find that $v_1^T r' \geq c$ if and only if $\alpha \geq 1$; that is, if r' belongs to the half-plane lying above the line (D_1AB_1) (see Figure 25). Hence the probability P_{12} is equal to the probability that r' falls within the wedge defined by the angle $\angle B_1AB_2$ (this is the shaded area in Figure 25). Karger et al. [125] bound this probability by the probability that r' lies on the right side of the vertical line through A , which is equal to $\text{prob}((v_1 + v_2)^T r' \geq 2c)$ and thus to $N(2c)$ (since $v_1 + v_2$ is a unit vector). The better bound $N(\sqrt{2}c)^2$ can be shown as follows. Let u_1, u_2 be orthogonal unit vectors in the plane forming each the angle $\frac{\pi}{4}$ with $v_1 + v_2$. Denote by E_i the intersection point of the line through the origin parallel to u_i with the line through A perpendicular

to u_i . One can easily verify that E_i is at distance $\sqrt{2}c$ from the origin. Now one can bound the probability P_{12} by the probability that r' falls within the wedge defined by the angle $\angle C_1AC_2$. The latter probability is just $\text{prob}(u_1^T r' \geq \sqrt{2}c \text{ and } u_2^T r' \geq \sqrt{2}c)$ which (by Lemma 24 (i) (ii)) is equal to $N(\sqrt{2}c)^2$. \blacksquare

We can now conclude the proof of Theorem 23. Lemma 26 implies that $E(m') \leq m \cdot N(\sqrt{2}c)^2$. As $m \leq \frac{n\Delta}{2}$, we obtain that

$$E(n' - m') \geq n \cdot N(c) - \frac{n\Delta}{2} N(\sqrt{2}c)^2 = n \left(N(c) - \frac{\Delta}{2} N(\sqrt{2}c)^2 \right).$$

Using Lemma 24 (iii) we find that

$$\frac{N(c)}{N(\sqrt{2}c)^2} \geq \frac{\left(\frac{1}{c} - \frac{1}{c^3}\right) \frac{1}{\sqrt{2\pi}} e^{-\frac{c^2}{2}}}{\frac{1}{4c^2\pi} e^{-2c^2}} = 2 \left(1 - \frac{1}{c^2}\right) \sqrt{2\pi} c e^{\frac{3}{2}c^2}.$$

As $c = \sqrt{\frac{2}{3} \ln \Delta - \frac{1}{3} \ln \ln \Delta}$, we have $e^{\frac{3}{2}c^2} = \frac{\Delta}{\sqrt{\ln \Delta}}$. One can verify that

$$2 \left(1 - \frac{1}{c^2}\right) \sqrt{2\pi} c e^{\frac{3}{2}c^2} > \sqrt{2\pi} c e^{\frac{3}{2}c^2} > \Delta.$$

(This holds for Δ large enough. However, one can colour G with $\Delta + 1$ colours in polynomial time (using a greedy algorithm) and thus find a stable set of size at least $\frac{n}{\Delta+1}$ which is $\Omega\left(\frac{n}{\Delta^{\frac{1}{3}}(\log \Delta)^{\frac{1}{3}}}\right)$ for bounded Δ .) This shows that $N(c) > \Delta \cdot N(\sqrt{2}c)^2$. Therefore, $E(n' - m') \geq \frac{n}{2} N(c)$ and, using again Lemma 24 (iii),

$$E(n' - m') \geq \frac{n}{2} \left(\frac{1}{c} - \frac{1}{c^3}\right) \frac{1}{\sqrt{2\pi}} e^{-\frac{c^2}{2}} = \Omega\left(\frac{n}{\Delta^{\frac{1}{3}}(\log \Delta)^{\frac{1}{3}}}\right).$$

This concludes the proof of Theorem 23.

We mention below the k -analogue of Theorem 23, whose proof is similar. (The analogue of Lemma 26 is that the probability P_{12} is bounded by $N(\sqrt{\frac{k-1}{k-2}}c)^2$, where $c = \sqrt{(1 - \frac{2}{k})(2 \ln \Delta - \ln \ln \Delta)}$.)

Theorem 27. *Let G be a vector k -colourable graph ($k \geq 2$) on n nodes with maximum degree Δ . Then an independent set of size $\Omega\left(\frac{n}{\Delta^{1-\frac{2}{k}}(\log \Delta)^{\frac{1}{k}}}\right)$ can be found in randomized polynomial time.*

Feige, Langberg and Schechtman [81] show that this result is in some sense best possible. They show that, for all $\epsilon > 0$ and $k > 2$, there are infinitely many graphs G that are vector k -colourable and satisfy $\alpha(G) \leq \frac{n}{\Delta^{1-\frac{2}{k}-\epsilon}}$, where n is the number of nodes and Δ is the maximum degree satisfying $\Delta > n^\delta$ for some constant $\delta > 0$.

The $O(n^{1-\frac{3}{k+1}}\sqrt{n})$ -colouring algorithm of Karger-Motwani-Sudan for vector k -colourable graphs. As before, it suffices to show that one can find in randomized polynomial time an independent set of size $\Omega\left(\frac{n^{\frac{3}{k+1}}}{\sqrt{\log n}}\right) = \Omega\left(\frac{n}{n^{1-\frac{3}{k+1}}\sqrt{\log n}}\right)$ in a vector k -colourable graph. (Indeed, using recursion,

one can then find in randomized polynomial time a semicolouring using $O(n^{1-\frac{3}{k+1}}\sqrt{\log n})$ colours and thus, using Lemma 19, a colouring using the same number of colours.) The result is shown by induction on k . Suppose the result holds for any vector $(k-1)$ -colourable graph. Set $\Delta_k(n) := n^{\frac{k}{k+1}}$ and let G be a vector k -colourable graph on n nodes. We distinguish two cases.

Suppose first that G has a node u of degree greater than $\Delta_k(n)$ and consider a subgraph H of G induced by a subset of $\Delta_k(n)$ nodes contained in the neighbourhood of u . Then H is vector $(k-1)$ -colourable (easy to verify; see [125]). By the induction assumption, we can find an independent set in H (and thus in G) of size $\Omega\left(\frac{\Delta_k(n)^{\frac{3}{k}}}{\sqrt{\log \Delta_k(n)}}\right) = \Omega\left(\frac{n^{\frac{3}{k+1}}}{\sqrt{\log n}}\right)$.

Suppose now that the maximum degree Δ of G is less than or equal to $\Delta_k(n)$. It follows from Theorem 27 that we can find an independent set in G of size $\Omega\left(\frac{n}{\Delta_k(n)^{1-\frac{2}{k}}\sqrt{\log \Delta_k(n)}}\right) = \Omega\left(\frac{n^{\frac{3}{k+1}}}{\sqrt{\log n}}\right)$. This concludes the proof.

6.5 Approximating the maximum stable set and vertex cover problems

The stable set problem. Determining the stability number of a graph is a hard problem. Arora et al. [21] show the existence of a constant $\epsilon > 0$ for which there is no polynomial time algorithm permitting to find a stable set in a graph G of size at least $n^{-\epsilon}\alpha(G)$ unless $P=NP$. We saw in Section 4.2 that the theta number $\vartheta(G)$ is a polynomially computable upper bound for $\alpha(G)$ which is tight for perfect graphs, in which case a maximum cardinality stable set can be found in polynomial time. For general graphs, the gap between $\alpha(G)$ and $\vartheta(G)$ can be arbitrarily large. Indeed, Feige [75] shows that, for all $\epsilon > 0$, there is a family of graphs for which $\vartheta(G) > n^{1-\epsilon}\alpha(G)$. The proof of Feige is nonconstructive; Alon and Kahale [9] give the following constructive proof for this result.

Theorem 28. *For every $\epsilon > 0$ one can construct a family of graphs on n nodes for which $\vartheta(G) \geq (\frac{1}{2} - \epsilon)n$ and $\alpha(G) = O(n^\delta)$ where $0 < \delta < 1$ is a constant depending on ϵ .*

PROOF. Given integers $0 < s < q$, let G_{qs} denote the graph on $n = \binom{2q}{q}$ nodes corresponding to all subsets A of $Q := \{1, \dots, 2q\}$ with cardinality $|A| = q$, where A, B are adjacent if $|A \cap B| = s$. We begin with evaluating the theta number of G_{qs} . For every vertex A of G_{qs} , set $d_A := (x+1)\chi^A - \chi^Q$, where x is the largest root of the quadratic polynomial $sx^2 - 2(q-s)x + s = 0$. Then, $d_A^T d_B = 0$ for all adjacent A, B . Therefore, the vectors $v_A := \frac{d_A}{\|d_A\|}$ form an orthonormal representation of \overline{G}_{qs} . Setting $d := \frac{1}{\sqrt{2q}}(1, \dots, 1)^T$ and using the definition from Theorem 12, we obtain:

$$\vartheta(G_{qs}) \geq \sum_A (d^T v_A)^2 = n \frac{(x-1)^2}{2(x^2+1)} = \frac{nq-2s}{2q-s}.$$

In order to evaluate the stability number of G_{qs} , one can use the following result of Frankl and Rödl [86]: For every $\gamma > 0$, there exists $0 < \delta < 1$ for which $\alpha(G_{qs}) \leq n^\delta$ if $\gamma q < s < (1-\gamma)q$.

We now indicate how to choose the parameters q, s in order to achieve the conclusion of the theorem. Let $\epsilon > 0$ be given. Define s as the largest integer for which $s < \frac{q}{2}$ and $\frac{q-2s}{2(q-s)} > \frac{1}{2} - \epsilon$ (i.e., $s < \frac{2q\epsilon}{1+2\epsilon}$). Choose γ such that $0 < \gamma < \frac{\epsilon}{q}$. Then $\gamma q < s < (1-\gamma)q$ and thus $\alpha(G_{qs}) \leq n^\delta$ for some $0 < \delta < 1$ by the Frankl-Rödl result. ■

On the positive side, Alon and Kahale [9] show the following two results; we present the second one without proof.

Theorem 29. *Let G be a graph on n nodes, $k \geq 3$, $m \geq 1$ be integers. If $\vartheta(G) \geq \frac{n}{k} + m$, then an independent set of cardinality $\Omega(m^{\frac{3}{k+1}} \log^{-\frac{1}{2}} m)$ can be found in randomized polynomial time.*

PROOF. Using the definition of $\vartheta(G)$ from Theorem 12, there exist unit vectors d, v_1, \dots, v_n where v_1, \dots, v_n form an orthonormal representation of \overline{G} . These vectors can be found in polynomial time since, as the proof of Theorem 12 shows, they can be computed from an optimum solution to the SDP program (58). Order the nodes in such a way that $(d^T v_1)^2 \geq \dots \geq (d^T v_n)^2$. As $\vartheta(G) \geq \frac{n}{k} + m$ and $(d^T v_i)^2 \leq 1$ for all i , we have $(d^T v_m)^2 \geq \frac{1}{k}$. Let H denote the subgraph of G induced by the nodes $1, \dots, m$. Then, v_1, \dots, v_m is an orthonormal representation of \overline{H} , the complementary graph of H . Using the definition of the theta number from Theorem 14, we deduce that

$$\vartheta(\overline{H}) \leq \max_{i=1, \dots, m} \frac{1}{(d^T v_i)^2} \leq k.$$

Therefore, H has a vector k -colouring. Applying the Karger-Motwani-Sudan results from the preceding subsection, one can find in randomized polynomial time a $O(m^{1-\frac{3}{k+1}} \sqrt{\log m})$ colouring of H . Then the largest colour class in this colouring has cardinality $\Omega(m^{\frac{3}{k+1}} \log^{-\frac{1}{2}} m)$. ■

Theorem 30. *If G is a graph on n nodes such that $\vartheta(G) > Mn^{1-\frac{2}{k}}$ for an appropriate absolute constant M , one can find in polynomial time a stable set in G of cardinality k .* ■

Halperin, Nathaniel and Zwick [110] show the following extension of Theorem 29.

Theorem 31. *Let G be a graph on n nodes that contains an independent set of size at least $\frac{n}{\alpha}$, where $\alpha \geq 1$, and set $k := \lfloor \alpha \rfloor$. Then an independent set of G of size $\tilde{\Omega}(n^{f(\alpha)})$ can be found in randomized polynomial time, where*

$$f(\alpha) = \frac{1}{k} \cdot \frac{\alpha(\alpha-1)}{\alpha(\alpha-k) + \frac{k^2-1}{3}}$$

(the notation $\tilde{\Omega}$ meaning that logarithmic factors are hidden). In particular, $f(\alpha) = 1$ for $1 \leq \alpha \leq 2$, $f(\alpha) = \frac{\alpha}{2(\alpha-1)}$ for $2 \leq \alpha \leq 3$, and $f(k) = \frac{3}{k+1}$ for every integer $k \geq 1$.

See, e.g., [106], [107], [108] for further results.

The vertex cover problem. We now turn to the vertex cover problem. A subset $X \subseteq V$ is a *vertex cover* if every edge is adjacent to a node in X ; that is, if $V \setminus X$ is a stable set. Denote by $vc(G)$ the minimum cardinality of a vertex cover in G . Thus $vc(G) = n - \alpha(G)$ and determining $vc(G)$ is therefore an NP-hard problem.

It is well known that $vc(G)$ can be approximated within a factor of 2 in polynomial time. An easy way to see it is to take a maximal matching M ; then the set C of vertices covered by M forms a vertex cover such that $vc(G) \leq |C| = 2|M| \leq 2 \cdot vc(G)$. Alternatively, this can be seen using an LP

relaxation of the problem. Indeed, consider the LP problem:

$$\begin{aligned} lp(G) := \min & \sum_{i \in V} x_i \\ \text{s.t.} & x_i + x_j \geq 1 \quad (ij \in E) \\ & 0 \leq x_i \leq 1 \quad (i \in V) \end{aligned} \quad (108)$$

which is a linear relaxation of the vertex cover problem:

$$\begin{aligned} vc(G) := \min & \sum_{i \in V} x_i \\ \text{s.t.} & x_i + x_j \geq 1 \quad (ij \in E) \\ & x_i \in \{0, 1\} \quad (i \in V). \end{aligned} \quad (109)$$

Obviously, $lp(G) \leq vc(G)$. Moreover, $vc(G) \leq 2 \cdot lp(G)$; indeed, given an optimum solution x to (108), the set $X := \{i \in V \mid x_i \geq \frac{1}{2}\}$ is a vertex cover whose cardinality satisfies $|X| \leq 2 \cdot lp(G)$.

On the negative side, it is known that the minimum vertex cover problem cannot be approximated in polynomial time within any factor smaller than $10\sqrt{5} - 21 \sim 1.36067$ if $P \neq NP$ [69]. The existence of a polynomial time approximation algorithm for the vertex cover problem with performance ratio $2 - \epsilon$ remains, however, open for any $\epsilon > 0$. Kleinberg and Goemans [132] propose to use the following semidefinite relaxation of the problem (109):

$$\begin{aligned} sd(G) := \min & \sum_{i=1}^n \frac{1 + v_0^T v_i}{2} \\ \text{s.t.} & (v_0 - v_i)^T (v_0 - v_j) = 0 \quad (ij \in E) \\ & v_0, v_1, \dots, v_n \text{ unit vectors.} \end{aligned} \quad (110)$$

They show that this semidefinite bound $sd(G)$ is equal to the obvious lower bound $n - \vartheta(G)$ for $vc(G)$, where $\vartheta(G)$ is the theta number bounding $\alpha(G)$. To see it, consider the matrix $X = (x_{ij})_{i,j=0}^n$ where $x_{ij} = v_i^T v_j$ and v_0, \dots, v_n satisfy (110); then X is constrained to be positive semidefinite with an all ones diagonal and to satisfy $1 + x_{ij} - x_{0i} - x_{0j} = 0$ for all edges ij of G . If we define the matrix $Y = (y_{ij})_{i,j=1}^n$ by

$$y_{ij} = \frac{1}{4}(1 + x_{ij} - x_{0i} - x_{0j}) \quad \text{for } i, j = 1, \dots, n,$$

then the objective function in (110) reads $n - \sum_{i=1}^n y_{ii}$ and X is feasible for (110) if and only if Y satisfies $Y - \text{diag}(Y)\text{diag}(Y)^T \succeq 0$ and $y_{ij} = 0$ ($ij \in E$); that is, if the vector $(y_{ii})_{i=1}^n$ belongs to the theta body $\text{TH}(G)$. (We use the definition of $\vartheta(G)$ from Theorem 11. See [155] for details on the above $X \mapsto Y$ mapping.)

A first observation is that this SDP bound is at least as good as the LP bound; namely,

$$sd(G) = n - \vartheta(G) \geq lp(G).$$

To see it, use the definition from Theorem 12. let d be a unit vector and v_1, \dots, v_n an orthonormal representation of \overline{G} such that $\vartheta(G) = \sum_{i \in V} (d^T v_i)^2$. Set $x_i := 1 - (d^T v_i)^2$ ($i \in V$). Then x is a feasible solution to the program (108) which shows that $lp(G) \leq \sum_i x_i = n - \vartheta(G)$.

Kleinberg and Goemans [132] construct a class of graphs G for which the ratio $\frac{vc(G)}{n - \vartheta(G)}$ converges to 2 as n goes to infinity, which shows that no improvement is made by using SDP instead of LP. (In fact, the class of graphs constructed in Theorem 28 displays the same behaviour.) They also propose to strengthen the semidefinite program (110) by adding to it the constraints

$$(v_0 - v_i)^T (v_0 - v_j) \geq 0 \quad (ij \in \overline{E});$$

the new semidefinite bound can be verified to be equal to $n - \vartheta'(G)$, where $\vartheta'(G)$ is the sharpening of $\vartheta(G)$ introduced in Section 4.4. Charikar [51] shows that the new integrality gap $\frac{vc(G)}{n - \vartheta'(G)}$ can again be made arbitrarily close to 2.

Improved approximation algorithms exist for graphs with bounded maximum degree Δ . Improving on earlier results, Halperin [108] shows that, for graphs with maximum degree Δ , the semidefinite relaxation (110) together with suitable randomized rounding permits to derive an approximation algorithm for the minimum vertex cover problem with performance ratio $2 - (1 - o(1)) \frac{2 \ln \ln \Delta}{\ln \Delta}$ for large Δ . We sketch this result below.

Halperin's algorithm is based on the following observation. Given a scalar $x \geq 0$, the set $C := \{i \in \{1, \dots, n\} \mid v_0^T v_i \geq -x\}$ is a vertex cover. Note that for $x = 0$, we have $|C| \leq 2 \cdot sd(G)$ and thus this gives again a 2-approximation algorithm. Moreover, if J is an independent set contained in the set $S_2 := \{i \in \{1, \dots, n\} \mid -x \leq v_0^T v_i < x\}$, then the set $C \setminus J$ is still a vertex cover. When x is small, nodes in S_2 correspond to vectors v_i that are approximatively orthogonal to v_0 and thus the endpoints of an edge contained in S_2 correspond to approximatively opposite vectors. Hence the set S_2 is likely to contain few edges and thus a large independent set J ; therefore, the set $C \setminus J$ is likely to be a small vertex cover.

More precisely, Halperin defines $x = \Theta(\frac{\ln \ln \Delta}{\ln \Delta})$ and the sets $S_1 = \{i \in \{1, \dots, n\} \mid v_0^T v_i \geq x\}$ and $S_2 = \{i \in \{1, \dots, n\} \mid -x \leq v_0^T v_i < x\}$ as above (thus $C = S_1 \cup S_2$). Then, $|S_1| \leq \frac{2}{x+1} sd(G)$ and $|S_2| \leq \frac{2}{1-x} sd(G)$. A large independent set J can be found in S_2 using the 'rounding via vector projections' technique from [125], exposed earlier in Section 6.4. Indeed, if ij is an edge contained in S_2 , then $v_i^T v_j = v_0^T v_i + v_0^T v_j - 1 < 2x - 1$. Hence, the subgraph of G induced by S_2 has a vector k -colouring for $k = \frac{2(1-x)}{1-2x}$. Therefore, Theorem 27 can be used for finding a large independent set in S_2 . These facts yield the desired performance ratio; see [108] for details.

As mentioned above, no polynomial time approximation algorithm is known for the vertex cover problem having a performance ratio $2 - \epsilon$ with $\epsilon > 0$. In fact, no tractable linear relaxation is known for (109), having an integrality gap lower than 2. Arora, Bollobás and Lovász [19] initiate a more systematic approach for proving non existence of tighter relaxations. They show an integrality gap of $2 - o(1)$ for some fairly general families of LP relaxations of (109). A first family consists of the LP relaxations in which each constraint has at most ϵn variables. A second family involves LP relaxations in which each constraint has defect at most ϵn ; the defect of an inequality $a^T x \geq b$ being $2b - \sum_i a_i$. A third family consists of the LP relaxations obtained after $O(1)$ iterations of the Lovász-Schrijver N operator applied to the LP in (108). It is an open question whether an analogue result holds for the N_+ operator.

6.6 Approximating MAX SAT

An instance of the MAX SAT problem in the Boolean variables x_1, \dots, x_n is composed of a collection \mathcal{C} of clauses C with nonnegative weights w_C associated to them. Each clause C is of the form $z_1 \vee \dots \vee z_k$ where each z_j is either a variable x_i or its negation \bar{x}_i (called a *literal*); k is its *length* and C is satisfied if at least one of the literals z_1, \dots, z_k is assigned value 1 (if a variable x_i is assigned value 1 then its negation \bar{x}_i is assigned value 0 and vice versa). The MAX SAT problem consists of finding an assignment of 0/1 values to the variables x_1, \dots, x_n so that the total weight of the satisfied clauses is maximized. Given an integer $k \geq 1$, the MAX k SAT problem is the special instance of MAX SAT where each clause has length at most k and MAX Ek SAT is the instance where all clauses have

length exactly k ; an instance of MAX SAT is said to be *satisfiable* if there is an assignment of the x_i 's satisfying all its clauses.

The MAX SAT and MAX k SAT problems are NP-hard. Moreover, Håstad [114] proved that, for any $\epsilon > 0$, there is no $(\frac{7}{8} + \epsilon)$ -approximation algorithm for MAX SAT, unless $P=NP$; his result also holds when restricted to satisfiable instances of MAX E3SAT. Håstad [114] also proved that, for any $\epsilon > 0$, there is no $(\frac{21}{22} + \epsilon)$ -approximation algorithm for MAX 2SAT unless $P=NP$.

A $\frac{3}{4}$ -approximation algorithm for MAX SAT. The first approximation algorithm for MAX SAT is the following $\frac{1}{2}$ -approximation algorithm due to Johnson [123]. Given $p_i \in [0, 1]$ ($i = 1, \dots, n$), set independently and randomly each variable x_i to 1 with probability p_i . Then the probability that a clause $C := \bigvee_{i \in I_C^+} x_i \bigvee \bigvee_{i \in I_C^-} \bar{x}_i$ is satisfied is equal to $1 - \prod_{i \in I_C^+} (1 - p_i) \prod_{i \in I_C^-} p_i$. If we set all p_i 's to $\frac{1}{2}$, then the total expected weight \hat{W}_1 of satisfied clauses satisfies:

$$\hat{W}_1 = \sum_{C \in \mathcal{C}} w_C (1 - \frac{1}{2^{k_C}}) \geq \frac{1}{2} \sum_{C \in \mathcal{C}} w_C$$

where k_C is the length of clause C . Therefore, this gives a randomized $\frac{1}{2}$ -approximation algorithm for MAX SAT or a $(1 - 2^{-k})$ -approximation algorithm for instances of MAX SAT where all clauses have length $\geq k$ (thus with performance ratio $\frac{3}{4}$ for MAX E2SAT and $\frac{7}{8}$ for MAX E3SAT); it can be derandomized using the method of conditional probabilities.

Goemans and Williamson [94] give an improved $\frac{3}{4}$ -approximation algorithm using linear programming. Consider the integer programming problem:

$$\begin{aligned} \max \quad & \sum_{C \in \mathcal{C}} w_C z_C \\ \text{s.t.} \quad & z_C \leq \sum_{i \in I_C^+} y_i + \sum_{i \in I_C^-} (1 - y_i) \quad (C \in \mathcal{C}) \\ & 0 \leq z_C \leq 1 \quad (C \in \mathcal{C}) \\ & y_i \in \{0, 1\} \quad (i = 1, \dots, n) \end{aligned} \tag{111}$$

and let Z_{LP}^* denote the optimum value of its linear programming relaxation obtained by relaxing the condition $y_i \in \{0, 1\}$ by $0 \leq y_i \leq 1$. If (y, z) is an optimum solution to (111), letting $x_i = 1$ if and only if $y_i = 1$, then clause C is satisfied precisely when $z_C = 1$; hence (111) solves the MAX SAT problem. The GW approximation algorithm goes as follows. First, solve the LP relaxation of (111) and let (y, z) be an optimum solution to it. Then, apply the Johnson's algorithm using the probabilities $p_i := y_i$; that is, set x_i to 1 with probability y_i . Setting $\beta_k := 1 - \left(1 - \frac{1}{k}\right)^k$ and using the fact⁴ that $1 - \prod_{i \in I_C^+} (1 - y_i) \prod_{i \in I_C^-} y_i \geq \beta_{k_C} z_C$, we find that the expected weight \hat{W}_2 of satisfied clauses satisfies:

$$\hat{W}_2 = \sum_{C \in \mathcal{C}} w_C \left(1 - \prod_{i \in I_C^+} (1 - y_i) \prod_{i \in I_C^-} y_i \right) \geq \sum_{C \in \mathcal{C}} w_C z_C \beta_{k_C}.$$

As β_k is a monotone decreasing function of k , this gives a randomized β_k -approximation algorithm for instances of MAX SAT where all clauses have at most k literals; thus a $(1 - \frac{1}{e})$ -approximation algorithm for MAX SAT, since $\lim_{k \rightarrow \infty} (1 - \frac{1}{k})^k = \frac{1}{e}$.

⁴The proof uses the arithmetic/geometric mean inequality: $\frac{a_1 + \dots + a_n}{n} \geq (a_1 \cdots a_n)^{\frac{1}{n}}$ for any nonnegative numbers a_1, \dots, a_n .

In order to obtain the promised $\frac{3}{4}$ performance ratio, it suffices to combine the above two algorithms. For this, note that $\frac{1}{2}(1 - \frac{1}{2^k} + \beta_k) \geq \frac{3}{4}$ for all $k \geq 1$. Therefore, $\frac{1}{2}(\hat{W}_1 + \hat{W}_2) \geq \frac{3}{4}Z_{LP}^*$. Hence the following is a $\frac{3}{4}$ -approximation algorithm for MAX SAT: with probability $\frac{1}{2}$, use the probabilities $p_i := \frac{1}{2}$ for determining the variables x_i and, with probability $\frac{1}{2}$, use instead the probabilities $p_i := y_i$.

Other $\frac{3}{4}$ -approximation algorithms for MAX SAT are given by Goemans and Williamson [94]. Instead of setting $x_i = 1$ with probability y_i , they set $x_i = 1$ with probability $f(y_i)$ for some suitably chosen function $f(\cdot)$.

Better approximation algorithms can be obtained using semidefinite relaxations instead of linear ones combined with adequate rounding techniques, as we now see.

The Goemans-Williamson α_0 -approximation algorithm for MAX 2SAT and their 0.7554-approximation algorithm for MAX SAT. Using a semidefinite relaxation for MAX SAT instead of a linear one and the hyperplane rounding technique, one can show a better approximation algorithm. It is convenient to introduce the new Boolean variables $x_{n+i} = \bar{x}_i$ for $i = 1, \dots, n$. Then a clause C can be expressed as a disjunction $C = \bigvee_{i \in I_C} x_i$, of the variables x_1, \dots, x_{2n} , with $I_C \subseteq \{1, \dots, 2n\}$. It is also convenient to work with ± 1 variables v_i (instead of $y_i \in \{0, 1\}$) and to introduce an additional ± 1 variable v_0 , the convention being to set x_i to 1 if $v_i = -v_0$ and to 0 if $v_i = v_0$. Hence the formulation (111) of MAX SAT can be rewritten as

$$\begin{aligned} \max \quad & \sum_{C \in \mathcal{C}} w_C z_C \\ \text{s.t.} \quad & z_C \leq \sum_{i \in I_C} \frac{1 - v_0 \cdot v_i}{2} \quad (C \in \mathcal{C}) \\ & 0 \leq z_C \leq 1 \quad (C \in \mathcal{C}) \\ & v_i \cdot v_{n+i} = -1 \quad (i = 1, \dots, n) \\ & v_0, v_1, \dots, v_{2n} \in \{\pm 1\}. \end{aligned} \tag{112}$$

For each clause $C = x_i \vee x_j$ of length 2, one can add the constraint:

$$z_C \leq 1 - \left(\frac{1 + v_0 \cdot v_i}{2} \right) \left(\frac{1 + v_0 \cdot v_j}{2} \right) = \frac{3 - v_0 \cdot v_i - v_0 \cdot v_j - v_i \cdot v_j}{4} \tag{113}$$

which, in fact, implies the constraint $z_C \leq \frac{1 - v_0 \cdot v_i}{2} + \frac{1 - v_0 \cdot v_j}{2}$.

Let (SDP) denote the semidefinite relaxation of the program (112) augmented with the constraints (113) for all clauses of length 2, which is obtained by introducing a matrix variable $X = (X_{ij})_{i,j=0}^{2n} \succeq 0$ and replacing each product $v_i \cdot v_j$ by X_{ij} . In other words, this amounts to replacing the constraint $v_0, \dots, v_{2n} \in \{\pm 1\}$ by the constraint $v_0, \dots, v_{2n} \in S^n$, S^n being the unit sphere in \mathbb{R}^{n+1} (the product $v_i \cdot v_j$ meaning then the inner product $v_i^T v_j$).

Goemans and Williamson [95] show that their basic α_0 -approximation algorithm for max-cut extends to MAX 2SAT. Namely, solve the relaxation (SDP) and let v_0, \dots, v_n be the optimum unit vectors solving it; select a random unit vector r and let H_r be the hyperplane with normal vector r ; set x_i to 1 if the hyperplane H_r separates v_0 and v_i and to 0 otherwise. Let θ_{ij} denote the angle (v_i, v_j) . Then the probability $\text{prob}(v_0, v_i)$ that the clause x_i is satisfied is equal to the probability that H_r separates v_0 and v_i and thus

$$\text{prob}(v_0, v_i) = \frac{\theta_{0i}}{\pi};$$

the probability $\text{prob}(v_0, v_i, v_j)$ that the clause $x_i \vee x_j$ is satisfied is equal to the probability that a random hyperplane separates v_0 from at least one of v_i and v_j which can be verified to be equal to

$$\text{prob}(v_0, v_i, v_j) = \frac{1}{2\pi}(\theta_{0i} + \theta_{0j} + \theta_{ij})$$

using the inclusion/exclusion principle. Therefore, for a clause $C = x_i \vee x_j$, we have

$$\frac{\text{prob}(v_0, v_i, v_j)}{z_C} \geq \frac{2}{\pi} \frac{\theta_{0i} + \theta_{0j} + \theta_{ij}}{3 - \cos \theta_{0i} - \cos \theta_{0j} - \cos \theta_{ij}} \geq \alpha_0,$$

where $\alpha_0 \simeq 0.87856$ is the Goemans-Williamson ratio from (84). The above relation also holds when $i = j$, i.e., when C is a clause of length 1, in which case one lets $\text{prob}(v_0, v_i, v_j) = \text{prob}(v_0, v_i)$. Hence the expected total weight of satisfied clauses is greater than or equal to α_0 times the optimum value of the relaxation (SDP); this gives therefore an α_0 -approximation algorithm for MAX 2SAT.

This improved MAX 2SAT algorithm leads to a slightly improved 0.7554-approximation algorithm for general MAX SAT. For this, one considers the following three algorithms: (1) set x_i to 1 independently with probability $p_i := \frac{1}{2}$; (2) set x_i to 1 independently with probability $p_i := \frac{1-v_0^T v_i}{2}$; (3) select a random hyperplane H_r and set x_i to 1 if H_r separates v_0 and v_i (the v_i 's being the optimum vectors to the relaxation (SDP)). One chooses algorithm (i) with probability q_i where $q_1 = q_2 = 0.4785$ and $q_3 = 1 - q_1 - q_2 = 0.0430$. Then the expected weight of the satisfied clauses is at least

$$\sum_{C|k_C \leq 2} w_C z_C \left(\frac{3}{2} q_1 + q_3 \cdot \alpha_0 \right) + \sum_{C|k_C \geq 3} w_C z_C \cdot q_1 \left(1 - \frac{1}{2^k} + 1 - \left(1 - \frac{1}{k} \right)^k \right)$$

which can be verified to be at least $0.7554 \cdot \sum_C w_C z_C$. A refinement of this algorithm is given in [94] with an improved performance ratio 0.7584.

The improved Feige-Goemans 0.931-approximation algorithm for MAX 2SAT. Feige and Goemans [77] show an improved performance ratio of about 0.931 for MAX 2SAT. For this, they strengthen the semidefinite relaxation (SDP) by adding to it the triangle inequalities:

$$X_{0i} + X_{0j} + X_{ij} \geq -1, \quad X_{0i} - X_{0j} - X_{ij} \geq -1, \quad -X_{0i} - X_{0j} + X_{ij} \geq -1 \quad (114)$$

for all $i, j \in \{1, \dots, 2n\}$. Moreover, they replace the vectors v_0, v_1, \dots, v_n (obtained from the optimum solution to the strengthened semidefinite program) by a new set of vectors v'_0, \dots, v'_n obtained by applying some *rotation* to the v_i 's. Then the assignment for the Boolean variables x_i are generated from the v'_i using as before the hyperplane rounding technique.

Let us explain how the vectors v'_i are generated from the v_i 's. Let $f : [0, \pi] \rightarrow [0, \pi]$ be a continuous function such that $f(0) = 0$ and $f(\pi - \theta) = \pi - f(\theta)$. As before, θ_{ij} denotes the angle (v_i, v_j) . The vector v_i is rotated in the plane spanned by v_0 and v_i until it forms an angle of $f(\theta_{0i})$ with v_0 ; the resulting vector is v'_i . If $v_i = v_0$ then $v'_i = v_i$. Moreover, let $v'_{n+i} = -v'_i$ for $i = 1, \dots, n$. Let θ'_{ij} be the angle (v'_i, v'_j) . Then $\theta'_{0i} = f(\theta_{0i})$ and Feige and Goemans [77] show the following equation permitting to express θ'_{ij} in terms of θ_{ij} :

$$\cos \theta'_{ij} = \cos \theta'_{0i} \cos \theta'_{0j} + \frac{\cos \theta_{ij} - \cos \theta_{0i} \cos \theta_{0j}}{\sin \theta_{0i} \sin \theta_{0j}} \sin \theta'_{0i} \sin \theta'_{0j}. \quad (115)$$

The probability that the clause $x_i \vee x_j$ is satisfied is now equal to

$$\text{prob}(v_0, v'_i, v'_j) = \frac{\theta'_{0i} + \theta'_{0j} + \theta'_{ij}}{2\pi}$$

while the contribution of this clause to the objective function of the semidefinite relaxation is

$$z_C \leq \frac{3 - \cos \theta_{0i} - \cos \theta_{0j} - \cos \theta_{ij}}{4}.$$

The performance ratio of the approximation algorithm using a rotation function f is, therefore, at least

$$\beta(f) := \min \frac{2}{\pi} \cdot \frac{\theta'_{01} + \theta'_{02} + \theta'_{12}}{3 - \cos \theta_{01} - \cos \theta_{02} - \cos \theta_{12}}$$

where the minimum is taken over all $\theta_{01}, \theta_{02}, \theta_{12} \in [0, \pi]$ for which $\cos \theta_{01}, \cos \theta_{02}, \cos \theta_{12}$ satisfy the triangle inequalities (114). Recall that $\theta'_{0i} = f(\theta_{0i})$ and relation (115) permits to express θ'_{12} in terms of θ_{01}, θ_{02} and θ_{12} .

Feige and Goemans [77] used a rotation function of the form

$$f_\lambda(\theta) = (1 - \lambda)\theta + \lambda \frac{\pi}{2}(1 - \cos \theta) \quad (116)$$

and, for the choice $\lambda = 0.806765$, they claim the lower bound 0.93109 for $\beta(f)$. Proving a correct evaluation of $\beta(f)$ is a non trivial task, since the minimization program defining $\beta(f)$ is too complicated to be handled analytically. Zwick [241] makes a detailed and rigorous analysis enabling him to prove a performance ratio of 0.931091 for MAX 2SAT.

The Matuura-Matsui 0.935-approximation algorithm for MAX 2SAT. Matuura and Matsui [172] design an approximation algorithm for MAX 2SAT with performance ratio 0.935. As in the Feige-Goemans algorithm, their starting point is to use the semidefinite relaxation (SDP') of MAX 2SAT obtained from (112) by adding the constraints (113) for the clauses of length 2 and the triangle inequalities (114); they fix v_0 to be equal to $(1, 0, \dots, 0)^T$. Let v_1, \dots, v_n be the unit vectors obtained from an optimum solution to the program (SDP'). No rotation is applied to the vectors v_i as in the Feige-Goemans algorithm. The new ingredient in the algorithm of Matuura-Matsui consists of selecting the random hyperplane using a distribution function f on the sphere which is skewed towards v_0 and uniform in any direction orthogonal to v_0 , instead of a uniform distribution.

Let \mathcal{F}_n denote the set of functions $f : S^n \rightarrow \mathbb{R}_+$ satisfying $\int_{S^n} f(v) dv = 1$, $f(v) = f(-v)$ for all $v \in S^n$, and $f(u) = f(v)$ for all $u, v \in S^n$ such that $u^T v_0 = v^T v_0$. Let $f \in \mathcal{F}_n$ and let the random unit vector r be now chosen according to the distribution function f . Then, $\text{prob}(v_i, v_j \mid f)$ denotes the probability that the clause $x_i \vee x_j$ is satisfied, i.e., as before, the probability that $\text{sign}(r^T v_0) \neq \text{sign}(r^T v_i)$ or $\text{sign}(r^T v_0) \neq \text{sign}(r^T v_j)$. Let P denote the linear subspace spanned by v_0, v_i, v_j and let \hat{f} denote the distribution on S^2 obtained by projecting onto P ; that is, $\hat{f}(v') := \int_{T(v')} f(v) dv$, where $T(v')$ is the set of all $v \in S^n$ whose projection on P is parallel to v' . Then the new approximation ratio of the algorithm is equal to

$$\alpha_{\hat{f}} := \min \frac{\text{prob}(v_i, v_j \mid \hat{f})}{1/4(3 - v_0^T v_i - v_0^T v_j - v_i^T v_j)}$$

where the minimum is taken over all $v_i, v_j \in S^2$ which together with $v_0 = (1, 0, 0)^T$ have their pairwise inner products satisfying the triangle inequalities (114).

The difficulty consists of constructing a distribution function $f \in \mathcal{F}_n$ for which $\alpha_{\hat{f}}$ is large. Matuura and Matsui [172] show the following. The function

$$g(v) := \cos^{1/1.3}(\theta) \text{ for all } v \in S^2 \text{ with } |v_0^T v| = \cos \theta, \quad (117)$$

is a distribution function on S^2 belonging to \mathcal{F}_2 ; it satisfies $\alpha_g \geq 0.935$ (this is proved numerically); and there exists $f \in \mathcal{F}_n$ for which $\hat{f} = g$.

The Lewin-Livnat-Zwick 0.940-approximation algorithm for MAX 2SAT. Lewin, Livnat and Zwick [157] achieve this improved performance ratio by combining the skewed hyperplane rounding technique exploited by Matuura and Matsui [172] with the pre-rounding rotation phase used by Feige and Goemans [77].

The Karloff-Zwick $\frac{7}{8}$ -approximation algorithm for MAX 3SAT. Karloff and Zwick [127] present an approximation algorithm for MAX 3SAT whose performance ratio they conjecture to be equal to $7/8 = 0.875$, thus the best possible since Håstad [114] proved the non existence of an approximation algorithm with performance ratio $> 7/8$ unless $P=NP$. Previous algorithms were using a reduction to the case of MAX 2SAT; for instance, Trevisan et al. [227] give a 0.801-approximation algorithm for MAX 3SAT using the Feige-Goemans 0.931 result for MAX 2SAT. Karloff and Zwick do not make such a reduction but consider instead the following direct semidefinite relaxation for MAX 3SAT:

$$\begin{aligned} \max \quad & \sum_{i,j,k \in \{1, \dots, 2n\}} w_{ijk} z_{ijk} \\ \text{s.t.} \quad & z_{ijk} \leq \text{relax}(v_0, v_i, v_j, v_k) \\ & v_i \cdot v_{n+i} = -1 \quad (i = 1, \dots, n) \\ & v_0, \dots, v_{2n} \in S^n, \quad z_{ijk} \in \mathbb{R}, \end{aligned}$$

where z_{ijk} is a scalar attached to the clause $x_i \vee x_j \vee x_k$ and

$$\text{relax}(v_0, v_i, v_j, v_k) := \min\left(1 - \frac{(v_0 + v_i)^T(v_j + v_k)}{4}, 1 - \frac{(v_0 + v_j)^T(v_i + v_k)}{4}, 1 - \frac{(v_0 + v_k)^T(v_i + v_j)}{4}, 1\right).$$

Note indeed that when the v_i 's are ± 1 scalars, then $\text{relax}(v_0, v_i, v_j, v_k)$ is equal to 0 precisely when $v_0 = v_i = v_j = v_k$ which corresponds to setting all variables x_i, x_j, x_k to 0 and thus to the clause $x_i \vee x_j \vee x_k$ not being satisfied.

Denote again by $\text{prob}(v_0, v_i, v_j, v_k)$ the probability that $x_i \vee x_j \vee x_k$ is satisfied and set

$$\text{ratio}(v_0, v_i, v_j, v_k) := \frac{\text{prob}(v_0, v_i, v_j, v_k)}{\text{relax}(v_0, v_i, v_j, v_k)}.$$

For a clause of length 1 or 2 (obtained by letting $j = k = 0$ or $k = 0$), it follows from the analysis of the GW algorithm that $\text{ratio}(v_0, v_i, v_j, v_k) \geq \alpha_0 > \frac{7}{8}$. For clauses of length 3, the analysis is technically much more involved and requires the computation of the volume of spherical tetrahedra as we now see.

Clearly, $\text{prob}(v_0, v_i, v_j, v_k)$ is equal to the probability that the random hyperplane H_r separates v_0 from at least one of v_i, v_j, v_k and thus to

$$1 - 2 \cdot \text{prob}(r^T v_h \geq 0 \quad \forall h = 0, i, j, k).$$

We may assume without loss of generality that v_0, v_i, v_j, v_k lie in \mathbb{R}^4 and, since we are only interested in the inner products $r^T v_h$, we can replace r by its normalized projection on \mathbb{R}^4 which is then uniformly distributed on the sphere S^3 . Define

$$T(v_0, v_i, v_j, v_k) := \{r \in S^3 \mid r^T v_h \geq 0 \quad \forall h = 0, i, j, k\}.$$

Then, $\text{prob}(v_0, v_i, v_j, v_k) = 1 - 2 \cdot \frac{\text{vol}(T(v_0, v_i, v_j, v_k))}{\text{vol}(S^3)}$, where $\text{vol}(\cdot)$ denotes the 3-dimensional spherical volume. As $\text{vol}(S^3) = 2\pi^2$, we find that

$$\text{prob}(v_0, v_i, v_j, v_k) = 1 - 2 \cdot \frac{\text{vol}(T(v_0, v_i, v_j, v_k))}{\pi^2}.$$

When the vectors v_0, v_i, v_j, v_k are linearly independent, $T(v_0, v_i, v_j, v_k)$ is a spherical tetrahedron, whose vertices are the vectors $v'_0, v'_i, v'_j, v'_k \in S^3$ satisfying $v_h^T v'_h > 0$ for all h and $v_{h_1}^T v'_{h_2} = 0$ for all distinct h_1, h_2 . That is,

$$T(v_0, v_i, v_j, v_k) = \left\{ \sum_{h=0,i,j,k} \alpha_h v'_h \mid \alpha_h \geq 0, \sum_h \alpha_h = 1 \right\}.$$

Therefore, evaluating the quantity $\text{ratio}(v_0, v_i, v_j, v_k)$ and thus the performance ratio of the algorithm relies on proving certain inequalities about volumes of spherical tetrahedra.

Karloff and Zwick [127] show that $\text{prob}(v_0, v_i, v_j, v_k) \geq \frac{7}{8}$ whenever $\text{relax}(v_0, v_i, v_j, v_k) = 1$, which shows a performance ratio $\frac{7}{8}$ for satisfiable instances of MAX 3SAT. Their proof is computer assisted as it involves one computation carried out with Mathematica. Zwick [242] can prove the performance ratio $\frac{7}{8}$ for general MAX 3SAT. Although his proof is again computer assisted, it can however be considered as a rigorous proof since it is carried out using a new system called RealSearch, written by Zwick, which involves only interval arithmetic (instead of floating point arithmetic). We refer to Zwick's paper for an interesting presentation and discussion.

Further extensions. Karloff and Zwick [127] describe a procedure for constructing strong semidefinite relaxations for general constraint satisfaction problems and thus for MAX k SAT. Halperin and Zwick [112] study approximation algorithms for MAX 4SAT using the semidefinite relaxation provided by the Karloff-Zwick recipe. The analysis of the classic hyperplane rounding technique necessitates now the evaluation of the probability $\text{prob}(v_0, \dots, v_4)$ that a random hyperplane separates v_0 from at least one of v_1, \dots, v_4 . Luckily, using the inclusion/exclusion formula, this probability can be expressed in terms of the probabilities $\text{prob}(v_i, v_j)$ and $\text{prob}(v_i, v_j, v_k, v_\ell)$ that were considered above. In this way, Halperin and Zwick can show a performance ratio of 0.845173 for MAX 4SAT, thus below the target ratio of $\frac{7}{8}$. They study in detail a variety of other possible rounding strategies which enable them to obtain some improved performance ratios, like 0.8721.

Asano and Williamson [22] present an improved approximation algorithm for MAX SAT with performance ratio 0.7846. For this, they use a new family of approximation algorithms extending the $\frac{3}{4}$ -approximation algorithm of Goemans and Williamson [94] (presented earlier in this section) combined with the semidefinite approaches for MAX 2SAT and MAX 3SAT of Karloff and Zwick [127] and Feige and Goemans [77].

Anjos [13] studies higher order semidefinite relaxations for the satisfiability problem.

6.7 Approximating the maximum directed cut problem

Given a directed graph $G = (V, A)$ and weights $w \in \mathbb{Q}_+^A$ associated to its arcs, the *maximum directed cut problem* asks for a directed cut $\delta^+(S)$ of maximum weight where, for $S \subseteq V$, the *directed cut* (or *dicut*) $\delta^+(S)$ is the set of arcs ij with $i \in S$ and $j \notin S$. This problem is NP-hard, since the max-cut problem in a undirected graph H reduces to the maximum dicut problem in the directed graph obtained by replacing each edge of H by two opposite arcs. Moreover, no approximation algorithm for the maximum dicut problem exists having a performance ratio $> \frac{12}{13}$ unless P=NP [114].

The simple random partition algorithm (which assigns each node to S independently with probability $\frac{1}{2}$) has a performance ratio $\frac{1}{4}$. Goemans and Williamson [95] show that their basic approximation algorithm for max-cut can be extended to the maximum dicut problem with performance ratio 0.79607. Feige and Goemans [77] prove an improved performance ratio of 0.859. These algorithms use the same ideas as the algorithms for MAX 2SAT presented in the same papers. Before presenting them, we

mention a simple $\frac{1}{2}$ -approximation algorithm of Halperin and Zwick [113] using a linear relaxation of the problem; this algorithm can in fact be turned into a purely combinatorial algorithm.

A $\frac{1}{2}$ -approximation algorithm by Halperin and Zwick. Consider the following linear program:

$$\begin{aligned} \max \quad & \sum_{ij \in A} w_{ij} z_{ij} \\ \text{s.t.} \quad & z_{ij} \leq x_i \quad (ij \in A) \\ & z_{ij} \leq 1 - x_j \quad (ij \in A) \\ & 0 \leq x_i \leq 1 \quad (i \in V). \end{aligned} \tag{118}$$

If we replace the linear constraint $0 \leq x \leq 1$ by the integer constraint $x \in \{0, 1\}^V$ then we obtain a formulation for the maximum dicut problem; the dicut $\delta^+(S)$ with $S = \{i \mid x_i = 1\}$ being an optimum dicut. Halperin and Zwick [113] show that the program (118) has a half-integer optimum solution. To see it, note first that (118) is equivalent to the program:

$$\begin{aligned} \max \quad & \sum_{ij \in A} w_{ij} z_{ij} \\ \text{s.t.} \quad & z_{ij} + z_{jk} \leq 1 \quad (ij \in A, jk \in A) \\ & 0 \leq z_{ij} \leq 1 \quad (ij \in A). \end{aligned} \tag{119}$$

Indeed, if (z, x) is feasible for (118), then z is feasible for (119); conversely, if z is feasible for (119) then (z, x) is feasible for (118), where $x_i := \max_{ij \in A} z_{ij}$ if $\delta^+(i) \neq \emptyset$ and $x_i := 0$ otherwise. Now, the constraints in (119) define in fact the fractional stable set polytope of the line graph of G (whose nodes are the arcs, with two arcs being adjacent if they form a path in G). Since the vertices of the fractional stable set polytope are half-integral, it follows that (119) and thus (118) has a half-integral optimum solution (x, z) . Then one constructs a directed cut $\delta^+(S)$ by putting node $i \in V$ in S with probability x_i . The expected weight of $\delta^+(S)$ is at least $\frac{1}{2} w^T z$. Therefore, this gives a $\frac{1}{2}$ -approximation algorithm. Moreover, this algorithm can be made purely combinatorial since a half-integral solution can be found using a bipartite matching algorithm (see [113]).

The Goemans-Williamson 0.796-approximation algorithm. One can alternatively model the maximum dicut problem in the following way. Given $v_0, v_1, \dots, v_n \in \{\pm 1\}$ and $S := \{i \in \{1, \dots, n\} \mid v_i = v_0\}$, the quantity

$$\frac{1}{4}(1 + v_0 \cdot v_i)(1 - v_0 \cdot v_j) = \frac{1}{4}(1 + v_0 \cdot v_i - v_0 \cdot v_j - v_i \cdot v_j)$$

is equal to 1 if $ij \in \delta^+(S)$ and to 0 otherwise. Therefore, the following program solves the maximum dicut problem:

$$\begin{aligned} \max \quad & \sum_{ij \in A} w_{ij} \frac{1}{4}(1 + v_0 \cdot v_i - v_0 \cdot v_j - v_i \cdot v_j) \\ \text{s.t.} \quad & v_0, v_1, \dots, v_n \in \{\pm 1\} \end{aligned} \tag{120}$$

Let (SDP) denote the relaxation of (120) obtained by replacing the condition $v_0, v_1, \dots, v_n \in \{\pm 1\}$ by the condition $v_0, v_1, \dots, v_n \in S^n$ and let z_{sdp} denote its optimum value. Goemans and Williamson propose the following analogue of their max-cut algorithm for solving the maximum dicut problem: Solve (SDP) and let v_0, \dots, v_n be an optimum solution to it; select a random unit vector r and let $S := \{i \in \{1, \dots, n\} \mid \text{sign}(v_0 \cdot r) = \text{sign}(v_i \cdot r)\}$. Let θ_{ij} denote the angle (v_i, v_j) . Then the expected weight $E(S)$ of the dicut $\delta^+(S)$ is equal to

$$E(S) = \sum_{ij \in A} w_{ij} \frac{1}{2\pi} (-\theta_{0i} + \theta_{0j} + \theta_{ij}).$$

In order to bound $\frac{E(S)}{z_{sdp}}$, one has to find lower bounds for the quantity

$$\frac{2}{\pi} \frac{-\theta_{0i} + \theta_{0j} + \theta_{ij}}{1 + \cos \theta_{0i} - \cos \theta_{0j} - \cos \theta_{ij}}.$$

Goemans and Williamson show the lower bound

$$\beta := \min_{0 \leq \theta < \arccos(-1/3)} \frac{2}{\pi} \frac{2\pi - 3\theta}{1 + 3 \cos \theta} > 0.79607$$

for it. Therefore, the above algorithm has performance ratio $\beta > 0.79607$.

The Feige-Goemans approximation algorithm. Feige and Goemans [77] propose an improved approximation algorithm for the maximum dicut problem analogue to their improved approximation algorithm for MAX 2SAT. Namely, strengthen the semidefinite program (SDP) by adding to it the triangle inequalities (114); replace the vectors v_0, \dots, v_n obtained as optimum solution of the strengthened SDP program by a new set of vectors v'_0, \dots, v'_n obtained by applying some rotation function to the v_i 's; generate from the v'_i 's the directed cut $\delta^+(S)$ where $S := \{i \in \{1, \dots, n\} \mid \text{sign}(v'_0 \cdot r) = \text{sign}(v'_i \cdot r)\}$. Thus one should now find lower bounds for the quantity

$$\frac{2}{\pi} \frac{-\theta'_{0i} + \theta'_{0j} + \theta'_{ij}}{1 + \cos \theta_{0i} - \cos \theta_{0j} - \cos \theta_{ij}}.$$

Using the rotation function f_λ from (116) with $\lambda = \frac{1}{2}$, Feige and Goemans claim a performance ratio of 0.857. Zwick [241] makes a detailed analysis of their algorithm enabling him to show a performance ratio of 0.859643 (using an adequate rotation function).

The Matuura-Matsui 0.863-approximation algorithm. Matuura and Matsui [171] propose an approximation algorithm for the maximum directed cut problem with performance ratio 0.863. Analogously to their algorithm for MAX 2SAT presented in the previous subsection, it relies on solving the semidefinite relaxation strengthened by the triangle inequalities (114) and applying the random hyperplane rounding phase using a distribution on the sphere which is skewed towards v_0 and uniform in any direction orthogonal to v_0 . As concrete choice, they propose to use the distribution function on S^2

$$g(v) = \cos^{1/1.8}(\theta) \quad \text{for all } v \in S^2 \text{ with } |v_0^T v| = \cos \theta \quad (121)$$

which can be realised as projection of a distribution on S^n and permits to show an approximation ratio of 0.863. (Compare (121) with the function g from (117) used for MAX 2SAT.)

The Lewin-Livnat-Zwick 0.874-approximation algorithm. Analogously to their improved algorithm for MAX 2SAT, Lewin, Livnat and Zwick [157] achieve this improved performance guarantee by combining the ideas of first suitably rotating the vectors obtained as solutions of the semidefinite program and of then using a skewed distribution function for choosing the random hyperplane.

7 Further Topics

7.1 Approximating polynomial programming using semidefinite programming

We come back in this section to the problem of approximating polynomial programs using semidefinite programming, which was already considered in Section 3.8. We present here the main ideas underlying this approach. They use results about representations of positive polynomials as sums of squares and moment sequences. Sums of squares will again be used in the next subsection for approximating the copositive cone. We then mention briefly some extensions to the general problem of testing whether a semi-algebraic set is empty.

Polynomial programs, sums of squares of polynomials, and moment sequences. Consider the following *polynomial programming problem*:

$$\min g(x) \quad \text{subject to} \quad g_\ell(x) \geq 0 \quad (\ell = 1, \dots, m) \quad (122)$$

where g, g_ℓ are polynomials in $x = (x_1, \dots, x_n)$. This is a very general problem which contains linear programming (when all polynomials have degree one) and 0/1 linear programming (since the integrality condition $x_i \in \{0, 1\}$ can be expressed as the polynomial equation: $x_i^2 - x_i = 0$). We mentioned in Section 3.8 that, under some technical assumption, the problem (122) can be approximated (getting arbitrarily close to its optimum) by the sequence of semidefinite programs (56). This result, due to Lasserre [141], relies on the fact that certain positive polynomials can be represented as sums of squares of polynomials. This idea of using sums of squares of polynomials for approximating polynomial programs has been introduced by Shor [217, 218, 219] and used by several other authors including Nesterov [182] and Parrilo [187, 188]; it seems to yield a more powerful method than other existing algebraic methods (see [189] for a comparison).

We would like to explain briefly here the main ideas underlying this approach. For simplicity, consider first the unconstrained problem:

$$p^* := \min g(x) \quad \text{subject to} \quad x \in \mathbb{R}^n \quad (123)$$

where $g(x) = \sum_{\alpha \in S_{2d}} g_\alpha x^\alpha$ is a polynomial of even degree $2d$; here S_k denotes the set of sequences $\alpha \in \mathbb{Z}_+^n$ with $|\alpha| := \sum_{i=1}^n \alpha_i \leq k$ for any integer k . One can assume w.l.o.g. that $g(0) = g_0 = 0$. In what follows the polynomial $g(x)$ is identified with its sequence of coefficients $g = (g_\alpha)_{\alpha \in S_{2d}}$. Obviously, (123) can be rewritten as

$$p^* = \max \lambda \quad \text{subject to} \quad g(x) - \lambda \geq 0 \quad \forall x \in \mathbb{R}^n. \quad (124)$$

Testing whether a polynomial is nonnegative is a hard problem, since it contains the problem of testing whether a matrix is copositive (see the next subsection). Lower bounds for p^* can be obtained by considering sufficient conditions for the polynomial $g(x) - \lambda$ to be nonnegative on \mathbb{R}^n . An obvious such sufficient condition being that $g(x) - \lambda$ be a sum of squares of polynomials. Therefore,

$$p^* \geq \max \lambda \quad \text{subject to} \quad g(x) - \lambda \text{ is a sum of squares.} \quad (125)$$

Testing whether a polynomial $p(x)$ is a sum of squares of polynomials amounts to testing feasibility of a semidefinite program (cf., e.g., [198]). Indeed, say $p(x)$ has degree $2d$, and let $z := (x^\alpha)_{\alpha \in S_d}$ be the vector consisting of all monomials of degree $\leq d$. Then one can easily verify that $p(x)$ is a sum of

squares if and only if $p(x) = z^T X z$ (identical polynomials) for some positive semidefinite matrix X . For $\gamma \in S_{2d}$, set

$$B_\gamma := \sum_{\alpha, \beta \in S_d | \alpha + \beta = \gamma} E_{\alpha, \beta},$$

where $E_{\alpha, \beta}$ is the elementary matrix with all zero entries except entries 1 at positions (α, β) and (β, α) .

Proposition 32. *A polynomial $p(x)$ of degree $2d$ is a sum of squares of polynomials if and only if the following semidefinite program:*

$$X \succeq 0, \langle B_\gamma, X \rangle = p_\gamma \quad (\gamma \in S_{2d}) \quad (126)$$

is feasible, where X is of order $\binom{n+d}{d}$ and with $\binom{n+2d}{2d}$ equations.

PROOF. As $z^T X z = \sum_{\alpha, \beta \in S_d} X_{\alpha, \beta} x^{\alpha + \beta} = \sum_{\gamma \in S_{2d}} x^\gamma \left(\sum_{\substack{\alpha, \beta \in S_d \\ \alpha + \beta = \gamma}} X_{\alpha, \beta} \right) = \sum_{\gamma \in S_{2d}} x^\gamma \langle B_\gamma, X \rangle$, $p(x) = z^T X z$ for some $X \succeq 0$ (which is equivalent to $p(x)$ being a sum of squares) if and only if the system (126) is feasible. \blacksquare

Note that the program (126) has a polynomial size for fixed n or d . Based on the result from Proposition 32, one can reformulate the lower bound for p^* from (125) as

$$p^* \geq \max_{\text{s.t. } g(x) - \lambda \text{ is a sum of squares}} \lambda = \max_{\text{s.t. } \langle B_\gamma, X \rangle = g_\gamma \quad (\gamma \in S_{2d} \setminus \{0\})} -\langle B_0, X \rangle \quad (127)$$

One can alternatively proceed in the following way for finding lower bounds for p^* . Obviously,

$$p^* = \min_{\mu} \int g(x) d\mu(x) \quad (128)$$

where the minimum is taken over all probability measures μ on \mathbb{R}^n . Define a sequence $y = (y_\alpha)_{\alpha \in S_{2d}}$ to be a *moment sequence* if $y_\alpha = \int x^\alpha d\mu(x)$ ($\alpha \in S_{2d}$) for some nonnegative measure μ on \mathbb{R}^n . Hence, (128) can be rewritten as

$$p^* = \min_{\alpha} \sum_{\alpha} g_\alpha y_\alpha \quad \text{s.t. } y \text{ is a moment sequence and } y_0 = 1. \quad (129)$$

Lower bounds for p^* can be obtained by replacing the condition that y be a moment sequence by a necessary condition for it. An obvious such necessary condition is that the moment matrix $M_d^{\mathbb{Z}}(y) = (y_{\alpha + \beta})_{\alpha, \beta \in S_d}$ (recall (54)) be positive semidefinite. Thus we find the following lower bound for p^* :

$$p^* \geq \min g^T y \quad \text{subject to } M_d^{\mathbb{Z}}(y) \succeq 0 \text{ and } y_0 = 1. \quad (130)$$

Note that the constraint in (130) is precisely condition (56) (when there are no constraints $g_\ell(x) \geq 0$). Since $M_d^{\mathbb{Z}}(y) = B_0 y_0 + \sum_{\gamma \in S_{2d} \setminus \{0\}} B_\gamma y_\gamma$, the semidefinite programs in (130) and in (127) are in fact dual of each other, which reflects the duality existing between the theories of nonnegative polynomials and of moment sequences.

The lower bound from (127) is equal to p^* if $g(x) - p^*$ is a sum of squares; this holds for $n = 1$ but not in general if $n \geq 2$. In general one can estimate p^* asymptotically by a sequence of SDP's

analogue to (127) if one assumes that an upper bound R is known a priori on the norm of a global minimizer x of $g(x)$, in which case

$$p^* = \min g(x) \text{ subject to } g_1(x) := R - \sum_{i=1}^n x_i^2 \geq 0.$$

Indeed, one can then use a result of Putinar [199] (quoted in Theorem 33 below) and conclude that, for any $\epsilon > 0$, the polynomial $g(x) - p^* + \epsilon$ is *positive* on $F := \{x \mid g_1(x) \geq 0\}$ and thus can be decomposed as $p(x) + p_1(x)g_1(x)$ for some polynomials $p(x)$ and $p_1(x)$ that are sums of squares. Testing for the existence of such decomposition can be expressed as a SDP program analogue to (127). Its dual (analogue to (130)) reads:

$$p_t^* := \min g^T y \text{ subject to } M_t(y) \geq 0, M_{t-1}(g_1 * y) \succeq 0, y_0 = 1.$$

Putinar's result permits to show the asymptotic convergence of p_t^* to p^* when t goes to infinity.

Theorem 33. [199] *Let g_1, \dots, g_m be polynomials and set $F := \{x \in \mathbb{R}^n \mid g_1(x) \geq 0, \dots, g_m(x) \geq 0\}$. Assume that F is compact and that there exists a polynomial u satisfying (i) the set $\{x \in \mathbb{R}^n \mid u(x) \geq 0\}$ is compact and (ii) u can be decomposed as $u_0 + \sum_{\ell=1}^m u_\ell g_\ell$ for some polynomials u_0, \dots, u_m that are sums of squares. Then every polynomial $p(x)$ which is positive on F can be decomposed as $p = p_0 + \sum_{\ell=1}^m p_\ell g_\ell$ for some polynomials p_0, \dots, p_m that are sums of squares.*

The above reasoning extends to the general program (122) if the assumption of Theorem 33 holds. This is the case, e.g., if the set $\{x \mid g_\ell(x) \geq 0\}$ is compact for one of the polynomials defining F . Then, Putinar's result permits to claim that, for any $\epsilon > 0$, the polynomial $g(x) - p^* + \epsilon$ can be decomposed as $p(x) + \sum_{\ell=1}^m p_\ell(x)g_\ell(x)$ for some polynomials $p(x)$, $p_\ell(x)$ that are sums of squares. Based on this, one can derive the asymptotic convergence to p^* of the minimum of $g^T y$ taken over all y satisfying (56) when t goes to ∞ . In the 0/1 case, when the constraints $x_i^2 - x_i = 0$ ($i = 1, \dots, n$) are part of the system defining F , there is in fact finite convergence in n steps [142] (see Section 3).

Semidefinite programming and the Positivstellensatz. Consider the following system:

$$\begin{aligned} f_j(x) &\geq 0 \quad (j = 1, \dots, s) \\ g_k(x) &\neq 0 \quad (k = 1, \dots, t) \\ h_\ell(x) &= 0 \quad (\ell = 1, \dots, u) \end{aligned} \tag{131}$$

where all f_j , g_k , h_ℓ are polynomials in the real variable $x = (x_1, \dots, x_n)$. The complexity of the problem of testing feasibility of this system has been the object of intensive research. Tarski [1951] showed that this problem is decidable and since then a number of other algorithms have been proposed, in particular, by Renegar [206] and Basu et al. [34].

We saw in Proposition 32 that testing whether a polynomial is a sum of squares can be formulated as a semidefinite program. Parrilo [187] showed that the general problem of testing infeasibility of the system (131) can also be formulated as a semidefinite programming problem (of very large size). This is based on the following result of real algebraic geometry, known as the 'Positivstellensatz'. The Positivstellensatz asserts that for a system of polynomial (in)equalities, either there is a solution in \mathbb{R}^n , or there is a polynomial identity giving a certificate that no real solution exists. This gives therefore a common generalization of Hilbert's 'Nullstellensatz' (in the complex case) and Farkas' lemma (for linear systems).

Theorem 34. ([222], [41]) *The system (131) is infeasible if and only if there exist polynomials f, g, h of the form*

$$\begin{aligned} f(x) &= \sum_{S \subseteq \{1, \dots, s\}} p_S \left(\prod_{j \in S} f_j \right) \quad \text{where all } p_S \text{ are sums of squares} \\ g(x) &= \prod_{k \in K} g_k \quad \text{where } K \subseteq \{1, \dots, t\} \\ h(x) &= \sum_{\ell=1}^u q_\ell h_\ell \quad \text{where all } q_\ell \text{ are polynomials} \end{aligned}$$

satisfying the equality $f + g^2 + h = 0$.

Bounds are known a priori for the degrees of the polynomials in the Positivstellensatz which make it possible to test infeasibility of the system (131) via semidefinite programming. However, these bounds are very large (triply exponential in n). Practically, one can use semidefinite programming for searching for infeasibility certificates of bounded degree.

7.2 Approximating combinatorial problems using copositive programming

We have seen throughout this chapter how semidefinite programming can be used for approximating combinatorial optimization problems. The idea of using the copositive cone and its dual, the cone of completely positive matrices, instead of the positive semidefinite cone has also been considered; cf., e.g., [43], [200]. We present below some results of de Klerk and Pasechnik [136] showing how the stability number of a graph can be computed using copositive relaxations.

Let us first recall some definitions. A symmetric matrix M of order n is *copositive* if $x^T M x \geq 0$ for all $x \in \mathbb{R}_+^n$ and M is *completely positive* if $M = \sum_{i=1}^k u_i u_i^T$ for some nonnegative vectors u_1, \dots, u_k . Let \mathcal{C}_n denote the set of symmetric copositive matrices of order n ; its dual cone \mathcal{C}_n^* is the set of completely positive matrices. Hence,

$$\mathcal{C}_n^* \subseteq \text{PSD}_n = \text{PSD}_n^* \subseteq \mathcal{C}_n.$$

Testing whether a matrix M is copositive is a co-NP-complete problem [178].

Let $G = (V, E)$ ($V = \{1, \dots, n\}$) be a graph and consider its theta number $\vartheta(G)$, defined by

$$\vartheta(G) = \max \langle J, X \rangle \text{ s.t. } X_{ij} = 0 \ (ij \in E), \text{Tr}(X) = 1, X \succeq 0 \quad (132)$$

(same as definition (58)). Then, $\vartheta(G)$ is an upper bound for the stability number of G , since for any stable set S in G , the matrix $X_S := \frac{1}{|S|} \chi^S (\chi^S)^T$ is feasible for the semidefinite program (132). Note that X_S is in fact completely positive. Therefore, one can define a tighter upper bound for $\alpha(G)$ by replacing in (132) the condition $X \succeq 0$ by the condition $X \in \mathcal{C}_n^*$. Letting A denote the adjacency matrix of G , we obtain:

$$\begin{aligned} \alpha(G) \leq \max \quad & \langle J, X \rangle & \leq \min \quad & \lambda \\ \text{s.t.} \quad & \text{Tr} X = 1 & \text{s.t.} \quad & \lambda I + yA - J \in \mathcal{C}_n \\ & X_{ij} = 0 \ (ij \in E) & & \lambda, y \in \mathbb{R} \\ & X \in \mathcal{C}_n^* & & \end{aligned} \quad (133)$$

where the right most program is obtained from the left most one using cone-LP duality. Using the following formulation for $\alpha(G)$ due to Motzkin and Straus [177]:

$$\frac{1}{\alpha(G)} = \min x^T(A + I)x \quad \text{subject to } x \geq 0 \text{ and } \sum_{i=1}^n x_i = 1,$$

one finds that the matrix $\alpha(G)(I + A) - J$ is copositive. This implies that the optimum value of the right most program in (133) is at most $\alpha(G)$. Therefore, equality holds throughout in (133). This shows again that copositive programming is not tractable.

Parrilo [187] proposes to approximate the copositive cone using sums of squares of polynomials. For this, note that a matrix M is copositive if and only if the polynomial

$$g_M(x) := \sum_{i,j=1}^n M_{ij}x_i^2x_j^2$$

is nonnegative on \mathbb{R}^n . Therefore, an obvious sufficient condition for M to be copositive is that $g_M(x)$ be a sum of squares or, more generally, that the polynomial $g_M(x) \left(\sum_{i=1}^n x_i^2 \right)^r$ be a sum of squares for some integer $r \geq 0$. A theorem of Polya asserts that, conversely, if M is strictly copositive (i.e., $x^T M x > 0$ for all $x \in \mathbb{R}_+^n \setminus \{0\}$), then $g_M(x) \left(\sum_{i=1}^n x_i^2 \right)^r$ is a sum of squares for some r . Powers and Reznick [197] give some upper bound for this integer r (depending only on M).

Let \mathcal{K}_n^r denote the set of symmetric matrices M of order n for which $g_M(x) \left(\sum_{i=1}^n x_i^2 \right)^r$ is a sum of squares. Thus

$$\text{PSD}_n \subseteq \mathcal{K}_n^0 \subseteq \dots \subseteq \mathcal{K}_n^r \subseteq \mathcal{C}_n.$$

We saw in the preceding subsection that testing whether a polynomial is a sum of squares can be solved via the semidefinite program (126). Therefore one can test membership in \mathcal{K}_n^r via semidefinite programming. For instance, Parrilo [187] shows that

$$M \in \mathcal{K}_n^0 \iff M = P + N \text{ for some } P \succeq 0, N \geq 0.$$

Moreover, $M \in \mathcal{K}_n^1$ if and only if the following system:

$$\begin{aligned} M - X^{(i)} &\succeq 0 & (i = 1, \dots, n) \\ X_{ii}^{(i)} &= 0 & (i = 1, \dots, n) \\ X_{ii}^{(j)} + 2X_{ij}^{(i)} &= 0 & (i \neq j = 1, \dots, n) \\ X_{jk}^{(i)} + X_{ik}^{(j)} + X_{ij}^{(k)} &\geq 0 & (1 \leq i < j < k \leq n) \end{aligned}$$

has a solution, where $X^{(1)}, \dots, X^{(n)}$ are symmetric $n \times n$ matrices ([187] and [44]).

Replacing in (133) the condition $\lambda I + yA - J \in \mathcal{C}_n$ by the condition $\lambda I + yA - J \in \mathcal{K}_n^r$, one can define the parameter

$$\vartheta^r(G) := \min \lambda \quad \text{subject to } \lambda I + yA - J \in \mathcal{K}_n^r.$$

Using the bound of Powers and Reznick [197], de Klerk and Pasechnik [136] show that

$$\alpha(G) = \lfloor \vartheta^r(G) \rfloor \quad \text{if } r \geq \alpha^2(G).$$

The same conclusion holds if we replace \mathcal{K}_n^r by the cone \mathcal{C}_n^r consisting of the matrices M for which $g_M(x) \left(\sum_{i=1}^n x_i^2 \right)^r$ has only nonnegative coefficients. Bomze and de Klerk [44] characterized the cone \mathcal{C}_n^r as the set of matrices M satisfying: $x^T M x - x^T \text{diag}(M) \geq 0$ for all $x \in \mathbb{Z}_+^n$ with $\sum_{i=1}^n x_i = r + 2$. It is also shown in [136] that $\vartheta^0(G) = \vartheta'(G)$, the Schrijver parameter from (65); $\vartheta^1(G) = \alpha(G)$ if G is an odd circuit, an odd wheel or their complement, or if $\alpha(G) = 2$. It is conjectured in [136] that $\vartheta^{\alpha(G)-1}(G) = \alpha(G)$.

Bomze and de Klerk [44] extend these ideas to standard quadratic optimization problems, of the form:

$$p^* := \min x^T Q x \quad \text{s.t. } x \in \Delta := \{x \in \mathbb{R}_+^n \mid e^T x = 1\} \quad (134)$$

where Q is a symmetric matrix. Problem (134) is equivalent to any of the following dual problems:

$$\begin{aligned} p^* &= \min \langle Q, X \rangle \quad \text{s.t. } \langle J, X \rangle = 1, X \in \mathcal{C}_n^* \\ &= \max \lambda \quad \text{s.t. } Q - \lambda J \in \mathcal{C}_n, \lambda \in \mathbb{R}. \end{aligned} \quad (135)$$

If we replace in (135) the cone \mathcal{C}_n by its approximation by \mathcal{C}_n^r (defined above), we obtain a lower bound p^r for p^* . Bomze and de Klerk [44] prove the following inequality for the quality of this approximation:

$$\frac{p^* - p^r}{Q_{max} - Q_{min}} \leq \frac{1}{r + 1}.$$

(Here, $Q_{max} = \max Q_{ij}$, $Q_{min} = \min Q_{ij}$, and Q, J are assumed to be linearly independent.) Bomze and de Klerk also show that p^r can be equivalently obtained as optimum value over a grid approximation of the simplex Δ ; namely,

$$p^r = \frac{r + 2}{r + 1} \min \left\{ y^T Q y - \frac{1}{r + 2} y^T \text{diag}(Q) \right\} \quad \text{s.t. } y \in \Delta, (r + 2)y \in \mathbb{Z}_+^n.$$

8 Semidefinite Programming and the Quadratic Assignment Problem

Quadratic problems in binary variables are the prime source for semidefinite models in combinatorial optimization. The simplest form, unconstrained quadratic programming in binary variables, corresponds to Max-Cut, and was described in detail in Section 5.

Assuming that the binary variables are the elements of a permutation matrix leads to the *Quadratic Assignment Problem (QAP)*. Formally, QAP consists in minimizing

$$\text{Tr}(AXB + C)X^T \quad (136)$$

over all permutation matrices X . One usually assumes that A and B are symmetric matrices of order n , while the linear term C is an arbitrary matrix of order n . There are many applications of this model problem, for instance in location theory. We refer to the recent monograph [48] for a description of published applications of QAP in Operations Research and combinatorial optimization.

The cost function (136) is quadratic in the matrix variable X . To rewrite this we use the vec-operator and (9). This leads to

$$\text{Tr}AXBX^T = \langle \text{vec}(X), \text{vec}(AXB) \rangle = x^T(B \otimes A)x, \quad (137)$$

because B is assumed to be symmetric. We can therefore express QAP equivalently as

$$\min\{x^T(B \otimes A)x + c^T x : x = \text{vec}(X), X \text{ permutation matrix}\}.$$

Here, $c = \text{vec}(C)$. To derive semidefinite relaxations of QAP we follow the generic pattern and linearize by introducing a new matrix variable for xx^T , leading to the study of

$$P = \text{conv}\{xx^T : x = \text{vec}(X), X \text{ permutation matrix}\}.$$

In section 3, we observed that any $Y \in P$ must satisfy the semidefiniteness condition (20), which in our present notation amounts to

$$Z = \begin{pmatrix} 1 & z^T \\ z & Y \end{pmatrix} \succeq 0, \quad \text{diag}(Y) = z.$$

The first question is to identify the smallest subcone of semidefinite matrices that contains P .

We use the following parametrization of matrices having row and column sums equal to e , the vector of all ones, see [104].

Lemma 35. [104] *Let V be an $n \times (n - 1)$ matrix with $V^T e = 0$ and $\text{rank}(V) = n - 1$. Then*

$$\mathcal{E} := \{X \in \mathbb{R}^{n \times n} : Xe = X^T e = e\} = \left\{ \frac{1}{n}ee^T + VMV^T : M \in \mathbb{R}^{(n-1) \times (n-1)} \right\} =: \mathcal{E}'.$$

PROOF. Let $Z = \frac{1}{n}ee^T + VMV^T \in \mathcal{E}'$. Then $Ze = Z^T e = e$, because $V^T e = 0$, hence $Z \in \mathcal{E}$. To see the other inclusion, let $V = QR$ be the QR-decomposition of V , i.e. $Q^T Q = I$, $QQ^T = I - \frac{1}{n}ee^T$ and $\text{rank}(R) = n - 1$. Let $X \in \mathcal{E}$ and set $M := R^{-1}Q^T X Q (R^{-1})^T$. Then $\frac{1}{n}ee^T + VMV^T = X \in \mathcal{E}'$. ■

We use this parametrization and define

$$W := \left(\frac{1}{n}e \otimes e, V \otimes V \right).$$

V can be any basis of e^\perp , as in the previous lemma. We can now describe the smallest subcone containing P .

Lemma 36. *Let $Y \in P$. Then there exists a symmetric matrix R of order $(n-1)^2 + 1$, indexed from 0 to $(n-1)^2$, such that*

$$R \succeq 0, \quad r_{00} = 1, \quad Y = WRW^T.$$

PROOF. (See also [239].) We first look at the extreme points of P , so let X be a permutation matrix. Thus we can write X as $X = \frac{1}{n}ee^T + VMV^T$, for some matrix M . Let $m = \text{vec}(M)$. Then, using (9),

$$x = \text{vec}(X) = \frac{1}{n}e \otimes e + (V \otimes V)m = Wz,$$

with $z = \begin{pmatrix} 1 \\ m \end{pmatrix}$. Now $xx^T = Wzz^TW^T = WRW^T$, with $r_{00} = 1$, $R \succeq 0$. The same holds for convex combinations formed from several permutation matrices. ■

To see that the set

$$\hat{P} := \{Y : \exists R \text{ such that } Y = WRW^T, \quad z = \text{diag}(Y), \quad \begin{pmatrix} 1 & z^T \\ z & Y \end{pmatrix} \succeq 0\} \quad (138)$$

is indeed the smallest subcone of positive semidefinite matrices containing P , it is sufficient to provide a positive definite matrix \hat{R} , such that $W\hat{R}W^T \in P$. In [239] it is shown that

$$\hat{R} = \begin{pmatrix} 1 & 0 \\ 0 & \frac{1}{n^2(n-1)}(nI_{n-1} - E_{n-1}) \otimes (nI_{n-1} - E_{n-1}) \end{pmatrix} \succ 0$$

gives

$$W\hat{R}W^T = \frac{1}{n!} \sum_{X \in \Pi} (xx^T),$$

the barycenter of P . Here $V = \begin{pmatrix} I_{n-1} \\ -e_{n-1}^T \end{pmatrix}$ has to be used in the definition of W .

Eliminating Y leaves the matrix variable R and $n^2 + 1$ equality constraints, fixing the first row equal to the main diagonal, and setting the first element equal to 1.

Thus we arrive at the following basic SDP relaxation of QAP:

$$(QAP_{R1}) \quad \min \text{Tr}(B \otimes A + \text{Diag}(c))Y \text{ such that } Y = WRW^T \in \hat{P}, \quad r_{00} = 1. \quad (139)$$

It is instructive to look at $W\hat{R}W^T$ for small values of n . For $n = 3$ we get

$$W\hat{R}W^T = \frac{1}{6} \left(\begin{array}{ccc|ccc|ccc} 2 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 1 \\ 0 & 2 & 0 & 1 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 2 & 1 & 1 & 0 & 1 & 1 & 0 \\ \hline 0 & 1 & 1 & 2 & 0 & 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 & 2 & 0 & 1 & 0 & 1 \\ 1 & 1 & 0 & 0 & 0 & 2 & 1 & 1 & 0 \\ \hline 0 & 1 & 1 & 0 & 1 & 1 & 2 & 0 & 0 \\ 1 & 0 & 1 & 1 & 0 & 1 & 0 & 2 & 0 \\ 1 & 1 & 0 & 1 & 1 & 0 & 0 & 0 & 2 \end{array} \right).$$

The zero pattern in this matrix is not incidental. In fact, any $X \in P$ will have entries equal 0 at positions corresponding to $x_{ij}x_{ik}$ and $x_{ji}x_{ki}$ for $j \neq k$. This corresponds to the off-diagonal elements of the main diagonal blocks, and the main-diagonal elements of the off diagonal blocks. To express these constraints, we introduce some more notation, and index the elements of matrices in P alternatively by $P = (p_{(i,j),(k,l)})$ for i, j, k, l between 1 and n .

Hence we can strengthen the above relaxation by asking that

$$y_{rs} = 0 \text{ for } r = (i, j), s = (i, k), \text{ or } r = (j, i), s = (k, j), j \neq k.$$

We collect all these equations in the constraint $G(Y) = 0$. Adding it to (139) results in a stronger relaxation. In [239] this model is called the 'Gangster model'. Aside from $n^2 + 1$ equality constraints from the basic model, we have $O(n^3)$ equations in this extended model. This amounts to serious computational work, but results in a very strong lower bound for QAP.

$$(QAP_{R2}) \quad \min \text{Tr}(B \otimes A + \text{Diag}(c))Y \text{ such that } Y = WRW^T \in \hat{P}, r_{00} = 1, G(Y) = 0. \quad (140)$$

Finally, one can include the constraints $y_{rs} \geq 0$ for all r, s , leading to

$$(QAP_{R3}) \quad \min \text{Tr}(B \otimes A + \text{Diag}(c))Y \text{ such that } Y = WTRW^T \in \hat{P}, r_{00} = 1, G(Y) = 0, Y \geq 0. \quad (141)$$

The resulting SDP has $O(n^4)$ constraints and can not be solved in a straightforward way by interior point methods for problems of interesting size ($n \geq 15$).

The Anstreicher-Brixius bound. Anstreicher et al. [16, 17] have recently achieved a breakthrough in solving several instances of QAP which could not be solved by previous methods. The size of these instances ranges from $n = 20$ to $n = 36$. The key to this breakthrough lies in the use of a bound for QAP that is both 'fast' to compute, and gives 'good' approximations to the exact value of QAP. This bounding procedure combines orthogonal, semidefinite and convex quadratic relaxations in a nontrivial way, starting from the Hoffman-Wielandt inequality, Theorem 5.

A simple way to derive this bound goes as follows. We use the parametrization

$$X = \frac{1}{n}ee^T + VYV^T \quad (142)$$

from Lemma 35, and assume in addition that $V^T V = I_{n-1}$. Substituting this into the cost of function of QAP results in

$$\text{Tr}(AXB + C)X^T = \text{Tr}\hat{A}Y\hat{B}Y^T + \text{Tr}(\hat{C} + \frac{2}{n}V^T Aee^T BV)Y^T + \frac{1}{n^2}s(A)s(B) + \frac{1}{n}s(C), \quad (143)$$

where $\hat{A} = V^T AV, \hat{B} = V^T BV, \hat{C} = V^T CV$ and $s(M) := e^T M e = \sum_{ij} m_{ij}$. The condition $V^T V = I$ implies that X in (142) is orthogonal if and only if Y is. Hadley et al. [104] use this to bound the quadratic term in Y by the minimal scalar product of the eigenvalues of \hat{A} and \hat{B} , see Theorem 5. Anstreicher and Brixius [16] use this observation as a starting point and observe that for any symmetric matrix \hat{S} , and any orthogonal Y , one has

$$0 = \text{Tr}\hat{S}(I - YY^T) = \text{Tr}\hat{S} - \text{Tr}\hat{S}YIY^T = \text{Tr}\hat{S} - \text{Tr}(I \otimes \hat{S})(yy^T).$$

This results in the following identity, true for any orthogonal Y and any symmetric \hat{S}, \hat{T} :

$$\text{Tr}\hat{A}Y\hat{B}Y^T = \text{Tr}(\hat{S} + \hat{T}) + \text{Tr}(\hat{B} \otimes \hat{A} - I \otimes \hat{S} - \hat{T} \otimes I)(yy^T). \quad (144)$$

We use $\hat{Q} = \hat{B} \otimes \hat{A} - I \otimes \hat{S} - \hat{T} \otimes I, \hat{D} = \hat{C} + \frac{2}{n}V^T A e e^T B V$ and substitute this into (143) to get

$$\text{Tr}(AXB + C)X^T = \text{Tr}(\hat{S} + \hat{T}) + y^T \hat{Q}y + \hat{d}^T y + \frac{1}{n^2}s(A)s(B) + \frac{1}{n}s(C), \quad (145)$$

This relation is true for any orthogonal X and Y related by (142) and symmetric \hat{S}, \hat{T} . It is useful to express the parts in (145) containing Y by the original matrix X . To do this we use the following identity:

$$0 = \text{Tr}\hat{S}(I - V^T V) = \text{Tr}\hat{S}(I - V^T X X^T V) = \text{Tr}\hat{S} - \text{Tr}(V\hat{S}V^T)XIX^T = \text{Tr}\hat{S} - \text{Tr}(I \otimes V\hat{S}V^T)(xx^T).$$

Hence, for any orthogonal X , and any symmetric \hat{S}, \hat{T} we also have

$$\text{Tr}(AXB + C)X^T = \text{Tr}(\hat{S} + \hat{T}) + x^T Qx + c^T x. \quad (146)$$

Here $Q = B \otimes A - I \otimes (V\hat{S}V^T) - (V\hat{T}V^T) \otimes I$. Comparing (145) and (146) we note that

$$y^T \hat{Q}y + \hat{d}^T y + \frac{1}{n^2}s(A)s(B) + \frac{1}{n}s(C) = x^T Qx + c^T x.$$

It should be observed that Q and \hat{Q} above depend on the specific choice of \hat{S}, \hat{T} . Anstreicher and Brixius use the optimal solution \hat{S}, \hat{T} from Theorem 6 and observe that dual feasibility yields $\hat{Q} \succeq 0$. Therefore the above problem is a convex quadratic programming problem. We denote its optimal solution as the Anstreicher-Brixius bound $ABB(A, B, C)$.

$$ABB(A, B, C) := \text{Tr}(\hat{S} + \hat{T}) + \min\{x^T Qx + c^T x : x = \text{vec}(X), X \text{ doubly stochastic}\}.$$

The interesting observation here is that \hat{S}, \hat{T} are obtained as a by-product of the Hoffman-Wielandt inequality, and that the resulting matrix Q is positive semidefinite over the set of doubly stochastic matrices (as a consequence of Theorem 6). These facts imply that the Anstreicher-Brixius bound is tractable.

To give a flavour of the quality of these bounds, we provide the following computational results on standard test sets from Nugent et al. [184]. These data sets have the following characteristics. The linear term C is equal to 0. The matrix B represents the rectilinear cell distance of a rectangular array of cells, hence there is some symmetry in these data. In case of $n = 12$, the resulting rectangular cell array has the following form:

1	2	3	4
5	6	7	8
9	10	11	12

We observe that the distance matrix B would not change, if the following cell array would have been used:

4	3	2	1
8	7	6	5
12	11	10	9

Mathematically speaking, there exist several permutation matrices X , such that $B = XBX^T$. Exploiting all these symmetries, it is sufficient to consider only the subproblems where the cells 1, 2, 5, 6 are assigned to some fixed location, say 1. All other permutations can be obtained by exploiting the automorphisms inherent in B .

We denote these subproblems by nug12.1, nug12.2, nug12.5, nug12.6 in Table 1. The instance $n = 15$ has a distance matrix B corresponding to a 5×3 rectangular grid, leading to subproblems nug15.1, nug15.2, nug15.3, nug15.6, nug15.7, nug15.8. The optimal values for these instances are contained in the column labeled 'exact'. These values can be computed routinely for $n \approx 15$. The biggest instance $n = 30$ was only recently solved to optimality, see [17]. The computational results for QAP_{R3} are from the forthcoming dissertation [221]. It is computationally infeasible to solve this relaxation by interior points. Sotirov [221] uses the bundle method to get approximate solutions of QAP_{R3} . Hence the values are only lower estimates of the true bound. The values of QAP_{R2} were obtained by Sotirov and Wolkowicz⁵ by making use of the NEOS distributed computing system. The bounds are obtained using interior point methods. The computational effort to get these values is prohibitively big. A more practical approach consists in using bundle methods to bargain computational efficiency against a slight decrease in the quality of the bound. Finally, the values of the Anstreicher-Brixius bound ABB are from [16].

These results indicate that the SDP models in combination with bundle methods may open the way to improved Branch and Bound approaches to solve larger QAP instances.

9 Epilogue: Semidefinite Programming and Algebraic Connectivity

An implicit message of all the preceding sections is that semidefinite programming relaxations have a high potential to significantly improve on purely polyhedral relaxations. This may give the wrong impression that semidefinite programming is a universal remedy to improve upon linear relaxations. This is in principle true, if we assume that some sort of semidefiniteness constraint is *added* to the polyhedral model.

If a model based on semidefinite programming is used *instead* of a linear model, it need not be true that the semidefinite model dominates the linear one. We conclude with an illustration of this perhaps not quite intuitive statement.

We consider the Traveling Salesman Problem (TSP), i.e. the problem of finding a shortest Hamiltonian cycle in an edge weighted graph. This problem is well known to be NP-hard, and has stimulated research since the late 1950's.

We need to recall some notation from graph theory. For an edge weighted graph, given by its weighted adjacency matrix X , with $X \geq 0$, $\text{diag}(X) = 0$ (setting to 0 the entries corresponding to nonedges), we consider vertex partitions $(S, V \setminus S)$ of its vertex set V and define

$$X(S, V \setminus S) := \sum_{i \in S, j \notin S} x_{ij}$$

⁵personal communication, 2001

problem	exact	QAP_{R2}	QAP_{R3}	ABB
nug12	578	529.3	552.1	482
nug12.1	586	550.7	573.6	-
nug12.2	586	550.6	571.3	-
nug12.5	578	551.8	572.2	-
nug12.6	600	555.8	578.8	-
nug15	1150	1070.5	1106.1	996
nug15.1	1150	1103.4	1131.6	-
nug15.2	1168	1116.3	1147.8	-
nug15.3	1164	1120.9	1148.4	-
nug15.6	1166	1113.6	1144.9	-
nug15.7	1182	1130.3	1161.9	-
nug15.8	1184	1134.1	1162.2	-
nug20	2570	2385.6	2441.9	2254
nug30	6124	5695.4	5803.2	5365

Table 1: Semidefinite relaxations and optimal value for some instances from the Nugent collection of test data. The column labeled QAP_{R3} gives lower estimates of the bound computed by the bundle method.

to be the weight of the cut, given by S . The *edge connectivity* $\mu(X)$ of X is defined as

$$\mu(X) := \min\{X(S, V \setminus S) : S \subseteq V, 1 \leq |S| \leq |V| - 1\}.$$

The polyhedral approach to TSP is based on approximating the convex hull of all Hamiltonian cycles by considering all two-edge connected graphs. Formally, this amounts to optimizing over the following set:

$$\{X : 0 \leq x_{ij} \leq 1, \text{diag}(X) = 0, Xe = 2e, \mu(X) = 2\}. \quad (147)$$

Even though there are $O(2^n)$ linear constraints defining this (polyhedral) set, it is possible to optimize over it in polynomial time, by using the ellipsoid method (because the separation problem amounts to a minimum capacity cut problem, which can thus be solved in polynomial time). It is also interesting to note that no combinatorial algorithm of provably polynomial running time exists for optimizing a linear function over this set.

Recently, Cvetcovic et al. [61] have proposed a model where 2-edge connectivity is replaced by the algebraic connectivity, leading to an SDP relaxation.

Fiedler [85] introduces the algebraic connectivity of a graph, given by its weighted adjacency matrix $X \geq 0, \text{diag}(X) = 0$, as follows. Let $L(X) := D - X$ be the Laplacian matrix corresponding to X , where $D := \text{Diag}(Xe)$, the diagonal matrix having the row sums of X on its main diagonal. Since $De = Xe$, it is clear that 0 is an eigenvalue of $L(X)$ corresponding to the eigenvector e . Moreover $X \geq 0$ implies by the Gersgorin disk theorem, that all eigenvalues of $L(X)$ are nonnegative, i.e., $L(X)$ is positive semidefinite in this case. Fiedler observed that the second smallest eigenvalue $\lambda_2(L(X)) = \min_{\|u\|=1, u^T e=0} u^T L(X)u$ is equal to 0 if and only if X is the adjacency matrix of a disconnected graph, otherwise $\lambda_2(L(X)) > 0$. Note also that $\lambda_2(L(X))$ is concave in X . Fiedler therefore denotes $\alpha(X) := \lambda_2(L(X))$ as the algebraic connectivity of the graph, given by the adjacency

matrix X . It is not difficult to calculate $\alpha(C_n)$, the algebraic connectivity of a cycle on n nodes,

$$\alpha(C_n) = 2(1 - \cos(\frac{2\pi}{n})) =: h_n$$

The concavity of $\alpha(X)$ therefore implies that

$$\alpha(X) \geq h_n$$

for any convex combination X of Hamiltonian cycles. We also note that the Taylor expansion of $\cos(x)$ gives $h_n \leq \frac{4\pi^2}{n^2}$. Cvetcovic et al. [61] propose to replace the polyhedral constraints $\mu(X) \geq 2$ by the nonlinear condition $\alpha(X) \geq h_n$, which can easily be shown to be equivalent to the semidefiniteness constraint

$$L(X) + ee^T - h_n I \succeq 0$$

on X . Replacing edge connectivity by algebraic connectivity in (147) leads to optimizing over

$$\{X : 0 \leq x_{ij} \leq 1, \text{diag}(X) = 0, Xe = 2e, L(X) + ee^T - h_n I \succeq 0\}. \quad (148)$$

This looks like a reasonable bargain, as we replace $O(2^n)$ linear constraints by a single semidefiniteness constraint. The crucial question of course is whether we can say anything about the relative strength of the two relaxations. Since $L(X) + ee^T \succeq 0$ it is clear that

$$\lambda_{\min}(L(X) + ee^T - h_n I) \geq -h_n \geq -\frac{4\pi^2}{n^2}.$$

Therefore the semidefiniteness constraint in (148) is nearly satisfied for any $X \geq 0$ as the dimension increases. We can say even more. Any matrix X feasible for (147) satisfies $\alpha(X) \geq h_n$, see [84] and the handbook [232], chapter 12 for further details. In other words, the simple semidefinite relaxation given by (148) is dominated by the polyhedral edge connectivity model (147).

10 Appendix: Surveys, Books and Software

Semidefinite Programming has undergone a rapid development in the last decade. We close with some practical information on semidefinite programming in connection with recent books, surveys, software and web-sites. The references given here are by no means complete. We apologize for any possible omissions.

Books and Survey papers: The proceedings volume [186] presents one of the first collection of papers devoted to semidefinite programming in connection with combinatorial optimization. The handbook [232] is currently a prime source for nearly all aspects of semidefinite optimization. It contains contributions from leading experts in the field, covering in 20 chapters algorithms, theory and applications. With nearly 900 references, it also reflects the state of the art up to about the year 1999. We also refer to [133] for a recent monograph on semidefinite programming, featuring also the development up to 2002.

The survey paper [229] has set the stage for many algorithmic and theoretical developments, that were to follow in the last few years. The surveys given by Lovász [166] and Goemans [91] focus on the interplay between semidefinite programming and NP-hard combinatorial optimization problems. We also refer to [205] and [226] for surveys focusing on algorithmic aspects and also the position of semidefinite programming in the context of general convex programming.

Software: The algorithmic machinery to solve semidefinite programs is rather sophisticated. It is therefore highly appreciated that many researchers offer their software to the scientific community for free use. The following two packages are currently considered state-of-the-art to deal with general semidefinite problems.

SEDUMI: <http://fewcal.kub.nl/software/sedumi.html>

SDPT3: <http://www.math.nus.edu.sg/mathtohkc/sdpt3.html>

Both packages use Matlab as the working horse and implement interior-point methods. The following package is written in C, and contains also specially tailored subroutines to compute the ϑ function.

CSDP: <http://www.nmt.edu/~borchers/csdp.html>

For large-scale problems, where interior-point methods are out of reach, the spectral bundle approach may be a possible alternative:

SBMethod: <http://www-user.tu-chemnitz.de/~helmberg/SBMethod.html>

Web-sites: Finally, we refer to the following two web-sites, which have been maintained over a long period of time, so we expect them to survive also in the future.

The Optimization-online web-site maintains an electronic library of technical reports in the field of optimization. A prominent part covers semidefinite programming and combinatorial optimization.

<http://www.optimization-online.org>

The semidefinite programming web-site maintained by C. Helmberg contains up-to-date information on various activities related to semidefinite programming (conferences, workshops, publications, software, people working in the field, etc) and pointers to other web-sites.

<http://www-user.tu-chemnitz.de/~helmberg/semidef.html>

Acknowledgements. We thank a referee for his careful reading and his suggestions that helped improve the presentation of this chapter.

References

- [1] N.E. Aguilera, M.S. Escalante, and G.L. Nasini. The disjunctive procedure and blocker duality. Preprint, 2001.
- [2] N.E. Aguilera, M.S. Escalante, and G.L. Nasini. A generalization of the perfect graph theorem under the disjunctive index. *Mathematics of Operations Research*, 27:460–469, 2002.
- [3] N.E. Aguilera, S.M. Bianchi, and G.L. Nasini. Lift and project relaxations for the matching polytope and related polytopes. Preprint, 2002.
- [4] A. Alfakih. Graph rigidity via Euclidean distance matrices. *Linear Algebra and its Applications*, 310:149–165, 2000.
- [5] A. Alfakih. On rigidity and realizability of weighted graphs. *Linear Algebra and its Applications*, 325:57–70, 2001.
- [6] A. Alfakih and H. Wolkowicz. On the embeddability of weighted graphs in Euclidean spaces. Technical Report, CORR 98-12, Department of Combinatorics and Optimization, University of Waterloo, 1998. Available at <http://orion.math.uwaterloo.ca/~hwolkowi/>
- [7] A. Alfakih, A. Khandani, and H. Wolkowicz. Solving Euclidean distance matrix completion problems via semidefinite programming. *Computational Optimization and Applications*, 12:13–30, 1999.
- [8] F. Alizadeh. Interior point methods in semidefinite programming with applications in combinatorial optimization. *SIAM Journal on Optimization*, 5:13–51, 1995.
- [9] N. Alon and N. Kahale. Approximating the independence number via the ϑ -function. *Mathematical Programming*, 80:253–264, 1998.
- [10] N. Alon and B. Sudakov. Bipartite subgraphs and the smallest eigenvalue. *Combinatorics, Probability and Computing*, 9:1–12, 2000.
- [11] N. Alon, B. Sudakov and U. Zwick. Constructing worst case instances for semidefinite programming based approximation algorithms. *SIAM Journal on Discrete Mathematics*, 15:58–72, 2002. [Preliminary version in *Proceedings of 12th ACM-SIAM Symposium on Discrete Algorithms*, pages 92–100, 2001.]
- [12] M.F. Anjos. *New Convex Relaxations for the Maximum Cut and VLSI Layout Problems*. PhD thesis, University of Waterloo, 2001.
- [13] M. Anjos. An improved semidefinite programming relaxation for the satisfiability problem. Technical Report UW-E&CE 2002-09, University of Waterloo, Canada, 2002.
- [14] M.F. Anjos and H. Wolkowicz. A strengthened SDP relaxation via a second lifting for the Max-Cut problem. *Discrete Applied Mathematics*, 119:79–106, 2002.

- [15] M.F. Anjos and H. Wolkowicz. Geometry of semidefinite Max-Cut relaxations via ranks. *Journal of Combinatorial Optimization*, 6:237–270, 2002.
- [16] K. Anstreicher, and N. Brixius. A lower bound for the Quadratic Assignment Problem based on Convex Quadratic Programming. *Mathematical Programming*, 89:341–357, 2001.
- [17] K. Anstreicher, N. Brixius, J.-P. Goux and J. Linderoth. Solving large quadratic assignment problems on computational grids. *Mathematical Programming B*, 91:563–588, 2002.
- [18] K. Anstreicher and H. Wolkowicz. On Lagrangian relaxation of quadratic matrix constraints. *SIAM Journal on Matrix Analysis and its Applications*, 22:41–55, 2000.
- [19] S. Arora, B. Bollobás and L. Lovász. Proving integrality gaps without knowing the linear program. In *Proceedings of the 43rd IEEE Symposium on Foundations of Computer Science*, IEEE Computer Science Press, Los Alamitos, CA, 2002.
- [20] S. Arora, D. Karger and M. Karpinski. Polynomial time approximation schemes for dense instances of NP-hard problems. In *Proceedings of the 27th Annual ACM Symposium on Theory of Computing*, ACM, New York, pages 284–293, 1995.
- [21] S. Arora, C. Lund, R. Motwani, M. Sudan, and M. Szegedy. Proof verification and intractability of approximation problems. In *Proceedings of the 33rd IEEE Symposium on Foundations of Computer Science*, IEEE Computer Science Press, Los Alamitos, CA, pages 14–23, 1992.
- [22] T. Asano and D.P. Williamson. Improved approximation algorithms for MAX SAT. In *Proceedings of 11th ACM-SIAM Symposium on Discrete Algorithms*, pages 96–115, 2000.
- [23] E. Balas. Disjunctive programming. *Annals of Discrete Mathematics*, 5:3–51, 1979.
- [24] E. Balas, S. Ceria, and G. Cornuéjols. A lift-and-project cutting plane algorithm for mixed 0-1 programs. *Mathematical Programming*, 58:295–324, 1993.
- [25] M.O. Ball, W. Liu, and W.R. Pulleyblank. Two terminal Steiner tree polyhedra. In *Contributions to Operations research and Economics*, B. Tulkens and H. Tulkens, eds., MIT Press, Cambridge, MA, pp. 251–284, 1989.
- [26] F. Barahona. On cuts and matchings in planar graphs. *Mathematical Programming*, 60:53–68, 1993.
- [27] F. Barahona. On the computational complexity of Ising spin glass models. *Journal of Physics A, Mathematical and General*, 15:3241–3253, 1982.
- [28] F. Barahona. The max-cut problem on graphs not contractible to K_5 . *Operations Research Letters*, 2:107–111, 1983.
- [29] F. Barahona and A.R. Mahjoub. On the cut polytope. *Mathematical Programming*, 36:157–173, 1986.
- [30] F. Barahona and A.R. Mahjoub. Compositions of graphs and polyhedra. II: Stable sets. *SIAM Journal on Discrete Mathematics*, 7:359–371, 1994.

- [31] A.I. Barvinok. Feasibility testing for systems of real quadratic equations. *Discrete and Computational Geometry*, 10:1–13, 1993.
- [32] A.I. Barvinok. Problems of distance geometry and convex properties of quadratic maps. *Discrete and Computational Geometry*, 13:189–202, 1995.
- [33] A.I. Barvinok. A remark on the rank of positive semidefinite matrices subject to affine constraints. *Discrete and Computational Geometry*, 25:23–31, 2001.
- [34] S. Basu, R. Pollack and M.-F. Roy. On the combinatorial and algebraic complexity of quantifier elimination. *Journal of the Association for Computing Machinery*, 43:1002–1045, 1996.
- [35] M. Bellare and P. Rogaway. The complexity of approximating a nonlinear program. *Mathematical Programming*, 69:429–441, 1995.
- [36] C. Berge. Sur une conjecture relative au problème des codes optimaux. Communication, 13ème assemblée générale de l’URSI, Tokyo, 1962.
- [37] P. Berman and M. Karpinski. On some tighter inapproximability results, further improvements. *Electronic Colloquium on Computational Complexity*, Report TR98-065, 1998.
- [38] D. Bienstock and M. Zuckerberg. Subset algebra lift operators for 0 – 1 integer programming. CORC Report 2002-01, Columbia University, NY, 2002.
- [39] A. Blum. New approximation algorithms for graph coloring. *Journal of the Association for Computing Machinery*, 41:470–516, 1994. [Preliminary version in *Proceedings of the 21st Annual ACM Symposium on Theory of Computing*, ACM, New York, pages 535–542, 1989 and in *Proceedings of the 31st IEEE Symposium on Foundations of Computer Science*, IEEE Computer Science Press, Los Alamitos, CA, pages 554–562, 1990.]
- [40] A. Blum and D. Karger. An $\tilde{O}(n^{3/14})$ -coloring algorithm for 3-colorable graphs. *Information Processing Letters*, 61:49-53, 1997.
- [41] J. Bochnak, M. Coste and M.-F. Roy. *Géométrie Algébrique Réelle*. Springer-Verlag, 1987.
- [42] A. Bockmayr, F. Eisenbrand, M. Hartmann, and A.S. Schulz. On the Chvátal rank of polytopes in the 0/1 cube. *Discrete Applied Mathematics*, 98:21–27, 1999.
- [43] I.M. Bomze, M. Dür, E. de Klerk, C. Roos, A.J. Quist, and T. Terlaky. On copositive programming and standard quadratic optimization problems. *Journal of Global Optimization*, 18:301–320, 2000.
- [44] I.M. Bomze and E. de Klerk. Solving standard quadratic optimization problems via semidefinite and copositive programming. *Journal of Global Optimization*, 24:163–185, 2002.
- [45] J.M. Borwein and H. Wolkowicz. Regularizing the abstract convex program. *Journal of Mathematical Analysis and Applications*, 83:495–530, 1981.
- [46] J. Bourgain. On Lipschitz embedding of finite metric spaces in Hilbert space. *Israel Journal of Mathematics*, 52:46–52, 1985.

- [47] A. Caprara and A.N. Letchford. On the separation of split cuts and related inequalities. Preprint, 2000. [*Mathematical Programming*, to appear.] Available at <http://www.lancs.ac.uk/staff/letchfoa/home.htm>
- [48] E. Cella. *The Quadratic Assignment Problem: Theory and Algorithms*. Kluwer Academic Publishers, USA, 1998.
- [49] S. Ceria. *Lift-and-Project Methods for Mixed 0-1 Programs*. PhD dissertation, Graduate School of Industrial Administration, Carnegie Mellon University, US, 1993.
- [50] S. Ceria and G. Pataki. Solving integer and disjunctive programs by lift-and-project. In R.E. Bixby, E.A. Boyd, and R.Z. Rios-Mercato, eds., *IPCO VI, Lecture Notes in Computer Science*, 1412:271–283, 1998.
- [51] M. Charikar. On semidefinite programming relaxations for graph colouring and vertex cover. In *Proceedings of 13th ACM-SIAM Symposium on Discrete Algorithms*, pages 616–620, 2002.
- [52] V. Chvátal. Edmonds polytopes and a hierarchy of combinatorial problems. *Discrete Mathematics*, 4:305–337, 1973.
- [53] V. Chvátal. On certain polytopes associated with graphs. *Journal of Combinatorial Theory B*, 18:138–154, 1975.
- [54] V. Chvátal, W. Cook, and M. Hartman. On cutting-plane proofs in combinatorial optimization. *Linear Algebra and its Applications*, 114/115:455–499, 1989.
- [55] W. Cook and S. Dash. On the matrix-cut rank of polyhedra. *Mathematics of Operations Research*, 26:19–30, 2001.
- [56] W. Cook, R. Kannan, and A. Schrijver. Chvátal closures for mixed integer programming problems. *Mathematical Programming*, 47:155–174, 1990.
- [57] G. Cornuéjols and Y. Li. Elementary closures for integer programs. *Operations Research Letters*, 28:1–8, 2001.
- [58] G. Cornuéjols and Y. Li. On the rank of mixed 0-1 polyhedra. In K. Aardal and A.M.H. Gerards, eds., *IPCO 2001, Lecture Notes in Computer Science*, 2081:71–77, 2001.
- [59] G. Cornuéjols and Y. Li. A connection between cutting plane theory and the geometry of numbers. *Mathematical Programming A*, 93:123–127, 2002.
- [60] G.M. Crippen and T.F. Havel. *Distance Geometry and Molecular Conformation*. Research Studies Press, Taunton, Somerset, England, 1988.
- [61] D. Cvetković, M. Cangalvić, and V. Kovačević-Vujčić. Semidefinite Programming Methods for the Symmetric Traveling Salesman Problem, *Proceedings of the 7th International IPCO Conference, Graz, Austria*, pp. 126–136, 1999.
- [62] S. Dash. *On the Matrix Cuts of Lovász and Schrijver and their Use in Integer Programming*. PhD thesis, Rice University, 2001.

- [63] S. Dash. An exponential lower bound on the length of some classes of branch-and-cut proofs. In W.J. Cook and A.S. Schulz, eds., IPCO 2002, *Lecture Notes in Computer Science*, 2337:145–160, 2002.
- [64] C. Delorme and S. Poljak. Laplacian eigenvalues and the maximum cut problem. *Mathematical Programming*, 62:557–574, 1993.
- [65] C. Delorme and S. Poljak. Combinatorial properties and the complexity of a max-cut approximation. *European Journal of Combinatorics*, 14:313–333, 1993.
- [66] C. Delorme and S. Poljak. The performance of an eigenvalue bound on the max-cut problem in some classes of graphs. *Discrete Mathematics*, 111:145–156, 1993.
- [67] P. Delsarte. An algebraic approach to the association schemes of coding theory. *Philips Research Reports Supplements*, No. 10, 1973.
- [68] M. Deza and M. Laurent. *Geometry of Cuts and Metrics*. Springer-Verlag, 1997.
- [69] I. Dinur and S. Safra. The importance of being biased. In *Proceedings of the 34th Annual ACM Symposium on Theory of Computing*, ACM, New York, pages 33–42, 2002.
- [70] R.J. Duffin. Infinite Programs, in: H.W. Kuhn and A.W. Tucker eds., *Linear Inequalities and Related Systems*, Annals of Mathematical Studies vol. 38, Princeton University Press, 1956, 157–170.
- [71] A. Eisenblätter. *Frequency Assignment in GSM Networks: Models, Heuristics, and Lower Bounds*. PhD Thesis, TU Berlin, Germany, 2001. Available at ftp://ftp.zib.de/pub/zib-publications/books/PhD_eisenblaetter.ps.Z
- [72] A. Eisenblätter. The semidefinite relaxation of the k -partition polytope is strong. In W.J. Cook and A.S. Schulz, eds., IPCO 2002, *Lecture Notes in Computer Science*, 2337:273–290, 2002.
- [73] F. Eisenbrand. On the membership problem for the elementary closure of a polyhedron. *Combinatorica*, 19:299–300, 1999.
- [74] F. Eisenbrand and A.S. Schulz. Bounds on the Chvátal rank of polytopes in the 0/1 cube. In G. Cornuéjols et al., eds., IPCO 1999, *Lecture Notes in Computer Science*, 1610:137–150, 1999.
- [75] U. Feige. Randomized graph products, chromatic numbers, and the Lovász ϑ -function. *Combinatorica*, 17: 79–90, 1997. [Preliminary version in *Proceedings of the 27th Annual ACM Symposium on Theory of Computing*, ACM, New York, pp. 635–640, 1995.]
- [76] U. Feige. Randomized rounding of semidefinite programs – variations on the MAX CUT example. Randomization, Approximation, and Combinatorial Optimization, Proceedings of Random-Approx’99, *Lecture Notes in Computer Science*, 1671:189–196, Springer-Verlag, 1999.
- [77] U. Feige and M. Goemans. Approximating the value of two prover proof systems, with applications to MAX 2SAT and MAX DICUT. *Proceedings of the 3rd Israel Symposium on the Theory of Computing and Systems*, ACM, New York, pages 182–189, 1995.
- [78] U. Feige, M. Karpinski, and M. Langberg. Improved approximation of max-cut on graphs of bounded degree. *Electronic Colloquium on Computational Complexity*, Report TR00-021, 2000.

- [79] U. Feige, M. Karpinski, and M. Langberg. A note on approximating max-bisection on regular graphs. *Electronic Colloquium on Computational Complexity*, Report TR00-043, 2000.
- [80] U. Feige and R. Krauthgamer. The probable value of the Lovász-Schrijver relaxations for maximum independent set. Preprint, 2002. [*SIAM Journal on Computing*, to appear.]
- [81] U. Feige, M. Langberg, and G. Schechtman. Graphs with tiny vector chromatic numbers and huge chromatic numbers. In *Proceedings of the 43rd Annual IEEE Symposium on Foundations of Computer Science*, IEEE Computer Science Press, Los Alamitos, CA, 2002.
- [82] U. Feige and G. Schechtman. On the optimality of the random hyperplane rounding technique for MAX CUT. *Random Structures and Algorithms*, 20:403–440, 2002.
- [83] U. Feige and G. Schechtman. On the integrality ratio of semidefinite relaxations of MAX CUT. In *Proceedings of the 33rd Annual ACM Symposium on Theory of Computing*, ACM, New York, pages 433–442, 2001.
- [84] M. Fiedler. Bounds for eigenvalues of doubly stochastic matrices. *Linear Algebra and its Applications*, 5:299–310, 1972.
- [85] M. Fiedler. Algebraic connectivity of graphs. *Czechoslovak Mathematical Journal*, 23:298–305, 1973.
- [86] P. Frankl and V. Rödl. Forbidden intersections. *Transactions of the American Mathematical Society*, 300:259–286, 1987.
- [87] A. Frieze and M. Jerrum. Improved approximation algorithms for MAX k -CUT and MAX BISECTION. *Algorithmica*, 18:67–81, 1997. [Preliminary version in *Proceedings of the 4th International IPCO Conference*, Copenhagen, *Lecture Notes in Computer Science*, 920:1–13, 1995.]
- [88] T. Fujie and M. Kojima. Semidefinite programming relaxation for nonconvex quadratic programs. *Journal of Global Optimization*, 10:367–380, 1997.
- [89] D.R. Fulkerson. Anti-blocking polyhedra. *Journal of Combinatorial Theory B*, 12:50–71, 1972.
- [90] M.R. Garey, D.S. Johnson and L. Stockmeyer. Some simplified NP-complete graph problems. *Theoretical Computer Science*, 1:237–267, 1976.
- [91] M.X. Goemans. Semidefinite programming in combinatorial optimization. *Mathematical Programming*, 143–161, 1997.
- [92] M. Goemans and F. Rendl. Semidefinite programs and association schemes. *Computing*, 63:331–340, 1999.
- [93] M.X. Goemans and L. Tunçel. When does the positive semidefiniteness constraint help in lifting procedures? *Mathematics of Operations Research*, 26:796–815, 2001.
- [94] M.X. Goemans and D.P. Williamson. New $3/4$ -approximation algorithms for the maximum satisfiability problem. *SIAM Journal on Discrete Mathematics*, 7:656–666, 1994.

- [95] M.X. Goemans and D.P. Williamson. Improved approximation algorithms for maximum cuts and satisfiability problems using semidefinite programming. *Journal of the Association for Computing Machinery*, 42:1115–1145, 1995. [Preliminary version in *Proceedings of the 26th Annual ACM Symposium on Theory of Computing*, ACM, New York, pp. 422–431, 1994.]
- [96] M.X. Goemans and D.P. Williamson. Approximation algorithms for MAX-3-CUT and other problems via complex semidefinite programming. In *Proceedings of the 33rd Annual ACM Symposium on Theory of Computing*, ACM, New York, pages 443–452, 2001.
- [97] D. Grigoriev, E.A. Hirsch, and D.V. Pasechnik. Complexity of semi-algebraic proofs. *Lecture Notes in Computer Science*, 2285:419–430, 2002.
- [98] D. Grigoriev, E. de Klerk, and D.V. Pasechnik. Finding optimum subject to few quadratic constraints in polynomial time. In preparation, 2002.
- [99] R. Grone, C.R. Johnson, E.M. Sá, and H. Wolkowicz. Positive definite completions of partial Hermitian matrices. *Linear Algebra and its Applications*, 58:109–124, 1984.
- [100] M. Grötschel, L. Lovász, and A. Schrijver. *Geometric Algorithms and Combinatorial Optimization*. Springer-Verlag, Berlin, New York, 1988.
- [101] M. Grötschel and W.R. Pulleyblank. Weakly bipartite graphs and the max-cut problem. *Operations Research Letters*, 1:23–27, 1981.
- [102] G. Gruber and F. Rendl. Computational experience with stable set relaxations. *SIAM Journal on Optimization*, to appear. Available at <http://www-sci.uni-klu.ac.at/math-or/home/publicat.htm>
- [103] B. Guenin. A characterization of weakly bipartite graphs. *Journal of Combinatorial Theory B*, 83:112–168, 2001.
- [104] S.W. Hadley, F. Rendl, and H. Wolkowicz. A new lower bound via projection for the quadratic assignment problem. *Mathematics of Operations Research*, 17:727–739, 1992.
- [105] M.M. Halldórsson. A still better performance guarantee for approximate graph coloring. *Information Processing Letters*, 45:19–23, 1993.
- [106] M.M. Halldórsson. Approximations of independent sets in graphs. In K. Jansen and J. Rolim, eds., APPROX '98, *Lecture Notes in Computer Science*, 1444:1–14, 1998.
- [107] M.M. Halldórsson. Approximations of weighted independent sets and hereditary subset problems. In T. Asano et al., eds., COCOON '99, *Lecture Notes in Computer Science*, 1627:261–270, 1999.
- [108] E. Halperin. Improved approximation algorithms for the vertex cover problem in graphs and hypergraphs. *SIAM Journal on Computing*, 31:1608–1623, 2002. [Preliminary version in *Proceedings of 11th ACM-SIAM Symposium on Discrete Algorithms*, pages 329–337, 2000.]
- [109] E. Halperin, D. Livnat, and U. Zwick. MAX-CUT in cubic graphs. In *Proceedings of 13th ACM-SIAM Symposium on Discrete Algorithms*, pages 506–513, 2002.

- [110] E. Halperin, R. Nathaniel, and U. Zwick. Coloring k -colorable graphs using relatively small palettes. In *Proceedings of 12th ACM-SIAM Symposium on Discrete Algorithms*, pages 319–326, 2001.
- [111] E. Halperin and U. Zwick. A unified framework for obtaining improved approximations algorithms for maximum graph bisection problems. In K. Aardal and A.M.H. Gerards, eds., *IPCO 2001, Lecture Notes in Computer Science*, 2081:210–225, 2001.
- [112] E. Halperin and U. Zwick. Approximation algorithms for MAX 4-SAT and rounding procedures for semidefinite programs. *Journal of Algorithms*, 40:184–211, 2001. [Preliminary version in *Proceedings of the 7th conference on Integer Programming and Combinatorial Optimization*, Graz, Austria, pp. 202–217, 1999.]
- [113] E. Halperin and U. Zwick. Combinatorial approximation algorithms for the maximum directed cut problem. In *Proceedings of 12th ACM-SIAM Symposium on Discrete Algorithms*, pages 1–7, 2001.
- [114] J. Håstad. Some optimal inapproximability results. In *Proceedings of the 29th Annual ACM Symposium on the Theory of Computing*, ACM, New York, pages 1–10, 1997. [Full version in *Electronic Colloquium on Computational Complexity*, Report TR97-037.]
- [115] C. Helmberg, F. Rendl, R.J. Vanderbei and H. Wolkowicz. An interior-point method for semidefinite programming. *SIAM Journal on Optimization*, 6:342–361, 1996.
- [116] C. Helmberg, F. Rendl and R. Weismantel. A semidefinite programming approach to the quadratic knapsack problem. *Journal of Combinatorial Optimization*, 4:197–215, 2000.
- [117] R.D. Hill and S.R. Waters. On the cone of positive semidefinite matrices. *Linear Algebra and its Applications*, 90:81–88, 1987.
- [118] A.J. Hoffman and H.W. Wielandt. The variation of the spectrum of a normal matrix. *Duke Mathematical Journal*, 20:37–39, 1953.
- [119] R.A. Horn, and C.R. Johnson. *Matrix Analysis*. Cambridge University Press, 1985.
- [120] K. Jansen, M. Karpinski, and A. Lingas. A polynomial time approximation scheme for MAX-BISECTION on planar graphs. *Electronic Colloquium on Computational Complexity*, Report TR00-064, 2000.
- [121] C.R. Johnson. Matrix completion problems: a survey. In C.R. Johnson, editor, *Matrix Theory and Applications*, volume 40 of *Proceedings of Symposia in Applied Mathematics*, pages 171–198. American Mathematical Society, Providence, Rhode Island, 1990.
- [122] C.R. Johnson, B. Kroschel, and H. Wolkowicz. An interior-point method for approximate positive semidefinite completions. *Computational Optimization and Applications*, 9:175–190, 1998.
- [123] D. Johnson. Approximation algorithms for combinatorial problems. *Journal of Computer and System Sciences*, 9:256–278, 1974.
- [124] V. Kann, S. Khanna, J. Lagergren, and A. Panconesi. On the hardness of approximating MAX k -CUT and its dual. *Chicago Journal of Theoretical Computer Science*, 2, 1997.

- [125] D. Karger, R. Motwani, and M. Sudan. Approximate graph colouring by semidefinite programming. *Journal of the Association for Computing Machinery*, 45:246–265, 1998. [Preliminary version in *Proceedings of 35th IEEE Symposium on Foundations of Computer Science*, IEEE Computer Science Press, Los Alamitos, CA, pages 2–13, 1994.]
- [126] H. Karloff. How good is the Goemans-Williamson max-cut algorithm? *SIAM Journal on Computing*, 29:336–350, 1999.
- [127] H. Karloff and U. Zwick. A 7/8-approximation algorithm for MAX 3SAT? In *Proceedings of the 38th Annual IEEE Symposium on Foundations of Computer Science*, IEEE Computer Science Press, Los Alamitos, CA, pages 406–415, 1997.
- [128] R.M. Karp. Reducibility among combinatorial problems. In *Complexity of Computer Computations*, Plenum Press, New York, pp. 85–103, 1972.
- [129] L. Khachiyan and L. Porkolab. Computing integral points in convex semi-algebraic sets. In *38th Annual Symposium on Foundations of Computer Science*, IEEE Computer Science Press, Los Alamitos, CA, pages 162–171, 1997.
- [130] L. Khachiyan and L. Porkolab. Integer optimization on convex semialgebraic sets. *Discrete and Computational Geometry*, 23:207–224, 2000.
- [131] S. Khanna, N. Linial and S. Safra. On the hardness of approximating the chromatic number. *Combinatorica*, 20:393–415, 2000. [Preliminary version in *Proceedings of the 2nd Israel Symposium on Theory and Computing Systems*, IEEE Computer Society Press, Los Alamos, CA, pp. 250–260, 1993.]
- [132] J. Kleinberg and M.X. Goemans. The Lovász theta function and a semidefinite programming relaxation of vertex cover. *SIAM Journal on Discrete Mathematics*, 11:196–204, 1998.
- [133] E. de Klerk. *Aspects of Semidefinite Programming: Interior Point Algorithms and Selected Applications*. Kluwer, 2002.
- [134] E. de Klerk, D.V. Pasechnik, and J.P. Warners. Approximate graph colouring and MAX- k -CUT algorithms based on the ϑ -function. Preprint, 2000. [*Journal of Combinatorial Optimization*, to appear.] Available at <http://ssor.twi.tudelft.nl/~deklerk/>
- [135] E. de Klerk and D.V. Pasechnik. On the performance guarantee of MAX 3-CUT approximation algorithms. Preprint, 2001.
- [136] E. de Klerk and D.V. Pasechnik. Approximation of the stability number of a graph via copositive programming. *SIAM Journal on Optimization*, 12:875–892, 2002.
- [137] D.E. Knuth. The sandwich theorem. *Electronic Journal of Combinatorics*, 1:1–48, 1994.
- [138] M. Kojima, S. Shindoh and S. Hara. Interior-point methods for the monotone semidefinite linear complementarity problem in symmetric matrices. *SIAM Journal on Optimization*, 7:86–125, 1997.
- [139] M. Kojima and L. Tunçel. Cones of matrices and successive convex relaxations of nonconvex sets. *SIAM Journal on Optimization*, 10:750–778, 2000.

- [140] J.B. Lasserre. Optimality conditions and LMI relaxations for 0 – 1 programs. Technical Report N. 00099, LAAS, Toulouse, 2000.
- [141] J.B. Lasserre. Global optimization with polynomials and the problem of moments. *SIAM Journal on Optimization*, 11:796–817, 2001.
- [142] J.B. Lasserre. An explicit exact SDP relaxation for nonlinear 0 – 1 programs. In K. Aardal and A.M.H. Gerards, eds., IPCO 2001, *Lecture Notes in Computer Science*, 2081:293–303, 2001. [See also: An explicit equivalent positive semidefinite program for nonlinear 0-1 programs. *SIAM Journal on Optimization*, 12:756–769, 2002.]
- [143] J.B. Lasserre. Semidefinite programming vs. LP relaxations for polynomial programming. *Mathematics of Operations Research*, 27:347–360, 2002.
- [144] M. Laurent. The real positive semidefinite completion problem for series-parallel graphs. *Linear Algebra and its Applications*, 252:347–366, 1997.
- [145] M. Laurent. A connection between positive semidefinite and Euclidean distance matrix completion problems. *Linear Algebra and its Applications*, 273:9–22, 1998.
- [146] M. Laurent. A tour d’horizon on positive semidefinite and Euclidean distance matrix completion problems. In P. Pardalos and H. Wolkowicz, editors, *Topics in Semidefinite and Interior-Point Methods*, vol. 18 of the Fields Institute for Research in Mathematical Science, Communication Series, Providence, Rhode Island, pages 51–76, 1998.
- [147] M. Laurent. Polynomial instances of the positive semidefinite and Euclidean distance matrix completion problems. *SIAM Journal on Matrix Analysis and its Applications*, 22:874–894, 2000.
- [148] M. Laurent. On the sparsity order of a graph and its deficiency in chordality. *Combinatorica*, 21:543–570, 2001.
- [149] M. Laurent. Tighter linear and semidefinite relaxations for max-cut based on the Lovász-Schrijver lift-and-project procedure. *SIAM Journal on Optimization*, 12:345–375, 2001.
- [150] M. Laurent. A comparison of the Sherali-Adams, Lovász-Schrijver and Lasserre relaxations for 0,1-programming. Report PNA-R0108, CWI, Amsterdam, 2001. Available at <http://www.cwi.nl/~monique>
- [151] M. Laurent. Semidefinite relaxations for Max-Cut. To appear in *The Sharpest Cut, Festschrift in Honour of M. Padberg’s 60th Birthday*, M. Grötschel, ed., SIAM. Available at <http://www.cwi.nl/~monique>
- [152] M. Laurent. Lower bound for the number of iterations in semidefinite hierarchies for the cut polytope. Preprint, 2002. Available at <http://www.cwi.nl/~monique>
- [153] M. Laurent and S. Poljak. On a positive semidefinite relaxation of the cut polytope. *Linear Algebra and its Applications*, 223/224:439–461, 1995.
- [154] M. Laurent and S. Poljak. On the facial structure of the set of correlation matrices. *SIAM Journal on Matrix Analysis and its Applications*, 17:530–547, 1996.

- [155] M. Laurent, S. Poljak, and F. Rendl. Connections between semidefinite relaxations of the max-cut and stable set problems. *Mathematical Programming*, 77:225–246, 1997.
- [156] H.W. Lenstra, Jr.. Integer programming with a fixed number of variables. *Mathematics of Operations Research*, 8:538–548, 1983.
- [157] M. Lewin, D. Livnat, and U. Zwick. Improved rounding techniques for the MAX 2-SAT and MAX DI-CUT problems. In W.J. Cook and A.S. Schulz, eds., IPCO 2002, *Lecture Notes in Computer Science*, 2337:67–82, 2002.
- [158] N. Linial, E. London, and Yu. Rabinovich. The geometry of graphs and some of its algorithmic consequences. *Combinatorica*, 15:215–245, 1995.
- [159] N. Linial, A. Magen, and A. Naor. Girth and Euclidean distortion. Preprint, 2001. Available at <http://www.cs.huji.ac.il/~nati/> [*Geometric and Functional Analysis*, to appear.]
- [160] N. Linial and M.E. Sachs. On the Euclidean distortion of complete binary trees. Preprint, 2001. [*Discrete and Computational Geometry*, to appear.] Available at <http://www.cs.huji.ac.il/~nati/>
- [161] L. Lipták and L. Tunçel. Stable set problem and the lift-and-project ranks of graphs. Research Report CORR 2002-13, Department of Combinatorics and Optimization, University of Waterloo, 2002. Available at <http://www.math.uwaterloo.ca/~ltuncel/publications/>
- [162] W. Liu. *Extended Formulations and Polyhedral Projection*. PhD thesis, Department of Combinatorics and Optimization, University of Waterloo, Canada, 1988.
- [163] L. Lovász. Normal hypergraphs and the perfect graph conjecture. *Discrete Mathematics*, 2:253–267, 1972.
- [164] L. Lovász. On the Shannon capacity of a graph. *IEEE Transactions on Information Theory*, IT-25:1–7, 1979.
- [165] L. Lovász. Stable sets and polynomials. *Discrete Mathematics*, 124:137–153, 1994.
- [166] L. Lovász. Semidefinite programs and combinatorial optimization. Lecture notes. Available at <http://www.research.microsoft.com/~lovasz>
- [167] L. Lovász and A. Schrijver. Cones of matrices and set-functions and 0-1 optimization. *SIAM Journal on Optimization*, 1:166–190, 1991.
- [168] C. Lund and M. Yannakakis. On the hardness of approximating minimization problems. In *Proceedings of the 25th Annual ACM Symposium on Theory of Computing*, ACM, New York, pages 286–293, 1993.
- [169] N. Maculan. The Steiner problem in graphs. *Annals of Discrete Mathematics*, 31:185–222, 1987.
- [170] S. Mahajan and H. Ramesh. Derandomizing semidefinite programming based approximation algorithms. In *Proceedings of the 36th Symposium on Foundations of Computer Science*, IEEE Computer Science Press, Los Alamitos, CA, pages 162–169, 1995.

- [171] S. Matuura and T. Matsui. 0.863-approximation algorithm for MAX DICUT. In M. Goemans et als., eds., APPROX 2001 and RANDOM 2001, *Lecture Notes in Computer Science*, 2129:138–146, 2001.
- [172] S. Matuura and T. Matsui. 0.935-approximation randomized algorithm for MAX 2SAT and its derandomization. Technical Report METR 2001-03, University of Tokyo, 2001. Available at <http://www.keisu.t.u-tokyo.ac.jp/METR.html>
- [173] R.J. McEliece, E.R. Rodemich and H.C. Rumsey, Jr. The Lovász bound and some generalizations. *Journal of Combinatorics and System Sciences*, 3:134–152, 1978.
- [174] P. Meurdesoif. Strengthening the Lovász $\vartheta(\overline{G})$ bound for graph colouring. Preprint, 2000. [*Mathematical Programming*, to appear.]
- [175] B. Mohar and S. Poljak. Eigenvalues and the max-cut problem. *Czechoslovak Mathematical Journal*, 40:343–352, 1990.
- [176] R.D.C. Monteiro. Primal-dual path-following algorithms for semidefinite programming. *SIAM Journal on Optimization*, 7:663–678, 1997.
- [177] T.S. Motzkin and E.G. Straus. Maxima for graphs and a new proof of a theorem of Túrán. *Canadian Journal of Mathematics*, 17:533–540, 1965.
- [178] K.G. Murty and S.N. Kabadi. Some NP-complete problems in quadratic and linear programming. *Mathematical Programming*, 39:117–129, 1987.
- [179] G. Nemhauser and L. Wolsey. *Integer and Combinatorial Optimization*. John Wiley and Sons, New York, 1988.
- [180] Y. Nesterov. Quality of semidefinite relaxation for nonconvex quadratic optimization. CORE Discussion Paper # 9719, Belgium, 1997.
- [181] Y. Nesterov. Semidefinite relaxation and nonconvex quadratic optimization. *Optimization Methods and Software*, 9:141–160, 1998.
- [182] Y. Nesterov. Squared functional systems and optimization problems. In J.B.G. Frenk, C. Roos, T. Terlaky, and S. Zhang, eds., *High Performance Optimization*, pages 405–440. Kluwer Academic Publishers, 2000.
- [183] J. von Neumann. Some matrix inequalities and metrization of matrix space. *Tomsk Univ. Rev.* 1:286–300, 1937. (reprinted in: John von Neumann: Collected works, vol. 4, A.H. Taub ed., MacMillan, 205–219, 1962.)
- [184] C.E. Nugent, T.E. Vollman, and J. Ruml. An experimental comparison of techniques for the assignment of facilities to locations. *Operations Research*, 16:150–173, 1968.
- [185] M.L. Overton and R.S. Womersley. On the sum of the largest eigenvalues of a symmetric matrix. *SIAM Journal on Matrix Analysis and its Applications*, 13:41–45, 1992.
- [186] P. M. Pardalos and H. Wolkowicz (eds.). Topics in semidefinite programming and interior point methods. *Fields Institute Communications* 18, American Mathematical Society, 1998.

- [187] P.A. Parrilo. *Structured Semidefinite Programs and Semialgebraic Geometry Methods in Robustness and Optimization*. PhD thesis, California Institute of Technology, May 2000.
- [188] P.A. Parrilo. Semidefinite programming relaxations for semialgebraic problems. Preprint, 2001. Available at <http://www.cds.caltech.edu/~pablo> [*Mathematical Programming*, to appear.]
- [189] P.A. Parrilo and B. Sturmfels. Minimizing polynomial functions. Preprint, 2001. Available at <http://www.arxiv.org/abs/math.OC/0031701>
- [190] G. Pataki. Cone-LP's and semidefinite programs: Geometry and a simplex-type method. In W.H. Cunningham, S.T. McCormick and M. Queyranne, eds., IPCO 1996, *Lecture Notes in Computer Science*, 1084:162–174, 1996.
- [191] G. Pataki. On the rank of extreme matrices in semidefinite programs and the multiplicity of optimal eigenvalues. *Mathematics of Operations Research*, 23:339–358, 1998.
- [192] S. Poljak. Polyhedral and eigenvalue approximations of the max-cut problem. In *Sets, Graphs, and Numbers*, vol. 60 of *Colloquia Mathematica Societatis János Bolyai*, pages 569–581. Budapest, Hungary, 1991.
- [193] S. Poljak and F. Rendl. Nonpolyhedral relaxation of graph-bisection problems. *SIAM Journal on Optimization*, 5:467–487, 1995.
- [194] S. Poljak and Z. Tuza. The expected relative error of the polyhedral approximation of the max-cut problem. *Operations Research Letters*, 16:191–198, 1994.
- [195] S. Poljak and Z. Tuza. Maximum cuts and largest bipartite subgraphs. In W. Cook, L. Lovász and P. Seymour, editors, *Combinatorial Optimization*, vol. 20 of *DIMACS Series in Discrete Mathematics and Theoretical Computer Science*, pages 181–244. American Mathematical Society, Providence, RI, 1995.
- [196] L. Porkolab and L. Khachiyan. On the complexity of semidefinite programs. *Journal of Global Optimization*, 10:351–365, 1997.
- [197] V. Powers and B. Reznick. A new bound for Pólya's Theorem with applications to polynomials positive on polyhedra. Preprint, 2000. [*Journal of Pure and Applied Algebra*, to appear.] Available at <http://www.mathcs.emory.edu/~vicki/preprint/>
- [198] V. Powers and T. Wörmann. An algorithm for sums of squares of real polynomials. *Journal of Pure and Applied Algebra*, 127:99–104, 1998.
- [199] M. Putinar. Positive polynomials on compact semi-algebraic sets. *Indiana University Mathematics Journal*, 42:969–984, 1993.
- [200] A.J. Quist, E. de Klerk, C. Roos and T. Terlaky. Copositive relaxation for general quadratic programming. *Optimization Methods and Software*, 9:185–209, 1998.
- [201] M.V. Ramana. An exact duality theory for semidefinite programming and its complexity implications. *Mathematical Programming*, 77:129–162, 1997.
- [202] M.V. Ramana and A. Goldman. Some geometric results in semidefinite programming. *Journal of Global Optimization*, 7:33–50, 1995.

- [203] M.V. Ramana, L. Tunçel, and H. Wolkowicz. Strong duality for semidefinite programming. *SIAM Journal on Optimization*, 7:641–662, 1997.
- [204] B.A. Reed and A.J.L. Ramirez. *Perfect Graphs*. Wiley, 2001.
- [205] F. Rendl. Semidefinite programming and combinatorial optimization. *Applied Numerical Mathematics*, 29:255–281, 1999.
- [206] J. Renegar. On the computational complexity and geometry of the first order theory of the reals. *Journal of Symbolic Computation*, 13(3):255–352, 1992.
- [207] A. Schrijver. A comparison of the Delsarte and Lovász bounds. *IEEE Transactions on Information Theory*, IT-25:425–429, 1979.
- [208] A. Schrijver. *Theory of Linear and Integer Programming*. John Wiley and Sons, New York, 1986.
- [209] A. Schrijver. A short proof of Guenin’s characterization of weakly bipartite graphs. *Journal of Combinatorial Theory B*, 85:255–260, 2002.
- [210] A. Schrijver. *Combinatorial Optimization - Polyhedra and Efficiency*. Springer-Verlag, Berlin, 2003.
- [211] P.D. Seymour. The matroids with the max-flow min-cut property. *Journal of Combinatorial Theory B*, 23:189–222, 1977.
- [212] H. Sherali and W. Adams. A hierarchy of relaxations between the continuous and convex hull representations for zero-one programming problems. *SIAM Journal on Discrete Mathematics*, 3:411–430, 1990.
- [213] H. Sherali and W. Adams. A hierarchy of relaxations and convex hull representations for mixed-integer zero-one programming problems. *Discrete Applied Mathematics*, 52:83–106, 1994.
- [214] H. Sherali and W. Adams. *A Reformulation-Linearization Technique (RLT) for Solving Discrete and Continuous Nonconvex Problems*. Kluwer, 1997.
- [215] H. Sherali and C.H. Tuncbilek. A global optimization algorithm for polynomial programming problems using a Reformulation-Linearization Technique. *Journal of Global Optimization*, 2:101–112, 1992.
- [216] H.D. Sherali and C.H. Tuncbilek. Reformulation-Linearization/Convexification relaxations for univariate and multivariate polynomial programming problems. *Operations Research Letters*, 21:1–10, 1997.
- [217] N.Z. Shor. An approach to obtaining global extremums in polynomial mathematical programming problems. *Kibernetika*, 5:102–106, 1987.
- [218] N.Z. Shor. Class of global minimum bounds of polynomial functions. *Cybernetics*, 6:731–734, 1987. [Translated from *Kibernetika*, 6:9–11, 1987.]
- [219] N.Z. Shor. *Nondifferentiable Optimization and Polynomial Problems*. Kluwer Academic Publishers, 1998.

- [220] M. Skutella. Convex quadratic and semidefinite programming relaxations in scheduling. *Journal of the Association for Computing Machinery*, 48:206–242, 2001.
- [221] R. Sotirov. Bundle methods in combinatorial optimization. Dissertation, University of Klagenfurt, forthcoming.
- [222] G. Stengle. A Nullstellensatz and a Positivstellensatz in semialgebraic geometry. *Mathematische Annalen*, 207:87–97, 1974.
- [223] T. Stephen and L. Tunçel. On a representation of the matching polytope via semidefinite liftings. *Mathematics of Operations Research*, 24:1–7, 1999.
- [224] M. Szegedy. A note on the ϑ number of Lovász and the generalized Delsarte bound. In *Proceedings of the 35th Annual IEEE Symposium on Foundations of Computer Science*, IEEE Computer Science Press, Los Alamitos, CA, pages 36–39, 1994.
- [225] M.J. Todd. A study of search directions directions in primal-dual interior-point methods for semidefinite programming. *Optimization Methods and Software*, 11:1–46, 1999.
- [226] M.J. Todd. Semidefinite Programming. *Acta Numerica* 10:515–560, 2001.
- [227] L. Trevisan, G.B. Sorkin, M. Sudan, and D.P. Williamson. Gadgets, approximation, and linear programming. In *Proceedings of the 37th Annual IEEE Symposium on Foundations of Computer Science*, IEEE Computer Science Press, Los Alamitos, CA, pages 617–626, 1996.
- [228] P. Tseng. Further results on approximating nonconvex quadratic optimization by semidefinite programming relaxation. Preprint, 2001. Available at http://www.optimization-online.org/DB_HTML/200/12/417.html
- [229] L. Vandenberghe and S. Boyd. Semidefinite Programming. *SIAM Review*, 38:49–95, 1996.
- [230] W.F. de la Vega. MAX-CUT has a randomized approximation scheme in dense graphs. *Random Structures and Algorithms*, 8:187–198, 1996.
- [231] A. Wigderson. Improving the performance guarantee for approximate graph colouring. *Journal of the Association for Computing Machinery*, 30:729–735, 1983.
- [232] H. Wolkowicz, R. Saigal, and L. Vandenberghe (eds.). *Handbook of Semidefinite Programming*. Kluwer, 2000.
- [233] M. Yannakakis. Expressing combinatorial optimization problems by linear programs. In *Proceedings of the 29th International IEEE Symposium on Foundations of Computer Science*, IEEE Computer Science Press, Los Alamitos, CA, pages 223–228, 1988.
- [234] M. Yannakakis. On the approximation of maximum satisfiability. *Journal of Algorithms*, 17:475–502, 1994.
- [235] Y. Ye. Approximating quadratic programming with bound and quadratic constraints. *Mathematical Programming*, 84:219–226, 1999.
- [236] Y. Ye. A 0.699-approximation algorithm for Max-Bisection. *Mathematical Programming*, 90:101–111, 2001.

- [237] Y. Zhang. On extending some primal-dual interior-point algorithms from linear programming to semidefinite programming. *SIAM Journal on Optimization*, 8:365–386, 1998.
- [238] S. Zhang. Quadratic minimization and semidefinite relaxation. *Mathematical Programming*, 87:453–465, 2000.
- [239] Q. Zhao, S.E. Karisch, F. Rendl, and H. Wolkowicz. Semidefinite programming relaxations for the Quadratic Assignment Problem. *Journal of Combinatorial Optimization*, 2:71-109, 1998.
- [240] U. Zwick. Outward rotations: a tool for rounding solutions of semidefinite programming relaxations, with applications to MAX CUT and other problems. In *Proceedings of the 31st Annual ACM Symposium on Theory of Computing*, ACM, New York, pp. 679–687, 1999.
- [241] U. Zwick. Analyzing the MAX 2-SAT and MAX DI-CUT approximation algorithms of Feige and Goemans. Preprint, 2000. Available at <http://www.math.tau.ac.il/~zwick/>
- [242] U. Zwick. Computer assisted proof of optimal approximability results. In *Proceedings of 13th ACM-SIAM Symposium on Discrete Algorithms*, pages 496–505, 2002.