

Research Reports on Mathematical and Computing Sciences

Large-scale semidefinite programs in electronic
structure calculation

Mituhiro Fukuda, Bastiaan J. Braams, Maho
Nakata, Michael L. Overton, Jerome K. Percus,
Makoto Yamashita, and Zhengji Zhao

February 2005, B-413

Department of
Mathematical and
Computing Sciences
Tokyo Institute of Technology

SERIES B: **Operations Research**

B-413 Large-scale semidefinite programs in electronic structure calculation

Mituhiko Fukuda, Bastiaan J. Braams, Maho Nakata, Michael L. Overton, Jerome K. Percus, Makoto Yamashita, and Zhengji Zhao

Abstract

Employing the variational approach having the two-body reduced density matrix (RDM) as variables to compute the ground state energies of atomic-molecular systems has been a long time dream in electronic structure theory in chemical physics/physical chemistry. Realization of the RDM approach has benefited greatly from recent developments in semidefinite programming (SDP). We present the actual state of this new application of SDP as well as the formulation of these SDPs, which can be arbitrarily large. Numerical results using parallel computation on high performance computers are given. The RDM method has several advantages including robustness and provision of high accuracy compared to traditional electronic structure methods, although its computational time and memory consumption are still extremely large.

Key words. Large-scale optimization – Computational chemistry – Semidefinite programming relaxation – Reduced Density Matrix – N-representability – Parallel computation

1 Introduction

Electronic structure theory is the source of some of the largest and most challenging problems in computational science. As the quantum mechanical basis for the computation of properties of molecules and solids it is also of immense practical importance.

Traditional formulations of the electronic structure problem give rise to large linear or nonlinear Hermitian eigenvalue problems, but using the reduced density matrix (RDM) method [5, 20], one is required instead to solve a very large semidefinite programming (SDP) problem. Until recently the RDM method could not compete either in accuracy or in speed with well-established electronic structure methods, but this is changing. Especially Nakata *et al.* [44, 42] showed that a well-established SDP code (*e.g.*, SDPA [17]) could be used to solve an SDP having the RDMs as variables with the basic conditions (the “ P ”, “ Q ”,

M. Fukuda: Department of Mathematical and Computing Sciences, Tokyo Institute of Technology, 2-12-1-W8-29 Oh-okayama, Meguro-ku, Tokyo, 152-8552 Japan. e-mail: mituhiko@is.titech.ac.jp. The work of this author was primarily conducted at Courant Institute of Mathematical Sciences, New York University

B. J. Braams: Department of Mathematics and Computer Science, Emory University, Atlanta, GA 30322. e-mail: braams@mathcs.emory.edu

M. Nakata: Department of Applied Chemistry, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo, 113-8656 Japan. e-mail: mnakata@qcl.t.u-tokyo.ac.jp

M. L. Overton: Department of Computer Science, Courant Institute of Mathematical Sciences, New York University, New York, NY 10012. e-mail: overton@cs.nyu.edu

J. K. Percus: Courant Institute of Mathematical Sciences and Department of Physics, New York University, New York, NY 10012. e-mail: percus@cims.nyu.edu

M. Yamashita: Department of Industrial Engineering and Management, Kanagawa University, 3-27-1 Rokkakubashi, Kanagawa-ku, Yokohama-shi, Kanagawa-ken, 221-8686 Japan. e-mail: makoto.yamashita@ie.kanagawa-u.ac.jp

Z. Zhao: High Performance Computing Research Department, Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Mail Stop 50F1650, Berkeley, CA 94720. e-mail: zzhao@lbl.gov

and “ G ” conditions, as will be clarified later) for a wide variety of interesting (although still small) molecules. Later, Zhao *et al.* [62] showed that with the inclusion of additional conditions (“ $T1$ ” and “ $T2$ ”), the accuracy that is obtained for small molecular systems compares favorably with the best widely used electronic structure methods. Very recently, Mazziotti [36, 37] announced results for larger molecular systems using the P , Q and G conditions.

For applied work, the main present challenge for the RDM approach is to develop the efficiency of the solution of the resulting large SDP problems to the stage where one has a method that is genuinely competitive in both accuracy and speed with traditional electronic structure methods. Accordingly, our objective in the present paper is to present the basics of the RDM method and the details of the resulting SDP, together with our computational experience, to experts in optimization theory and semidefinite programming. Very briefly: the RDM method gives rise to arbitrarily large SDP problems. Each of the data matrices of the SDP has a small number of diagonal blocks and is extremely sparse. On the other hand, the linear combination of these matrices, *i.e.*, the dual slack matrix, is mostly dense, but its largest block has density only about 20%. One of the keys to successfully and drastically reduce the size of the SDP is to formulate it as a dual SDP problem. The dual formulation has many fewer dual variables (primal constraints) than the original primal formulation and can therefore be solved more efficiently. The SDP problems must be solved to high accuracy – typically 7 digits for the optimal value – and this is an extremely important consideration in the choice of solution methods and codes. We have experience solving these problems with both the SeDuMi code [50] and with the SDPA/SDPARA codes [17, 58].

In the next section, Section 2, we present the electronic structure problem and the RDM theory. In Section 2.1, we show the basic equations and the general form of the RDM reformulation, and we explain the concept of N -representability conditions. In Section 2.2, we exhibit the principal N -representability conditions: P , Q , G , $T1$ and $T2$, respectively, and in Section 2.3, we give a chronological overview of applications of SDP to these equations.

In Section 3, we present the precise formulation of the RDM equations in dual SDP form using inequality and equality constraints. This is an improvement over the previous result [62] where equality constraints were split into a slightly relaxed pair of inequalities. The quality of this modification is also discussed in Section 4.3. We also consider the computational advantages of the dual SDP formulation compared to the primal one in terms of both number of floating point operations and memory usage.

Section 4 gives our main results. Section 4.1 discusses the sizes and the sparsity of this class of SDPs. Section 4.2 gives the ground state energies and the dipole moments of small atomic-molecular systems solving small- and medium-scale SDPs by SeDuMi, which can handle inequality and equality constraints in the dual SDP problem. The numerical results confirm that the RDM approach employing the P , Q , G , $T1$ and $T2$ conditions provides accurate, robust and most of the time better values for the ground state energy and the dipole moment than the traditional electronic structure methods. Section 4.3 gives the same results for large-scale SDPs using the parallel SDPARA-SMP code, which has a better memory storage scheme than SDPARA [58]. Only inequality constraints are considered in the dual SDP formulation here. We also discuss essential techniques to solve large-scale problems in a high performance parallel environment. Possibly, we solved the largest SDP reported with 20709 dual variables (primal constraints) and largest block matrices with size 3211×3211 with such density and accuracy. This size was not exceeded because of lack

of available hours at the computational provider. Finally, Section 4.4 briefly reports our particular experience in using alternative formulations and methods for this problem. We conclude with an outlook for further work.

To finalize this introduction, we indicate briefly the key ideas and terminology of the RDM method in a pure linear algebra setting. Let $\mathbf{A} \in \mathbb{S}^n$ (the space of real symmetric matrices of dimension $n \times n$) and let E_0 be its smallest eigenvalue. By the Rayleigh-Ritz variational principle, E_0 is equal to the minimum of $\langle \mathbf{x}, \mathbf{A}\mathbf{x} \rangle$ over $\mathbf{x} \in \mathbb{R}^n$ subject to $\langle \mathbf{x}, \mathbf{x} \rangle = 1$, and by convexity this has an optimization formulation:

$$E_0 = \begin{cases} \min & \langle \mathbf{A}, \mathbf{X} \rangle \\ \text{subject to} & \langle \mathbf{X}, \mathbf{I} \rangle = 1, \\ & \mathbf{X} \succeq \mathbf{O}, \mathbf{X} \in \mathbb{S}^n. \end{cases} \quad (1)$$

We are interested in the case where the dimension of \mathbf{A} is exponentially large in some natural parameter, but the matrix \mathbf{A} has a special structure: $\mathbf{A} = M^\dagger(\mathbf{B})$ where $\mathbf{B} \in \mathbb{S}^m$, $M : \mathbb{S}^n \rightarrow \mathbb{S}^m$ is a linear map, and M^\dagger is its adjoint. The dimension of \mathbf{B} is only polynomially large in the same natural parameter. The map M has another important property: the $n \times n$ identity matrix is in the range of M^\dagger ; more specifically, $M^\dagger(\mathbf{I}_m) = d\mathbf{I}_n$ where d is a nonzero constant. Also, M preserves positive semidefiniteness. Thus, letting $\mathbf{Y} = M(\mathbf{X})$, the problem (1) can be reformulated as a minimization problem in \mathbb{S}^m :

$$E_0 = \begin{cases} \min & \langle \mathbf{B}, \mathbf{Y} \rangle \\ \text{subject to} & \langle \mathbf{Y}, \mathbf{I} \rangle = d, \mathbf{Y} \in \mathbb{S}^m, \text{ and} \\ & \text{“representability”}. \end{cases} \quad (2)$$

Here, “representability” means: there exists a positive semidefinite matrix $\mathbf{X} \in \mathbb{S}^n$ such that $\mathbf{Y} = M(\mathbf{X})$. An obvious necessary condition for representability, since M preserves semidefiniteness, is $\mathbf{Y} \succeq \mathbf{O}$. The success of this approach might seem to rely now on being able to specify concrete necessary and sufficient conditions for representability that do not require the reconstruction of the large matrix \mathbf{X} , but this is understood to be intractable. Instead the conditions that are known are necessary but not sufficient, and so they serve to define an approximation – a lower-bound approximation – to the original exponentially large problem. The conditions that have turned out to be most effective are all of semidefinite kind.

In the language of electronic structure, \mathbf{A} is the Hamiltonian, E_0 is the ground state energy, \mathbf{x} is a wavefunction, and \mathbf{X} is a (full) density matrix. The dimension of the matrix \mathbf{A} is exponentially large in the natural parameter N (the number of electrons) or in another natural parameter r (the size of the one-electron basis), assuming the ratio r/N is kept fixed. A matrix of the form $\mathbf{Y} = M(\mathbf{X})$ is called a reduced density matrix, \mathbb{S}^m is the space of RDMs, and the dimension m scales as r^2 .

2 The electronic structure problem and reduced density matrices

2.1 Basic formalism

The electronic structure problem is to determine the ground state energy of a many-electron system in a given external potential [51]. For an N -electron system this ground state energy is the smallest eigenvalue of a Hermitian operator (the Schrödinger operator or Hamiltonian) that acts on a space of N -electron *wavefunctions*, which are complex-valued square-integrable functions of N single-electron coordinates simultaneously that are totally antisymmetric under the interchange of any pair of electrons. The single-electron coordinates are three space coordinates and a discrete *spin* coordinate, $\mathbb{R}^3 \times \{-\frac{1}{2}, +\frac{1}{2}\}$. A wavefunction is often denoted by Ψ and if we let $\mathbf{z}_1, \dots, \mathbf{z}_N$ denote single-electron coordinates then the antisymmetry is expressed by

$$\Psi(\mathbf{z}_1, \dots, \mathbf{z}_a, \dots, \mathbf{z}_b, \dots, \mathbf{z}_N) = -\Psi(\mathbf{z}_1, \dots, \mathbf{z}_b, \dots, \mathbf{z}_a, \dots, \mathbf{z}_N)$$

for all distinct pairs of indices (a, b) and for all arguments of the wavefunction.

In our work we follow the usual approach of discretizing the many-electron space of wavefunctions by way of a discretization of the single-electron space of wavefunctions, and for purpose of exposition, we let that single-electron basis be orthonormal. If the single-electron basis functions are ψ_i for $1 \leq i \leq r$, then the discretized many-electron wavefunctions are of the form

$$\Psi(\mathbf{z}_1, \mathbf{z}_2, \dots, \mathbf{z}_N) = \sum_{i_1, i_2, \dots, i_N} c(i_1, i_2, \dots, i_N) \psi_{i_1}(\mathbf{z}_1) \psi_{i_2}(\mathbf{z}_2) \cdots \psi_{i_N}(\mathbf{z}_N), \quad (3)$$

where i_1, i_2, \dots, i_N are distinct indices from 1 to r . The antisymmetry requirement on Ψ is carried over to the discrete coefficients \mathbf{c} , so we require

$$c(i_1, \dots, i_a, \dots, i_b, \dots, i_N) = -c(i_1, \dots, i_b, \dots, i_a, \dots, i_N)$$

for all distinct pairs (a, b) and all values of the arguments.

If single-electron wavefunctions are discretized using a basis of size r then the associated basis for N -electron wavefunctions has size $r!/(N!(r-N)!)$ (taking into account the antisymmetry requirement). Even under a rather coarse discretization the basis size r is a small multiple of N , say $r = 2N$, and so the number of coefficients scales exponentially with N and quickly becomes astronomical.

This discrete formulation of the electronic structure problem as an exponentially large eigenvalue problem is called full configuration-interaction (FCI), and it is intractable except for very small systems. More practical approaches [51] involve truncating the many electron basis in some systematic way. In the most direct versions (for example in the SDCI approach: singly and doubly substituted configuration interaction) this leads to a smaller Hermitian eigenvalue problem and in other versions (for example in the CCSD approach: coupled cluster expansion using single and double excitations) one obtains a nonlinear eigenvalue problem.

An entirely different conceptual approach to the ground state electronic structure problem relies on the concept of the two-body reduced density matrix (2-RDM) of a many-electron system. This approach, first articulated in detail in two papers in the early 1960's

[5, 20] (but note as well the earlier refs. [26, 30, 31]), was the subject of active theoretical [8, 39, 13] and computational [28, 29, 19, 40, 48, 18] investigations through the 1970's, but because of limited success interest somewhat waned. The basic ideas of the RDM method were indicated in the Introduction, and we now describe them again in the specific setting of electronic structure theory.

We assume that the space of wavefunctions has been discretized as just discussed, and so we obtain a discrete Hamiltonian \mathbf{H} . The discretized ground state problem then asks for the minimum eigenvalue E_0 for the problem $\mathbf{H}\mathbf{c} = E_0\mathbf{c}$, where \mathbf{c} is the discretized wavefunction (3). The corresponding matrix variational problem (1) then becomes

$$\begin{cases} \min & \langle \mathbf{H}, \mathbf{\Gamma}^{\text{full}} \rangle \\ \text{subject to} & \langle \mathbf{\Gamma}^{\text{full}}, \mathbf{I} \rangle = 1, \\ & \mathbf{\Gamma}^{\text{full}} \succeq \mathbf{O}. \end{cases} \quad (4)$$

Here $\mathbf{\Gamma}^{\text{full}}$ denotes the *full density matrix*: a function of two pairs of N -electron variables, so of the form $\mathbf{\Gamma}^{\text{full}}(i_1, \dots, i_N; i'_1, \dots, i'_N)$ which is, like the wavefunction, antisymmetric under interchange of any pair of indices in i_1, \dots, i_N and also under interchange of any pair of indices in i'_1, \dots, i'_N . In the general case where wavefunctions are assumed to be complex, then $\mathbf{\Gamma}^{\text{full}}$ is complex Hermitian under interchange of (i_1, \dots, i_N) with (i'_1, \dots, i'_N) , but in the present work we consider only the real case and then $\mathbf{\Gamma}^{\text{full}}$ is real symmetric under that operation. $\mathbf{\Gamma}^{\text{full}}$ is an exponentially large object that is not suitable as ingredient of an effective computational method. However, a reduction to the form described in (2) is possible.

Given a full density matrix $\mathbf{\Gamma}^{\text{full}}$, the corresponding p -body RDM $\mathbf{\Gamma}_p$ is a function of two pairs of p -electron variables defined as a (scaled) partial trace over the remaining $N - p$ variables:

$$\mathbf{\Gamma}_p(i_1, \dots, i_p; i'_1, \dots, i'_p) = \frac{N!}{(N-p)!} \sum_{i_{p+1}, \dots, i_N} \mathbf{\Gamma}^{\text{full}}(i_1, \dots, i_p, i_{p+1}, \dots, i_N; i'_1, \dots, i'_p, i_{p+1}, \dots, i_N). \quad (5)$$

The linear space of $\mathbf{\Gamma}_p$ is defined by antisymmetry conditions (which are obviously inherited from $\mathbf{\Gamma}^{\text{full}}$): $\mathbf{\Gamma}_p(i_1, \dots, i_p; i'_1, \dots, i'_p)$ is antisymmetric under the interchange of any two unprimed coordinates and is also antisymmetric under the interchange of any two primed coordinates. Furthermore, $\mathbf{\Gamma}_p$ is Hermitian or real symmetric (following $\mathbf{\Gamma}^{\text{full}}$) under interchange of the vector of unprimed coordinates with the vector of primed coordinates.

The key property for RDM theory is described in the language of physics and chemistry by saying that the Hamiltonian involves – for the case of nonrelativistic electronic structure – one-body and two-body interaction terms only. The mathematical description is that the energy depends only on the one-body and two-body RDMs. Thus we have discrete operators \mathbf{H}_1 and \mathbf{H}_2 – the one-body and two-body parts of the Hamiltonian – such that on the space of density matrices

$$\langle \mathbf{H}, \mathbf{\Gamma}^{\text{full}} \rangle = \langle \mathbf{H}_1, \mathbf{\Gamma}_1 \rangle + \langle \mathbf{H}_2, \mathbf{\Gamma}_2 \rangle.$$

It is easily seen that $\langle \mathbf{\Gamma}_p, \mathbf{I} \rangle = N!/(N-p)!$ and also that the mapping $\mathbf{\Gamma}^{\text{full}} \rightarrow \mathbf{\Gamma}_p$ preserves the positive semidefiniteness property. Now a formulation of the electronic structure problem is obtained in the form of (2):

$$\begin{cases} \min & \langle \mathbf{H}_1, \mathbf{\Gamma}_1 \rangle + \langle \mathbf{H}_2, \mathbf{\Gamma}_2 \rangle \\ \text{subject to} & \langle \mathbf{\Gamma}_1, \mathbf{I} \rangle = N, \langle \mathbf{\Gamma}_2, \mathbf{I} \rangle = N(N-1), \text{ and} \\ & \text{“}N\text{-representability”}. \end{cases} \quad (6)$$

In this equation, “ N -representability” has yet to be defined – or rather, approximated – by a tractable family of necessary conditions for N -representability of the matrices $\mathbf{\Gamma}_1$ and $\mathbf{\Gamma}_2$ in the form of (5) for a positive semidefinite matrix $\mathbf{\Gamma}^{\text{full}}$.

2.2 Specific N -representability conditions

The linear space of $\mathbf{\Gamma}_1$ is the space of real symmetric $r \times r$ matrices, \mathbb{S}^r . As defined in (5), $\mathbf{\Gamma}_2$ depends on two pairs of indices, $\mathbf{\Gamma}_2(i_1, i_2; i'_1, i'_2)$, and is antisymmetric in both pairs and real symmetric under interchange of the pairs. The antisymmetry in (i_1, i_2) and in (i'_1, i'_2) will be built into the basis, and so $\mathbf{\Gamma}_2 \in \mathbb{S}^{r(r-1)/2}$. It is also clear from (5) that $\mathbf{\Gamma}_1$ is itself a scaled partial trace of $\mathbf{\Gamma}_2$:

$$\mathbf{\Gamma}_1(i, i') = \frac{1}{N-1} \sum_{j=1}^r \mathbf{\Gamma}_2(i, j; i', j). \quad (7)$$

$\mathbf{\Gamma}_1$ could therefore be eliminated entirely from the problem. However, both the objective function and the N -representability conditions are more conveniently formulated if $\mathbf{\Gamma}_1$ is retained and if the trace condition (7) is used as a set of linear constraints on the pair $(\mathbf{\Gamma}_1, \mathbf{\Gamma}_2)$. We follow this approach.

The trace conditions on $\mathbf{\Gamma}_1$ and $\mathbf{\Gamma}_2$ were specified in (6). The remaining conditions are in the form of convex inequalities. Moreover, all conditions that we have used are of semidefinite form.

For the 1-RDM the remaining necessary and sufficient N -representability conditions [5] are:

$$\mathbf{I} \succeq \mathbf{\Gamma}_1 \succeq \mathbf{O}. \quad (8)$$

For the 2-RDM, a complete family of constructive necessary and sufficient conditions is not available. On a smaller subspace of matrices (the “diagonal” 2-RDM’s), the N -representability problem is well understood: this diagonal N -representability problem is equivalent to characterization of the Correlation Polytope, also known as the Boolean Quadric Polytope and equivalent via a linear bijection to the Cut Polytope [9, p. 54]. Optimization over the Boolean Quadric Polytope is \mathcal{NP} -hard (it is the same as the unconstrained 0-1 quadratic programming problem), and as is pointed out in [9, p. 397], it follows from a result of Karp and Papadimitriou [27] that a polynomially concise description of all the facets of this polytope is not available unless $\mathcal{NP} = \text{co-}\mathcal{NP}$. For earlier investigations into the diagonal N -representability problem, we note [8, 39, 13]. As the original problem (4) is exponentially large, this complexity barrier should not deter us – the RDM method is to be viewed as an approximation method and one works with necessary conditions for N -representability that are known not to be sufficient.

The basic well known convex inequalities for the 2-RDM are the P and the Q conditions (so named in [20], but they are also found in [5]) and the G condition [20]. In our previous work [62] we added to this a $T1$ and a $T2$ condition, which as we pointed out are implied by

a much earlier paper of Erdahl [13]. All these conditions are of semidefinite form: $\mathbf{P} \succeq \mathbf{O}$, $\mathbf{Q} \succeq \mathbf{O}$, $\mathbf{G} \succeq \mathbf{O}$, $\mathbf{T1} \succeq \mathbf{O}$, and $\mathbf{T2} \succeq \mathbf{O}$, where the matrices \mathbf{P} , \mathbf{Q} , \mathbf{G} , $\mathbf{T1}$ and $\mathbf{T2}$ are defined by linear combinations of the entries of the basic matrices $\mathbf{\Gamma}_1$ and $\mathbf{\Gamma}_2$. Specifically (all indices range over $1, \dots, r$ and δ is the Kronecker delta):

$$\mathbf{P} \equiv \mathbf{\Gamma}_2, \quad (9)$$

$$Q(i, j; i', j') \equiv \Gamma_2(i, j; i', j') - \delta(i, i')\Gamma_1(j, j') - \delta(j, j')\Gamma_1(i, i') + \delta(i, j')\Gamma_1(j, i') \\ + \delta(j, i')\Gamma_1(i, j') + \delta(i, i')\delta(j, j') - \delta(i, j')\delta(j, i'). \quad (10)$$

The matrices \mathbf{P} and \mathbf{Q} are of the same size as $\mathbf{\Gamma}_2$ and have the same antisymmetry property, so they belong to $\mathbb{S}^{r(r-1)/2}$. Also,

$$G(i, j; i', j') = \Gamma_2(i, j'; j, i') + \delta(i, i')\Gamma_1(j', j). \quad (11)$$

In the matrix \mathbf{G} there is no antisymmetry in (i, j) or in (i', j') , so \mathbf{G} belongs to \mathbb{S}^{r^2} . Also,

$$T1(i, j, k; i', j', k') = \delta(i, i')\Gamma_2(k', j'; k, j) - \delta(i, j')\Gamma_2(k', i'; k, j) \\ + \delta(i, k')\Gamma_2(j', i'; k, j) - \delta(j, i')\Gamma_2(k', j'; k, i) \\ + \delta(j, j')\Gamma_2(k', i'; k, i) - \delta(j, k')\Gamma_2(j', i'; k, i) \\ + \delta(k, i')\Gamma_2(k', j'; j, i) - \delta(k, j')\Gamma_2(k', i'; j, i) \\ + \delta(k, k')\Gamma_2(j', i'; j, i) \\ + (\delta(j, k')\delta(k, j') - \delta(j, j')\delta(k, k'))\Gamma_1(i', i) \\ + (\delta(i, j')\delta(k, k') - \delta(i, k')\delta(k, j'))\Gamma_1(i', j) \\ + (\delta(i, k')\delta(j, j') - \delta(i, j')\delta(j, k'))\Gamma_1(i', k) \\ + (\delta(j, i')\delta(k, k') - \delta(j, k')\delta(k, i'))\Gamma_1(j', i) \\ + (\delta(i, k')\delta(k, i') - \delta(i, i')\delta(k, k'))\Gamma_1(j', j) \\ + (\delta(i, i')\delta(j, k') - \delta(i, k')\delta(j, i'))\Gamma_1(j', k) \\ + (\delta(j, j')\delta(k, i') - \delta(j, i')\delta(k, j'))\Gamma_1(k', i) \\ + (\delta(i, i')\delta(k, j') - \delta(i, j')\delta(k, i'))\Gamma_1(k', j) \\ + (\delta(i, j')\delta(j, i') - \delta(i, i')\delta(j, j'))\Gamma_1(k', k) \\ + \delta(i, i')\delta(j, j')\delta(k, k') - \delta(i, j')\delta(j, i')\delta(k, k') \\ - \delta(i, i')\delta(j, k')\delta(k, j') + \delta(i, j')\delta(j, k')\delta(k, i') \\ + \delta(i, k')\delta(j, i')\delta(k, j') - \delta(i, k')\delta(j, j')\delta(k, i'). \quad (12)$$

$\mathbf{T1}$ is fully antisymmetric in both its index triples, so it belongs to $\mathbb{S}^{r(r-1)(r-2)/6}$. Finally,

$$T2(i, j, k; i', j', k') = \delta(i, i')\Gamma_2(j', k'; j, k) - \delta(j, j')\Gamma_2(k', i; k, i') \\ - \delta(k, k')\Gamma_2(j', i; j, i') + \delta(j, k')\Gamma_2(j', i; k, i') \\ + \delta(k, j')\Gamma_2(k', i; j, i') + \delta(k, k')\delta(j, j')\Gamma_1(i, i') \\ - \delta(j, k')\delta(k, j')\Gamma_1(i, i'). \quad (13)$$

$T2(i, j, k; i', j', k')$ is antisymmetric in (j, k) and in (j', k') , so it belongs to $\mathbb{S}^{r^2(r-1)/2}$.

Let us briefly discuss spin symmetries, which can be used to reduce the above matrices to a block-diagonal form. In order to make use of spin symmetry the basis functions ψ_i must be organized in pairs; each function is a product of one of $r/2$ orthogonal spatial

basis functions (functions of the spatial coordinates) and one of two orthogonal spin states (functions of the spin coordinate). Each index i can then be further refined by a bijection to a pair of indices $n(i)$ (spatial orbitals) and $\sigma(i)$ (spin states). The spatial orbitals $n(i)$ can take the indices $1, 2, \dots, \frac{r}{2}$ while the spin states $\sigma(i)$ can take the values $+1/2$ (α spin, spin up) or $-1/2$ (β spin, spin down).

Then our variable matrices and the matrix conditions will vanish for these values of the indices:

$$\begin{aligned}
\Gamma_1(n(i)\sigma(i), n(i')\sigma(i')) &= 0 && \text{for } \sigma(i) \neq \sigma(i'), \\
A(n(i)\sigma(i), n(j)\sigma(j); n(i')\sigma(i'), n(j')\sigma(j')) &= 0 && \text{where } \mathbf{A} = \mathbf{\Gamma}_2 \text{ or } \mathbf{Q} \\
&&& \text{for } \sigma(i) + \sigma(j) \neq \sigma(i') + \sigma(j'), \\
G(n(i)\sigma(i), n(j)\sigma(j); n(i')\sigma(i'), n(j')\sigma(j')) &= 0 && \text{for } \sigma(i) + \sigma(j') \neq \sigma(j) + \sigma(i'), \\
T1(n(i)\sigma(i), n(j)\sigma(j), n(k)\sigma(k); n(i')\sigma(i'), n(j')\sigma(j'), n(k')\sigma(k')) &= 0 && (14) \\
&&& \text{for } \sigma(i) + \sigma(j) + \sigma(k) \neq \sigma(i') + \sigma(j') + \sigma(k'), \\
T2(n(i)\sigma(i), n(j)\sigma(j), n(k)\sigma(k); n(i')\sigma(i'), n(j')\sigma(j'), n(k')\sigma(k')) &= 0 && \\
&&& \text{for } \sigma(i) + \sigma(j') + \sigma(k') \neq \sigma(j) + \sigma(k) + \sigma(i').
\end{aligned}$$

In addition to the trace conditions (6), there are trace conditions corresponding to the number of electrons with α spin:

$$\sum_{n(i)=1}^{r/2} \Gamma_1(n(i)\alpha, n(i)\alpha) = N_\alpha, \quad (15)$$

$$\sum_{n(i), n(j)=1}^{r/2} \Gamma_2(n(i)\alpha, n(j)\alpha; n(i)\alpha, n(j)\alpha) = N_\alpha(N_\alpha - 1), \quad (16)$$

where $N = N_\alpha + N_\beta$ ($\beta = -\alpha$).

Finally there is a linear constraint for the given total spin S ,

$$\begin{aligned}
&\sum_{n(i), n(j)=1}^{r/2} (\Gamma_2(n(i)\alpha, n(j)\alpha; n(i)\alpha, n(j)\alpha) + \Gamma_2(n(i)\beta, n(j)\beta; n(i)\beta, n(j)\beta)) \\
&- 2 \sum_{n(i), n(j)=1}^{r/2} \Gamma_2(n(i)\alpha, n(j)\beta; n(i)\alpha, n(j)\beta) \\
&- 4 \sum_{n(i), n(j)=1}^{r/2} \Gamma_2(n(i)\alpha, n(j)\beta; n(j)\alpha, n(i)\beta) + 3N = 4S(S + 1).
\end{aligned} \quad (17)$$

2.3 Previous numerical computations using the RDM method

Following the clear statement of the RDM approach and of the most important N -representability conditions [5, 20], the first significant computational results came in the 1970s. Kijewski [28, 29] applied the RDM method to doubly ionized carbon ($N = 4$), C^{++} , using a basis of 10 spin orbitals ($r = 10$). Garrod and co-authors were the first ones to actually solve the SDP imposing the P , Q and G conditions, by which they obtained very accurate results for atomic beryllium ($N = 4$ and $r = 10$) [19, 48, 18]. Mihailović and Rosina also considered the RDM method for nuclear physics [40], but reported rather poor accuracy.

This early work belongs firmly to semidefinite programming, although that name was not yet in use. The analytical work [20] is focused on semidefinite conditions (for fermion and also for boson systems), and the subsequent computational methods would be recognized by anyone working in semidefinite programming today. Rosina and Garrod [48] described two main algorithms to solve the SDP. One successively added cutting planes into the linear programming relaxation of the problem, and the other minimized the objective function incorporating a barrier function for the cone of positive semidefinite matrices!

Because of the high computational cost and the lack of progress on the N -representability problem interest in the computational aspects of the RDM approach fell off during the 1980s, but it has been rekindled in recent years. Nakata et al. [44] showed that the RDM method with the P , Q and G conditions provides ground state energies that compare very favorably to Hartree-Fock results for a wide variety of small molecules (r up to 16). In subsequent work [42], they demonstrated that the method maintains its accuracy when molecular dissociation is modeled – a test that is failed by many of the traditional methods of electronic structure calculation. In a previous paper [62] several of us confirmed and extended the results of [44, 42] for the accuracy of the RDM method with P , Q and G conditions relative to the Hartree-Fock approximation. We further showed that by adding two additional N -representability conditions, which we called $T1$ and $T2$, one obtains for small molecular systems (r up to 20) an accuracy that compares favorably not just with Hartree-Fock but with the best standard methods of quantum chemistry. Although the cost of the RDM method is still very high compared to traditional methods, Mazziotti [36, 37] recently announced results for considerably larger systems (r up to 36) for the RDM approach imposing only the P , Q and G conditions.

In the present paper we discuss in detail only our chosen approach of optimizing the 2-RDM subject to semidefinite N -representability conditions ($P, Q, G, T1, T2$), without invoking 3-body or higher RDMs. We note here, however, a related approach being actively pursued that employs 2-body and higher reduced density matrices. In this other approach, under the name of Density Equation (DE) or Contracted Schrödinger Equation (CSE) [4, 45, 55, 7, 32, 59, 60], the primary unknown is the 1-RDM or 2-RDM and the equations involve an approximate reconstructed 3-RDM or 4-RDM. An excellent survey can be found in the edited volume [3] that includes contributions by Coleman [6], Erdahl [15], Nakatsuji [46], Valdemoro [56] and Mazziotti [33]. Applications of the DE/CSE approach to quantum chemistry include [60, 12]. In its original form the DE/CSE method does not impose the basic positivity conditions on the 2-RDM, but Erdahl and Jin [14, 15] and Mazziotti [38, 34] set up and solve equations closely related to the DE/CSE ones in which positivity conditions are imposed on the 2-RDM and on higher-order reconstructed density matrices [35].

3 The SDP formulation of the RDM method

Let \mathbf{C}, \mathbf{A}_p ($p = 1, 2, \dots, m$) be given block-diagonal symmetric matrices with prescribed block sizes, and $\mathbf{c}, \mathbf{a}_p \in \mathbb{R}^s$ ($p = 1, 2, \dots, m$) be given s -dimensional real vectors. We denote by $\mathbf{Diag}(\mathbf{a})$ a diagonal matrix with the elements of \mathbf{a} on its diagonal.

The primal SDP is defined as

$$\begin{cases} \max & \langle \mathbf{C}, \mathbf{X} \rangle + \langle \mathbf{Diag}(\mathbf{c}), \mathbf{Diag}(\mathbf{x}) \rangle \\ \text{subject to} & \langle \mathbf{A}_p, \mathbf{X} \rangle + \langle \mathbf{Diag}(\mathbf{a}_p), \mathbf{Diag}(\mathbf{x}) \rangle = b_p, \quad (p = 1, 2, \dots, m) \\ & \mathbf{X} \succeq \mathbf{O}, \quad \mathbf{x} \in \mathbb{R}^s, \end{cases} \quad (18)$$

and its dual

$$\begin{cases} \min & \mathbf{b}^T \mathbf{y} \\ \text{subject to} & \mathbf{S} = \sum_{p=1}^m \mathbf{A}_p y_p - \mathbf{C} \succeq \mathbf{O}, \\ & \sum_{p=1}^m \mathbf{Diag}(\mathbf{a}_p) y_p = \mathbf{Diag}(\mathbf{c}), \\ & \mathbf{y} \in \mathbb{R}^m, \end{cases} \quad (19)$$

where (\mathbf{X}, \mathbf{x}) are the primal variables and (\mathbf{S}, \mathbf{y}) are the dual variables.

Primal-dual interior-point methods and their variants are the most established and efficient algorithms to solve general SDPs. Details on how these iterative methods work can be found in [57, 52, 41].

In this section, we formulate the RDM method with the $(P, Q, G, T1, T2)$ N -representability conditions as an SDP. Observe that the 1-RDM variational variable $\mathbf{\Gamma}_1$ and its corresponding Hamiltonian \mathbf{H}_1 is a two index matrix (see (6)), but the 2-RDM variational variable $\mathbf{\Gamma}_2$, the corresponding Hamiltonian \mathbf{H}_2 , as well as \mathbf{Q} and \mathbf{G} are four index matrices, and moreover, $\mathbf{T1}$ and $\mathbf{T2}$ are six index matrices. We map each pair i, j or triple i, j, k of indices to a composite index for these matrices, resulting in symmetric matrices of order $r(r-1)/2 \times r(r-1)/2$ for $\mathbf{\Gamma}_2$, \mathbf{H}_2 and \mathbf{Q} , a symmetric matrix of order $r(r-1)(r-2)/6 \times r(r-1)(r-2)/6$ for $\mathbf{T1}$, and a symmetric matrix of order $r^2(r-1)/2 \times r^2(r-1)/2$ for $\mathbf{T2}$. For example, the four-index element $\Gamma_2(i, j; i', j')$, with $1 \leq i < j \leq r$, $1 \leq i' < j' \leq r$, can be associated with the two-index element $\widetilde{\Gamma}_2(j-i + (2r-i)(i-1)/2, j'-i' + (2r-i')(i'-1)/2)$. We assume henceforth that all matrices have their indices mapped to two indices, and we keep the same notation for simplicity. Furthermore, due to the antisymmetry property of the 2-RDM $\mathbf{\Gamma}_2$ and of the N -representability conditions $Q, T1$ and $T2$, and also due to the spin symmetry, all these matrices reduce to block-diagonal matrices of size specified in Table 1.

Now, let us define a linear transformation $\text{svec} : \mathbb{S}^n \rightarrow \mathbb{R}^{n(n+1)/2}$ as

$$\text{svec}(\mathbf{U}) = (U_{11}, \sqrt{2}U_{12}, U_{22}, \sqrt{2}U_{13}, \sqrt{2}U_{23}, U_{33}, \dots, \sqrt{2}U_{1n}, \dots, U_{nn})^T, \quad \mathbf{U} \in \mathbb{S}^n.$$

To formulate the RDM method with the $(P, Q, G, T1, T2)$ conditions in (6) as the dual SDP (19), define

$$\mathbf{y} = (\text{svec}(\mathbf{\Gamma}_1)^T, \text{svec}(\mathbf{\Gamma}_2)^T)^T \in \mathbb{R}^m \quad \text{and} \quad \mathbf{b} = (\text{svec}(\mathbf{H}_1)^T, \text{svec}(\mathbf{H}_2)^T)^T \in \mathbb{R}^m.$$

It is now relatively straightforward to express the N -representability conditions (8) through (13) as the dual slack matrix variable \mathbf{S} by defining it to have the following diagonal blocks: $\mathbf{\Gamma}_1, \mathbf{I} - \mathbf{\Gamma}_1, \mathbf{\Gamma}_2, \mathbf{Q}, \mathbf{G}, \mathbf{T1}, \mathbf{T2}$ taking into account the spin symmetry (14) and making suitable definitions for the matrices \mathbf{C}, \mathbf{A}_p ($p = 1, 2, \dots, m$). The equalities in (6) and (7), and (15) through (17) will define the vectors \mathbf{c}, \mathbf{a}_p ($p = 1, 2, \dots, m$).

The required number of floating point operations when solving these problems for instance using the parallel code SDPARA [58] are as follows. The computational flops per

iteration when using SDPARA (Section 4.3) can be estimated as $\mathcal{O}(m^2 f^2/q + m^3/q + mn_{\max}^2 + n_{\max}^3)$, where n_{\max} is the size of the largest block matrix, f is the maximum number of nonzero elements in each data matrix \mathbf{A}_p ($p = 1, 2, \dots, m$), and q is the number of used processors. In our case, $m = \mathcal{O}(r^4)$, $n_{\max} = \mathcal{O}(r^3)$ and $f = \mathcal{O}(r^2)$, and therefore, the computational flops per iteration is $\mathcal{O}(r^{12}/q)$, while the total memory usage becomes $\mathcal{O}(m^2) = \mathcal{O}(r^8)$.

The formulation of the RDM method as a dual SDP, as considered here, has a clear advantage over the primal SDP formulation [44, 34, 42, 36, 37] as detailed in [62]. When using the primal SDP formulation with the $(P, Q, G, T1, T2)$ conditions, we have $m = \mathcal{O}(r^6)$, $n_{\max} = \mathcal{O}(r^3)$ and $f = \mathcal{O}(1)$, and then, the computational flops per iterations becomes $\mathcal{O}(r^{18}/q)$, while the total memory usage becomes $\mathcal{O}(m^2) = \mathcal{O}(r^{12})$.

The formulation (19) proposed here is novel in the sense that it now includes equality constraints that were previously absent [62]. The implications of these two different formulations are discussed in Section 4.3.

4 Numerical results for the RDM method

First, we present the sizes of the SDPs which arise from this application, and briefly analyze their sparsity. In Section 4.2, we solve small- and medium-scale SDPs with discretization basis size r up to 18 using SeDuMi. The ground state energies and dipole moments obtained by the RDM method imposing the $(P, Q, G, T1, T2)$ conditions are compared with the values obtained by the traditional electronic structure methods from the specialized packages Gamess and Gaussian 98. Section 4.3 gives similar results for large-scale SDPs with r up to 26 using the parallel code SDPARA. The SDP formulation is slightly changed here and we discuss several issues regarding large-scale optimization. Finally, in Section 4.4, we briefly comment on the limitations of alternative optimization methods to solve these problems.

4.1 Sizes and sparsity of SDPs

Table 1 shows the typical size of the SDP relaxation problem (19) as a function of the discretization basis size r , itemizing the sizes of block matrices for each of the N -representability conditions.

Observe that the number of equality constraints in the primal SDP (18) grows as $m \approx 3r^4/64$, while the size of the largest block matrices corresponding to the $T2$ condition grows as approximately $3r^3/16$, and they do not depend on the number of electrons N of the system.

As one can observe from the N -representability conditions given in Section 2.2 and the actual formulation (19) as an SDP, all data matrices for our problem have integral values, excepting the diagonal matrices $\mathbf{Diag}(\mathbf{c})$, $\mathbf{Diag}(\mathbf{a}_p)$ ($p = 1, 2, \dots, m$) which have rational values, and the objective function vector \mathbf{b} which has real values. Also, if we have two different systems with a common discretization basis size r , only the diagonal matrices and the objective function vector differ, and the entries corresponding to the semidefinite conditions of the 1-RDM $\mathbf{\Gamma}_1$ and the $(P, Q, G, T1, T2)$ conditions will be exactly the same. This fact can eventually be explored to re-solve a new system with the same discretization basis size r once we have the results from a previous one.

Table 1: Size of the SDP relaxation problem as a function of the discretization basis size r .

# constraints m in primal SDP (18)	$\frac{r}{4}(\frac{3r^3}{16} - \frac{r^2}{4} + \frac{9r}{4} + 1)$
N -representability conditions	Size of block matrices
dimension of the free variable \mathbf{x}	$\frac{r}{2}(\frac{r}{2} + 1) + 5$
$\mathbf{\Gamma}_1 \succeq \mathbf{O}$	$\frac{r}{2}, \frac{r}{2}$
$\mathbf{I} - \mathbf{\Gamma}_1 \succeq \mathbf{O}$	$\frac{r}{2}, \frac{r}{2}$
$\mathbf{P} \equiv \mathbf{\Gamma}_2 \succeq \mathbf{O}$	$\frac{r^2}{4}, \frac{r}{4}(\frac{r}{2} - 1), \frac{r}{4}(\frac{r}{2} - 1)$
$\mathbf{Q} \succeq \mathbf{O}$	$\frac{r^2}{4}, \frac{r}{4}(\frac{r}{2} - 1), \frac{r}{4}(\frac{r}{2} - 1)$
$\mathbf{G} \succeq \mathbf{O}$	$\frac{r^2}{2}, \frac{r^2}{4}, \frac{r^2}{4}$
$\mathbf{T}_1 \succeq \mathbf{O}$	$\frac{r^2}{8}(\frac{r}{2} - 1), \frac{r^2}{8}(\frac{r}{2} - 1), \frac{r}{12}(\frac{r}{2} - 1)(\frac{r}{2} - 2), \frac{r}{12}(\frac{r}{2} - 1)(\frac{r}{2} - 2)$
$\mathbf{T}_2 \succeq \mathbf{O}$	$\frac{r^2}{8}(\frac{3r}{2} - 1), \frac{r^2}{8}(\frac{3r}{2} - 1), \frac{r^2}{8}(\frac{r}{2} - 1), \frac{r^2}{8}(\frac{r}{2} - 1)$

Figure 1 shows the number of nonzero elements of *each* data matrix. We counted the nonzero elements of each data matrix $(\mathbf{C}, \mathbf{Diag}(\mathbf{c}))$, $(\mathbf{A}_p, \mathbf{Diag}(\mathbf{a}_p))$ ($p = 1, 2, \dots, m$) and sorted them in non-increasing order for $r = 10$ with $m = 465$, and $r = 26$ with $m = 20709$. The maximum number of nonzero elements which occurred for each case are 245 out of 76985 possible occurrences, and 3325 out of 25151105, respectively, which correspond to the data matrix $(\mathbf{C}, \mathbf{Diag}(\mathbf{c}))$. After that, the data matrices $(\mathbf{A}_p, \mathbf{Diag}(\mathbf{a}_p))$ with the second most nonzero elements correspond to the coefficients of the variables $\mathbf{\Gamma}_1$ and $\mathbf{\Gamma}_2$ in this order. Therefore, each data matrix is extremely sparse.




Figure 1: Number of nonzero elements of each data matrix in the SDP for $r = 10$ with $m = 465$ (left) and $r = 26$ with $m = 20709$ (right).

A more interesting sparsity characterization of the problem can be observed by analyzing the density rate of the dual slack matrix variable $\mathbf{S} = \sum_{p=1}^m \mathbf{A}_p y_p - \mathbf{C}$, which has 21 block matrices as itemized in Table 1, for a random nonzero vector $\mathbf{y} \in \mathbb{R}^m$. From the definition and the dual SDP formulation (19) we used, one can see that the block matrices corresponding to the 1-RDM characterization, the P condition, and the Q condition are fully dense. In addition, the two smallest block matrices of the G condition are fully dense,

too. Figure 2 (left) depicts the density of the other block matrices as a function of the discretization basis size r . More specifically, this figure shows the density of the largest block matrix of the G condition, and the block matrices corresponding to the $T1$ and $T2$ conditions (see Table 1). The density rate of the two largest block matrices of the $T1$ condition coincides with the two smallest block matrices of the $T2$ condition here.




Figure 2: Density rates of the sparse block matrices as a function of the discretization basis size r (left), and sparsity structure for the two largest block matrices of the $T2$ condition for $r = 12$ (right).

A very positive aspect of the density rates is that for all the block matrices corresponding to the $T1$ and $T2$ conditions, the density decreases as r increases. In particular, the crucial block matrix corresponding to the two largest block matrices of the $T2$ condition are the sparsest ones due to the product of Kronecker deltas (13), although they are still rather dense: 19.3% for $r = 26$.

Figure 2 (right) shows the sparsity structure corresponding to the two largest block matrices of the $T2$ condition from \mathbf{S} for $r = 12$. These block matrices are still very dense (37.7 %) and apparently do not have an obvious sparsity structure which could be exploited.

4.2 Numerical results for small- and medium-scale problems

We utilized SeDuMi 1.05 [50] for small- and medium-scale SDPs on a Pentium Xeon 2.4GHz with 6GB of memory, and a level two cache of size 512KB. SDPT3 3.1 [54] is the only other software package that can solve SDPs with inequality and equality constraints in the dual SDP (19), but our experiments showed that SeDuMi provides much more accurate solutions.

Table 2 shows the actual sizes, the typical time and memory usage of the SDPs we picked for each discretization basis size r . We only listed the sizes of the largest block matrices among the 21 block matrices and one diagonal matrix. Here, 306x2 for instance means that there are two block matrices of sizes 306×306 each.

Table 3 shows our main result, the ground state energies calculated by the RDM method, imposing the (P,Q,G) , $(P,Q,G,T1)$, $(P,Q,G,T2)$, and $(P,Q,G,T1,T2)$ conditions (columns 7–10) to verify numerically the effectiveness of each N -representability condition. For all the

Table 2: Sizes, required time and memory to solve the SDPs (imposing the $(P,Q,G,T1,T2)$ conditions) as a function of the discretization basis size r for small- and medium-scale problems using SeDuMi.

Basis size r	conditions	# constraints m	Sizes of the largest block matrices	Time	Memory (GB)
10	P,Q,G	465	50x1,25x4,10x4,5x4	11s	0.0
	$P,Q,G,T1$	465	50x3,25x4,10x6,5x4	10s	0.0
	$P,Q,G,T1,T2$	465	175x2,50x5,25x4,10x6	86s	0.1
12	P,Q,G	948	72x1,36x4,15x4,6x4	2.3min	0.1
	$P,Q,G,T1$	948	90x2,72x1,36x4,20x2	2.8min	0.1
	$P,Q,G,T1,T2$	948	306x2,90x4,72x1,36x4	17min	0.1
14	P,Q,G	1743	98x1,49x4,21x4,7x4	13min	0.1
	$P,Q,G,T1$	1743	147x2,98x1,49x4,35x2	14min	0.1
	$P,Q,G,T1,T2$	1743	490x2,147x4,98x1,49x4	1.4h	0.2
16	P,Q,G	2964	128x1,64x4,28x4,8x4	41min	0.3
	$P,Q,G,T1$	2964	224x2,128x1,64x4,56x2	1.4h	0.3
	$P,Q,G,T1,T2$	2964	736x2,224x4,128x1,64x4	6.4h	0.4
18	P,Q,G	4743	162x1,81x4,36x4,9x4	1.9h	0.6
	$P,Q,G,T1$	4743	324x2,162x1,84x2,81x4	2.7h	0.7
	$P,Q,G,T1,T2$	4743	1053x2,324x4,162x1,84x2	12h	1.0

tables that follow, “ r ” is the discretization basis size, “basis” is the spin orbital (one-electron) basis, “state” is the equilibrium state of the system, “ $N(N_\alpha)$ ” is the electron (α spin electron) number, and “ $2S + 1$ ” is the spin multiplicity. For non-atomic systems, it is also necessary to add the repulsion energies to the optimal values of SDPs to obtain the ground state energies. These results are compared with the mainstream electronic structure methods: coupled cluster singles and doubles with perturbational treatment of triples (CCSD(T)) (from Gaussian 98 [16] – column 11), singly and doubly substituted configuration interaction (SDCI) (from Gamess [49] – column 12), and Hartree-Fock (HF) (from Gamess – column 13). The standard for these comparisons is the Full Configuration Interaction method (FCI) (from Gamess – column 14) which essentially consists in computing the minimum eigenvalue of a symmetric matrix with size $\mathcal{O}(r!/N!(r - N!))$. All of the energies are given as a difference between them and the FCI values. Also, in all the tables that follow, the actual discretization basis is from [10, 11, 23, 24, 63], and the experimental geometries for these systems are from [21, 22, 25]. In all calculations using Gaussian 98 and Gamess, we unfroze the core orbitals which are frozen by default. The entry “F/C” means fail to converge.

The RDM method with the (P,Q,G) conditions gives better results than the classic HF. With the $(P,Q,G,T1)$ conditions we get improvements, but imposing the $(P,Q,G,T1,T2)$ conditions, the results are clearly better than the best traditional electronic structure method CCSD(T) (from Gaussian 98). One of the great advantages of the RDM method compared to the traditional electronic structure methods is that it is more numerically robust in the sense that the SDPs can be solved without tuning or sensitive parameter setting

Table 3: The ground state energies (in difference from that of FCI) calculated by the RDM method adding the (P,Q,G) , $(P,Q,G,T1)$, $(P,Q,G,T2)$, and $(P,Q,G,T1,T2)$ conditions (columns 7–10), and those obtained by CCSD(T), SDCI, and HF (columns 11–13) from Gamess and Gaussian 98. The last column shows the FCI results. The energy and the energy differences are in Hartree ($= 4.3598 \times 10^{-18}$ Joules). SDPs solved by SeDuMi.

r	System	Basis	State	$N(N_\alpha)$	$2S+1$	ΔE_{PQG}	ΔE_{PQGT1}	ΔE_{PQGT2}	$\Delta E_{PQGT1T2}$	$\Delta E_{CCSD(T)}$	ΔE_{SDCI}	ΔE_{HF}	E_{FCI}
10	Li	STO-6G	2S	3(2)	2	-0.0000	-0.0000	-0.0000	-0.0000	+0.0000	-0.0000	+0.0003	-7.4002
10	Be	STO-6G	1S	4(2)	1	-0.0000	-0.0000	-0.0000	-0.0000	+0.0000	+0.0000	+0.0527	-14.5561
12	H ₃	double- ζ	$^2A'_1$	3(2)	2	-0.0007	-0.0005	-0.0000	-0.0000	F/C	†+0.0001	+0.0314	-1.4861
12	BeH ⁺	STO-6G	$^1\Sigma^+$	4(2)	1	-0.0000	-0.0000	-0.0000	-0.0000	+0.0000	+0.0000	+0.0204	-14.8433
14	NH ₂ ⁻	STO-6G	1A_1	10(5)	1	-0.0020	-0.0013	-0.0000	-0.0000	+0.0000	+0.0007	+0.0454	-55.1607
14	FH ₂ ⁺	STO-6G	1A_1	10(5)	1	-0.0011	-0.0005	-0.0000	-0.0000	+0.0001	+0.0006	+0.0416	-99.8294
16	CH ₃ ⁺	STO-6G	$^1E'$	8(4)	1	-0.0135	-0.0038	-0.0002	-0.0002	+0.0002	+0.0016	+0.0596	-39.2147
16	CH ₃	STO-6G	$^2A'_2$	9(5)	2	-0.0105	-0.0018	-0.0001	-0.0001	F/C	+0.0016	+0.0631	-39.5178
16	NH ₃ ⁺	STO-6G	$^2A'_2$	9(5)	2	-0.0098	-0.0018	-0.0002	-0.0002	F/C	+0.0015	+0.0618	-55.7924
18	Be	split-valence	1S	4(2)	1	-0.0001	-0.0000	-0.0000	-0.0000	+0.0000	+0.0000	+0.0447	-14.6156
18	CH ₄	STO-6G	1A_1	10(5)	1	-0.0195	-0.0041	-0.0002	-0.0002	+0.0001	+0.0027	+0.0802	-40.1906
18	NH ₄ ⁺	STO-6G	1A_1	10(5)	1	-0.0170	-0.0041	-0.0002	-0.0002	+0.0001	+0.0028	+0.0829	-56.4832
18	Na	STO-6G	2S	11(6)	2	-0.0010	-0.0004	-0.0000	-0.0000	-0.0001	+0.0014	+0.0430	-161.0770

† from Gaussian 98 since Gamess did not converge, “F/C” fail to converge.

required by the traditional electronic structure methods. We are mostly interested in solving highly-correlated systems, and CCSD(T) is frequently used for such systems. However, CCSD(T) and its family of methods have severe limitations for such systems and may not reflect a physical situation, for example, due to the spin contaminations. CCSD(T) solves a nonlinear eigenvalue problem so that there are systems which are hard to solve or do not converge (H₃, CH₃, NH₃⁺ in Tables 3 and 6), or due to its non-variational nature, the energy can get lower than the FCI energy (Na, LiOH in Tables 3 and 6). Unfortunately, the RDM method is not competitive in terms of time since heuristic based electronic structure methods provide results in a few seconds.

The RDM method with $(P,Q,G,T1,T2)$ conditions provides a more reliable approximation of the ground state energy than using only the (P,Q,G) conditions if we pay a price for the computational time and memory as shown in Table 2. However their complexity in terms of floating point operations per iteration (of the interior-point method) and total memory usage are the same: $\mathcal{O}(r^{12})$ and $\mathcal{O}(r^8)$, respectively (see Section 3).

It is interesting to comment here that the RDM method, through an SDP relaxation, can always derive an extremely good lower bound for the ground state energy in polynomial time in r , while the targeting value from the FCI is only computable in factorial time in N and in a fixed discretization basis r . At the same time, though, it is quite impressive that some electronic structure methods like CCSD(T) can often provide comparably good values in a much shorter time.

Observe from Table 3 that we usually require at least 7 digits of accuracy for the optimal value of the SDP for systems with less than -100.0 Hartrees of energy. This means that, adding to the difficulty of solving large-scale SDPs, we need highly precise optimal values and solutions. This particular requirement apparently excludes the possibility of using methods such as the bundle method, Krylov iterative methods or nonlinear formulations (see refs. in [57, 52, 53, 2]).

The *dipole moment* $\langle \hat{\mu} \rangle$ is defined as the norm of $(\langle \hat{\mu}_x \rangle, \langle \hat{\mu}_y \rangle, \langle \hat{\mu}_z \rangle)$, *i.e.*,

$$\langle \hat{\mu} \rangle = \sqrt{\langle \hat{\mu}_x \rangle^2 + \langle \hat{\mu}_y \rangle^2 + \langle \hat{\mu}_z \rangle^2},$$

where

$$\langle \hat{\mu}_x \rangle = \langle \boldsymbol{\mu}_x, \boldsymbol{\Gamma}_1 \rangle,$$

$$[\boldsymbol{\mu}_x]_{ij} = \int \psi_i(\mathbf{z}) x \psi_j(\mathbf{z}) d\mathbf{z}, \quad (i, j = 1, 2, \dots, r)$$

and ψ_i ($i = 1, 2, \dots, r$) are the basis function for the discretization. $\langle \hat{\mu}_y \rangle$ and $\langle \hat{\mu}_z \rangle$ are also defined in a similar way.

Table 4: The (nonzero) dipole moments in Debye ($= 3.356 \times 10^{-30}$ Coulomb meters) calculated by the RDM method adding the (P, Q, G) , $(P, Q, G, T1)$, $(P, Q, G, T2)$, and $(P, Q, G, T1, T2)$ conditions (columns 7–10), and those obtained by SDCI, HF and FCI (columns 11–13) from Gamess. SDPs solved by SeDuMi.

r	System	Basis	State	$N(N_\alpha)$	$2S+1$	D_{PQG}	D_{PQGT1}	D_{PQGT2}	$D_{PQGT1T2}$	D_{SDCI}	D_{HF}	D_{FCI}
12	H ₃	double- ζ	${}^2A'_1$	3(2)	2	0.000000	0.000000	0.000000	0.000000	†0.8908	†0.9211	0.859481
12	BeH ⁺	STO-6G	${}^1\Sigma_+$	4(2)	1	3.730358	3.730202	3.729455	3.729455	3.729713	3.979810	3.729456
14	NH ₂ ⁻	STO-6G	1A_1	10(5)	1	1.179561	1.173888	1.178982	1.178952	1.186210	1.190041	1.178952
14	FH ₂ ⁺	STO-6G	1A_1	10(5)	1	2.296445	2.299526	2.303632	2.303690	2.295315	2.465680	2.303915

† from Gaussian 98 instead, since Gamess calculated a higher energy.

In Table 4, we show (only) the nonzero dipole moments $\langle \hat{\mu} \rangle$ of H₃, BeH⁺, NH₂⁻, and FH₂⁺ in Debye. H₃ is a regular triangle shaped molecule, whose dipole moment should be 0. Surprisingly, the RDM method reproduces this, while traditional electronic structure methods calculated nonzero dipole moments. The H₃ system has many local minima, and HF gets trapped at them due to the spatial symmetry, as well as SDCI and FCI which use HF as a reference. In contrast, the RDM method computes the global minimum. In general, the dipole moments from the RDM method with (P, Q, G) conditions are better than from HF, and worse than from SDCI. But with $(P, Q, G, T1, T2)$ conditions, they almost reproduce the FCI results. The dipole moment results for NH₂⁻ have a noteworthy feature. When more conditions are added to constrain the variational space, then the result for the energy must become better and one expects that normally the result for the dipole moment will become better at the same time. However, when the $T1$ condition was added to the (P, Q, G) conditions, the dipole moment result for NH₂⁻ became worse. This suggests that for that system the (P, Q, G) dipole result was in some sense accidentally very good. Summing up, the RDM with $(P, Q, G, T1, T2)$ conditions reproduces good dipole moments and it is a stable and a robust method even when traditional electronic structure methods fail to work.

Finally, the error measures for the approximate optimal solution $(\hat{\mathbf{X}}, \hat{\mathbf{x}}, \hat{\mathbf{S}}, \hat{\mathbf{y}})$ of the SDPs are as follows:

(I) duality gap $\equiv \mathbf{b}^T \hat{\mathbf{y}} - \langle \mathbf{C}, \hat{\mathbf{X}} \rangle - \langle \mathbf{Diag}(\mathbf{c}), \mathbf{Diag}(\hat{\mathbf{x}}) \rangle,$

(II) primal feasibility error $\equiv \max_{p=1,2,\dots,m} |\langle \mathbf{A}_p, \hat{\mathbf{X}} \rangle + \langle \mathbf{Diag}(\mathbf{a}_p), \mathbf{Diag}(\hat{\mathbf{x}}) \rangle - b_p|,$

$$(III) \text{ dual feasibility error} \equiv \max \left\{ \max_{i,j=1,2,\dots,n} \left| [\hat{\mathbf{S}} - \sum_{p=1}^m \mathbf{A}_p \hat{y}_p + \mathbf{C}]_{ij} \right|, \max_{i=1,2,\dots,s} \left| \left[\sum_{p=1}^m \mathbf{a}_p y_p - \mathbf{c} \right]_i \right| \right\},$$

(IV) minimum eigenvalue of $\hat{\mathbf{X}}$,

(V) minimum eigenvalue of $\hat{\mathbf{S}}$.

The largest errors obtained for the instances solved in this section, not necessarily for the same problem, are (I) 6.86×10^{-7} , (II) 2.16×10^{-7} , (III) 0, (IV) 1.93×10^{-9} , and (V) 3.51×10^{-9} . Since they are small values, they guarantee that we are very close to the optimal solution (see [57, 52, 41] for optimality criteria).

Basically, there are two reasons we could not solve larger SDPs by SeDuMi. First, lack of memory caused by the use of MATLAB. Second, the computational time becomes very large for a serial code. Therefore, we solved large-scale SDPs by the parallel code SDPARA [58] using high performance computers in the next subsection.

4.3 Numerical results for large-scale problems

SDPARA [58] is a C++ open source parallel code for solving general SDPs under GNU General Public License. It is an implementation of the primal-dual predictor-corrector infeasible interior-point method. The main ways that SDPARA benefits from parallel computation are the following two routines. In the framework of primal-dual interior-point methods for general SDPs, the most computationally intense routines involve the construction and the solution of a linear equation whose coefficient matrix is known as the *Schur complement matrix* (SCM). A close look at this matrix [58] reveals that each element can be evaluated on a different processor, independently from the others, if each of them stores the input data matrices \mathbf{A}_p ($p = 1, 2, \dots, m$) and the variable matrices \mathbf{X} and \mathbf{S} in their own memory space. This characteristic is well suited for parallel computation. In addition to the evaluation of the SCM, its parallel Cholesky factorization can be done efficiently by a routine provided by ScaLAPACK [1].

We installed SDPARA on two IBM RS/6000 SPs, *seaborg* ($16 \times 375\text{MHz}$ Power3+ with level two cache of size 8MB, and a maximum of 64GB of memory per Nighthawk node) at the National Energy Research Scientific Computing Center (NERSC), and *eagle* ($4 \times 375\text{MHz}$ Power3-II with level two cache of size 8MB, and 2GB of memory per Winterhawk-II thin node) at Oak Ridge National Laboratory. We also installed SDPARA on an IBM pSeries 690, *cheetah* ($32 \times 1.3\text{GHz}$ Power4 with level two cache of size 1.5MB per chip, level three cache of size 32MB, and maximum of 128GB memory per Regatta node) at Oak Ridge National Laboratory. We chose to report the time and the total memory usage for *seaborg* since we performed most of the computation there.

SDPARA was compiled with IBM C++ using the 64-bit addressing option which allows handling more than 2GB of data. We also made two modifications to SDPARA 0.90 [58], which limited the size of SDPs that could be solved to $r = 20$ with $m = 7230$ and $n_{\max} = 1450$ [62]. First, a check point was introduced, permitting a re-start of SDPARA after a certain number of iterations. This was due to a technical restriction on the running time of twelve hours at these multiple-user facilities. Second, the memory storage was changed. SDPARA 0.90 keeps duplicate copies of three type of matrices: the input

data matrices \mathbf{C}, \mathbf{A}_p ($p = 1, 2, \dots, m$), the variable matrices \mathbf{X} and \mathbf{S} , and a considerable number of auxiliary matrices such as $\mathbf{X}^{-1}, \mathbf{S}^{-1}$, various matrix products, and the search direction at each processor. See [58] for details. Storing the input data matrices and the variable matrices at each processor is essential for constructing the SCM elements by parallel processing. The advantage of also storing the auxiliary matrices at each processor is that this reduces communication time, but the disadvantage is the excessive use of local memory. We modified the code to just keep a single copy of the auxiliary matrices at a specific processor. Before evaluating the SCM elements at each iteration of the interior-point method, we transmit copies of only the updated variable matrices from the specific processor to all other processors. We will call this version of the code SDPARA-SMP.

Table 5 shows the great reduction in total memory usage that resulted from this change, where the last column indicates the number of processors used. Furthermore, a reduction in the running time was also achieved, especially for problems with (P, Q, G) and $(P, Q, G, T1)$ conditions, by making a minor improvement in handling zero block matrices. Fortunately, the computational time was not increased by these modifications, mostly because most of communications were done within the node, which shares a common memory space between several processors, and not between different nodes.

Table 5: Sizes, required time, memory, and number of processors to solve the SDP (imposing the $(P, Q, G, T1, T2)$ conditions) as a function of the discretization basis size r using SDPARA 0.90 and SDPARA-SMP.

Basis size r	conditions	# constraints m	Sizes of the largest block matrices	SDPARA 0.90		SDPARA-SMP		# proc.
				Time	Memory (GB)	Time	Memory (GB)	
10	P, Q, G	465	50x1, 25x4, 10x4, 5x4	5.8s	0.2	5.3s	0.2	16
	$P, Q, G, T1$	465	50x3, 25x4, 10x6, 5x4	9.7s	0.2	8.0s	0.2	16
	$P, Q, G, T1, T2$	465	175x2, 50x5, 25x4, 10x6	37s	0.5	36s	0.2	16
12	P, Q, G	948	72x1, 36x4, 15x4, 6x4	16s	0.3	13s	0.2	16
	$P, Q, G, T1$	948	90x2, 72x1, 36x4, 20x2	26s	0.3	21s	0.2	16
	$P, Q, G, T1, T2$	948	306x2, 90x4, 72x1, 36x4	3.2min	1.1	3.1min	0.4	16
14	P, Q, G	1743	98x1, 49x4, 21x4, 7x4	45s	0.4	37s	0.3	16
	$P, Q, G, T1$	1743	147x2, 98x1, 49x4, 35x2	1.7min	0.5	1.3min	0.4	16
	$P, Q, G, T1, T2$	1743	490x2, 147x4, 98x1, 49x4	15min	2.6	15min	0.8	16
16	P, Q, G	2964	128x1, 64x4, 28x4, 8x4	2.1min	0.6	1.7min	0.5	16
	$P, Q, G, T1$	2964	224x2, 128x1, 64x4, 56x2	4.3min	1.0	3.6min	0.6	16
	$P, Q, G, T1, T2$	2964	736x2, 224x4, 128x1, 64x4	55min	5.6	54min	1.5	16
18	P, Q, G	4743	162x1, 81x4, 36x4, 9x4	6.9min	1.0	5.7min	0.9	16
	$P, Q, G, T1$	4743	324x2, 162x1, 84x2, 81x4	15min	1.9	12min	1.1	16
	$P, Q, G, T1, T2$	4743	1053x2, 324x4, 162x1, 84x2	3.3h	11.2	3.3h	2.9	16
20	P, Q, G	7230	200x1, 100x4, 45x4, 10x4	19min	1.8	16min	1.6	16
	$P, Q, G, T1$	7230	450x2, 200x1, 120x2, 100x4	37min	3.5	34min	2.0	16
	$P, Q, G, T1, T2$	7230	1450x2, 450x4, 200x1, 120x2	14h	27.2	13h	5.7	16
22	P, Q, G	10593	242x1, 121x4, 55x4, 11x4	1.3h	3.3	56min	2.9	16
	$P, Q, G, T1$	10593	605x2, 242x1, 165x2, 121x4	2.3h	6.3	2.0h	3.6	16
	$P, Q, G, T1, T2$	10593	1936x2, 605x4, 242x1, 165x2	*	48.4	2.0days	10.2	16
24	P, Q, G	15018	288x1, 144x4, 66x4, 12x4	3.2h	5.8	2.3h	5.3	16
	$P, Q, G, T1$	15018	792x2, 288x1, 220x2, 144x4	7.5h	10.9	6.9h	6.4	16
	$P, Q, G, T1, T2$	15018	2520x2, 792x4, 288x1, 220x2	*	*	3.3days	26.3	32
26	P, Q, G	20709	338x1, 169x4, 78x4, 13x4	8.3h	10.2	6.2h	9.3	16
	$P, Q, G, T1$	20709	1014x2, 338x1, 286x2, 169x4	21h	18.5	21h	11.2	16
	$P, Q, G, T1, T2$	20709	3211x2, 1014x4, 338x1, 286x2	*	*	5.4days	73.9	64

“*” memory was exceeded or the running time would have been excessive.

Another limitation in using SDPARA-SMP is that it does not handle equality constraints in the dual SDP (19) as SeDuMi does. Therefore, we introduced a small perturbation into the formulation which is equivalent to a further relaxation of the problem (19) [62]. Equalities like $\langle \mathbf{\Gamma}_1, \mathbf{I} \rangle = N$ were all replaced by $-\epsilon \leq \langle \mathbf{\Gamma}_1, \mathbf{I} \rangle - N$, and $\langle \mathbf{\Gamma}_1, \mathbf{I} \rangle - N \leq \epsilon$, where ϵ was fixed to 10^{-5} for SDP relaxations with $(P, Q, G, T2)$ or $(P, Q, G, T1, T2)$ conditions and $r \geq 16$, and 10^{-7} otherwise.

Table 6: The ground state energies (in difference from that of the FCI) calculated by the RDM method adding the (P,Q,G) , $(P,Q,G,T1)$, $(P,Q,G,T2)$, and $(P,Q,G,T1,T2)$ conditions (columns 7–10), and those obtained by CCSD(T), SDCI, and HF (columns 11–13) from Gamess and Gaussian 98. The last column shows the FCI results. The energy and the energy differences are in Hartree ($= 4.3598 \times 10^{-18}$ Joules). SDPs solved by SDPARA-SMP.

r	System	Basis	State	$N(N_\alpha)$	$2S+1$	ΔE_{PQG}	ΔE_{PQGT1}	ΔE_{PQGT2}	$\Delta E_{PQGT1T2}$	$\Delta E_{CCSD(T)}$	ΔE_{SDCI}	ΔE_{HF}	E_{FCI}
10	Li	STO-6G	2S	3(2)	2	-0.0000	-0.0000	-0.0000	-0.0000	+0.0000	-0.0000	+0.0003	-7.4002
10	Be	STO-6G	1S	4(2)	1	-0.0000	-0.0000	-0.0000	-0.0000	+0.0000	+0.0000	+0.0527	-14.5561
12	H ₃	double- ζ	$^2A'_1$	3(2)	2	-0.0008	-0.0006	-0.0000	-0.0000	F/C	†+0.0001	+0.0314	-1.4861
12	BeH ⁺	STO-6G	$^1\Sigma^+$	4(2)	1	-0.0000	-0.0000	-0.0000	-0.0000	+0.0000	+0.0000	+0.0204	-14.8433
14	NH ₂ ⁻	STO-6G	1A_1	10(5)	1	-0.0020	-0.0013	-0.0000	-0.0000	+0.0000	+0.0007	+0.0454	-55.1607
14	FH ₂ ⁺	STO-6G	1A_1	10(5)	1	-0.0011	-0.0005	-0.0000	-0.0000	+0.0001	+0.0006	+0.0416	-99.8294
16	CH ₃ ⁺	STO-6G	$^1E'$	8(4)	1	-0.0135	-0.0038	-0.0003	-0.0003	+0.0002	+0.0016	+0.0596	-39.2147
16	CH ₃	STO-6G	$^2A'_2$	9(5)	2	-0.0106	-0.0018	-0.0006	-0.0006	F/C	+0.0016	+0.0631	-39.5178
16	NH ₃ ⁺	STO-6G	$^2A'_2$	9(5)	2	-0.0098	-0.0018	-0.0005	-0.0005	F/C	+0.0015	+0.0618	-55.7924
18	Be	split-valence	1S	4(2)	1	-0.0001	-0.0001	-0.0001	-0.0000	+0.0000	+0.0000	+0.0447	-14.6156
18	CH ₄	STO-6G	1A_1	10(5)	1	-0.0195	-0.0041	-0.0004	-0.0004	+0.0001	+0.0027	+0.0802	-40.1906
18	NH ₃ ⁺	STO-6G	1A_1	10(5)	1	-0.0171	-0.0041	-0.0004	-0.0004	+0.0001	+0.0028	+0.0829	-56.4832
18	Na	STO-6G	2S	11(6)	2	-0.0010	-0.0005	-0.0001	-0.0001	-0.0001	+0.0014	+0.0430	-161.0770
20	C	double- ζ	3P	6(4)	3	-0.0039	-0.0031	-0.0010	-0.0009	+0.0002	+0.0011	+0.0520	-37.7365
20	O	double- ζ	1D	8(4)	1	-0.0187	-0.0140	-0.0021	-0.0019	+0.0028	+0.0144	+0.1088	-74.7873
20	Ne	double- ζ	1S	10(5)	1	-0.0067	-0.0026	-0.0010	-0.0007	-0.0000	+0.0042	+0.1165	-128.6388
22	HLi ₂	STO-6G	$^2A'_1$	7(4)	2	-0.0010	-0.0007	-0.0002	-0.0002	+0.0002	+0.0005	+0.0235	-15.4055
22	LiOH	STO-6G	$^1\Sigma^+$	12(6)	1	-0.0086	-0.0040	-0.0007	-0.0007	-0.0005	+0.0109	+0.0940	-82.6484
22	HN ₂ ⁺	STO-6G	$^1\Sigma^+$	14(7)	1	-0.0259	-0.0119	-0.0023	-0.0023	+0.0014	+0.0144	+0.1712	-108.9307
22	HNO	STO-6G	$^1A'$	16(8)	1	-0.0190	-0.0136	-0.0010	-0.0010	+0.0012	+0.0093	+0.1499	-129.4479
24	LiH	double- ζ	$^1\Sigma^+$	4(2)	1	-0.0003	-0.0002	-0.0001	—	+0.0000	+0.0002	+0.0276	-8.0087
24	BH	double- ζ	$^1\Sigma^+$	6(3)	1	-0.0065	-0.0047	-0.0006	—	+0.0003	+0.0034	+0.0740	-25.1877
24	HF	double- ζ	$^1\Sigma^+$	10(5)	1	-0.0116	-0.0058	-0.0003	-0.0003	+0.0003	+0.0134	+0.1383	-100.1603
26	CH ₃ N	STO-6G	1A_1	16(8)	1	-0.0385	-0.0164	-0.0013	-0.0013	+0.0007	+0.0113	+0.1574	-93.8845

† from Gaussian 98 since Gamess did not converge, “F/C” fail to converge, “—” not computed.

Table 6 gives the ground state energy for all systems we solved using SDPARA-SMP, including the small- and medium-scale ones we solved previously using SeDuMi. The basic conclusions about the quality of the results of the RDM method compared to the traditional electronic structure methods are the same as previously stated. A comparison between this table and Table 3 shows that the small perturbations we included in the formulation can lower the energy in some cases as much as 0.0005 Hartrees (CH₃ with $(P,Q,G,T1,T2)$ conditions), which is still acceptable but not desirable. On the other hand, this means that the actual energies obtained by the SDPs especially imposing the $(P,Q,G,T1,T2)$ conditions with equality constraints should be slightly higher than shown in Table 6, and they still must give comparably better results than CCSD(T).

In particular, we believe that we solved the largest SDP found in the literature so far ($m = 20709$, largest block matrix $n_{\max} = 3211$) with this density and accuracy. Larger problems could not be solved because we had limited access to these high performance computers.

Table 7 shows the nonzero dipole moments for the corresponding molecules. We derive the same conclusions as in Section 4.2. Here, NH₂⁻ and HNO with $(P,Q,G,T1)$ conditions and LiOH with $(P,Q,G,T2)$ conditions contradict our expectation that adding more N -representability conditions into the variational space, we get better approximations for the dipole moments (from FCI).

In Table 8, we show the occupation number calculated by the RDM method with (P,Q,G) , $(P,Q,G,T1)$, and $(P,Q,G,T1,T2)$ conditions and compared to HF’s and FCI’s ones for CH₃ and CH₄. The *occupation number* is defined by the eigenvalue of the density

Table 7: The (nonzero) dipole moments in Debye ($= 3.3356 \times 10^{-30}$ Coulomb meters) calculated by the RDM method adding the (P,Q,G) , $(P,Q,G,T1)$, $(P,Q,G,T2)$, and $(P,Q,G,T1,T2)$ conditions (columns 7–10), and those obtained by SDCl, HF and FCI (columns 11–13) from Gamess. SDPs solved by SDPARA-SMP.

r	System	Basis	State	$N(N_\alpha)$	$2S+1$	D_{PQG}	D_{PQGT1}	D_{PQGT2}	$D_{PQGT1T2}$	D_{SDCl}	D_{HF}	D_{FCI}
12	H ₃	double- ζ	${}^2A'_1$	3(2)	2	0.000000	0.000000	0.000000	0.000000	†0.8908	†0.9211	0.859481
12	BeH ⁺	STO-6G	${}^1\Sigma_+$	4(2)	1	3.730328	3.730157	3.729477	3.729459	3.729713	3.979810	3.729456
14	NH ₂ ⁻	STO-6G	1A_1	10(5)	1	1.179587	1.173864	1.178975	1.178942	1.186210	1.190041	1.178952
14	FH ₂ ⁺	STO-6G	1A_1	10(5)	1	2.296407	2.299562	2.303633	2.303682	2.295315	2.465680	2.303915
22	HLi ₂	STO-6G	2A_1	7(4)	2	0.561751	0.570898	0.574802	0.574843	0.582754	0.529818	0.575991
22	LiOH	STO-6G	${}^1\Sigma_+$	12(6)	1	0.214317	0.224562	0.318120	0.318062	0.277222	1.994491	0.330520
22	HN ₃ ⁺	STO-6G	${}^1\Sigma_+$	14(7)	1	3.086371	3.130303	3.168295	3.168292	3.290407	3.161953	3.170068
22	HNO	STO-6G	${}^1A'$	16(8)	1	1.212286	1.211767	1.249927	1.250139	1.286756	1.467859	1.254617
24	LiH	double- ζ	${}^1\Sigma_+$	4(2)	1	5.537220	5.541333	5.547012	—	5.572690	5.937220	5.548159
24	BH	double- ζ	${}^1\Sigma_+$	6(3)	1	1.558380	1.565279	1.593296	—	1.662481	2.030271	1.594179
24	HF	double- ζ	${}^1\Sigma_+$	10(5)	1	2.255452	2.260646	2.281847	2.281844	2.283955	2.378904	2.282259
26	CH ₃ N	STO-6G	1A_1	16(8)	1	1.625011	1.662642	1.701920	1.701928	1.715631	1.871751	1.706174

† from Gaussian 98 instead, since Gamess calculated a higher energy, “—” not computed.

matrix $\mathbf{D}_1(i, i')$:

$$\mathbf{D}_1(i, i') = \sum_{\sigma=\alpha,\beta} \Gamma_1(i\sigma, i'\sigma), \quad (i, i' = 1, 2, \dots, r/2) \quad (20)$$

where σ is the spin variable. We chose CH₃ because it is an open shell systems, and CH₄ because unusually bad results were obtained by previous investigations [42, 43]. The occupation number of HF is always 0, 1 or 2. Apparently, the RDM method overestimates the occupation numbers when the FCI’s ones are close to zero, and underestimates otherwise. Adding the $T1$ and $T2$ conditions to the (P,Q,G) conditions, we recover the FCI’s occupation numbers. Observe that the core orbitals (close to integral values) are almost inactive and unaffected by the approximation levels.

Finally, we give the error measures for the approximate optimal solution $(\hat{\mathbf{X}}, \hat{\mathbf{S}}, \hat{\mathbf{y}})$ for the SDPs. Now that we do not have the equality constraints in the dual SDP (19), the errors (I), (II) and (III) can be restated as follows:

$$(I') \text{ duality gap} \equiv \mathbf{b}^T \hat{\mathbf{y}} - \langle \mathbf{C}, \hat{\mathbf{X}} \rangle,$$

$$(II') \text{ primal feasibility error} \equiv \max_{p=1,2,\dots,m} |\langle \mathbf{A}_p, \hat{\mathbf{X}} \rangle - b_p|,$$

$$(III') \text{ dual feasibility error} \equiv \max_{i,j=1,2,\dots,n} |[\hat{\mathbf{S}} - \sum_{p=1}^m \mathbf{A}_p \hat{y}_p + \mathbf{C}]_{ij}|.$$

The largest errors obtained for the instances solved in this section, not necessary for the same problem, are (I') 1.73×10^{-5} , (II') 1.28×10^{-6} , (III') 4.48×10^{-13} , (IV) 2.27×10^{-10} , and (V) 3.85×10^{-12} .

4.4 Considerations on alternative methods

The small perturbations we introduced into the formulation, splitting one equality constraint into two inequality constraints, as explained at Section 4.3, are not desirable. Instead, we

Table 8: Occupation number for CH_3 and CH_4 computed from the RDM method with (P,Q,G) , $(P,Q,G,T1)$, $(P,Q,G,T1,T2)$ conditions compared to those from HF and FCI. SDPs solved by SDPARA-SMP.

r	System	Basis	State	$N(N_\alpha)$	$2S+1$	PQG	$PQGT1$	$PQGT1T2$	HF	FCI
16	CH_3	STO-6G	${}^2A_2''$	9(5)	2	0.026261	0.021318	0.020702	0	0.0205
						0.026261	0.021318	0.020702	0	0.0205
						0.028585	0.028945	0.028186	0	0.0279
						0.999989	1.000492	1.000042	1	1.0000
						1.972254	1.975063	1.976044	2	1.9763
						1.972254	1.975063	1.976044	2	1.9763
						1.974420	1.977839	1.978330	2	1.9784
						1.999976	1.999963	1.999959	2	2.0000
18	CH_4	STO-6G	1A_1	10(5)	1	0.025979	0.021372	0.020305	0	0.0201
						0.027267	0.023940	0.022851	0	0.0227
						0.027267	0.023941	0.022852	0	0.0227
						0.027267	0.023941	0.022852	0	0.0227
						1.972743	1.974314	1.975499	2	1.9757
						1.972743	1.974314	1.975499	2	1.9757
						1.972743	1.974315	1.975499	2	1.9757
						1.974026	1.983911	1.984701	2	1.9848
	1.999963	1.999952	1.999952	2	2.0000					

tried to eliminate some variables (at \mathbf{y} in (19)) using these equalities as equations, producing an equivalent SDP with fewer variables and no equality constraints. Preliminary numerical experiments demonstrated, however, that these linear transformations introduce undesirable numerical properties into the problem and SDPARA was not able to get enough accuracy [61, Section 5.3.3]. Therefore, incorporation of equality constraints as a standard option, as done in SeDuMi and SDPT3, certainly is a desirable addition to SDPARA’s capability.

Alternative methods such as discussed by [41] may be worth considering, but we have felt up until now that they are not able to deliver the accuracy that we require for this application. This is certainly our experience with the spectral bundle method; early experiments reported in [47] indicated that is very difficult to obtain satisfactory accuracy. We also experimented with the new code SDPLR 1.01 [2] which combines an augmented Lagrangian technique with limited-memory BFGS. However, we even could not solve the smallest problems to the accuracy that we need since the number of internal limited-memory BFGS iterations increases prohibitively as the optimal solution is approached. Surprisingly, Mazziotti [36, 37] very recently announced some results for larger systems ($r = 36$, with $m = 468684$ and $n_{\max} = 648$, using only the (P,Q,G) conditions), for which he solved the SDPs by a method similar to that used in SDPLR.

The use of the conjugate gradient method to solve the SCM system or other iterative methods to solve the related indefinite “augmented system” (see [53]) could be a further alternative, but the extreme ill-conditioning of these linear systems makes it very difficult to obtain the accuracy that we need. It is possible that eliminating some of the degeneracies in the system could lead to improved performance of these methods.

5 Conclusion and further directions

The RDM method, which provides a lower bound for the ground state energy of a many-electron system subject to a given external potential, can be formulated as an SDP problem through the known $(P, Q, G, T1, T2)$ N -representability conditions. The new formulation presented here as a dual SDP (19) seems the most suitable one for the state-of-art software to solve general SDPs. The numerical experiments carried out since 2001, including the ones reported here, demonstrate for the first time the quality, the strength, and the actual effectiveness of the N -representability conditions known for more than forty years in electronic structure calculation. In fact, they demonstrate that the RDM method with the $(P, Q, G, T1, T2)$ conditions can give better ground state energies than the current electronic structure methods, although it is not competitive in terms of time at least at present. It also has the advantage of robust convergence which is not the case for the traditional electronic structure methods. In addition, our results for the dipole moment confirm that the RDM itself is computed with excellent accuracy compared with traditional wavefunction-based methods.

We also report results for the largest problems in literature using the $(P, Q, G, T1, T2)$ conditions with discretization basis size $r = 26$ ($m = 20709$, $n_{\max} = 3211$), while the previous ones were $r = 20$ ($m = 7230$, $n_{\max} = 1450$) [62]. The SDPs which arise from this application can be arbitrarily large, and may require special techniques for their solution. Parallel computation and large memory management are indispensable. In fact, it seems that we will always face a dual hardware limitation in solving large-scale problems: time and memory, both of which depend on the number of available processors and physical memory.

The recent series of numerical results for this application opens up a whole research field which was once very active, and at the same time raises many questions for future investigations.

Some fundamental questions for physicists are the search for new N -representability conditions and understanding the role of the known conditions. Chemists might be interested in understanding why the same $(P, Q, G, T1, T2)$ conditions provide very good approximations for some systems and not for others, and also in studying many desirable properties obtainable by this unique method, like dissociation processes of highly-correlated systems having multiple bonds and high spin states, which are difficult to calculate. And finally, optimizers have the challenge of solving larger SDPs with $m > 20000$ and $n > 3000$ with high accuracy. However, it is certain that novel algorithms and exploration of new physical properties of the N -representability conditions are necessary in order for the RDM method becomes practical.

As a final observation, we recognize that there is a need to provide physicists/chemists easy-to-use black box SDP solvers based on their own terminology.

Acknowledgments

This work would not have been possible without the support of the National Science Foundation (NSF) via Grant No. ITR-DMS 0113852, which provided essential funding for M.F. and B.J.B, summer support for Z.Z., and short-term visiting support to New York University for M.N. and M.Y. The authors M.F. and M.N. were also supported by fellowships from the Japan Society for the Promotion of Science and Grant-in-Aids for Scientific Research

from the Japanese Ministry of Education, Culture, Sports, Science and Technology, B.J.B by Department of Energy (DOE) Grant No. DE-FG02-86ER53223, M.L.O. by NSF Grant No. DMS-0412049, J.K.P. by DOE Grant No. DE-FG02-02ER15292, and Z.Z. by a teaching fellowship at the Department of Physics, New York University. The numerical experiments used resources of NERSC (National Energy Research Scientific Computing Center), which is supported by the Office of Science of the Department of Energy under Contract No. DE-AC03-76SF00098, and also of the Center for Computational Sciences at Oak Ridge National Laboratory, which is supported by the Office of Science of the Department of Energy under Contract DE-AC05-00OR22725. We also thank Masakazu Kojima from Tokyo Institute of Technology for allowing us to use the Pentium Xeon machine.

References

- [1] Blackford, L.S., Choi, J., Cleary, A., D’Azevedo E., Demmel, J., Dhillon, I., Dongarra, J., Hammarling, S., Henry, G., Petitet, A., Stanley, K., Walker, D., Whaley, R.C.: ScaLAPACK User’s Guide. SIAM, Philadelphia, 1997
- [2] Burer, S., Monteiro, R.D.C., Choi, C.: SDPLR version 1.01 user’s guide (short). Department of Management Sciences, University of Iowa, Iowa City, IA 52242-1000, 2004. <http://dollar.biz.uiowa.edu/~burer/software/SDPLR/>
- [3] Cioslowski, J.: Many-Electron Densities and Reduced Density Matrices. Kluwer Academic/Plenum Publishers, New York, 2000
- [4] Cohen, L., Frishberg, C.: Hierarchy equations for reduced density matrices. *Phys. Rev.* **A13**, 927–930 (1976)
- [5] Coleman, A.J.: Structure of fermion density matrices. *Rev. Mod. Phys.* **35**, 668–687 (1963)
- [6] Coleman, A.J.: RDMs: How did we get here? In: [3], pp. 1–17
- [7] Colmenero, F., Valdemoro, C.: Self-consistent approximate solution of the 2nd-order contracted Schrödinger-equation. *Int. J. Quantum Chem.* **51**, 369–388 (1994)
- [8] Davidson, E.R.: Linear inequalities for density matrices. *J. Math. Phys.* **10**, 725–734 (1969)
- [9] Deza, M.M., Laurent, M.: *Geometry of Cuts and Metrics*. Springer-Verlag, Berlin, 1997
- [10] Dunning, T.H.Jr., Gaussian basis functions for use in molecular calculations. I. Contraction of ($9s5p$) atomic basis sets for the first-row atoms. *J. Chem. Phys.* **53**, 2823–2833 (1970)
- [11] Dunning, T.H.Jr., Hay, P.J.: Gaussian basis sets for molecular calculations. In: *Modern Theoretical Chemistry Vol. 3: Methods of Electronic Structure Theory*, Schaefer III, H.F. (ed.), Plenum, New York, 1977, pp. 1–28

- [12] Ehara, M., Nakata, M., Kou, H., Yasuda, K., Nakatsuji, H.: Direct determination of the density matrix using the density equation: Potential energy curves of HF, CH₄, BH₃, NH₃, and H₂O. *Chem. Phys. Lett.* **305**, 483–488 (1999)
- [13] Erdahl, R.M.: Representability. *Int. J. Quantum Chem.* **13**, 697–718 (1978)
- [14] Erdahl, R.M., Jin, B.: The lower bound method for reduced density matrices. *Journal of Molecular Structure: THEOCHEM* **527**, 207–220 (2000)
- [15] Erdahl, R.M., Jin, B.: On calculating approximate and exact density matrices. In: [3], pp. 57–84
- [16] Frisch, M.J., Trucks, G.W., Schlegel, H.B., *et al.*: Gaussian 98, Revision A.11.3, Gaussian, Inc., Pittsburgh PA, 2002
- [17] Fujisawa, K., Nakata, K., Kojima, M.: SDPA (SemiDefinite Programming Algorithm) – user’s manual – version 5.00. Research Report B-308, Department of Mathematical and Computing Sciences, Tokyo Institute of Technology, Tokyo, Japan, 1999
- [18] Garrod, C., Fusco, M.A.: A density matrix variational calculation for atomic Be. *Int. J. Quantum Chem.* **10**, 495–510 (1976)
- [19] Garrod, C., Mihailović, M.V., Rosina, M.: The variational approach to the two-body density matrix. *J. Math. Phys.* **16**, 868–874 (1975)
- [20] Garrod, C., Percus, J.K.: Reduction of the N -particle variational problem. *J. Math. Phys.* **5**, 1756–1776 (1964)
- [21] Graner, G., Hirota, E., Iijima, T., Kuchitsu, K., Ramsay, D.A., Vogt, J., Vogt, N.: Landolt-Börnstein - Group II Molecules and Radicals, Volume 25, Subvolume A. Springer-Verlag, Heidelberg, 1998
- [22] Graner, G., Hirota, E., Iijima, T., Kuchitsu, K., Ramsay, D.A., Vogt, J., Vogt, N.: Landolt-Börnstein - Group II Molecules and Radicals, Volume 25, Subvolume B. Springer-Verlag, Heidelberg, 1999
- [23] Hehre, W.J., Ditchfield, R., Stewart, R.F., Pople, J.A.: Self-consistent molecular orbital methods. IV. Use of Gaussian expansions of slater-type orbitals. Extension to second-row molecules. *J. Chem. Phys.* **52**, 2769–2773 (1970)
- [24] Hehre, W.J., Stewart, R.F., Pople, J.A.: Self-consistent molecular-orbital methods. I. Use of Gaussian expansions of slater-type atomic orbitals. *J. Chem. Phys.* **51**, 2657–2664 (1969)
- [25] Huber, K.P., Herzberg, G.: *Molecular Spectra and Molecular Structure IV, Electronic Constants of Diatomic Molecules*. Van Nostrand Reinhold, New York, 1979
- [26] Husimi, K.: Some formal properties of the density matrix. *Proc. Phys. Math. Soc. Jpn.* **22**, 264–314 (1940)

- [27] Karp, R.M., Papadimitriou C.H.: On linear characterization of combinatorial optimization problems. *SIAM J. Comput.* **11**, 620–632 (1982)
- [28] Kijewski, L.J.: Effectiveness of symmetry and the Pauli condition on the 1 matrix in the reduced-density-matrix variational principle. *Phys. Rev.* **A6**, 31–35 (1972)
- [29] Kijewski, L.J.: Strength of the G -matrix condition in the reduced-density-matrix variational principle. *Phys. Rev.* **A9**, 2263–2266 (1974)
- [30] Löwdin, P.-O.: Quantum theory of many-particle systems. I. Physical interpretations by means of density matrices, natural spin-orbitals, and convergence problems in the method of configurational iteration. *Phys. Rev.* **97**, 1474–1489 (1955)
- [31] Mayer, J.E.: Electron correlation. *Phys. Rev.* **100**, 1579–1586 (1955)
- [32] Mazziotti, D.A.: Contracted Schrödinger equation: Determining quantum energies and two-particle density matrices without wave functions. *Phys. Rev.* **A57** 4219–4234 (1998)
- [33] Mazziotti, D.A.: Cumulants and the contracted Schrödinger equation. In: [3], pp. 139–163
- [34] Mazziotti, D.A.: Variational minimization of atomic and molecular ground-state energies via the two-particle reduced density matrix. *Phys. Rev.* **A65**, 062511 (2002)
- [35] Mazziotti, D.A.: Solution of the 1,3-contracted Schrödinger equation through positivity conditions on the two-particle reduced density matrix. *Phys. Rev.* **A66**, 062503 (2002)
- [36] Mazziotti, D.A.: Realization of quantum chemistry without wave functions through first-order semidefinite programming. *Phys. Rev. Lett.* **93**, 213001 (2004)
- [37] Mazziotti, D.A.: First-order semidefinite programming for the direct determination of two-electron reduced density matrices with application to many-electron atoms and molecules. *J. Chem. Phys.* **121**, 10957–10966 (2004)
- [38] Mazziotti, D.A., Erdahl, R.M.: Uncertainty relations and reduced density matrices: Mapping many-body quantum mechanics onto four particles. *Phys. Rev.* **A63**, 042113 (2001)
- [39] McRae, W.B., Davidson, E.R.: Linear inequalities for density matrices II. *J. Math. Phys.* **13**, 1527–1538 (1972)
- [40] Mihailović, M.V., Rosina, M.: The variational approach to the density matrix for light nuclei. *Nucl. Phys.* **A237**, 221–228 (1975)
- [41] Monteiro, R.D.C.: First- and second-order methods for semidefinite programming. *Math. Prog.* **B97**, 209–244 (2003)
- [42] Nakata, M., Ehara, M., Nakatsuji, H.: Density matrix variational theory: Application to the potential energy surfaces and strongly correlated systems. *J. Chem. Phys.* **116**, 5432–5439 (2002)

- [43] Nakata, M., Ehara, M., Nakatsuji, H.: Density matrix variational theory: Strength of Weinhold-Wilson inequalities. In: *Fundamental World of Quantum Chemistry*, Löwdin, P.-O., Kryachko, E.S., (eds.), Kluwer Academic, Boston, 2003, pp. 543–557
- [44] Nakata, M., Nakatsuji, H., Ehara, M., Fukuda, M., Nakata, K., Fujisawa, K.: Variational calculations of fermion second-order reduced density matrices by semidefinite programming algorithm. *J. Chem. Phys.* **114**, 8282–8292 (2001)
- [45] Nakatsuji, H.: Equation for direct determination of density matrix. *Phys. Rev.* **A14**, 41–50 (1976)
- [46] Nakatsuji, H.: Density equation theory in chemical physics. In: [3], pp. 85–116
- [47] Nayakkankuppam, M.V.: Optimization over symmetric cones. PhD thesis, Department of Computer Science, New York University, New York, 1999
- [48] Rosina, M., Garrod C.: The variational calculation of reduced density matrices. *J. Comput. Phys.* **18**, 300-310 (1975)
- [49] Schmidt, M.W., Baldridge, K.K., Boatz, J.A., Elbert, S.T., Gordon, M.S., Jensen, J.H., Koseki, S., Matsunaga, N., Nguyen, K.A., Su, S.J., Windus, T.L., Dupuis, M., Montgomery, J.A.: General atomic and molecular electronic structure system. *J. Comput. Chem.* **14**, 1347-1363 (1993)
- [50] Sturm, J.F.: Using SeDuMi 1.02, a MATLAB toolbox for optimization over symmetric cones. *Optim. Methods Softw.* **11–12**, 625–653 (1999). <http://sedumi.mcmaster.ca/>
- [51] Szabo, A., Ostlund, N.S.: *Modern Quantum Chemistry: Introduction to Advanced Electronic Structure Theory*. Dover Publications, Inc., Mineola, New York, 1996
- [52] Todd, M.J.: Semidefinite optimization. *Acta Numer.* **10**, 515–560 (2001)
- [53] Toh, K.-C.: Solving large scale semidefinite programs via an iterative solver on the augmented systems. *SIAM J. Optim.* **14**, 670–698 (2003)
- [54] Toh, K.-C., Tütüncü, R.H., Todd, M.J.: On the implementation of SDPT3 (version 3.1) – a Matlab software package for semidefinite-quadratic-linear programming. *IEEE Conference on Computer-Aided Control System Design*, 2004
- [55] Valdemoro, C.: Approximating the 2nd-order reduced density-matrix in terms of the 1st-order one. *Phys. Rev.* **A45**, 4462–4467 (1992)
- [56] Valdemoro, C., Tel, L.M., Pérez-Romero, E.: Critical questions concerning iterative solution of the contracted Schrödinger equation. In: [3], pp. 117–138
- [57] Wolkowicz, H., Saigal, R., Vandenberghe, L.: *Handbook of Semidefinite Programming: Theory, Algorithms, and Applications*. Kluwer Academic, Boston, 2000
- [58] Yamashita, M., Fujisawa, K., Kojima, M.: SDPARA : SemiDefiniteProgramming Algorithm paRAllel version. *Parallel Comput.* **29**, 1053–1067 (2003). <http://grid.r.dendai.ac.jp/sdpa/>

- [59] Yasuda, K.: Direct determination of the quantum-mechanical density matrix: Parquet theory. *Phys. Rev.* **A59**, 4133–4149 (1999)
- [60] Yasuda, K., Nakatsuji, H.: Direct determination of the quantum-mechanical density matrix using the density equation II. *Phys. Rev.* **A56**, 2648–2657 (1997)
- [61] Zhao, Z.: The reduced density matrix method for electronic structure calculation – application of semidefinite programming to N -fermion systems. PhD thesis, Department of Physics, New York University, New York, 2004
- [62] Zhao, Z., Braams, B.J., Fukuda, M., Overton, M.L., Percus, J.K.: The reduced density matrix method for electronic structure calculations and the role of three-index representability conditions. *J. Chem. Phys.* **120**, 2095–2104 (2004)
- [63] Basis sets were obtained from the Extensible Computational Chemistry Environment Basis Set Database, Version 02/25/04, as developed and distributed by the Molecular Science Computing Facility, Environmental and Molecular Sciences Laboratory which is part of the Pacific Northwest Laboratory, P.O. Box 999, Richland, Washington 99352, USA, and funded by the U.S. Department of Energy. The Pacific Northwest Laboratory is a multi-program laboratory operated by Battelle Memorial Institute for the U.S. Department of Energy under contract DE-AC06-76RLO 1830. <http://www.emsl.pnl.gov/forms/basisform.html>