

The Strong Second Order Sufficient Condition and Constraint Nondegeneracy in Nonlinear Semidefinite Programming and Their Implications *

Defeng Sun

Department of Mathematics

National University of Singapore, Singapore

Email: matsundf@nus.edu.sg

May 15, 2005

Abstract

For a locally optimal solution to the nonlinear semidefinite programming problem, under Robinson's constraint qualification, the following conditions are proved to be equivalent: the strong second order sufficient condition and constraint nondegeneracy; the nonsingularity of Clarke's Jacobian of the Karush-Kuhn-Tucker system; the strong regularity of the Karush-Kuhn-Tucker point; and others.

1 Introduction

Consider the optimization problem

$$\begin{aligned} (OP) \quad & \min \quad f(x) \\ & \text{s.t.} \quad G(x) \in K, \\ & \quad \quad x \in X, \end{aligned} \tag{1}$$

where $f : X \rightarrow \Re$ and $G : X \rightarrow Y$ are twice continuously differentiable functions, X and Y are two finite dimensional real vector spaces each equipped with a scalar product denoted by $\langle \cdot, \cdot \rangle$ and its induced norm denoted by $\| \cdot \|$, and K is a closed convex set in Y . We reserve Z to represent an arbitrary real vector space with a scalar product $\langle \cdot, \cdot \rangle$. We denote by $\mathcal{J}_x f(x)$ and $\mathcal{J}_{xx}^2 f(x)$ the derivative and the second order derivative of f with respect to

*This research was partially supported by Grant R146-000-061-112 of the National University of Singapore.

$x \in X$, respectively. For any given linear operator A , we denote its adjoint by A^* . The first order optimality condition, namely the Karush-Kuhn-Tucker (KKT) condition, for (OP) takes the following form:

$$\mathcal{J}_x L(x, \mu) = 0 \quad \text{and} \quad \mu \in \mathcal{N}_K(G(x)), \quad (2)$$

where the Lagrangian function $L : X \times Y \rightarrow \Re$ is defined by

$$L(x, \mu) := f(x) + \langle \mu, G(x) \rangle, \quad (x, \mu) \in X \times Y, \quad (3)$$

$\mathcal{J}_x L(x, \mu)$ is the derivative of $L(x, \mu)$ at (x, μ) with respect to $x \in X$, and $\mathcal{N}_K(y)$ denotes the normal cone of K at y in convex analysis [33]:

$$\mathcal{N}_K(y) = \begin{cases} \{d \in Y : \langle d, z - y \rangle \leq 0 \quad \forall z \in K\} & \text{if } y \in K, \\ \emptyset & \text{if } y \notin K. \end{cases}$$

For any (x, μ) satisfying (2), we call x a *stationary point* and (x, μ) a *KKT point* of (OP) , respectively.

During the last three decades, tremendous progress has been achieved towards sensitivity and stability analysis of solutions to the optimization problem (OP) subject to data perturbation [4, 10, 16, 34]. When K is a polyhedral set, the corresponding theory is quite complete. This is especially the case for the conventional nonlinear programming

$$\begin{aligned} (NLP) \quad & \min \quad f(x) \\ & \text{s.t.} \quad h(x) = 0, \\ & \quad \quad g(x) \leq 0, \\ & \quad \quad x \in X, \end{aligned} \quad (4)$$

where $f : X \rightarrow \Re$, $h : X \rightarrow \Re^m$ and $g : X \rightarrow \Re^p$ are twice continuously differentiable functions.

For the case that K is a general nonpolyhedral set, much less has been known. However, when K is \mathcal{C}^2 -cone reducible in the sense of Bonnans and Shapiro [4, Definition 3.135], the full picture of sensitivity and stability of solutions for problem (OP) is emerging [1, 2, 3, 4]. The class of \mathcal{C}^2 -cone reducible sets is rich. It includes notably the polyhedral set, the second order cone (ice-cream cone or Lorentz cone), the cone of symmetric positive semidefinite matrices, and their Cartesian product [4, 36].

Compared to the conventional nonlinear programming (NLP) , the theory for (OP) with \mathcal{C}^2 -cone reducible sets is yet to complete. Let \bar{x} be a feasible solution to (OP) . Robinson's constraint qualification (CQ) [27] is said to hold at \bar{x} if

$$0 \in \text{int}\{G(\bar{x}) + \mathcal{J}_x G(\bar{x})X - K\}, \quad (5)$$

where “int” denotes the topological interior part of a given set. If \bar{x} is a locally optimal solution to (OP) and Robinson’s CQ holds at \bar{x} , then there exists a Lagrangian multiplier $\bar{\mu} \in Y$, together with \bar{x} , satisfying the KKT condition:

$$\mathcal{J}_x L(\bar{x}, \bar{\mu}) = 0 \quad \text{and} \quad \bar{\mu} \in \mathcal{N}_K(G(\bar{x})). \quad (6)$$

For any closed (not necessary convex) set $D \subseteq Y$ and $y \in Y$, denote

$$\text{dist}(y, D) := \inf\{\|y - d\| : d \in D\}.$$

For any closed set $D \subseteq Y$, we write $\mathcal{T}_D^i(y)$ and $\mathcal{T}_D(y)$ for the *inner tangent* cone and the *contingent* (Bouligand) cone of D at y , respectively. That is,

$$\mathcal{T}_D^i(y) = \{d \in Y : \text{dist}(y + td, D) = o(t), t \geq 0\}$$

and

$$\mathcal{T}_D(y) = \{d \in Y : \exists t_k \downarrow 0, \text{dist}(y + t_k d, D) = o(t_k)\}.$$

When D is a closed convex set, the inner tangent cone and the contingent cone are equal:

$$\mathcal{T}_D(y) = \mathcal{T}_D^i(y) = \{d \in Y : \text{dist}(y + td, D) = o(t), t \geq 0\}, \quad y \in D$$

and will be called *tangent cones* directly. Since Y is assumed to be a finite dimensional space and K is a closed convex set, Robinson’s CQ (5) can be equivalently written as

$$\mathcal{J}_x G(\bar{x})X + \mathcal{T}_K(G(\bar{x})) = Y, \quad (7)$$

which reduces to the well known Mangasarian-Fromovitz constraint qualification (MFCQ) for the conventional nonlinear programming (NLP) [20]:

$$\left\{ \begin{array}{l} \mathcal{J}_x h_i(\bar{x}), \quad i = 1, \dots, m, \text{ are linearly independent,} \\ \exists d \in X : \mathcal{J}_x h_i(\bar{x})d = 0, i = 1, \dots, m, \mathcal{J}_x g_j(\bar{x})d < 0, j \in \mathcal{I}(\bar{x}), \end{array} \right. \quad (8)$$

where the active set $\mathcal{I}(\bar{x})$ of $g(\cdot)$ at \bar{x} is defined by

$$\mathcal{I}(\bar{x}) := \{j : g_j(\bar{x}) = 0, j = 1, \dots, p\}.$$

For a proof on this equivalence, see [26, Theorem 3]. A stronger notion than the MFCQ in (NLP) is the linear independence constraint qualification (LICQ):

$$\{\mathcal{J}_x h_i(\bar{x})\}_{i=1}^m \text{ and } \{\mathcal{J}_x g_j(\bar{x})\}_{j \in \mathcal{I}(\bar{x})} \text{ are linearly independent.} \quad (9)$$

Let $\mathcal{M}(\bar{x})$ denote the set of Lagrangian multipliers satisfying (6). Then $\mathcal{M}(\bar{x})$ is nonempty and bounded if and only if Robinson’s CQ holds at \bar{x} [4, Proposition 2.71], which generalizes an analogue of Gauvin [11] for (NLP): $\mathcal{M}(\bar{x})$ is nonempty and bounded if and only if the MFCQ holds \bar{x} . For (NLP), the LICQ implies that $\mathcal{M}(\bar{x})$ is a singleton.

In one of his seminal papers, Robinson [28] introduced an important concept called *strong regularity* for generalized equations, which include the KKT system (6) as a special case, and defined a *strong second order sufficient condition* for (NLP) . He also showed that the strong second order sufficient condition and the LICQ implies the strong regularity of the solution to the KKT system (6). Interestingly, the converse is also true, see [14, 5, 8] and [4, Proposition 5.38].

The primary objective of this paper is to build up the connections between the strong sufficient second order condition and strong regularity for the nonlinear semidefinite programming

$$\begin{aligned}
(NLSDP) \quad & \min \quad f(x) \\
& \text{s.t.} \quad h(x) = 0, \\
& \quad \quad g(x) \in \mathcal{S}_+^p, \\
& \quad \quad x \in X,
\end{aligned} \tag{10}$$

where $f : X \rightarrow \mathbb{R}$, $h : X \rightarrow \mathbb{R}^m$, and $g : X \times \mathcal{S}^p \rightarrow \mathcal{S}^p$ are twice continuously differentiable, \mathcal{S}^p is the linear space of all $p \times p$ real symmetric matrices, and \mathcal{S}_+^p is the cone of all $p \times p$ positive semidefinite matrices. Problem $(NLSDP)$ is a special case of (OP) with

$$G(x) := (h(x), g(x)), \quad x \in X, \quad Y := \mathbb{R}^m \times \mathcal{S}^p, \quad \text{and} \quad K := \{0\} \times \mathcal{S}_+^p. \tag{11}$$

We achieve this via the study of the nonsingularity of generalized Jacobian of the system of nonsmooth equations reformulated from (2). Consequently, we show that if \bar{x} is a locally optimal solution to $(NLSDP)$ and Robinson's CQ holds at \bar{x} , then the nonsingularity of Clarke's Jacobian of the corresponding nonsmooth system is not only sufficient but also necessary for the strong regularity. Since the nonsingularity of Clarke's Jacobian is a stronger condition than many other conditions posed for general nonsmooth equations [18, 24], this actually establishes the equivalence of many conditions discussed in a wide range of literatures for $(NLSDP)$.

The organization of this paper is as follows. In Section 2, we study some useful properties of Clarke's Jacobian for Lipschitz functions, in particular for the metric projector over \mathcal{S}_+^p . We propose a strong second order sufficient condition for the nonlinear semidefinite programming $(NLSDP)$ in Section 3. It is shown that this strong second order sufficient condition and constraint nondegeneracy imply the nonsingularity of Clarke's Jacobian of the corresponding nonsmooth system. The promised equivalent conditions are discussed in Section 4. We conclude this paper by pointing out some possible research topics in Section 5.

2 Jacobian Properties

Let X , Y , and Z be three arbitrary finite dimensional real vector spaces each equipped with a scalar product $\langle \cdot, \cdot \rangle$ and its induced norm $\| \cdot \|$. Let \mathcal{O} be an open set in Y and $\Xi : \mathcal{O} \subseteq Y \rightarrow Z$ be a locally Lipschitz continuous function on the open set \mathcal{O} . Rademacher's theorem says that Ξ is almost everywhere F(réchet)-differentiable in \mathcal{O} . We denote by \mathcal{O}_Ξ the set of points in \mathcal{O} where Ξ is F-differentiable. Then Clarke's generalized Jacobian of Ξ at y is well defined [7]:

$$\partial \Xi(y) := \text{conv}\{\partial_B \Xi(y)\},$$

where “conv” denotes the convex hull and

$$\partial_B \Xi(y) := \{V : V = \lim_{k \rightarrow \infty} \mathcal{J}_y \Xi(y^k), y^k \in \mathcal{O}_\Xi\}.$$

The next lemma is about the generalized Jacobian for composite functions.

Lemma 1 *Let $\Psi : X \rightarrow Y$ be a continuously differentiable function on an open neighborhood \widehat{N} of \bar{x} and $\Xi : \mathcal{O} \subseteq Y \rightarrow Z$ be a locally Lipschitz continuous function on an open set \mathcal{O} containing $\bar{y} := \Psi(\bar{x})$. Suppose that Ξ is directionally differentiable at every point in \mathcal{O} and that $J_x \Psi(\bar{x}) : X \rightarrow Y$ is onto. Then it holds that*

$$\partial_B \Phi(\bar{x}) = \partial_B \Xi(\bar{y}) \mathcal{J}_x \Psi(\bar{x}), \quad (12)$$

where $\Phi : \widehat{N} \rightarrow Z$ is defined by $\Phi(x) := \Xi(\Psi(x))$, $x \in \widehat{N}$.

Proof. By shrinking \widehat{N} if necessary, we may assume that $\Psi(\widehat{N}) \subseteq \mathcal{O}$. Then Φ is Lipschitz continuous and directionally differentiable on \widehat{N} . By further shrinking \widehat{N} if necessary, we may also assume that for each $x \in \widehat{N}$, $\mathcal{J}_x \Psi(x)$ is onto.

We shall first show that Φ is F-differentiable at $x \in \widehat{N}$ if and only Ξ is F-differentiable at $\Psi(x)$, which ensures that

$$\partial_B \Phi(\bar{x}) \subseteq \partial_B \Xi(\bar{y}) \mathcal{J}_x \Psi(\bar{x}).$$

Certainly, Φ is F-differentiable at $x \in \widehat{N}$ if Ξ is F-differentiable at $\Psi(x)$. Now, suppose that Φ is F-differentiable at $x \in \widehat{N}$. Then, since Ξ is directionally differentiable at $\Psi(x)$, for any $d \in X$ we have

$$\mathcal{J}_x \Phi(x) d = \Xi'(\Psi(x); \mathcal{J}_x \Psi(x) d),$$

which implies that for any $s, t \in \Re$ and $u, v \in X$,

$$\begin{aligned} \Xi'(\Psi(x); s \mathcal{J}_x \Psi(x) u + t \mathcal{J}_x \Psi(x) v) &= \Xi'(\Psi(x); \mathcal{J}_x \Psi(x) (su + tv)) \\ &= \mathcal{J}_x \Phi(x) (su + tv) \\ &= s \mathcal{J}_x \Phi(x) u + t \mathcal{J}_x \Phi(x) v \\ &= s \Xi'(\Psi(x); \mathcal{J}_x \Psi(x) u) + t \Xi'(\Psi(x); \mathcal{J}_x \Psi(x) v). \end{aligned}$$

By the surjectivity of $\mathcal{J}_x\Psi(x)$, we can conclude that $\Xi'(\Psi(x); \cdot)$ is a linear operator and so Ξ is Gâteaux differentiable at $\Psi(x)$. Since Ξ is assumed to be locally Lipschitz continuous on \mathcal{O} , Ξ is F-differentiable at $\Psi(x)$.

Next, we show that the second half inclusion holds:

$$\partial_B\Phi(\bar{x}) \supseteq \partial_B\Xi(\bar{y})\mathcal{J}_x\Psi(\bar{x}).$$

Let $W \in \partial_B\Xi(\bar{y})$. Then there exists a sequence $\{y^k\}$ in \mathcal{O} converging to \bar{y} such that Ξ is F-differentiable at y^k and $W = \lim_{k \rightarrow \infty} \mathcal{J}_y\Xi(y^k)$. By applying the classical Inverse Function Theorem to

$$\Psi(\bar{x} + \mathcal{J}_x\Psi(\bar{x})^*(y - \bar{y})) - \Psi(\bar{x}) = 0,$$

we obtain that there exists a sequence $\{\tilde{y}^k\}$ in \mathcal{O} converging to \bar{y} such that

$$\Psi\left(\bar{x} + \mathcal{J}_x\Psi(\bar{x})^*(\tilde{y}^k - \bar{y})\right) - \Psi(\bar{x}) = y^k - \Psi(\bar{x})$$

for all k sufficiently large. Let $\tilde{x}^k := \bar{x} + \mathcal{J}_x\Psi(\bar{x})^*(\tilde{y}^k - \bar{y})$. Then $y^k = \Psi(\tilde{x}^k)$ and Φ is F-differentiable at \tilde{x}^k with

$$\mathcal{J}_x\Phi(\tilde{x}^k) = \mathcal{J}_y\Xi(y^k)\mathcal{J}_x\Psi(\tilde{x}^k).$$

By using the fact that $\tilde{y}^k \rightarrow \bar{y}$ implies $\tilde{x}^k \rightarrow \bar{x}$, we know that there exists a $V \in \partial_B\Phi(\bar{x})$ such that

$$W\mathcal{J}_x\Psi(\bar{x}) = \lim_{k \rightarrow \infty} \mathcal{J}_y\Xi(y^k) \lim_{k \rightarrow \infty} \mathcal{J}_x\Psi(\tilde{x}^k) = \lim_{k \rightarrow \infty} \mathcal{J}_x\Phi(\tilde{x}^k) = V \in \partial_B\Phi(\bar{x}).$$

This completes the proof. Q.E.D.

Let D be a closed convex set in Z . Let $\Pi_D : Z \rightarrow Z$ denote the metric projector over D . That is, for any $y \in Z$, $\Pi_D(y)$ is the unique optimal solution to the convex quadratic programming problem:

$$\begin{aligned} \min \quad & \frac{1}{2}\langle z - y, z - y \rangle \\ \text{s.t.} \quad & z \in D. \end{aligned} \tag{13}$$

It is well known [39] that the metric projector $\Pi_D(\cdot)$ is contractive, i.e., for any two vectors $y, z \in Z$,

$$\|\Pi_D(y) - \Pi_D(z)\| \leq \|y - z\|.$$

Hence, $\Pi_D(\cdot)$ is F-differentiable almost everywhere in Z and for any $y \in Z$, $\partial\Pi_D(y)$ is well defined.

Lemma 2 [21, Proposition 1] *Let $D \subseteq Z$ be a closed convex set. Then, for any $y \in Z$ and $V \in \partial\Pi_D(y)$,*

(a) *V is self-adjoint.*

$$(b) \quad \langle d, Vd \rangle \geq 0 \quad \forall d \in Z.$$

$$(c) \quad \langle Vd, d - Vd \rangle \geq 0 \quad \forall d \in Z.$$

Lemma 2 provides general properties about $\partial\Pi_D(\cdot)$. In our analysis, we need a finer characterization about Clarke's Jacobian of $\Pi_{\mathcal{S}_+^p}(\cdot)$. We write $A \succeq 0$ and $A \succ 0$ to mean that A is a symmetric positive semidefinite matrix and a symmetric positive definite matrix, respectively. For any two matrices A and B in \mathcal{S}^p , we write

$$\langle A, B \rangle := \text{Tr}(A^T B)$$

for the *Frobenius inner product* between A and B , where “Tr” denotes the trace of a matrix. Under the Frobenius inner product, the projection $A_+ := \Pi_{\mathcal{S}_+^p}(A)$ of a matrix $A \in \mathcal{S}^p$ onto the cone \mathcal{S}_+^p satisfies the following complementarity condition:

$$\mathcal{S}_+^p \ni A_+ \perp (A_+ - A) \in \mathcal{S}_+^p, \quad (14)$$

where for any two matrices B and S in \mathcal{S}^p , $B \perp S \iff \langle B, S \rangle = 0$. Let A have the following spectral decomposition

$$A = P\Lambda P^T, \quad (15)$$

where Λ is the diagonal matrix of eigenvalues of A and P is a corresponding orthogonal matrix of orthonormal eigenvectors. Then

$$A_+ = P\Lambda_+ P^T,$$

where Λ_+ is the diagonal matrix whose diagonal entries are the nonnegative parts of the respective diagonal entries of Λ [13, 38]. Define three index sets of positive, zero, and negative eigenvalues of A , respectively, as

$$\alpha := \{i : \lambda_i > 0\}, \quad \beta := \{i : \lambda_i = 0\}, \quad \gamma := \{i : \lambda_i < 0\}.$$

Write

$$\Lambda = \begin{bmatrix} \Lambda_\alpha & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \Lambda_\gamma \end{bmatrix} \quad \text{and} \quad P = [P_\alpha \quad P_\beta \quad P_\gamma]$$

with $P_\alpha \in \mathbb{R}^{p \times |\alpha|}$, $P_\beta \in \mathbb{R}^{p \times |\beta|}$, and $P_\gamma \in \mathbb{R}^{p \times |\gamma|}$. Define the matrix $U \in \mathcal{S}^p$ with entries

$$U_{ij} := \frac{\max\{\lambda_i, 0\} + \max\{\lambda_j, 0\}}{|\lambda_i| + |\lambda_j|}, \quad i, j = 1, \dots, p,$$

where $0/0$ is defined to be 1. In [1, 2], Bonnans, Cominetti and Shapiro showed, among many other important things, that $\Pi_{\mathcal{S}_+^p}$ is directionally differentiable everywhere in \mathcal{S}^p . Sun

and Sun showed in [37] that $\Pi_{\mathcal{S}_+^p}$ is a strongly semismooth matrix-valued function and for any $H \in \mathcal{S}^p$, gave an explicit formula for the directional derivative of $\Pi'_{\mathcal{S}_+^p}(A; H)$:

$$\Pi'_{\mathcal{S}_+^p}(A; H) = P \begin{bmatrix} \tilde{H}_{\alpha\alpha} & \tilde{H}_{\alpha\beta} & U_{\alpha\gamma} \circ \tilde{H}_{\alpha\gamma} \\ \tilde{H}_{\alpha\beta}^T & \Pi_{\mathcal{S}_+^{|\beta|}}(\tilde{H}_{\beta\beta}) & 0 \\ \tilde{H}_{\alpha\gamma}^T \circ U_{\alpha\gamma}^T & 0 & 0 \end{bmatrix} P^T, \quad (16)$$

where $\tilde{H} := P^T H P$ and \circ denotes the Hadamard product. Hence, we have

$$\mathcal{T}_{\mathcal{S}_+^p}(A_+) = \{B \in \mathcal{S}^p : B = \Pi'_{\mathcal{S}_+^p}(A_+; B)\} = \{B \in \mathcal{S}^p : P_{\bar{\alpha}}^T B P_{\bar{\alpha}} \succeq 0\}, \quad (17)$$

where $\bar{\alpha} := \{1, \dots, p\} \setminus \alpha$ and $P_{\bar{\alpha}} := [P_{\beta} \ P_{\gamma}]$. The lineality space of $\mathcal{T}_{\mathcal{S}_+^p}(A_+)$, i.e., the largest linear space in $\mathcal{T}_{\mathcal{S}_+^p}(A_+)$, denoted by $\text{lin}(\mathcal{T}_{\mathcal{S}_+^p}(A_+))$, takes the following form:

$$\text{lin}(\mathcal{T}_{\mathcal{S}_+^p}(A_+)) = \{B \in \mathcal{S}^p : P_{\bar{\alpha}}^T B P_{\bar{\alpha}} = 0\}. \quad (18)$$

The *critical cone* of \mathcal{S}_+^p at $A \in \mathcal{S}^p$, associated with the complementarity problem (14), is defined as

$$C(A; \mathcal{S}_+^p) := \mathcal{T}_{\mathcal{S}_+^p}(A_+) \cap (A_+ - A)^\perp,$$

which can be completely described:

$$C(A; \mathcal{S}_+^p) = \{B \in \mathcal{S}^p : P_{\beta}^T B P_{\beta} \succeq 0, \ P_{\beta}^T B P_{\gamma} = 0, \ P_{\gamma}^T B P_{\gamma} = 0\}. \quad (19)$$

The affine hull of $C(A; \mathcal{S}_+^p)$, which we denote $\text{aff}(C(A; \mathcal{S}_+^p))$, can thus be written as

$$\text{aff}(C(A; \mathcal{S}_+^p)) = \{B \in \mathcal{S}^p : P_{\beta}^T B P_{\gamma} = 0, \ P_{\gamma}^T B P_{\gamma} = 0\}. \quad (20)$$

We summarize some differential properties of $\Pi_{\mathcal{S}_+^p}(\cdot)$ in the following proposition. For details, see [24, Corollary 10 & Lemma 11].

Proposition 3 *The following three statements are true.*

- (a) $\Pi_{\mathcal{S}_+^p}(\cdot)$ is F-differentiable at $A \in \mathcal{S}^p$ if and only if A is nonsingular.
- (b) For any $A \in \mathcal{S}^p$, the directional derivative $\Pi'_{\mathcal{S}_+^p}(A; \cdot)$ is F-differentiable at $H \in \mathcal{S}^p$ if and only if $\tilde{H}_{\beta\beta}$ is nonsingular, where $\tilde{H} := P^T H P$.
- (c) Let $A \in \mathcal{S}^p$ be arbitrary and $\Phi(\cdot) := \Pi'_{\mathcal{S}_+^p}(A; \cdot)$. It holds that

$$\partial_B \Pi_{\mathcal{S}_+^p}(A) = \partial_B \Phi(0).$$

Proposition 3 and Lemma 1 allow us to prove the following useful result on $\partial \Pi_{\mathcal{S}_+^p}(\cdot)$.

Proposition 4 Suppose that $A \in \mathcal{S}^p$ has the spectral decomposition as in (15). Then for any $V \in \partial_B \Pi_{\mathcal{S}_+^p}(A)$ (respectively, $\partial \Pi_{\mathcal{S}_+^p}(A)$), there exists a $W \in \partial_B \Pi_{\mathcal{S}_+^{|\beta|}}(0)$ (respectively, $\partial \Pi_{\mathcal{S}_+^{|\beta|}}(0)$) such that

$$V(H) = P \begin{bmatrix} \tilde{H}_{\alpha\alpha} & \tilde{H}_{\alpha\beta} & U_{\alpha\gamma} \circ \tilde{H}_{\alpha\gamma} \\ \tilde{H}_{\alpha\beta}^T & W(\tilde{H}_{\beta\beta}) & 0 \\ \tilde{H}_{\alpha\gamma}^T \circ U_{\alpha\gamma}^T & 0 & 0 \end{bmatrix} P^T \quad \forall H \in \mathcal{S}^p, \quad (21)$$

where $\tilde{H} := P^T H P$. Conversely, for any $W \in \partial_B \Pi_{\mathcal{S}_+^{|\beta|}}(0)$ (respectively, $\partial \Pi_{\mathcal{S}_+^{|\beta|}}(0)$), there exists a $V \in \partial_B \Pi_{\mathcal{S}_+^p}(A)$ (respectively, $\partial \Pi_{\mathcal{S}_+^p}(A)$) such that (21) holds.

Proof. We only need to prove that (21) holds for $V \in \partial_B \Pi_{\mathcal{S}_+^p}(A)$ and $W \in \partial_B \Pi_{\mathcal{S}_+^{|\beta|}}(0)$.

Let $\Phi(\cdot) := \Pi'_{\mathcal{S}_+^p}(A; \cdot)$. Define $\Psi : \mathcal{S}^p \rightarrow \mathcal{S}^p$ by $\Psi(H) := P^T H P$, $H \in \mathcal{S}^p$ and $\Xi : \mathcal{S}^p \rightarrow \mathcal{S}^p$ by

$$\Xi(B) := P \begin{bmatrix} B_{\alpha\alpha} & B_{\alpha\beta} & U_{\alpha\gamma} \circ B_{\alpha\gamma} \\ B_{\alpha\beta}^T & \Pi_{\mathcal{S}_+^{|\beta|}}(B_{\beta\beta}) & 0 \\ B_{\alpha\gamma}^T \circ U_{\alpha\gamma}^T & 0 & 0 \end{bmatrix} P^T, \quad B \in \mathcal{S}^p.$$

Then, by (16), we have

$$\Phi(H) = \Xi(\Psi(H)), \quad H \in \mathcal{S}^p.$$

Since $\Pi_{\mathcal{S}_+^{|\beta|}}$ is directionally differentiable everywhere and $\mathcal{J}_H \Psi(H) : \mathcal{S}^p \rightarrow \mathcal{S}^p$ is onto, we know from Lemma 1 that

$$\partial_B \Phi(0) = \partial_B \Xi(0) \mathcal{J}_H \Psi(0).$$

This, together with (c) of Proposition 3, completes the proof. Q.E.D.

Remark 5 Relation (21) in Proposition 4 holds for any orthogonal matrix P such that the spectral decomposition (15) holds. One may further characterize $\partial_B \Pi_{\mathcal{S}_+^{|\beta|}}(0)$ as in [24, Lemma 11]. In this paper, we do not need the structure of $\partial_B \Pi_{\mathcal{S}_+^{|\beta|}}(0)$.

Motivated by Shapiro [35, p. 313] and Bonnans and Shapiro [4, p. 487], for any given $B \in \mathcal{S}^p$ we introduce a linear-quadratic function $\Upsilon_B : \mathcal{S}^p \times \mathcal{S}^p \rightarrow \mathfrak{R}$ in the next definition.

Definition 6 For any given $B \in \mathcal{S}^p$, define the linear-quadratic function $\Upsilon_B : \mathcal{S}^p \times \mathcal{S}^p \rightarrow \mathfrak{R}$, which is linear in the first argument and quadratic in the second argument, by

$$\Upsilon_B(\Gamma, A) := 2 \left\langle \Gamma, A B^\dagger A \right\rangle, \quad (\Gamma, A) \in \mathcal{S}^p \times \mathcal{S}^p,$$

where B^\dagger is the Moore-Penrose pseudo-inverse of B .

The following result plays an important role in our subsequent analysis.

Proposition 7 *Suppose that $B \in \mathcal{S}_+^p$ and $\Gamma \in \mathcal{N}_{\mathcal{S}_+^p}(B)$. Then for any $V \in \partial\Pi_{\mathcal{S}_+^p}(B + \Gamma)$ and $\Delta B, \Delta\Gamma \in \mathcal{S}^p$ such that $\Delta B = V(\Delta B + \Delta\Gamma)$, it holds that*

$$\langle \Delta B, \Delta\Gamma \rangle \geq -\Upsilon_B(\Gamma, \Delta B). \quad (22)$$

Proof. Let $A := B + \Gamma$. Then we know from [9] that

$$B = \Pi_{\mathcal{S}_+^p}(B + \Gamma) = \Pi_{\mathcal{S}_+^p}(A) \quad \text{and} \quad B\Gamma = \Gamma B = 0.$$

Thus, we can assume that A has the spectral decomposition as in (15),

$$B = P \begin{bmatrix} \Lambda_\alpha & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} P^T, \quad \text{and} \quad \Gamma = P \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \Lambda_\gamma \end{bmatrix} P^T.$$

Let $\tilde{B} := P^T B P$, $\tilde{\Gamma} := P^T \Gamma P$, $\Delta\tilde{B} := P^T \Delta B P$, and $\Delta\tilde{\Gamma} := P^T \Delta\Gamma P$. Then, by Proposition 4, there exists a $W \in \partial\Pi_{\mathcal{S}_+^{|\beta|}}(0)$ such that

$$V(\Delta B + \Delta\Gamma) = P \begin{bmatrix} \Delta\tilde{B}_{\alpha\alpha} + \Delta\tilde{\Gamma}_{\alpha\alpha} & \Delta\tilde{B}_{\alpha\beta} + \Delta\tilde{\Gamma}_{\alpha\beta} & U_{\alpha\gamma} \circ (\Delta\tilde{B}_{\alpha\gamma} + \Delta\tilde{\Gamma}_{\alpha\gamma}) \\ (\Delta\tilde{B}_{\alpha\beta} + \Delta\tilde{\Gamma}_{\alpha\beta})^T & W(\Delta\tilde{B}_{\beta\beta} + \Delta\tilde{\Gamma}_{\beta\beta}) & 0 \\ (\Delta\tilde{B}_{\alpha\gamma} + \Delta\tilde{\Gamma}_{\alpha\gamma})^T \circ U_{\alpha\gamma}^T & 0 & 0 \end{bmatrix} P^T,$$

which, together with the assumption that $\Delta B = V(\Delta B + \Delta\Gamma)$, implies that

$$\Delta\tilde{\Gamma}_{\alpha\alpha} = 0, \quad \Delta\tilde{\Gamma}_{\alpha\beta} = 0, \quad \Delta\tilde{B}_{\beta\gamma} = 0, \quad \Delta\tilde{B}_{\gamma\gamma} = 0, \quad (23)$$

$$\Delta\tilde{B}_{\beta\beta} = W(\Delta\tilde{B}_{\beta\beta} + \Delta\tilde{\Gamma}_{\beta\beta}), \quad (24)$$

and

$$\Delta\tilde{B}_{\alpha\beta} - U_{\alpha\gamma} \circ \Delta\tilde{B}_{\alpha\beta} = U_{\alpha\gamma} \circ \Delta\tilde{\Gamma}_{\beta\beta}. \quad (25)$$

By (c) of Lemma 2 and equation (24), we obtain that

$$\langle \Delta\tilde{B}_{\beta\beta}, \Delta\tilde{\Gamma}_{\beta\beta} \rangle = \langle W(\Delta\tilde{B}_{\beta\beta} + \Delta\tilde{\Gamma}_{\beta\beta}), (\Delta\tilde{B}_{\beta\beta} + \Delta\tilde{\Gamma}_{\beta\beta}) - W(\Delta\tilde{B}_{\beta\beta} + \Delta\tilde{\Gamma}_{\beta\beta}) \rangle \geq 0. \quad (26)$$

Hence, by equations (23), (25), and (26),

$$\begin{aligned} \langle \Delta B, \Delta\Gamma \rangle &= \langle \Delta\tilde{B}, \Delta\tilde{\Gamma} \rangle = 2 \operatorname{Tr} \left((\Delta\tilde{B}_{\alpha\gamma})^T \Delta\tilde{\Gamma}_{\alpha\gamma} \right) + \operatorname{Tr} \left(\Delta\tilde{B}_{\beta\beta} \Delta\tilde{\Gamma}_{\beta\beta} \right) \\ &\geq 2 \operatorname{Tr} \left((\Delta\tilde{B}_{\alpha\gamma})^T \Delta\tilde{\Gamma}_{\alpha\gamma} \right) = 2 \sum_{i \in \alpha, j \in \gamma} (\Delta\tilde{B})_{ij} (\Delta\tilde{\Gamma})_{ij} \\ &= 2 \sum_{i \in \alpha, j \in \gamma} \frac{|\lambda_j|}{\lambda_i} \left((\Delta\tilde{B})_{ij} \right)^2 = -2 \sum_{i \in \alpha, j \in \gamma} \frac{\lambda_j}{\lambda_i} \left((\Delta\tilde{B})_{ij} \right)^2. \end{aligned} \quad (27)$$

On the other hand, since

$$B^\dagger = P \begin{bmatrix} (\Lambda_\alpha)^{-1} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} P^T,$$

we obtain from (23) and the spectral decomposition of Γ that

$$\begin{aligned} \Upsilon_B(\Gamma, \Delta B) &= 2 \langle \Gamma, (\Delta B) B^\dagger (\Delta B) \rangle = 2 \langle (\Delta B) \Gamma, B^\dagger (\Delta B) \rangle \\ &= 2 \langle (\Delta \tilde{B}) \tilde{\Gamma}, (P^T B^\dagger P) (\Delta \tilde{B}) \rangle \\ &= 2 \operatorname{Tr} \left((\Delta \tilde{B}_{\alpha\gamma} \Lambda_\gamma)^T (\Lambda_\alpha)^{-1} \Delta \tilde{B}_{\alpha\gamma} \right) \\ &= 2 \sum_{i \in \alpha, j \in \gamma} \frac{\lambda_j}{\lambda_i} \left((\Delta \tilde{B})_{ij} \right)^2. \end{aligned} \quad (28)$$

By combining (27) and (28), we get (22).

Q.E.D.

3 Strong Second Order Sufficient Condition and Constraint Nondegeneracy

Let D be an arbitrary closed set in a given finite dimensional real vector space Z . The *inner* and *out second order tangent sets* to the set D at the point $y \in D$ and in the direction $d \in Z$ can be defined, respectively, by

$$\mathcal{T}_D^{i,2}(y, d) := \{w \in Z : \operatorname{dist}(y + td + \frac{1}{2}t^2w, D) = o(t^2), t \geq 0\} \quad (29)$$

and

$$\mathcal{T}_D^2(y, d) := \{w \in Z : \exists t_k \downarrow 0 \text{ such that } \operatorname{dist}(y + t_k d + \frac{1}{2}t_k^2w, D) = o(t_k^2)\}. \quad (30)$$

From the definitions of inner and out second order tangent sets, we can see directly that $\mathcal{T}_D^{i,2}(z, d) \subseteq \mathcal{T}_D^2(y, d)$ and $\mathcal{T}_D^{i,2}(z, d) = \emptyset$ (respectively, $\mathcal{T}_D^2(z, d) = \emptyset$) if $d \notin \mathcal{T}_D^i(y)$ (respectively, $d \notin \mathcal{T}_D(y)$). In general, $\mathcal{T}_D^{i,2}(z, d) \neq \mathcal{T}_D^2(z, d)$ even if D is convex [4, Section 3.3]. However, if D is \mathcal{C}^2 -cone reducible, the equality always holds [4, Proposition 3.136]. In particular, when $K := \{0\} \times \mathcal{S}_+^p \subset Y := \mathbb{R}^m \times \mathcal{S}^p$,

$$\mathcal{T}_K^{i,2}(y, d) = \mathcal{T}_K^2(y, d) \quad \forall y, d \in Y.$$

Let \bar{x} be a feasible solution to the nonlinear semidefinite programming (*NLSDP*). The *critical cone* $C(\bar{x})$ of (*NLSDP*) at \bar{x} is defined by

$$C(\bar{x}) := \{d : \mathcal{J}_x G(\bar{x})d \in \mathcal{T}_K(G(\bar{x})), \mathcal{J}_x f(\bar{x})d \leq 0\}, \quad (31)$$

i.e.,

$$C(\bar{x}) = \left\{ d : \mathcal{J}_x h(\bar{x})d = 0, \mathcal{J}_x g(\bar{x})d \in \mathcal{T}_{\mathcal{S}_+^p}(g(\bar{x})), \mathcal{J}_x f(\bar{x})d \leq 0 \right\}. \quad (32)$$

If \bar{x} is a stationary point of (NLSDP), i.e., if $\mathcal{M}(\bar{x})$ is nonempty, then

$$C(\bar{x}) = \left\{ d : \mathcal{J}_x h(\bar{x})d = 0, \mathcal{J}_x g(\bar{x})d \in \mathcal{T}_{\mathcal{S}_+^p}(g(\bar{x})), \mathcal{J}_x f(\bar{x})d = 0 \right\}. \quad (33)$$

Let \bar{x} be a stationary point of (NLSDP). Then there exists $(\bar{\zeta}, \bar{\Gamma}) \in \mathcal{M}(\bar{x})$ such that

$$\mathcal{J}_x L(\bar{x}, \bar{\zeta}, \bar{\Gamma}) = 0, \quad -h(\bar{x}) = 0, \quad \text{and} \quad \bar{\Gamma} \in \mathcal{N}_{\mathcal{S}_+^p}(g(\bar{x})).$$

By using the fact that

$$\bar{\Gamma} \in \mathcal{N}_{\mathcal{S}_+^p}(g(\bar{x})) \iff \mathcal{S}_+^p \ni (-\bar{\Gamma}) \perp g(\bar{x}) \in \mathcal{S}_+^p,$$

we may assume that $A := g(\bar{x}) + \bar{\Gamma}$ has the spectral decomposition as in (15),

$$g(\bar{x}) = P \begin{bmatrix} \Lambda_\alpha & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} P^T, \quad \text{and} \quad \bar{\Gamma} = P \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \Lambda_\gamma \end{bmatrix} P^T. \quad (34)$$

Then, by (17) and (18), we have

$$\mathcal{T}_{\mathcal{S}_+^p}(g(\bar{x})) = \{ B \in \mathcal{S}^p : [P_\beta \ P_\gamma]^T B [P_\beta \ P_\gamma] \succeq 0 \},$$

$$\mathcal{T}_{\mathcal{S}_+^p}(g(\bar{x})) \cap \bar{\Gamma}^\perp = \{ B \in \mathcal{S}^p : P_\beta^T B P_\beta \succeq 0, \quad P_\beta^T B P_\gamma = 0, \quad P_\gamma^T B P_\gamma = 0 \}, \quad (35)$$

and

$$\text{lin} \left(\mathcal{T}_{\mathcal{S}_+^p}(g(\bar{x})) \right) = \{ B \in \mathcal{S}^p : [P_\beta \ P_\gamma]^T B [P_\beta \ P_\gamma] = 0 \}. \quad (36)$$

Furthermore, since $\mathcal{M}(\bar{x})$ is nonempty,

$$\begin{aligned} C(\bar{x}) &= \{ d : \mathcal{J}_x h(\bar{x})d = 0, \quad [P_\beta \ P_\gamma]^T (\mathcal{J}_x g(\bar{x})d) [P_\beta \ P_\gamma] \succeq 0, \quad P_\gamma^T (\mathcal{J}_x g(\bar{x})d) P_\gamma = 0 \} \\ &= \left\{ d : \mathcal{J}_x h(\bar{x})d = 0, \quad P_\beta^T (\mathcal{J}_x g(\bar{x})d) P_\beta \succeq 0, \quad P_\beta^T (\mathcal{J}_x g(\bar{x})d) P_\gamma = 0, \right. \\ &\quad \left. P_\gamma^T (\mathcal{J}_x g(\bar{x})d) P_\gamma = 0 \right\}. \end{aligned} \quad (37)$$

However, it is not easy to give an explicit formula to the affine hull of $C(\bar{x})$, which we denote $\text{aff}(C(\bar{x}))$. Instead, we define the following outer approximation set to $\text{aff}(C(\bar{x}))$ with respect to $(\bar{\zeta}, \bar{\Gamma})$ by

$$\text{app}(\bar{\zeta}, \bar{\Gamma}) := \{ d : \mathcal{J}_x h(\bar{x})d = 0, \quad \mathcal{J}_x g(\bar{x})d \in \text{aff}(C(A; \mathcal{S}_+^p)) \}, \quad (38)$$

where $C(A; \mathcal{S}_+^p)$ is the critical cone of \mathcal{S}_+^p at $A = g(\bar{x}) + \bar{\Gamma}$. By (20), it holds that

$$\text{app}(\bar{\zeta}, \bar{\Gamma}) = \{ d : \mathcal{J}_x h(\bar{x})d = 0, \quad P_\beta^T (\mathcal{J}_x g(\bar{x})d) P_\gamma = 0, \quad P_\gamma^T (\mathcal{J}_x g(\bar{x})d) P_\gamma = 0 \}. \quad (39)$$

Then by the definition of $\text{aff}(C(\bar{x}))$, we have for any $(\bar{\zeta}, \bar{\Gamma}) \in \mathcal{M}(\bar{x})$ that

$$\text{aff}(C(\bar{x})) \subseteq \text{app}(\bar{\zeta}, \bar{\Gamma}). \quad (40)$$

Obviously, the two sets in (40) coincide if the *strict complementary* condition holds at $(\bar{x}, \bar{\zeta}, \bar{\Gamma})$:

$$\text{rank}(g(\bar{x})) + \text{rank}(\bar{\Gamma}) = p,$$

where “rank” denotes the rank of a square matrix. In general, these two sets may be different even if $\mathcal{M}(\bar{x})$ is a singleton as in the case for the conventional nonlinear programming (NLP).

The next proposition shows that the equality in (40) holds if $(\bar{\zeta}, \bar{\Gamma}) \in \mathcal{M}(\bar{x})$ satisfies a stronger constraint qualification than Robinson’s CQ (7) at \bar{x} , which, in the context of (NLSDP), is equivalent to

$$\begin{cases} \mathcal{J}_x h(\bar{x})X = \mathfrak{R}^m, \\ \mathcal{J}_x g(\bar{x})X + \mathcal{T}_{\mathcal{S}_+^p}(g(\bar{x})) = \mathcal{S}^p. \end{cases} \quad (41)$$

Proposition 8 *Let \bar{x} be a feasible solution to the nonlinear semidefinite programming (NLSDP) and $(\bar{\zeta}, \bar{\Gamma}) \in \mathcal{M}(\bar{x})$. Suppose that $(\bar{\zeta}, \bar{\Gamma})$ satisfies the following strict constraint qualification:*

$$\begin{cases} \mathcal{J}_x h(\bar{x})X = \mathfrak{R}^m, \\ \mathcal{J}_x g(\bar{x})X + \mathcal{T}_{\mathcal{S}_+^p}(g(\bar{x})) \cap \bar{\Gamma}^\perp = \mathcal{S}^p. \end{cases} \quad (42)$$

Then $\mathcal{M}(\bar{x})$ is a singleton, i.e., $\mathcal{M}(\bar{x}) = \{(\bar{\zeta}, \bar{\Gamma})\}$, and $\text{aff}(C(\bar{x})) = \text{app}(\bar{\zeta}, \bar{\Gamma})$.

Proof. The uniqueness of $(\bar{\zeta}, \bar{\Gamma})$ follows from [4, Proposition 4.50]. We only need to show

$$\text{app}(\bar{\zeta}, \bar{\Gamma}) \subseteq \text{aff}(C(\bar{x})).$$

Let d be an arbitrary vector in $\text{app}(\bar{\zeta}, \bar{\Gamma})$. Let $A := g(\bar{x}) + \bar{\Gamma}$. We may assume that A has the spectral decomposition as in (15) and the two matrices $g(\bar{x})$ and $\bar{\Gamma}$ satisfy (34). Let S be any matrix in \mathcal{S}^p such that

$$P^T S P = \begin{bmatrix} P_\alpha^T S P_\alpha & P_\alpha^T S P_\beta & P_\alpha^T S P_\gamma \\ P_\beta^T S P_\alpha & P_\beta^T S P_\beta & 0 \\ P_\gamma^T S P_\alpha & 0 & 0 \end{bmatrix} \quad \text{with } P_\beta^T S P_\beta \succ 0.$$

By the strict constraint qualification (42), we know that there exist a vector $\bar{d} \in X$ and a matrix $U \in \mathcal{T}_{\mathcal{S}_+^p}(g(\bar{x})) \cap \bar{\Gamma}^\perp$ such that

$$\begin{cases} \mathcal{J}_x h(\bar{x})(-\bar{d}) = 0, \\ \mathcal{J}_x g(\bar{x})(-\bar{d}) + U = -S, \end{cases} \quad (43)$$

which, together with (35), implies that

$$P^T(\mathcal{J}_x g(\bar{x})\bar{d})P = U + S \in \mathcal{T}_{S_+^p}(g(\bar{x})) \cap \bar{\Gamma}^\perp, \quad P_\beta^T(\mathcal{J}_x g(\bar{x})\bar{d})P_\beta \succ 0, \quad \text{and} \quad \bar{d} \in C(\bar{x}).$$

Let $\bar{\tau} > 0$ be sufficiently large such that

$$P_\beta^T[\mathcal{J}_x g(\bar{x})(\bar{\tau}\bar{d} - d)]P_\beta = \bar{\tau}P_\beta^T(\mathcal{J}_x g(\bar{x})\bar{d})P_\beta - P_\beta^T(\mathcal{J}_x g(\bar{x})d)P_\beta \succeq 0.$$

Furthermore, since

$$P_\beta^T(\mathcal{J}_x g(\bar{x})\bar{d})P_\gamma = P_\beta^T(\mathcal{J}_x g(\bar{x})d)P_\gamma = 0 \quad \text{and} \quad P_\gamma^T(\mathcal{J}_x g(\bar{x})\bar{d})P_\gamma = P_\gamma^T(\mathcal{J}_x g(\bar{x})d)P_\gamma = 0,$$

it holds that

$$\bar{\tau}\bar{d} - d \in C(\bar{x}).$$

Therefore, by using the facts that $d = \bar{\tau}\bar{d} - (\bar{\tau}\bar{d} - d)$ and both $\bar{\tau}\bar{d}$ and $\bar{\tau}\bar{d} - d$ are in the critical cone $C(\bar{x})$, we complete the proof. Q.E.D.

Before we state the second order conditions for (NLSDP), we need below the concept of \mathcal{C}^2 -cone reducibility, which is adapted from Bonnans and Shapiro [4, Definition 3.315].

Definition 9 *A closed (not necessarily convex) set $D \subseteq Y$ is called \mathcal{C}^2 -cone reducible at a point $\bar{y} \in D$ if there exist a neighborhood $\mathcal{V} \subseteq Y$ of \bar{y} , a pointed closed convex cone Q (a cone is said to be pointed if and only its lineality space is the origin) in a finite dimensional space Z and a twice continuously differentiable mapping $\Xi : \mathcal{V} \rightarrow Z$ such that: (i) $\Xi(\bar{y}) = 0 \in Z$, (ii) the derivative mapping $J_y \Xi(\bar{y}) : Y \rightarrow Z$ is onto, and (iii) $D \cap \mathcal{V} = \{y \in \mathcal{V} \mid \Xi(y) \in Q\}$. We say that D is \mathcal{C}^2 -cone reducible if D is \mathcal{C}^2 -cone reducible at every point $\bar{y} \in Y$ (possibly to a different pointed cone Q).*

Many interesting sets such as the polyhedral convex set, the second-order cone, and the cone S_+^p are all \mathcal{C}^2 -cone reducible, and the Cartesian product of \mathcal{C}^2 -cone reducible sets is again \mathcal{C}^2 -cone reducible [4, 36]. In particular, $K = \{0\} \times S_+^p$ is \mathcal{C}^2 -cone reducible.

Recall that for any set $D \subseteq Z$, the support function of the set D is defined as

$$\sigma(y, D) := \sup_{z \in D} \langle z, y \rangle, \quad y \in Z.$$

Combining Theorem 3.45 and Proposition 3.316 with Theorem 3.317 in [4], we can state in the following theorem the second order necessary condition and the second order sufficient condition for nonlinear semidefinite programming (NLSDP). See also [2].

Theorem 10 *(Second order conditions.) Let $K = \{0\} \times S_+^p \subset \mathbb{R}^m \times S^p$. Suppose that \bar{x} is a locally optimal solution to the nonlinear semidefinite programming (NLSDP) and Robinson's CQ holds at \bar{x} . Then the following inequality holds:*

$$\sup_{\mu \in \mathcal{M}(\bar{x})} \left\{ \langle d, \mathcal{J}_{xx}^2 L(\bar{x}, \mu)d \rangle - \sigma(\mu, \mathcal{T}_K^2(G(\bar{x}), \mathcal{J}_x G(\bar{x})d)) \right\} \geq 0 \quad \forall d \in C(\bar{x}). \quad (44)$$

Conversely, let \bar{x} be a feasible solution to (NLSDP) such that $\mathcal{M}(\bar{x})$ is nonempty. Suppose that Robinson's CQ holds at \bar{x} . Then the following condition

$$\sup_{\mu \in \mathcal{M}(\bar{x})} \{ \langle d, \mathcal{J}_{xx}^2 L(\bar{x}, \mu) d \rangle - \sigma(\mu, \mathcal{T}_K^2(G(\bar{x}), \mathcal{J}_x G(\bar{x})d)) \} > 0 \quad \forall d \in C(\bar{x}) \setminus \{0\} \quad (45)$$

is necessary and sufficient for the quadratic growth condition at the point \bar{x} :

$$f(x) \geq f(\bar{x}) + c\|x - \bar{x}\|^2 \quad \forall x \in \hat{N} \text{ such that } G(x) \in K \quad (46)$$

for some constant $c > 0$ and a neighborhood \hat{N} of \bar{x} in X .

Obviously, when the second order growth condition (46) holds, \bar{x} is a strictly local solution of (NLSDP). So there exists no gap between the above second order sufficient condition (45) and the second order necessary condition (44).

We write $\mu = (\zeta, \Gamma) \in \mathbb{R}^m \times \mathcal{S}^p$ for any $\mu \in \mathcal{M}(\bar{x})$. Then for $\mu \in \mathcal{M}(\bar{x})$ and $d \in C(\bar{x})$ the ‘‘sigma term’’ in (44) and (45) can be written as

$$\begin{aligned} \sigma(\mu, \mathcal{T}_K^2(G(\bar{x}), \mathcal{J}_x G(\bar{x})d)) &= \sigma\left(\zeta, \mathcal{T}_{\{0\}}^2(h(\bar{x}), \mathcal{J}_x h(\bar{x})d)\right) + \sigma\left(\Gamma, \mathcal{T}_{\mathcal{S}_+^p}^2(g(\bar{x}), \mathcal{J}_x g(\bar{x})d)\right) \\ &= 0 + \sigma\left(\Gamma, \mathcal{T}_{\mathcal{S}_+^p}^2(g(\bar{x}), \mathcal{J}_x g(\bar{x})d)\right) \\ &= \sigma\left(\Gamma, \mathcal{T}_{\mathcal{S}_+^p}^2(g(\bar{x}), \mathcal{J}_x g(\bar{x})d)\right), \end{aligned}$$

which becomes $-\infty$ for any $d \in X$ such that $\mathcal{J}_x g(\bar{x})d \notin \mathcal{T}_{\mathcal{S}_+^p}(g(\bar{x}))$. This means that in order to define a stronger second order sufficient condition over a set larger than $C(\bar{x})$ one needs to find a substitute for this sigma term. The following lemma, due to Shapiro [35, p. 313] and Bonnans and Shapiro [4, p. 487], makes it possible.

Lemma 11 *Let \bar{x} be a feasible solution to (NLSDP) such that $\mathcal{M}(\bar{x})$ is nonempty. Then for any $(\zeta, \Gamma) \in \mathcal{M}(\bar{x})$ with $\zeta \in \mathbb{R}^m$ and $\Gamma \in \mathcal{S}^p$, one has*

$$\Upsilon_{g(\bar{x})}(\Gamma, \mathcal{J}_x g(\bar{x})d) = \sigma\left(\Gamma, \mathcal{T}_{\mathcal{S}_+^p}^2(g(\bar{x}), \mathcal{J}_x g(\bar{x})d)\right) \quad \forall d \in C(\bar{x}).$$

Now, we are ready to define a strong second order sufficient condition, which extends an analogue for the conventional nonlinear programming (NLP) introduced by Robinson [28] to the nonlinear semidefinite programming (NLSDP).

Definition 12 *Let \bar{x} be a stationary point of the nonlinear semidefinite programming (NLSDP). We say that the strong second order sufficient condition holds at \bar{x} if*

$$\sup_{(\zeta, \Gamma) \in \mathcal{M}(\bar{x})} \{ \langle d, \mathcal{J}_{xx}^2 L(\bar{x}, \zeta, \Gamma) d \rangle - \Upsilon_{g(\bar{x})}(\Gamma, \mathcal{J}_x g(\bar{x})d) \} > 0 \quad \forall d \in \hat{C}(\bar{x}) \setminus \{0\}, \quad (47)$$

where for any $(\zeta, \Gamma) \in \mathcal{M}(\bar{x})$, $(\zeta, \Gamma) \in \mathbb{R}^m \times \mathcal{S}^p$ and

$$\hat{C}(\bar{x}) := \bigcap_{(\zeta, \Gamma) \in \mathcal{M}(\bar{x})} \text{app}(\zeta, \Gamma).$$

Next, we define a nondegeneracy condition for $(NLSDP)$, which is an analogue of the LICQ for the conventional nonlinear programming (NLP) . The concept of nondegeneracy originally appeared in Robinson [29] for the general optimization problem (OP) . Here we adopt a somewhat slightly different version from Robinson's original definition.

Definition 13 *We say that a feasible point \bar{x} to optimization problem (OP) is constraint nondegenerate if*

$$\mathcal{J}_x G(\bar{x})X + \text{lin}(\mathcal{T}_K(\bar{y})) = Y, \quad (48)$$

where $\bar{y} := G(\bar{x})$.

The name “constraint nondegeneracy” was coined by Robinson in [32]. The nondegeneracy condition (48) given here is consistent with the version given in [30] and has been extensively used in [4, 36] for sensitivity and stability analysis in optimization and variational inequalities. See [4, 36] for various equivalent forms. Certainly, the constraint nondegenerate condition (48) implies Robinson's CQ (7). For the conventional (NLP) , as observed in Robinson [29] and Shapiro [36], the LICQ is equivalent to the constraint nondegeneracy. For the nonlinear semidefinite programming $(NLSDP)$, the constraint nondegeneracy takes the following form:

$$\begin{cases} \mathcal{J}_x h(\bar{x})X = \mathbb{R}^m, \\ \mathcal{J}_x g(\bar{x})X + \text{lin}(\mathcal{T}_{\mathcal{S}_+^p}(g(\bar{x}))) = \mathcal{S}^p. \end{cases} \quad (49)$$

Let x be a feasible point to $(NLSDP)$ such that $\mathcal{M}(x)$ is nonempty. Then there exists $(\zeta, \Gamma) \in \mathbb{R}^m \times \mathcal{S}^p$, together with x , satisfying the following KKT condition:

$$\mathcal{J}_x L(x, \zeta, \Gamma) = 0, \quad -h(x) = 0, \quad \text{and} \quad \Gamma \in \mathcal{N}_{\mathcal{S}_+^p}(g(x)), \quad (50)$$

where

$$L(x, \zeta, \Gamma) = f(x) + \langle \zeta, h(x) \rangle + \langle \Gamma, g(x) \rangle.$$

Since, from Eaves [9],

$$\Gamma \in \mathcal{N}_{\mathcal{S}_+^p}(g(x)) \iff g(x) = \Pi_{\mathcal{S}_+^p}(g(x) + \Gamma),$$

we can write the KKT condition (50) equivalently as

$$F(x, \zeta, \Gamma) := \begin{bmatrix} \mathcal{J}_x L(x, \zeta, \Gamma) \\ -h(x) \\ -g(x) + \Pi_{\mathcal{S}_+^p}(g(x) + \Gamma) \end{bmatrix} = \begin{bmatrix} \mathcal{J}_x L(x, \zeta, \Gamma) \\ -h(x) \\ \Gamma - \Pi_{\mathcal{S}_+^p}(\Gamma + g(x)) \end{bmatrix} = 0, \quad (51)$$

where \mathcal{S}_-^p is the cone of negative semidefinite symmetric matrices in \mathcal{S}^p , i.e., $\mathcal{S}_-^p = -\mathcal{S}_+^p$. Both (50) and (51) are equivalent to

$$0 \in \begin{bmatrix} \mathcal{J}_x L(x, \zeta, \Gamma) \\ -h(x) \\ -g(x) \end{bmatrix} + \begin{bmatrix} \mathcal{N}_X(x) \\ \mathcal{N}_{\mathbb{R}^m}(\zeta) \\ \mathcal{N}_{\mathcal{S}_-^p}(\Gamma) \end{bmatrix}. \quad (52)$$

Problem (52) is in the form of the following generalized equation:

$$0 \in \phi(z) + \mathcal{N}_D(z), \quad (53)$$

where ϕ is a continuously differentiable mapping from a given finite dimensional real vector space Z to itself and D is a closed convex set in Z .

Robinson in [28] introduced the far reaching concept of *strong regularity* for a solution of the generalized equation (53).

Definition 14 *Let \bar{z} be a solution of the generalized equation (53). We say that \bar{z} is a strongly regular solution of the generalized equation (53) if there exist neighborhoods \mathcal{B} of the origin $0 \in Z$ and \mathcal{V} of \bar{z} such that for every $\delta \in \mathcal{B}$, the following linearized generalized equation*

$$\delta \in \phi(\bar{z}) + \mathcal{J}_z \phi(\bar{z})(z - \bar{z}) + \mathcal{N}_D(z)$$

has a unique solution in \mathcal{V} , denoted by $z_{\mathcal{V}}(\delta)$, and the mapping $z_{\mathcal{V}} : \mathcal{B} \rightarrow \mathcal{V}$ is Lipschitz continuous.

Remark 15 *Recall that a function $\Xi : \mathcal{O} \subseteq Z \rightarrow Z$ is said to be a locally Lipschitz homeomorphism near $\bar{z} \in \mathcal{O}$ if there exists an open neighborhood $\mathcal{V} \subseteq \mathcal{O}$ of \bar{z} such that the restricted mapping $\Xi|_{\mathcal{V}} : \mathcal{V} \rightarrow \Xi(\mathcal{V})$ is Lipschitz continuous and bijective, and its inverse is also Lipschitz continuous. Define two mappings $\hat{\Xi}, \Xi : Z \rightarrow Z$ by*

$$\hat{\Xi}(z) := z - \Pi_D(z - \hat{\phi}(z)) \quad \text{and} \quad \Xi(z) := z - \Pi_D(z - \phi(z)),$$

where $\hat{\phi}(z) := \phi(\bar{z}) + \mathcal{J}_z \phi(\bar{z})(z - \bar{z})$, $z \in Z$. From Lemma 3.1 in Robinson [31] or Theorem 3.1 in Kummer [18] we know that $\hat{\Xi}$ is a locally Lipschitz homeomorphism near \bar{z} if and only if Ξ is so. See also Theorem 5.2.8 in Facchinei and Pang [10]. Thus, \bar{z} is a strongly regular solution of the generalized equation (53) is equivalent to say that $\hat{\Xi}$ or Ξ is a locally Lipschitz homeomorphism near \bar{z} .

The next proposition relates the strong second order sufficient condition and constraint nondegeneracy to the nonsingularity of Clarke's Jacobian of the mapping F and the strong regularity of a solution to the generalized equation (52).

Proposition 16 *Let \bar{x} be a feasible solution to the nonlinear semidefinite programming (NLSDP). Let $\bar{\zeta} \in \mathbb{R}^m$ and $\bar{\Gamma} \in \mathcal{S}^p$ be such that $(\bar{\zeta}, \bar{\Gamma}) \in \mathcal{M}(\bar{x})$, i.e., let $(\bar{x}, \bar{\zeta}, \bar{\Gamma})$ be a KKT point of (NLSDP). Consider the following three statements:*

- (a) *The strong second order sufficient condition (47) holds at \bar{x} and \bar{x} is constraint nondegenerate.*
- (b) *Any element in $\partial F(\bar{x}, \bar{\zeta}, \bar{\Gamma})$ is nonsingular.*

(c) The KKT point $(\bar{x}, \bar{\zeta}, \bar{\Gamma})$ is a strongly regular solution of the generalized equation (52).

It holds that (a) \implies (b) \implies (c).

Proof. “(a) \implies (b)” Since the constraint nondegeneracy condition (49) is assumed to hold at \bar{x} , $(\bar{\zeta}, \bar{\Gamma})$ satisfies the strict constraint qualification (42). Thus, by Proposition 8, $\mathcal{M}(\bar{x}) = \{(\bar{\zeta}, \bar{\Gamma})\}$ and $\text{aff}(C(\bar{x})) = \text{app}(\bar{\zeta}, \bar{\Gamma})$. The strong second order sufficient condition (47) then takes the following form:

$$\langle d, \mathcal{J}_{xx}^2 L(\bar{x}, \bar{\zeta}, \bar{\Gamma})d \rangle - \Upsilon_{g(\bar{x})}(\bar{\Gamma}, \mathcal{J}_x g(\bar{x})d) > 0 \quad \forall d \in \text{aff}(C(\bar{x})) \setminus \{0\}. \quad (54)$$

Let W be an arbitrary element in $\partial F(\bar{x}, \bar{\zeta}, \bar{\Gamma})$. We shall show that W is nonsingular. Let $(\Delta x, \Delta \zeta, \Delta \Gamma) \in X \times \Re^m \times \mathcal{S}^p$ be such that

$$W(\Delta x, \Delta \zeta, \Delta \Gamma) = 0.$$

Let $A := g(\bar{x}) + \bar{\Gamma}$. Without loss of generality, we assume that A has the spectral decomposition as in (15) and $g(\bar{x})$ and $\bar{\Gamma}$ satisfy (34). Then, by Lemma 1, we know that there exists a $V \in \partial \Pi_{\mathcal{S}_+^p}(A)$ such that

$$W(\Delta x, \Delta \zeta, \Delta \Gamma) = \begin{bmatrix} \mathcal{J}_{xx}^2 L(\bar{x}, \bar{\zeta}, \bar{\Gamma})\Delta x + \mathcal{J}_x h(\bar{x})^* \Delta \zeta + \mathcal{J}_x g(\bar{x})^* \Delta \Gamma \\ -\mathcal{J}_x h(\bar{x})\Delta x \\ -\mathcal{J}_x g(\bar{x})\Delta x + V(\mathcal{J}_x g(\bar{x})\Delta x + \Delta \Gamma) \end{bmatrix} = 0. \quad (55)$$

From Proposition 4, (39), and the second and the third equations of (55) we know that

$$\Delta x \in \text{app}(\bar{\zeta}, \bar{\Gamma}) = \text{aff}(C(\bar{x})). \quad (56)$$

By the first and second equations of (55), we obtain that

$$\begin{aligned} 0 &= \langle \Delta x, \mathcal{J}_{xx}^2 L(\bar{x}, \bar{\zeta}, \bar{\Gamma})\Delta x + \mathcal{J}_x h(\bar{x})^* \Delta \zeta + \mathcal{J}_x g(\bar{x})^* \Delta \Gamma \rangle \\ &= \langle \Delta x, \mathcal{J}_{xx}^2 L(\bar{x}, \bar{\zeta}, \bar{\Gamma})\Delta x \rangle + \langle \Delta x, \mathcal{J}_x h(\bar{x})^* \Delta \zeta \rangle + \langle \Delta x, \mathcal{J}_x g(\bar{x})^* \Delta \Gamma \rangle \\ &= \langle \Delta x, \mathcal{J}_{xx}^2 L(\bar{x}, \bar{\zeta}, \bar{\Gamma})\Delta x \rangle + \langle \Delta \zeta, \mathcal{J}_x h(\bar{x})\Delta x \rangle + \langle \Delta \Gamma, \mathcal{J}_x g(\bar{x})\Delta x \rangle \\ &= \langle \Delta x, \mathcal{J}_{xx}^2 L(\bar{x}, \bar{\zeta}, \bar{\Gamma})\Delta x \rangle + \langle \mathcal{J}_x g(\bar{x})\Delta x, \Delta \Gamma \rangle, \end{aligned}$$

which, together with the third equation of (55) and Proposition 7, implies that

$$0 \geq \langle \Delta x, \mathcal{J}_{xx}^2 L(\bar{x}, \bar{\zeta}, \bar{\Gamma})\Delta x \rangle - \Upsilon_{g(\bar{x})}(\bar{\Gamma}, \mathcal{J}_x g(\bar{x})\Delta x). \quad (57)$$

Hence, we can conclude from (56), (57), and the strong second order sufficient condition (54) that

$$\Delta x = 0.$$

Thus, (55) reduces to

$$\begin{bmatrix} \mathcal{J}_x h(\bar{x})^* \Delta \zeta + \mathcal{J}_x g(\bar{x})^* \Delta \Gamma \\ V(\Delta \Gamma) \end{bmatrix} = 0. \quad (58)$$

From Proposition 4 and $V(\Delta \Gamma) = 0$, we obtain that

$$P_\alpha^T \Delta \Gamma P_\alpha = 0, \quad P_\alpha^T \Delta \Gamma P_\beta = 0, \quad \text{and} \quad P_\alpha^T \Delta \Gamma P_\gamma = 0. \quad (59)$$

By the constraint nondegeneracy condition (49), there exist a vector $d \in X$ and a matrix $S \in \text{lin}(\mathcal{T}_{\mathcal{S}_+^p}(g(\bar{x})))$ such that

$$\mathcal{J}_x h(\bar{x})d = \Delta \zeta \quad \text{and} \quad \mathcal{J}_x g(\bar{x})d + S = \Delta \Gamma. \quad (60)$$

Hence, by (60) and the first equation of (58), we obtain

$$\begin{aligned} \langle \Delta \zeta, \Delta \zeta \rangle + \langle \Delta \Gamma, \Delta \Gamma \rangle &= \langle \mathcal{J}_x h(\bar{x})d, \Delta \zeta \rangle + \langle \mathcal{J}_x g(\bar{x})d + S, \Delta \Gamma \rangle \\ &= \langle d, \mathcal{J}_x h(\bar{x})^* \Delta \zeta \rangle + \langle d, \mathcal{J}_x g(\bar{x})^* \Delta \Gamma \rangle + \langle S, \Delta \Gamma \rangle \\ &= \langle d, \mathcal{J}_x h(\bar{x})^* \Delta \zeta + \mathcal{J}_x g(\bar{x})^* \Delta \Gamma \rangle + \langle S, \Delta \Gamma \rangle \\ &= \langle S, \Delta \Gamma \rangle \\ &= \langle P^T S P, P^T \Delta \Gamma P \rangle, \end{aligned}$$

which, together with (59) and (36), implies that

$$\langle \Delta \zeta, \Delta \zeta \rangle + \langle \Delta \Gamma, \Delta \Gamma \rangle = \langle P^T S P, P^T \Delta \Gamma P \rangle = 0.$$

Thus,

$$\Delta \zeta = 0 \quad \text{and} \quad \Delta \Gamma = 0.$$

This, together with $\Delta x = 0$, shows that W is nonsingular.

“(b) \implies (c)” By Clarke’s inverse function theorem [6, 7], F is a locally Lipschitz homeomorphism near $(\bar{x}, \bar{\zeta}, \bar{\Gamma})$. Thus, by Remark 15, $(\bar{x}, \bar{\zeta}, \bar{\Gamma})$ is a strongly regular solution of the generalized equation (52). Q.E.D.

In Proposition 16, it is shown that (a) \implies (b) \implies (c). In the next section, we shall show that if \bar{x} is a locally optimal solution to the nonlinear semidefinite programming (NLSDP) and Robinson’s CQ holds at \bar{x} , then these three statements are actually equivalent to each other.

4 Equivalent Conditions

We first introduce a uniform version of the second order growth condition defined in [4, Definition 5.16]. Let U be a Banach space and $f : X \times U \rightarrow \Re$ and $G : X \times U \rightarrow Y$. We

say that $(f(x, u), G(x, u))$, with $u \in U$, is a \mathcal{C}^2 -smooth parameterization of the optimization problem (OP) if $f(\cdot, \cdot)$ and $g(\cdot, \cdot)$ are twice continuously differentiable and there exists a $\bar{u} \in U$ such that $f(\cdot, \bar{u}) = f(\cdot)$ and $G(\cdot, \bar{u}) = G(\cdot)$. The corresponding parameterized problem takes the form:

$$\begin{aligned} (OP_u) \quad & \min \quad f(x, u) \\ & \text{s.t.} \quad G(x, u) \in K, \\ & \quad \quad x \in X. \end{aligned} \tag{61}$$

We say that a parameterization is *canonical* if $U := X \times Y$, $\bar{u} = (0, 0) \in X \times Y$, and

$$(f(x, u), G(x, u)) := (f(x) - \langle u_1, x \rangle, G(x) + u_2), \quad x \in X, \quad u := (u_1, u_2) \in X \times Y.$$

Definition 17 Let \bar{x} be a stationary point of the optimization problem (OP). We say that the uniform second order (quadratic) growth condition holds at \bar{x} with respect to a \mathcal{C}^2 -smooth parameterization $(f(x, u), G(x, u))$ if there exist $c > 0$ and neighborhoods \mathcal{V}_X of \bar{x} and \mathcal{V}_U of \bar{u} such that for any $u \in \mathcal{V}_U$ and any stationary point $x(u) \in \mathcal{V}_X$ of the corresponding parameterized problem (OP_u) , the following holds:

$$f(x, u) \geq f(x(u), u) + c\|x - x(u)\|^2 \quad \forall x \in \mathcal{V}_X \text{ such that } G(x, u) \in K. \tag{62}$$

We say that the uniform second order growth condition holds at \bar{x} if (62) holds for every \mathcal{C}^2 -smooth parameterization of (OP).

The next lemma shows that for the nonlinear semidefinite programming (NLSDP) the uniform second order growth condition implies the strong second order sufficient condition.

Lemma 18 Let \bar{x} be a stationary point of the nonlinear semidefinite programming (NLSDP). Suppose that Robinson's CQ holds at \bar{x} . If the uniform second order growth condition holds at \bar{x} with respect to the canonical parameterization, then the strong second order sufficient condition (47) holds at \bar{x} .

Proof. Let $(\bar{\zeta}, \bar{\Gamma}) \in \mathcal{M}(\bar{x})$. We may assume that $A := g(\bar{x}) + \bar{\Gamma}$ has the spectral decomposition as in (15) and $g(\bar{x})$ and $\bar{\Gamma}$ satisfy (34). Consider the following parameterized nonlinear semidefinite programming problem:

$$\begin{aligned} \min \quad & f(x) \\ \text{s.t.} \quad & h(x) = 0, \\ & g(x) + \tau P_\beta P_\beta^T \in \mathcal{S}_+^p, \\ & x \in X, \end{aligned} \tag{63}$$

where $\tau \in \Re$ is a parameter. Then, for any $\tau > 0$, $(\bar{\zeta}, \bar{\Gamma})$, together with \bar{x} , satisfies the following KKT condition of the parameterized problem (63):

$$\mathcal{J}_x L_\tau(\bar{x}, \zeta, \Gamma) = \mathcal{J}_x L(\bar{x}, \zeta, \Gamma) = 0, \quad -h(\bar{x}) = 0, \quad \text{and} \quad \Gamma \in \mathcal{N}_{\mathcal{S}_+^p}(g(\bar{x}) + \tau P_\beta P_\beta^T), \tag{64}$$

where for each $\tau \in \mathbb{R}$,

$$L_\tau(x, \zeta, \Gamma) := L(x, \zeta, \Gamma) + \tau \langle \Gamma, P_\beta P_\beta^T \rangle, \quad (x, \zeta, \Gamma) \in X \times \mathbb{R}^m \times \mathcal{S}^p.$$

Let $\mathcal{M}_\tau(\bar{x})$ be the set consisting of all $(\zeta, \Gamma) \in \mathbb{R}^m \times \mathcal{S}^p$ that satisfy (64). Thus, since $\text{rank}(g(\bar{x}) + \tau P_\beta P_\beta^T) + \text{rank}(\bar{\Gamma}) = p$ for any $\tau > 0$, the critical cone $C_\tau(\bar{x})$ of the parameterized problem (63) at \bar{x} for $\tau > 0$ takes the form:

$$C_\tau(\bar{x}) = \{d : \mathcal{J}_x h(\bar{x})d = 0, \quad P_\gamma^T(\mathcal{J}_x g(\bar{x})d)P_\gamma = 0\} \supseteq \text{app}(\bar{\zeta}, \bar{\Gamma}), \quad (65)$$

where $\text{app}(\bar{\zeta}, \bar{\Gamma})$ satisfies (39). Therefore, by Lemma 11 and the second part of Theorem 10, we have for all $\tau > 0$ that

$$\sup_{(\zeta, \Gamma) \in \mathcal{M}_\tau(\bar{x})} \left\{ \langle d, \mathcal{J}_{xx}^2 L_\tau(\bar{x}, \zeta, \Gamma)d \rangle - \Upsilon_{(g(\bar{x}) + \tau P_\beta P_\beta^T)}(\Gamma, \mathcal{J}_x g(\bar{x})d) \right\} > 0 \quad \forall d \in C_\tau(\bar{x}) \setminus \{0\},$$

which, together with the fact that for any $(\zeta, \Gamma) \in \mathcal{M}_\tau(\bar{x})$,

$$\Upsilon_{(g(\bar{x}) + \tau P_\beta P_\beta^T)}(\Gamma, \mathcal{J}_x g(\bar{x})d) = \Upsilon_{g(\bar{x})}(\Gamma, \mathcal{J}_x g(\bar{x})d) \quad \forall d \in \text{app}(\bar{\zeta}, \bar{\Gamma}),$$

$\mathcal{J}_{xx}^2 L_\tau(\bar{x}, \zeta, \Gamma) = \mathcal{J}_{xx}^2 L(\bar{x}, \zeta, \Gamma)$, and (65), implies

$$\sup_{(\zeta, \Gamma) \in \mathcal{M}_\tau(\bar{x})} \left\{ \langle d, \mathcal{J}_{xx}^2 L(\bar{x}, \zeta, \Gamma)d \rangle - \Upsilon_{g(\bar{x})}(\Gamma, \mathcal{J}_x g(\bar{x})d) \right\} > 0 \quad \forall d \in \text{app}(\bar{\zeta}, \bar{\Gamma}) \setminus \{0\}. \quad (66)$$

Since for any $\tau > 0$, $\mathcal{M}_\tau(\bar{x}) \subseteq \mathcal{M}(\bar{x})$, we derive from (66) that

$$\sup_{(\zeta, \Gamma) \in \mathcal{M}(\bar{x})} \left\{ \langle d, \mathcal{J}_{xx}^2 L(\bar{x}, \zeta, \Gamma)d \rangle - \Upsilon_{g(\bar{x})}(\Gamma, \mathcal{J}_x g(\bar{x})d) \right\} > 0 \quad \forall d \in \text{app}(\bar{\zeta}, \bar{\Gamma}) \setminus \{0\}.$$

This shows that the strong second order sufficient condition (47) holds. Q.E.D.

Another important concept in sensitivity and stability analysis of the optimization problem (OP) is the *strong stability* of a stationary point, introduced by Kojima in [17]. Below the definition of strong stability is from Bonnans and Shapiro [4, Definition 5.3].

Definition 19 *Let \bar{x} be a stationary point of the optimization problem (OP). We say that \bar{x} is strongly stable with respect to a \mathcal{C}^2 -smooth parameterization $(f(x, u), G(x, u))$ if there exist neighborhoods \mathcal{V}_X of \bar{x} and \mathcal{V}_U of \bar{u} such that for any $u \in \mathcal{V}_U$, the corresponding perturbed problem (OP_u) has a unique stationary point $x(u) \in \mathcal{V}_X$ and $x(\cdot)$ is continuous on \mathcal{V}_U . If this holds for any \mathcal{C}^2 -smooth parameterization, we say that \bar{x} is strongly stable.*

Let $F : X \times \mathbb{R}^m \times \mathcal{S}^p \rightarrow X \times \mathbb{R}^m \times \mathcal{S}^p$ be defined as in (51) and $(\bar{x}, \bar{\zeta}, \bar{\Gamma}) \in X \times \mathbb{R}^m \times \mathcal{S}^p$ be a KKT point of (NLSDP). Then

$$F(\bar{x}, \bar{\zeta}, \bar{\Gamma}) = \begin{bmatrix} \mathcal{J}_x L(\bar{x}, \bar{\zeta}, \bar{\Gamma}) \\ -h(\bar{x}) \\ -g(\bar{x}) + \Pi_{\mathcal{S}_+^p}(g(\bar{x}) + \bar{\Gamma}) \end{bmatrix} = 0.$$

Let $A := g(\bar{x}) + \bar{\Gamma}$. Assume that A has the spectral decomposition as in (15) and $g(\bar{x})$ and $\bar{\Gamma}$ satisfy (34). Then, F is directionally differentiable at $(\bar{x}, \bar{\zeta}, \bar{\Gamma})$ and for any $\delta := (\delta_1, \delta_2, \delta_3) \in X \times \mathfrak{R}^m \times \mathcal{S}^p$,

$$\Phi(\delta) := F'(\bar{x}, \bar{\zeta}, \bar{\Gamma}; \delta) = \begin{bmatrix} \mathcal{J}_{xx}^2 L(\bar{x}, \bar{\zeta}, \bar{\Gamma}) \delta_1 + \mathcal{J}_x h(\bar{x})^* \delta_2 + \mathcal{J}_x g(\bar{x})^* \delta_3 \\ -\mathcal{J}_x h(\bar{x}) \delta_1 \\ -\mathcal{J}_x g(\bar{x}) \delta_1 + \Pi'_{\mathcal{S}_+^p}(A; \mathcal{J}_x g(\bar{x}) \delta_1 + \delta_3) \end{bmatrix}, \quad (67)$$

where $\Pi'_{\mathcal{S}_+^p}(A; \cdot)$ is given by (16). Since $\Phi(\cdot)$ is Lipschitz continuous, $\partial_B \Phi(0)$ is well defined.

Lemma 20 *Let $A = g(\bar{x}) + \bar{\Gamma}$ and Φ be defined by (67). It holds that*

$$\partial_B \Phi(0) = \partial_B F(\bar{x}, \bar{\zeta}, \bar{\Gamma}).$$

Proof. Define $\Xi : X \times \mathfrak{R}^m \times \mathcal{S}^p \rightarrow \mathcal{S}^p$ by

$$\Xi(\delta) := \Pi'_{\mathcal{S}_+^p}(A; \Psi(\delta)),$$

where $\Psi(\delta) := \mathcal{J}_x g(\bar{x}) \delta_1 + \delta_3$, $\delta := (\delta_1, \delta_2, \delta_3) \in X \times \mathfrak{R}^m \times \mathcal{S}^p$. Then, by Lemma 1 and (c) of Proposition 3, we obtain that

$$\partial_B \Xi(0) = \partial_B \Pi_{\mathcal{S}_+^p}(A) \mathcal{J}_\delta \Psi(0),$$

which, together with Lemma 1, proves the conclusion of this lemma. Q.E.D.

Let $\text{ind}(\phi, \bar{z})$ denote the index of a continuous function $\phi : Z \rightarrow Z$ at its zero $\bar{z} \in Z$ used in degree theory [19, 23]. Now, we are ready to state the main result of this paper.

Theorem 21 *Let \bar{x} be a locally optimal solution to the nonlinear semidefinite programming (NLSDP). Suppose that Robinson's CQ (41) holds at \bar{x} so that \bar{x} is necessarily a stationary point of (NLSDP). Let $(\bar{\zeta}, \bar{\Gamma}) \in \mathfrak{R}^m \times \mathcal{S}^p$ be such that $(\bar{x}, \bar{\zeta}, \bar{\Gamma})$ is a KKT point of (NLSDP). Then the following statements are equivalent:*

- (a) *The strong second order sufficient condition (47) holds at \bar{x} and \bar{x} is constraint nondegenerate.*
- (b) *Any element in $\partial F(\bar{x}, \bar{\zeta}, \bar{\Gamma})$ is nonsingular.*
- (c) *The KKT point $(\bar{x}, \bar{\zeta}, \bar{\Gamma})$ is a strongly regular solution of the generalized equation (52).*
- (d) *The uniform second order growth condition holds at \bar{x} and \bar{x} is constraint nondegenerate.*
- (e) *The point \bar{x} is strongly stable and \bar{x} is constraint nondegenerate.*
- (f) *F is a locally Lipschitz homeomorphism near the KKT point $(\bar{x}, \bar{\zeta}, \bar{\Gamma})$.*

(g) For every $V \in \partial_B F(\bar{x}, \bar{\zeta}, \bar{\Gamma})$, $\text{sgn det } V = \text{ind}(F, (\bar{x}, \bar{\zeta}, \bar{\Gamma})) = \pm 1$.

(h) Φ is a globally Lipschitz homeomorphism.

(i) For every $V \in \partial_B \Phi(0)$, $\text{sgn det } V = \text{ind}(\Phi, 0) = \pm 1$.

(j) Any element in $\partial \Phi(0)$ is nonsingular.

Proof. We have already known from Proposition 16 that (a) \implies (b) \implies (c) and from Remark 15 that (c) \iff (f). The relations (c) \iff (d) \iff (e) follow from Bonnans and Shapiro [4, Theorems 5.24 & 5.35]. Since $\Pi_{\mathcal{S}_+^p}(\cdot)$ is strongly semismooth everywhere [37], F is a semismooth function (see [22, 25] for discussions on semismooth functions). Then, by Gowda [12, Theorem 3 & Corollary 4], we know that (f) \iff (g). Furthermore, by the semismoothness of $\Pi_{\mathcal{S}_+^p}(\cdot)$, Lemma 20, and Theorem 6 in [24], it holds that (g) \iff (h) \iff (i). By Lemma 20, the relation (b) \iff (j) holds. The proof of this theorem will be completed if one can show that (d) \implies (a). The latter, however, is implied by Lemma 18. Q.E.D.

Remark 22 As mentioned in Section 1, the equivalence between (a) and (c) has already been known for the conventional nonlinear programming (NLP). This equivalence for (NLP) follows from Jongen et al. [14, Theorem 3.1], Robinson [28, Theorem 4.1] and Kojima [17, Corollary 6.6]. For different proofs, see [4, Proposition 5.38], [5, Theorem 4.10], and [8, Theorem 6]. By assuming \bar{x} to be a stationary point (not necessarily a local optimal solution), for (NLP), Jongen, Klatte, and Tammer in [15] proved (b) \iff (e) for a different but equivalent KKT system. By focusing on the local optimal solution case only, we extend these results in Theorem 21 from (NLP) to (NLSDP).

5 Conclusions

In this paper, we discussed a strong second order sufficient condition for the nonlinear semidefinite programming (NLSDP). This strong second order sufficient condition, together with constraint nondegeneracy, is shown to be equivalent to many conditions, notably the strong regularity of the KKT point and the nonsingularity of Clarke's Jacobian of the mapping F at the KKT point. There are many important questions not addressed in this paper. For example, it would be interesting to know whether these equivalent results given in Theorem 21 can be generalized to other \mathcal{C}^2 -cone reducible sets. Another possibility is to see which of these conditions are still equivalent to (b) or (j) if \bar{x} is assumed to be a stationary point only.

References

- [1] J.F. BONNANS, R. COMINETTI, AND A. SHAPIRO. Sensitivity analysis of optimization problems under second order regularity constraints. *Mathematics of Operations Research* 23 (1998) 803–832.
- [2] J.F. BONNANS, R. COMINETTI, AND A. SHAPIRO. Second order optimality conditions based on parabolic second order tangent sets. *SIAM Journal on Optimization* 9 (1999) 466–493.
- [3] J.F. BONNANS AND A. SHAPIRO. Optimization problems with perturbations: A guided tour. *SIAM Review* 40 (1998) 202–227.
- [4] J.F. BONNANS AND A. SHAPIRO. *Perturbation Analysis of Optimization Problems*, Springer (New York, 2000).
- [5] J.F. BONNANS AND A. SULEM. Pseudopower expansion of solutions of generalized equations and constrained optimization problems. *Mathematical Programming* 70 (1995) 123–148.
- [6] F.H. CLARKE. On the inverse function theorem. *Pacific Journal of Mathematics* 64 (1976) 97–102.
- [7] F.H. CLARKE. *Optimization and Nonsmooth Analysis*, John Wiley and Sons (New York, 1983).
- [8] A.L. DONTCHEV AND R.T. ROCKAFELLAR. Characterizations of strong regularity for variational inequalities over polyhedral convex sets. *SIAM Journal on Optimization* 6 (1996) 1087–1105.
- [9] B.C. EAVES. On the basic theorem of complementarity. *Mathematical Programming* 1 (1971) 68–75.
- [10] F. FACCHINEI AND J.S. PANG. *Finite-Dimensional Variational Inequalities and Complementarity Problems: Volume I*, Springer (New York, 2003).
- [11] J. GAUVIN. A necessary and sufficient regularity condition to have bounded multipliers in nonconvex programming. *Mathematical Programming* 12 (1977) 136–138.
- [12] M.S. GOWDA. Inverse and implicit function theorems for H-differentiable and semismooth functions. *Optimization Methods and Software* 19 (2004) 443–461.
- [13] N.J. HIGHAM. Computing a nearest symmetric positive semidefinite matrix. *Linear Algebra and Applications* 103 (1988) 103–118.

- [14] H.TH. JONGEN, T. MOBERT, J. RÜCKMANN, AND K. TAMMER. On inertia and Schur complement in optimization. *Linear Algebra and its Applications* 95 (1987) 97–109.
- [15] H.TH. JONGEN, J. RÜCKMANN, AND K. TAMMER. Implicit functions and sensitivity of stationary points. *Mathematical Programming* 49 (1990) 123–138.
- [16] D. KLATTE AND B. KUMMER. *Nonsmooth Equations in Optimization: Regularity, Calculus, Methods and Applications*, Kluwer Academic Publishers (Boston, 2002).
- [17] M. KOJIMA. Strongly stable stationary solutions in nonlinear programs. In: S.M. Robinson, editor, *Analysis and Computation of Fixed Points*, Academic Press (New York, 1980), pp. 93–138.
- [18] B. KUMMER. Lipschitzian inverse functions, directional derivatives, and applications in $C^{1,1}$ -optimization. *Journal of Optimization Theory and Applications* 70 (1991) 559–580.
- [19] N.G. LLOYD. *Degree Theory*, Cambridge University Press (Cambridge, 1978).
- [20] O.L. MANGASARIAN AND S. FROMOVITZ. The Fritz-John necessary optimality conditions in the presence of equality and inequality constraints. *Journal of Mathematical Analysis and Applications* 7 (1967) 37–47.
- [21] F. MENG, D. SUN, AND G. ZHAO. Semismoothness of solutions to generalized equations and the Moreau-Yosida regularization. *Mathematical Programming, Series B*, to appear.
- [22] R. MIFFLIN. Semismooth and semiconvex functions in constrained optimization. *SIAM Journal on Control and Optimization* 15 (1977) 957–972.
- [23] J.M. ORTEGA AND W.C. RHEINBOLDT. *Iterative Solution of Nonlinear Equations in Several Variables*, Academic Press (New York, 1970).
- [24] J.S. PANG, D. SUN, AND J. SUN. Semismooth homeomorphisms and strong stability of semidefinite and Lorentz complementarity problems. *Mathematics of Operations Research* 28 (2003) 39–63.
- [25] L. QI AND J. SUN. A nonsmooth version of Newton’s method. *Mathematical Programming* 58 (1993) 353–367.
- [26] S.M. ROBINSON. Stability theory for systems of inequalities, Part I: Linear systems. *SIAM Journal on Numerical Analysis* 12 (1975) 754–769.
- [27] S.M. ROBINSON. First order conditions for general nonlinear optimization. *SIAM Journal on Applied Mathematics* 30 (1976) 597–607.

- [28] S.M. ROBINSON. Strongly regular generalized equations. *Mathematics of Operations Research* 5 (1980) 43–62.
- [29] S.M. ROBINSON. Local structure of feasible sets in nonlinear programming, Part II: Nondegeneracy. *Mathematical Programming Study* 22 (1984) 217–230.
- [30] S.M. ROBINSON. Local structure of feasible sets in nonlinear programming, Part III: Stability and sensitivity. *Mathematical Programming Study* 30 (1987) 45–66.
- [31] S.M. ROBINSON. An implicit-function theorem for a class of nonsmooth functions. *Mathematics of Operations Research* 16 (1991) 292–309.
- [32] S.M. ROBINSON. Constraint nondegeneracy in variational analysis. *Mathematics of Operations Research* 28 (2003) 201–232.
- [33] R.T. ROCKAFELLAR. *Convex Analysis*, Princeton (New Jersey, 1970).
- [34] R.T. ROCKAFELLAR AND R.J.-B. WETS. *Variational Analysis*, Springer (New York, 1998).
- [35] A. SHAPIRO. First and second order analysis of nonlinear semidefinite programs. *Mathematical Programming, Series B* 77 (1997) 301–320.
- [36] A. SHAPIRO. Sensitivity analysis of generalized equations. *Journal of Mathematical Sciences* 115 (2003) 2554–2565.
- [37] D. SUN AND J. SUN. Semismooth matrix valued functions. *Mathematics of Operations Research* 27 (2002) 150–169.
- [38] P. TSENG, Merit functions for semi-definite complementarity problems. *Mathematical Programming* 83 (1998) 159–185.
- [39] E.H. ZARANTONELLO. Projections on convex sets in Hilbert space and spectral theory I and II. In E.H. Zarantonello, editor, *Contributions to Nonlinear Functional Analysis*, Academic Press (New York, 1971) pp. 237–424.