

Dynamic updates of the barrier parameter in primal-dual methods for nonlinear programming

Paul ARMAND,[†] Joël BENOIST[†] and Dominique ORBAN[‡]

Abstract. We introduce a framework in which updating rules for the barrier parameter in primal-dual interior-point methods become dynamic. The original primal-dual system is augmented to incorporate explicitly an updating function. A Newton step for the augmented system gives a primal-dual Newton step and also a step in the barrier parameter. Based on local information and a linesearch, the decrease of the barrier parameter is automatically adjusted. We analyze local convergence properties, report numerical experiments on a standard collection of nonlinear problems and compare our results to a state-of-the-art interior-point implementation. In many instances, the adaptive algorithm reduces the number of iterations and of function evaluations. Its design guarantees a better fit between the magnitudes of the primal-dual residual and of the barrier parameter along the iterations.

Key words. constrained optimization, interior point method, nonlinear programming, primal-dual method, barrier method

AMS subject classification. 65K05, 90C06, 90C26, 90C30, 90C51

1 Introduction

We consider nonlinear minimization problems of the form

$$\begin{aligned} \min \quad & f(x), \\ \text{s.t.} \quad & c(x) = 0, \\ & x \geq 0, \end{aligned} \tag{1.1}$$

where $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and $c : \mathbb{R}^n \rightarrow \mathbb{R}^m$ are twice continuously differentiable. Let $(y, z) \in \mathbb{R}^m \times \mathbb{R}^n$ denote a vector of Lagrange multipliers associated to the constraints, $w = (x, y, z)$, $v = (x, z)$ and define the Lagrangian function $\ell(w) = f(x) + y^\top c(x) - z^\top x$ associated to problem (1.1). The first order optimality conditions of problem (1.1) can be written

$$F(w) = 0 \quad \text{and} \quad v \geq 0 \tag{1.2}$$

where $F : \mathbb{R}^{2n+m} \rightarrow \mathbb{R}^{2n+m}$ is defined by

$$F(w) = \begin{pmatrix} \nabla_x \ell(w) \\ c(x) \\ XZe \end{pmatrix}, \tag{1.3}$$

[†]Université de Limoges (France); e-mail: Paul.Armand@unilim.fr, Joel.Benoist@unilim.fr.

[‡]Ecole Polytechnique de Montréal (Canada); e-mail: Dominique.Orban@polymtl.ca.

where $e = (1 \cdots 1)$ is the vector of all ones. We use X to denote the diagonal matrix $\text{diag}(x_1, \dots, x_n)$ and employ similar notation for other quantities.

Primal-dual interior point methods apply a Newton-type method to identify an approximate solution w_k to the *primal-dual system*

$$F(w) = \mu \tilde{e}, \quad (1.4)$$

where $\tilde{e} = (0 \ 0 \ e^\top)^\top$ and μ is held fixed at some value $\mu_k > 0$, while enforcing $v_k > 0$ —a process referred to as *inner* iterations. The value of μ_k is then decreased to ensure that $\{\mu_k\} \downarrow 0$ and a new series of inner iterations begins. The effort involved in an inner iteration can be significant, especially when far from a solution to (1.1). Typically, a sequence of inner iterations terminates as soon as

$$\|F(w_k) - \mu_k \tilde{e}\| \leq \varepsilon_k, \quad (1.5)$$

for some tolerance $\varepsilon_k > 0$ converging to zero as $\mu_k \downarrow 0$. In order to find an appropriate starting point for the next sequence of inner iterations and to guarantee fast local convergence, an extrapolation step is often performed from w_k . A possibility, originally devised in a purely primal framework, is to linearize the primal optimality conditions about (w_k, μ_k) [4]. It is now known that, in the primal-dual framework, this is equivalent to a Newton step for (1.4) using the *new* value μ_{k+1} [8]

$$F'(w_k)d_k + F(w_k) = \mu_{k+1}\tilde{e}. \quad (1.6)$$

If w_k satisfies (1.5) and if μ_k and ε_k are appropriately updated, this scheme preserves strict feasibility and guarantees asymptotic Q-subquadratic convergence [2, 8, 9].

In this paper, we study the possibility of adjusting μ dynamically as the iterations proceed based on local information at the current point. This proposal is motivated by considering the augmented system

$$\begin{pmatrix} F(w) - \mu \tilde{e} \\ \theta(\mu) \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad (1.7)$$

where θ is a scalar function defined on \mathbb{R}_+ , differentiable, increasing, strictly convex and such that $\theta(0) = 0$. Clearly, the vector (w^*, μ^*) is a solution of (1.7) if and only if $\mu^* = 0$ and $F(w^*) = 0$. The Newton step for (1.7) from (w_k, μ_k) solves

$$\begin{pmatrix} F'(w_k) & -\tilde{e} \\ 0 & \theta'(\mu_k) \end{pmatrix} \begin{pmatrix} d_k^w \\ d_k^\mu \end{pmatrix} + \begin{pmatrix} F(w_k) - \mu_k \tilde{e} \\ \theta(\mu_k) \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}. \quad (1.8)$$

The first equation of (1.8) may be rewritten

$$F'(w_k)d_k^w + F(w_k) = (\mu_k + d_k^\mu)\tilde{e}, \quad (1.9)$$

which is just the system (1.6) with $\mu_{k+1} = \mu_k + d_k^\mu$. The value of d_k^μ is obtained from the second equation of (1.8)

$$d_k^\mu = -\frac{\theta(\mu_k)}{\theta'(\mu_k)}. \quad (1.10)$$

This point of view shows that both the decrease of the barrier parameter and the extrapolation step can be interpreted as one and only Newton step.

A first consequence of this formulation is that the tools of the convergence analysis of Newton's method can be directly applied to a local convergence analysis of the interior point algorithm. This point of view is developed in Section 4.

A second consequence is that a control of the decrease of the barrier parameter can be introduced in a natural manner; decreasing μ rather conservatively when far from a solution where the extrapolation step can be a poor direction of improvement, and more enthusiastically when closer to a solution. The algorithmic scheme that we consider is based on the application of a sequence of steps (1.8) while maintaining positivity of v . We introduce a steplength, computed by a linesearch, in the updating formula for the barrier parameter. A first candidate steplength is given by the so-called *fraction to the boundary* rule: the largest $\alpha \in (0, 1]$ ensuring

$$v_k + \alpha d_k^v \geq (1 - \tau_k)v_k, \quad (1.11)$$

where $\tau_k \in (0, 1)$. A natural choice for the next value of the barrier parameter is then

$$\mu_{k+1} = \mu_k + \alpha_k d_k^\mu. \quad (1.12)$$

A second candidate steplength is given by a sufficient decrease condition on a merit function. Finally, we explore the possibility of updating μ at each inner iteration. This results in an algorithm which is not warranted by a global convergence theory, however we illustrate that it possesses some stabilizing properties. The benefits of damping the update of μ are illustrated by some numerical experiments in Section 7.

Besides offering a framework in which μ is naturally updated adaptively, the scheme (1.9)–(1.10) attempts to resolve two numerical difficulties inherent to interior-point methods. The first is that if μ decreases too fast too early, the iteration may not finish within the allowed time limits. The reason for this is a worsening discrepancy between the magnitude of μ and the magnitude of the residual $\|F(w) - \mu\bar{e}\|$, causing (1.5) to require many inner iterations. An illustration of this behaviour is given in Fig. 7.1 with a state-of-the-art implementation. We will show how an algorithm based on (1.9)–(1.10) does not suffer from such a discrepancy. The second numerical difficulty is that it is nearly impossible to know in practice when μ may be allowed to decrease faster. If μ decreases fast too early, the first difficulty may arise and if it decreases fast too late, we may be spending too much time and effort solving (1.4).

Under standard assumptions on (1.1), as $\mu \downarrow 0$, exact solutions to (1.4) may be expressed as an implicit function $w(\mu)$ and trace a smooth trajectory, called the central path, converging to a solution w^* of (1.1) as $\mu \downarrow 0$ [6]. For schemes based on proximity to the central path as enforced by (1.5), under standard regularity conditions, the local convergence rate of $\{w_k\}$ to w^* is directly tied to the convergence rate of $\{\mu_k\}$ to zero. The positivity requirement $v > 0$ imposes that μ cannot decrease too fast in the sense that

$$\mu_{k+1} \geq \kappa \mu_k^{2-\epsilon}$$

for some constant $\kappa > 0$ and some $\epsilon > 0$ as small as desired. Various rules for updating μ can be found in the literature. They range from the simplest, but globally convergent, monotonic decreases of μ , rules that draw inspiration from linear programming to rules of a different nature which are based on a measure of centrality. Most algorithms decrease μ linearly and will attempt to recover fast local convergence by switching to a rule which decreases μ superlinearly as soon as it is declared sufficiently small [15]. However, the notion *sufficiently small* is problem dependent and, it remains unclear how to wisely switch from one rule to the other. In [7], the authors use a rule which mimics rules based on the duality gap in linear and convex programming but enforces that $\{\mu_k\}$ be decreasing. Numerically, the monotonicity of $\{\mu_k\}$ can be a source of difficulties and more dynamic rules are investigated, often at the expense of global convergence. An example is given by [5] where the barrier parameter is allowed to increase occasionally. Recently, the authors of [12] incorporated such a rule into a safeguarded globally convergent framework with a heuristic switch. At variance, the authors of [13] note the importance of keeping individual complementarity pairs clustered. They measure deviation from centrality and update μ accordingly in a manner similar to [11]. Unfortunately, this practice is not warranted by any global convergence theory and can cause failure if μ_k diverges or becomes too small prematurely.

The scheme (1.9)–(1.10), although fairly general, does not necessarily impose that new updating functions $\theta(\cdot)$ be devised. Rather, known updates may be used and the linesearch, by means of (1.11)–(1.12), will adjust d_μ as the iterations proceed. Two suitable updates will be given in §2. At variance with many traditional implementations, the sequence $\{\mu_k\}$ is not fixed ahead of time and moreover, is also adjusted dynamically even in the early iterations. The present framework alleviates the need for an artificial “switch” between a conservative—usually linear—update of μ in the early iterations and a more enthusiastic—usually superlinear—update as μ becomes small. In the literature, μ_k is only chosen on-the-fly in the late iterations using rules such as, e.g., $\mu_{k+1} = O(\|F(w_k) - \mu_k \bar{e}\|^2)$ [14, 17].

The paper is structured as follows. Section 2 describes the notation and assumptions used throughout the text. Section 3 presents a number of preliminary results concerning local solutions to equation (1.4) and properties of a Newton iteration. Section 4 studies local convergence properties of the sequences defined by (1.8) but

ignores the positivity constraints. The latter are addressed in Section 5 where conditions on the tolerance ε_k and on the decreasing rate of μ_k are derived to ensure that a single inner iteration is asymptotically sufficient. Finally, Section 6 presents three algorithm schemes and gives a general convergence result. Results of some numerical experiments are discussed in Section 7.

2 Notation and assumptions

Vector inequalities are understood componentwise. Given two vectors $x, y \in \mathbb{R}^n$, their Euclidean scalar product is denoted by $x^\top y$ and the associated ℓ_2 norm is $\|x\| = (x^\top x)^{1/2}$. The open Euclidean ball centered at x with radius $r > 0$ is denoted by $B(x, r)$, that is $B(x, r) := \{y : \|x - y\| < r\}$.

For two nonnegative scalar sequences $\{a_k\}$ and $\{b_k\}$ converging to zero, we use the Landau symbols $a_k = o(b_k)$ if $\lim_{k \rightarrow \infty} a_k/b_k = 0$ and $a_k = O(b_k)$ if there exists a constant $c > 0$, such that $a_k \leq cb_k$ for all sufficiently large k . We use similar symbols with vector arguments, in which case they are understood normwise. We write $a_k = \Theta(b_k)$ if both $a_k = O(b_k)$ and $b_k = O(a_k)$ hold. We also write $a_k \sim b_k$ if $\lim_{k \rightarrow \infty} a_k/b_k = 1$. For a positive function $\epsilon : \mathbb{R}_+ \rightarrow \mathbb{R}_+$, we write $\epsilon(t) = o(t)$ if $\lim_{t \downarrow 0} \epsilon(t)/t = 0$, $\epsilon(t) = O(t)$ if $\limsup_{t \downarrow 0} \epsilon(t)/t < +\infty$ and $\epsilon(t) = \Theta(t)$ if both $0 < \liminf_{t \downarrow 0} \epsilon(t)/t$ and $\limsup_{t \downarrow 0} \epsilon(t)/t < +\infty$ hold.

We denote by e_i the vectors of the canonical basis of \mathbb{R}^n . For all $x \in \mathbb{R}^n$, $\nabla c(x)$ denotes the transpose of the Jacobian matrix of c at x , i.e., the $n \times m$ matrix whose i -th column is $\nabla c_i(x)$. Let $x^* \in \mathbb{R}^n$ be a local solution of problem (1.1). Let $\mathcal{A} := \{i : x_i^* = 0\}$ be the set of indices of active inequality constraints. Throughout the paper, we assume that the following assumptions are satisfied.

Assumption 2.1 The functions f and c are twice continuously differentiable and their second derivatives are Lipschitz continuous over an open neighborhood of x^* .

Assumption 2.2 The linear independence constraint qualification holds at x^* , i.e., $\{\nabla c_i(x^*), i = 1, \dots, m\} \cup \{e_i, i \in \mathcal{A}\}$ is a linearly independent set of vectors.

Note that Assumptions 2.1 and 2.2 imply that there exists a strictly feasible point, i.e., $\bar{x} \in \mathbb{R}$ such that $c(\bar{x}) = 0$ and $\bar{x} > 0$, and that there exists a unique vector of Lagrange multipliers $(y^*, z^*) \in \mathbb{R}^{m+n}$ such that $w^* = (x^*, y^*, z^*)$ is a solution of (1.2).

Assumption 2.3 The strong second-order sufficiency condition is satisfied at w^* , i.e., $u^\top \nabla_{xx}^2 \ell(w^*) u > 0$ for all $u \neq 0$ satisfying $\nabla c(x^*)^\top u = 0$ and $u_i = 0$ for all $i \in \mathcal{A}$.

Assumption 2.4 Strict complementarity holds at w^* , that is

$$\min\{x_i^* + z_i^* : i = 1 \dots n\} > 0.$$

Under Assumptions 2.1-2.4, the Jacobian of F , defined by

$$F'(w) = \begin{pmatrix} \nabla_{xx}^2 \ell(w) & \nabla c(x) & -I \\ \nabla c(x)^\top & 0 & 0 \\ Z & 0 & X \end{pmatrix},$$

is uniformly nonsingular over a neighbourhood of w^* . This fact will allow us to state strong results on the solution to (1.4) as $\mu \downarrow 0$.

We end this section by giving some examples of function θ that can be used in equation (1.7). Our basic assumption on θ is

Assumption 2.5 The function $\theta : \mathbb{R}_+ \rightarrow \mathbb{R}$ is continuously differentiable, increasing, strictly convex and such that $\theta(0) = 0$.

A popular update is the linear update with ratio σ where $0 < \sigma < 1$. It is easy to verify that this update is given by

$$\theta_L(\mu) = \mu^{\frac{1}{1-\sigma}}, \quad \text{and} \quad \mu_{k+1} = \sigma \mu_k. \quad (2.1)$$

A second very common update is the q -superlinear update given by

$$\theta_S(\mu) = \frac{\mu}{(b^\gamma - \mu^\gamma)^{\frac{1}{\gamma}}} \text{ for } \mu \in [0, b) \quad \text{and} \quad \mu_{k+1} = \frac{1}{b^\gamma} \mu_k^{1+\gamma} \quad (2.2)$$

for some parameters $b > 0$ and $0 < \gamma < 1$. Such a rule is used in, e.g., [2, p. 52].

3 Preliminary results

In this section we give some useful results about the solution of equation (1.4).

Lemma 3.1 *Under Assumptions 2.1–2.2, the Jacobian F' is locally Lipschitzian about w^* , i.e., there exist $L > 0$ and $\delta_L > 0$ such that, for all $w_1, w_2 \in B(w^*, \delta_L)$,*

$$\|F'(w_1) - F'(w_2)\| \leq L \|w_1 - w_2\|.$$

The next two lemmas are consequences of the nonsingularity of $F'(w^*)$ (see e.g. [6]).

Lemma 3.2 *Under Assumptions 2.1–2.4, there exist $\delta_K > 0$ and $K > 0$ such that, for all $w \in B(w^*, \delta_K)$ the Jacobian $F'(w)$ is nonsingular and*

$$\frac{1}{K} \leq \|F'(w)^{-1}\| \leq K.$$

Lemma 3.3 *Under Assumptions 2.1–2.4, there exist $C > 0$, $\delta_C > 0$, $\mu_C > 0$ and a continuously differentiable function $w(\cdot) : (-\mu_C, \mu_C) \rightarrow \mathbb{R}^{n+m+n}$ such that*

1. for all $w \in B(w^*, \delta_C)$ and $|\mu| < \mu_C$,

$$F(w) = \mu \tilde{e} \quad \text{if and only if} \quad w(\mu) = w,$$

2. the following expansion holds

$$w(\mu) = w^* + w'(0)\mu + o(\mu)$$

where the tangent vector to the central path $w(\cdot)$ at $\mu = 0$ is given by $w'(0) = F'(w^*)^{-1}\tilde{e} \neq 0$,

3. for all $\mu_1, \mu_2 \in (-\mu_C, \mu_C)$,

$$\|w(\mu_1) - w(\mu_2)\| \leq C|\mu_1 - \mu_2|.$$

The following result states a property of the central path and its tangent vector. It is a consequence of the strict complementarity assumption.

Lemma 3.4 *Let Assumptions 2.1–2.4 hold. Let $w(\cdot)$ be the implicit function defined in Lemma 3.3. Let $w(\mu) := (x(\mu), y(\mu), z(\mu))$, then*

$$x_i^* z_i'(0) + x_i'(0) z_i^* = 1. \quad (3.1)$$

Proof. By virtue of Lemma 3.3, for all $i = 1, \dots, n$, we have

$$x_i(\mu) = x_i^* + \mu x_i'(0) + o(\mu) \quad \text{and} \quad z_i(\mu) = z_i^* + \mu z_i'(0) + o(\mu).$$

By using $x_i^* z_i^* = 0$ and $F(w(\mu)) = \mu \tilde{e}$, we deduce that

$$x_i(\mu) z_i(\mu) = (x_i^* z_i'(0) + x_i'(0) z_i^*) \mu + o(\mu) = \mu.$$

Dividing both sides by μ and taking the limit $\mu \rightarrow 0$, we obtain (3.1). \square

The next two lemmas analyse some basic properties of the Newton step generated by the primal-dual method.

Lemma 3.5 *Under Assumptions 2.1–2.4, there exist $\delta^* > 0$, $\mu^* > 0$ and $M_1 > 0$ such that, for all $w \in B(w^*, \delta^*)$ and $|\mu| < \mu^*$, the Newton iterate, defined by*

$$N(w, \mu) = w - F'(w)^{-1}(F(w) - \mu \tilde{e}),$$

satisfies

$$\|N(w, \mu) - w(\mu)\| \leq M_1 \|w - w(\mu)\|^2. \quad (3.2)$$

and

$$\|N(w, \mu) - w^*\| \leq \frac{1}{2} \|w - w^*\| + \frac{\mu}{\mu^*} \frac{\delta^*}{2}. \quad (3.3)$$

In particular, $N(w, \mu) \in B(w^*, \delta^*)$.

Proof. Let us define

$$M_1 := \frac{1}{2}KL, \quad \delta^* := \min\{\delta_L, \delta_K, \delta_C, \frac{1}{4M_1}\} \text{ and } \mu^* := \min\{\mu_C, \frac{\sqrt{1+4M_1\delta^*}-1}{4M_1C}\},$$

where the constants are defined in Lemmas 3.1-3.3.

Let $(w, \mu) \in B(w^*, \delta^*) \times (-\mu^*, \mu^*)$. By Lemma 3.3, one has $F(w(\mu)) - \mu\tilde{e} = 0$. It follows that

$$\begin{aligned} N(w, \mu) - w(\mu) &= w - w(\mu) - F'(w)^{-1}(F(w) - \mu\tilde{e}) \\ &= -F'(w)^{-1}(F(w) - \mu\tilde{e} - F'(w)(w - w(\mu))) \\ &= -F'(w)^{-1} \int_0^1 (F'(w(\mu) + t(w - w(\mu))) - F'(w))(w - w(\mu)) dt. \end{aligned}$$

Taking the norm on both sides, then using Lemmas 3.1 and 3.2, we obtain (3.2).

Using (3.2), Lemma 3.3 and the definition of the constants M_1 , δ^* and μ^* , we have

$$\begin{aligned} \|N(w, \mu) - w^*\| &\leq \|N(w, \mu) - w(\mu)\| + \|w(\mu) - w^*\| \\ &\leq M_1\|w - w(\mu)\|^2 + \|w(\mu) - w^*\| \\ &\leq 2M_1(\|w - w^*\|^2 + \|w(\mu) - w^*\|^2) + \|w(\mu) - w^*\| \\ &\leq 2M_1\delta^*\|w - w^*\| + (2M_1C\mu^* + 1)C\mu \\ &\leq \frac{1}{2}\|w - w^*\| + \frac{\delta^*}{2\mu^*}\mu \\ &< \delta^*, \end{aligned}$$

which proves (3.3) and concludes the proof. \square

Lemma 3.6 *Let Assumptions 2.1-2.4 hold. Let δ^* and μ^* be the radii of convergence defined in Lemma 3.5. Then, there exists $M_2 > 0$ such that, for all $w \in B(w^*, \delta^*)$ and $0 \leq \mu^+ \leq \mu < \mu^*$, the Newton iterate*

$$N(w, \mu^+) = w - F'(w)^{-1}(F(w) - \mu^+\tilde{e}),$$

satisfies

$$\|N(w, \mu^+) - w(\mu^+)\| \leq M_2(\|w - w(\mu)\|^2 + \mu^2). \quad (3.4)$$

Proof. Let C , M_1 , δ^* and μ^* be the constants defined in Lemmas 3.3 and 3.5. Using Lemmas 3.3 and 3.5, one has

$$\begin{aligned} \|N(w, \mu^+) - w(\mu^+)\| &\leq M_1\|w - w(\mu^+)\|^2 \\ &\leq M_1(\|w - w(\mu)\| + \|w(\mu) - w(\mu^+)\|)^2 \\ &\leq M_1(\|w - w(\mu)\| + C(\mu - \mu^+))^2 \\ &\leq 2M_1 \max\{1, C^2\}(\|w - w(\mu)\|^2 + \mu^2). \end{aligned}$$

The result follows with $M_2 = 2M_1 \max\{1, C^2\}$. \square

4 Local convergence analysis

In this section we give convergence results that are direct consequences of the properties of the Newton method. An important point to note is that it is not assumed that the iterates remain strictly feasible during the iterations. The fraction to the boundary rule is not used to maintain the positivity of v_k . A second point to note is that the results are stated without any assumption on the proximity of the iterates to the central path.

Theorem 4.1 *Let Assumptions 2.1–2.5 hold. Consider the sequences defined by the recurrence*

$$\mu_{k+1} = \mu_k + d_k^\mu \quad \text{and} \quad w_{k+1} = w_k + d_k^w,$$

where d_k^μ is defined by (1.10) and d_k^w is the solution of (1.9). Let δ^* and μ^* be as in Lemma 3.5. For all $w_0 \in B(w^*, \delta^*)$ and all $0 < \mu_0 < \mu^*$, $\{\mu_k\}$ is decreasing and $\{(w_k, \mu_k)\}$ converges to $(w^*, 0)$. Moreover, if $\theta'(0) \neq 0$, then the rate of convergence of both sequences is q -superlinear. In addition, if θ' is locally Lipschitzian at 0, the rate of convergence of both sequences is quadratic.

Proof. It follows from Assumption 2.5 that $\theta(\mu) > 0$ and $\theta'(\mu) > 0$ for all $\mu > 0$, therefore the sequence $\{\mu_k\}$ is decreasing. Using the strict convexity of θ , one has

$$\mu_{k+1}\theta'(\mu_k) = \theta'(\mu_k)\mu_k - \theta(\mu_k) > -\theta(0) = 0.$$

By induction, we then obtain $\mu_k > 0$ for all k and thus the sequence $\{\mu_k\}$ converges. Taking the limit in the above equality and using the fact that zero is the unique solution of $\theta(\mu) = 0$, we deduce that $\{\mu_k\}$ converges to zero.

From (3.3) we have

$$\|w_{k+1} - w^*\| \leq \frac{1}{2}\|w_k - w^*\| + \frac{\delta^*}{2\mu^*}\mu_{k+1},$$

and therefore $w_k \rightarrow w^*$ as $k \rightarrow \infty$.

Assume $\theta'(0) \neq 0$. By Lemmas 3.1 and 3.2 the Jacobian matrix

$$\begin{pmatrix} F'(w) & -\tilde{e} \\ 0 & \theta'(\mu) \end{pmatrix}$$

is nonsingular and locally Lipschitzian at $(w^*, 0)$. The sequence $\{(w_k, \mu_k)\}$ is then generated by the Newton method with a step defined by (1.8). The convergence and the rates of convergence of the sequences follow from the local convergence properties of the Newton method (see for example [1, Proposition 1.17]). \square

This result allows to consider a globally convergent version of a primal-dual method with an update of the barrier parameter at each iteration. For example it

suffices to apply the globally convergent version of the Newton method proposed in [10, Chapter 8]. Since $\theta(\mu) > 0$ for $\mu > 0$, we can use the merit function

$$(w, \mu) \mapsto \|F(w) - \mu\tilde{e}\| + \theta(\mu)$$

(where $\|\cdot\|$ is any norm) and apply an Armijo backtracking linesearch technique. It follows from [10, Theorem 8.2.1] that if the sequence $\{w_k\}$ is bounded and if F' is uniformly Lipschitz continuous and bounded away from zero on a neighborhood of $\{w_k\}$, then the resulting algorithm is globally convergent. But an important feature of primal-dual methods is to maintain strict feasibility of the iterates and if we add the fraction to the boundary rule to a globally convergent version of the algorithm, then the analysis is far from straightforward and needs a full study. So we do not go further with this idea in this paper.

Theorem 4.1 shows also that, whenever $\theta'(0) \neq 0$, the sequence $\{w_k\}$ converges to w^* with a r -superlinear or r -quadratic rate of convergence, depending on whether θ' is locally Lipschitzian at zero or not. It would be interesting to exhibit conditions on the decrease of $\{\mu_k\}$ so that the sequence $\{w_k\}$ converges with a q -rate of convergence. The next result shows that, provided that $\{\mu_k\}$ does not converge too fast to zero, both sequences $\{w_k\}$ and $\{\mu_k\}$ have the same rate of convergence.

Theorem 4.2 *Let Assumptions 2.1–2.4 hold. Let δ^* and μ^* be the threshold values defined in Lemma 3.5. Assume $\{\mu_k\}$ is a sequence of positive scalars converging to zero such that $0 < \mu_0 < \mu^*$ and*

$$\liminf_{k \rightarrow \infty} \frac{\mu_{k+1}}{\mu_k^p} > 0, \quad (4.1)$$

for some parameter $p \in (1, 2)$. Consider the sequence defined by the recurrence

$$w_{k+1} = w_k + d_k^w,$$

where d_k^w is the solution of (1.9) with d_k^μ defined by $d_k^\mu = \mu_{k+1} - \mu_k$. Then for all $w_0 \in B(w^*, \delta^*)$, the sequence $\{w_k\}$ converges to w^* and

$$w_k = w(\mu_k) + o(\mu_k). \quad (4.2)$$

In particular, $v_k > 0$ for sufficiently large k and both sequences $\{w_k\}$ and $\{\mu_k\}$ have the same rate of convergence.

Proof. As in the proof of Theorem 4.1, from (3.3) we have

$$\|w_{k+1} - w^*\| \leq \frac{1}{2} \|w_k - w^*\| + \frac{\delta^*}{2\mu^*} \mu_{k+1},$$

and thus $w_k \rightarrow w^*$ when $k \rightarrow \infty$.

Let us prove now that (4.2) holds. Let M_2 be the constant defined in Lemma 3.6. Let us define $s_k := M_2 \|w_k - w(\mu_k)\|$ and $\epsilon_k := M_2 \mu_k$. Condition (4.1) implies that $\mu_k^2 = o(\mu_{k+1})$ and that there exist $c > 0$, $\beta \in (0, 1)$ and \hat{k} such that

$$c\beta^{p^k} \leq \mu_k \quad \text{for all } k \geq \hat{k} \quad (4.3)$$

(see for example [1, Proposition 1.2]). From (3.4), one has

$$s_{k+1} \leq s_k^2 + \epsilon_k^2. \quad (4.4)$$

Since $\epsilon_k^2 = o(\epsilon_{k+1})$, there exists an index $\bar{k} \geq 0$ such that $\epsilon_k^2 \leq \frac{1}{2}\epsilon_{k+1}$ for all $k \geq \bar{k}$. Let us prove that there exists $k_0 \geq \bar{k}$ such that $s_{k_0} \leq \epsilon_{k_0}$. For the purpose of obtaining a contradiction, suppose that $s_k > \epsilon_k$ for all $k \geq \bar{k}$. From (4.4) we would deduce that $s_{k+1} \leq 2s_k^2$, therefore $\{s_k\}$ would converge quadratically to zero, which should imply that $s_k = O(\alpha^{2^k})$ for some $\alpha > 0$. Since we have supposed that $s_k > \epsilon_k$, we would then have $\epsilon_k = O(\alpha^{2^k})$, a contradiction with (4.3) knowing that $1 < p < 2$.

Let us prove now by induction on k that $s_k \leq \epsilon_k$ for $k \geq k_0$. Our claim holds for $k = k_0$. Assume that it is true for a given $k \geq k_0$. By (4.4) and the induction hypothesis one has $s_{k+1} \leq 2\epsilon_k^2 \leq \epsilon_{k+1}$, and thus our claim is also true for $k + 1$.

From (4.4), we can then deduce that $s_{k+1} \leq 2\epsilon_k^2 = o(\epsilon_{k+1})$, which implies (4.2) and proves the first part of the theorem.

Let us show that $v_k > 0$ for sufficiently large k . From (4.2) and Lemma 3.3 we have

$$(x_k)_i = x_i^* + x_i'(0)\mu_k + o(\mu_k),$$

for all index i . By (3.1) we have either $x_i^* > 0$ or $x_i'(0) > 0$. It follows that $(x_k)_i > 0$ for sufficiently large k . The reasoning is similar for the components $(z_k)_i$.

At last, using (4.2) and Lemma 3.3, we have $w_k - w^* = w_k - w(\mu_k) + w(\mu_k) - w^* = w'(0)\mu_k + o(\mu_k)$, and thus $\|w_k - w^*\| \sim \|w'(0)\|\mu_k$. \square

As we said at the beginning of this section, it is interesting to note that unlike other local convergence analyses [2, 7, 18], Theorem 4.2 assumes neither strict feasibility of the iterates— w_0 might very well have a component $v_0 \not\geq 0$ —nor proximity of the iterates to the central path. Strict feasibility and proximity however follow, as specified by (4.2), from the sub-quadratic rate of convergence of the barrier parameter to zero given by (4.1). The iterates w_k asymptotically approach the central path tangentially since from (4.2) and Lemma 3.3,

$$\lim_{k \rightarrow \infty} \frac{w_k - w^*}{\mu_k} = \lim_{k \rightarrow \infty} \frac{w(\mu_k) - w^*}{\mu_k} = w'(0).$$

Asymptotic results having a related yet different flavour are given in [5, 14, 16, 17]. There, the update of μ is quadratic or faster, can be based on the size of the current residual, proximity to the central path is not enforced and feasibility

is eventually not satisfied. However, the stabilized primal-dual method in [14] may produce iterates with nonpositive components. Quadratic convergence follows but takes place in a restrictive neighbourhood around w^* .

5 Maintaining the positivity and the proximity to the central path

Assume that at the current iteration, the iterates w and μ satisfy the approximate optimality condition

$$\|F(w) - \mu\tilde{e}\| \leq \varepsilon(\mu), \quad (5.1)$$

for some tolerance $\varepsilon(\mu) > 0$. The Newton iterate for (1.7) is

$$w^+ = w - F'(w)^{-1}(F(w) - \mu^+\tilde{e}), \quad (5.2)$$

where $\mu^+ = \mu - \theta(\mu)/\theta'(\mu)$. We propose to analyse the conditions under which w^+ will satisfy the fraction to the boundary rule and the next stopping test, that is

$$v^+ \geq (1 - \tau(\mu))v, \quad (5.3)$$

for some parameter $\tau(\mu) \in (0, 1)$ and

$$\|F(w^+) - \mu^+\tilde{e}\| \leq \varepsilon(\mu^+). \quad (5.4)$$

The analysis proposed below uses some tools developed in [2].

Lemma 5.1 *Let Assumptions 2.1–2.4 hold. Assume also that $\varepsilon(\mu) = O(\mu)$. Let δ^* be defined in Lemma 3.5. Then, there exists $M_3 > 0$ such that, for all $w \in B(w^*, \delta^*)$ satisfying (5.1) and all $\mu > 0$ sufficiently small,*

$$\|w - w^*\| \leq M_3\mu.$$

Proof. Let $w \in B(w^*, \delta^*)$. Since $F(w^*) = 0$ and $F'(w^*)$ is invertible (Lemma 3.2), we have

$$w - w^* = F'(w^*)^{-1}(F(w) - \int_0^1 (F'(w^* + t(w - w^*)) - F'(w^*))(w - w^*) dt).$$

Taking the norm on both sides, using the Cauchy-Schwarz inequality and Lemmas 3.1 and 3.2, we obtain

$$\|w - w^*\| \leq K(\|F(w)\| + \frac{L}{2}\|w - w^*\|^2).$$

Recall that $\delta^* \leq \frac{1}{2KL}$ (see the definition of δ^* in the proof of Lemma 3.5), it follows that

$$\|w - w^*\| \leq \frac{4}{3}K\|F(w)\|.$$

Now using $\|\tilde{e}\| = \sqrt{n}$, (5.1) and $\varepsilon(\mu) = O(\mu)$, we then have

$$\begin{aligned}\|w - w^*\| &\leq \frac{4}{3}K(\|F(w) - \mu\tilde{e}\| + \sqrt{n}\mu) \\ &\leq \frac{4}{3}K(\varepsilon(\mu) + \sqrt{n}\mu) \\ &\leq M_3\mu.\end{aligned}$$

□

Lemma 5.2 *Let Assumptions 2.1–2.4 hold. Assume also that $\varepsilon(\mu) = O(\mu)$. Let δ^* and μ^* be defined in Lemma 3.5. Then, there exists $M_4 > 0$ such that, for all $w \in B(w^*, \delta^*)$ satisfying (5.1) and all $0 < \mu^+ \leq \mu$ sufficiently small, the Newton iterate w^+ defined by (5.2) satisfies*

$$\|w^+ - w(\mu^+)\| \leq M_4\mu^2. \quad (5.5)$$

Proof. By using Lemmas 5.1 and 3.3 we obtain

$$\begin{aligned}\|w - w(\mu)\| &\leq \|w - w^*\| + \|w^* - w(\mu)\| \\ &\leq (M_3 + C)\mu\end{aligned}$$

Using this last inequality in (3.4), we obtain (5.5) with a constant M_4 defined by $M_4 = M_2((M_3 + C)^2 + 1)$. □

Regarding positivity of the iterates, we distinguish the case where no fraction to the boundary rule is enforced, to the case where an explicit fraction to the boundary rule is present.

Theorem 5.3 *Let Assumptions 2.1–2.4 hold. Assume also that the parameters μ^+ and $\varepsilon(\mu)$ are chosen such that*

$$\mu^2 = o(\mu^+) \quad \text{and} \quad \varepsilon(\mu) = O(\mu).$$

Let δ^ be defined in Lemma 3.5. Then for all $w \in B(w^*, \delta^*)$ satisfying (5.1) and such that $v > 0$, the Newton iterate w^+ defined by (5.2) is such that $v^+ > 0$ and $F(w^+) - \mu^+\tilde{e} = o(\mu^+)$ for sufficiently small $\mu > 0$. Moreover, if $\tau(\mu)$ satisfies*

$$\limsup_{\mu \downarrow 0} (1 - \tau(\mu)) \frac{\mu + \varepsilon(\mu)}{\mu^+} < 1, \quad (5.6)$$

then (5.3) is satisfied for sufficiently small $\mu > 0$. In addition, if $\varepsilon(\mu) = \Theta(\mu)$, then (5.4) is satisfied for sufficiently small $\mu > 0$.

Proof. Let $0 < \kappa < 1$. We begin by proving that, for all index i

$$x_i^+ \geq \kappa \frac{\mu^+}{\mu + \varepsilon(\mu)} x_i.$$

Consider first the case where $x_i^* > 0$. Lemmas 5.1 and 5.2 imply that by taking $\mu > 0$ sufficiently small, w and w^+ can be arbitrarily close to w^* . It follows that

$$\lim_{\mu \downarrow 0} \frac{x_i^+}{x_i} = \frac{x_i^*}{x_i^*} = 1.$$

Using $\mu^+ < \mu + \varepsilon(\mu)$ and $0 < \kappa < 1$ we deduce that

$$\frac{x_i^+}{x_i} \geq \kappa \frac{\mu^+}{\mu + \varepsilon(\mu)}, \quad (5.7)$$

for sufficiently small $\mu > 0$.

Consider now the case where $x_i^* = 0$. By Lemma 3.4, we have $x_i'(0) > 0$. By using Lemma 5.2, Lemma 3.3 and $\mu^2 = o(\mu^+)$, we have

$$\begin{aligned} x_i^+ &= (x_i^+ - x_i(\mu^+)) + x_i(\mu^+) \\ &\geq -M_4 \mu^2 + x_i'(0) \mu^+ + o(\mu^+) \\ &\geq x_i'(0) \mu^+ + o(\mu^+). \end{aligned}$$

Dividing both sides by x_i we obtain

$$\frac{x_i^+}{x_i} \geq \frac{x_i'(0) \mu^+ + o(\mu^+)}{x_i}.$$

From the definition of F and (5.1), we deduce that $x_i z_i \leq \mu + \varepsilon(\mu)$. Multiplying and dividing the previous equation by z_i yields

$$\frac{x_i^+}{x_i} \geq \frac{\mu^+}{\mu + \varepsilon(\mu)} (x_i'(0) z_i + o(1)).$$

Using Lemma 3.4, the term in parentheses tends to $x_i'(0) z_i^* = 1 - x_i^* z_i'(0) = 1$ as μ goes to zero. We then deduce that (5.7) also holds for sufficiently small $\mu > 0$ and for all i such that $x_i^* = 0$.

By the symmetric role of the variables x and z , the reasoning is the same for the components z_i . We then have

$$v^+ \geq \kappa \frac{\mu^+}{\mu + \varepsilon(\mu)} v. \quad (5.8)$$

Since we have assumed that $v > 0$, (5.8) implies that $v^+ > 0$.

If condition (5.6) is satisfied, it implies that there exists $\kappa \in (0, 1)$ such that, for sufficiently small $\mu > 0$, we have

$$1 - \tau(\mu) \leq \kappa \frac{\mu^+}{\mu + \varepsilon(\mu)}.$$

From this last inequality and (5.8) we deduce that (5.3) holds.

Turning now to the residual at the updated iterate, by using Lemma 5.2 and $\mu^2 = o(\mu^+)$, we have

$$\begin{aligned} \|F(w^+) - \mu^+ \tilde{e}\| &= \|F(w^+) - F(w(\mu^+))\| \\ &= O(\|w^+ - w(\mu^+)\|) \\ &= O(\mu^2) \\ &= o(\mu^+). \end{aligned}$$

If $\varepsilon(\mu) = \Theta(\mu)$, this last equation implies that for sufficiently small μ , (5.4) is satisfied. \square

6 Practical algorithms

In this section, we are concerned with ways to incorporate the ideas presented above into practical algorithms. We first present two algorithms which embed steps in (w, μ) into a globally-convergent framework and give a convergence theorem. Next, we give a third algorithm, whose global convergence has not been established, but which is conceptually closer to intuition and possesses desirable numerical properties.

We choose a scalar function θ satisfying Assumption 2.5 and for all $\mu > 0$ we define

$$\mu^+ = \mu - \frac{\theta(\mu)}{\theta'(\mu)}.$$

We select two tolerance functions $\varepsilon : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ and $\tau : \mathbb{R}_+ \rightarrow (0, 1)$, such that

$$\mu^2 = o(\mu^+), \quad \varepsilon(\mu) = \Theta(\mu) \quad \text{and} \quad \limsup_{\mu \downarrow 0} (1 - \tau(\mu)) \frac{\mu + \varepsilon(\mu)}{\mu^+} < 1. \quad (6.1)$$

The functions θ given by examples at the end of Section 2 meet the requirement $\mu^2 = o(\mu^+)$. The other conditions in (6.1) are satisfied if, for instance, we choose $\varepsilon(\mu) = \rho\mu$ for some parameter $\rho > 0$ and $\tau(\mu) = \max\{\bar{\tau}, 1 - \mu\}$ for some $0 < \bar{\tau} < 1$. We describe our first algorithm as Algorithm 6.1.

ALGORITHM 6.1

Given an initial barrier parameter $\mu_0 > 0$, a tolerance $\varepsilon_0 := \varepsilon(\mu_0)$ and $w_0 := (x_0, y_0, z_0) \in \mathbb{R}^{2n+m}$ such that $v_0 > 0$ and $\|F(w_0) - \mu_0 \tilde{e}\| \leq \varepsilon_0$, set $k = 0$ and perform the following steps:

1. Compute d_k^μ from (1.10) and d_k^w from (1.9).
2. Choose $\tau_k := \tau(\mu_k) \in (0, 1)$ and compute α_k the largest value of $\alpha \in (0, 1]$ such that (1.11) holds. Set $\mu_{k+1} = \mu_k + \alpha_k d_k^\mu$ and $\bar{w}_k = w_k + \alpha_k d_k^w$.
3. Choose $\varepsilon_{k+1} := \varepsilon(\mu_{k+1}) < \varepsilon_k$. Starting from \bar{w}_k , apply a sequence of inner iterations to find w_{k+1} satisfying

$$\|F(w_{k+1}) - \mu_{k+1} \tilde{e}\| \leq \varepsilon_{k+1}.$$

4. Set $k := k + 1$ and go to Step 1.
-

Upon denoting the Newton iterates by

$$w_k^+ = w_k + d_k^w \quad \text{and} \quad \mu_k^+ = \mu_k + d_k^\mu.$$

where d_k^μ and d_k^w are defined by (1.10) and (1.9), equation (1.9) can be rewritten

$$F'(w_k)(w_k^+ - w_k) + F(w_k) = \mu_k^+ \tilde{e}.$$

We also have

$$\bar{w}_k = \alpha_k w_k^+ + (1 - \alpha_k) w_k \quad \text{and} \quad \mu_{k+1} = \alpha_k \mu_k^+ + (1 - \alpha_k) \mu_k.$$

Theorem 6.1 *Let Assumptions 2.1–2.4 and 2.5 hold. Assume also that the function $\theta(\cdot)$, $\varepsilon(\cdot)$ and $\tau(\cdot)$ satisfy the properties (6.1). Let $\{w_k\}$ and $\{\mu_k\}$ be the sequences generated by Algorithm 6.1. Assume finally that $\{\mu_k\}$ converges to zero and that a subsequence of $\{w_k\}$ converges to w^* . The following conclusions hold:*

- (i) $w_k \rightarrow w^*$ when $k \rightarrow \infty$,
- (ii) $\alpha_k = 1$ and $w_{k+1} = w_k^+ = \bar{w}_k$ for sufficiently large k ,
- (iii) $\frac{\|w_{k+1} - w^*\|}{\|w_k - w^*\|} \sim \frac{\mu_{k+1}}{\mu_k}$.

Proof. Let δ^* and μ^* be defined in Lemma 3.5. Let $\bar{\mu}$ be a threshold value such that all the conclusions of Theorem 5.3 are satisfied. Let us define $\hat{\mu} = \min\{1, \mu^*, \bar{\mu}, \frac{\delta^*}{M_4 + C}\}$, where M_4 and C are defined in Lemmas 5.2 and 3.3. By assumption, there exists an index $k_0 \geq 0$ such that, $w_{k_0} \in B(w^*, \delta^*)$ and $\mu_{k_0} < \hat{\mu}$. Let us prove that $w_k \in B(w^*, \delta^*)$ for all $k \geq k_0$. We proceed by induction on k .

Our claim is true for $k = k_0$. Assume that it is true for a given $k \geq k_0$. Considering the assumptions that have been made and the induction hypothesis, Theorem 5.3 applies, therefore $\alpha_k = 1$ and $w_{k+1} = w_k^+$. By Lemmas 5.2 and 3.3, we have

$$\begin{aligned} \|w_{k+1} - w^*\| &\leq \|w_{k+1} - w(\mu_{k+1})\| + \|w(\mu_{k+1}) - w^*\| \\ &\leq M_4\mu_k^2 + C\mu_{k+1} \\ &< (M_4 + C)\hat{\mu} \\ &< \delta^*, \end{aligned}$$

and thus our claim is also true for $k + 1$.

By applying Lemma 5.1 one has $\|w_k - w^*\| \leq M_3\mu_k$ for all $k \geq k_0$, which proves conclusion (i).

Applying again Theorem 5.3, we have $\alpha_k = 1$ and $w_{k+1} = \bar{w}_k = w_k^+$ for all $k \geq k_0$, which proves point (ii).

Lemma 5.2 and the fact that $\mu_k^2 = o(\mu_{k+1})$ imply that $\|w_k - w(\mu_k)\| = o(\mu_k)$. Using again Lemma 3.3, it follows that $w_k - w^* = w_k - w(\mu_k) + w(\mu_k) - w^* = \mu_k w'(0) + o(\mu_k)$, from which conclusion (iii) follows. \square

We will refer to the steps described in Step 2 of Algorithm 6.1 as *extrapolation* steps. Note that since the inner iteration in Step 3 of Algorithm 6.1 relies on a globally convergent method, global convergence of Algorithm 6.1 is ensured once the generated sequence $\{\mu_k\}$ converges to zero. This will be the case provided the sequence of steplengths $\{\alpha_k\}$ along the extrapolation is bounded away from zero. Once the iterates reach a region where the central path exists and once μ is sufficiently small, it is easy to show that there exists $\alpha_{\min} > 0$ such that $\alpha_k \geq \alpha_{\min} > 0$ for all sufficiently large k . Theorem 5.3 confirms this and goes even further to show that the unit step is asymptotically accepted. In a practical implementation, preventing α_k from converging to zero when far from a solution, and possibly in regions where no central path exists, can be achieved by imposing a minimal decrease in μ at each outer iteration. This has, however, not been necessary in the tests of §7.

From a numerical point of view, a danger with Algorithm 6.1 is that \bar{w}_k may lie too close to the boundary, causing ill conditioning and potentially leading to a large number of inner iterations. Although this situation did not arise in the tests of §7, it could certainly occur. To circumvent this possibility, we suggest performing a linesearch on a merit function along extrapolation steps in order to keep \bar{w}_k safely away from the boundary. This results in the following algorithm, for which we only specify the difference with Algorithm 6.1.

ALGORITHM 6.2

2. Choose $\tau_k := \tau(\mu_k) \in (0, 1)$ and compute $\tilde{\alpha}_k$ the greatest value of $\alpha \in (0, 1]$ such that (1.11) holds. Starting with $\alpha = \tilde{\alpha}_k$ and from $\tilde{w}_k = w_k + \tilde{\alpha}_k d_k^w$, perform a backtracking linesearch on a merit function for which d_k^w is a descent direction from w_k . Let $\alpha_k > 0$ be the resulting steplength. Set $\mu_{k+1} = \mu_k + \alpha_k d_k^\mu$ and $\bar{w}_k = w_k + \alpha_k d_k^w$.
-

In our tests, the inner iterations are globalized by an exact penalty function and we use this merit function to identify μ_{k+1} and \bar{w}_k at Step 2 of Algorithm 6.2. It will be made explicit in Section 7.

We now present a third algorithm which only takes steps in (w, μ) but is not embedded into any globally-convergent framework. It simply eliminates the sequences of inner iterations and only takes extrapolation steps. Quite surprisingly, this algorithm turned out to be more robust in practice. We describe it as Algorithm 6.3.

ALGORITHM 6.3

Given an initial barrier parameter $\mu_0 > 0$, a tolerance $\varepsilon_0 := \varepsilon(\mu_0)$ and $w_0 := (x_0, y_0, z_0) \in \mathbb{R}^{2n+m}$ such that $v_0 > 0$ and $\|F(w_0) - \mu_0 \tilde{e}\| \leq \varepsilon_0$, set $k = 0$ and perform the following steps:

1. Compute d_k^μ from (1.10) and d_k^w from (1.9).
 2. Choose $\tau_k := \tau(\mu_k) \in (0, 1)$ and compute $\tilde{\alpha}_k$ the greatest value of $\alpha \in (0, 1]$ such that (1.11) holds. Starting with $\alpha = \tilde{\alpha}_k$ and from $\tilde{w}_k = w_k + \tilde{\alpha}_k d_k^w$, perform a backtracking linesearch on a merit function for which d_k^w is a descent direction from w_k . Let α_k be the resulting steplength. Set $\mu_{k+1} = \mu_k + \alpha_k d_k^\mu$ and $w_{k+1} = w_k + \alpha_k d_k^w$.
 3. Set $k := k + 1$ and go to Step 1.
-

Used as such, the merit function in Algorithm 6.3 mostly serves the purpose of staying safely away from the boundary of the feasible set. Obviously, this algorithm would be globally convergent using for instance a linesearch on the residual (1.4). However, this abandons the minimization aspect of the problem and merely attempts to identify a first-order critical point. Research on a global convergence framework for Algorithm 6.3 is under way.

7 Numerical experiments

In this section we describe our framework for numerical experiments and summarize the results. The three algorithms in Section 6 were implemented as modifications

of the IPOPT software [15]¹, an implementation of a globally-convergent algorithm for general smooth nonlinear programming. Nonlinear inequality constraints are transformed into equalities by the addition of slack variables, resulting in a problem of the form (1.1). Bounds on the variables and slacks are treated by means of a logarithmic barrier. The IPOPT implementation fixes a pre-determined sequence $\{\mu_k\}$ of barrier parameters which decreases to zero and each set of inner iterations consists in an SQP algorithm applied to the barrier subproblem. Global convergence is ensured in one of several ways, which one to choose is left as an option to the user. IPOPT was chosen for its excellent robustness, demonstrated numerical efficiency and availability.

Algorithms 6.1 and 6.2 depart from the IPOPT reference implementation mainly in the extrapolation step and the fact that cutting back this step influences the next value of μ . Step 3 of both algorithms is identical to the globalization implemented in IPOPT. We had to choose particular parameters in IPOPT and for completeness, we now describe them and mention what options differ from their default values and why. We stress that the following modifications were applied to *all* algorithms compared in this section.

1. Globalization in Step 3 consisted of a linesearch ensuring sufficient decrease in the non-differentiable exact merit function

$$\psi(x; \mu, \nu) = f(x) - \mu \sum_{i=1}^n \log x_i + \nu \|c(x)\|_2, \quad (7.1)$$

in which $\mu > 0$ is fixed and $\nu > 0$ is dynamically adjusted.

2. We disabled second-order correction steps after an extrapolation step since the latter is not concerned with descent. All other second-order corrections settings were left unchanged.
3. After a barrier subproblem is solved, IPOPT resets ν to 10^{-6} in the merit function (7.1). We also ensured that it was reset after an extrapolation step.
4. We allowed for a barrier subproblem to be deemed solved if w_k readily satisfies the next barrier stopping test. No extrapolation step was imposed in such a case. Similarly, if \bar{w}_k , resulting from an extrapolation step, satisfied the barrier stopping test with μ_{k+1} , no inner iteration step was imposed and we moved on to the next barrier subproblem. By default, IPOPT *imposes* at least one step in each sequence of inner iterations². For purposes of comparison, we relaxed IPOPT so as to not impose such a step, which improved the

¹version 2.3.x

²Contrary to what is stated in [15], imposing a step is sufficient but not necessary to guarantee fast local convergence [2, 8, 9]

reference implementation noticeably. This exposed a problem in the original IPOPT implementation where optimality for a barrier subproblem with the smallest allowed value of μ did not imply optimality for the original problem [15, § 2.1, Equation (7)]. Fixing this inconsistency noticeably increased the reliability of the reference implementation.

5. For consistency with Newton’s method and the usual theory of interior-point methods, we elected to take equal primal and dual steplengths.

We compared the three algorithms on all the problems from the COPS 3.0 collection [3] which possess at least one inequality constraint and/or at least one bound constraint. This results in 28 problems, all of which were used in their default dimension. In all cases, the initial barrier parameter was left at its default value in IPOPT, $\mu_0 = 0.1$. Each problem was given a limit of 1000 iterations and 10 minutes of CPU time.

As mentioned in §1, a common occurrence of numerical failure in practical interior point methods is the situation where, at the beginning of a sequence of inner iterations, μ_k and $\|F(w_k) - \mu_k \tilde{e}\|$ differ by several orders of magnitude. This situation may lead to a large number of inner iterations and eventually, failure may occur due to limited allowed CPU time or a maximal number of iterations being reached. Using the reference implementation of IPOPT, this situation occurs, among others, on instances of problem `dirichlet`. The behaviour of μ_k and $\|F(w_k) - \mu_k \tilde{e}\|$ is depicted in Fig. 7.1. After a few iterations, the two differ sufficiently to cause a large number of inner iterations and, in one of the two cases, the maximum CPU time to be reached.




Figure 7.1: Log-plot of the evolution of μ_k (dashed line) and $\|F(w_k) - \mu_k \tilde{e}\|$ (solid line) using the reference implementation on two instances of problem `dirichlet`. On the left plot, the discrepancy is moderate and the method manages to identify an optimal solution. On the right plot, the discrepancy is too large and causes failure due to CPU time.

it	IPOPT			1-STEP-1.1		
	$\ F(w_k) - \mu_k \tilde{e}\ $	μ_k	α_k	$\ F(w_k) - \mu_k \tilde{e}\ $	μ_k	α_k
0	1.00e+01	1.00e-01	0.00e+00	1.00e+01	1.00e-01	0.00e+00
1	1.07e+00	1.00e-01	6.01e-01	1.07e+00	8.76e-02	6.01e-01
2	5.28e-01	1.00e-01	5.17e-01	4.89e-01	7.72e-02	5.53e-01
3	1.98e-01	3.16e-02	6.31e-01	3.13e-01	7.07e-02	3.70e-01
4	1.67e-02	5.62e-03	9.66e-01	1.34e-01	6.13e-02	5.71e-01
5	3.16e-03	4.22e-04	9.45e-01	6.56e-02	5.37e-02	5.12e-01
6	2.86e-03	8.66e-06	1.09e-01	4.29e-02	4.40e-02	7.08e-01
7	1.21e-02	8.66e-06	4.07e-01	3.27e-02	3.39e-02	8.58e-01
8	1.18e-02	8.66e-06	4.39e-01	1.23e-02	2.42e-02	1.00e+00
9	6.53e-02	8.66e-06	2.57e-01	1.42e+00	1.67e-02	1.00e+00
10	5.48e-02	8.66e-06	3.31e-01	6.34e-02	1.11e-02	1.00e+00
11	3.10e-01	8.66e-06	7.94e-02	8.08e-02	7.05e-03	1.00e+00

Table 7.1: Comparison of the first few iterations of the reference algorithm and Algorithm 6.3 on problem `dirichlet3`. In the reference algorithm, the current inner iteration will not finish within the time limits. In the case of Algorithm 6.3, the barrier parameter and the primal-dual residual remain of comparable magnitudes.

In contrast, Algorithm 6.3, by means of the steplength in the variable μ , appears to introduce a regularization which works towards avoiding a discrepancy such as that of Fig. 7.1. This is illustrated in Fig. 7.2. Table 7.1 gives a detailed look at the first few iterations of the reference algorithm and Algorithm 6.3 on problem `dirichlet3`, and exposes the influence of the steplength α_k on the value of μ_k .

Note that on the two plots of Fig. 7.2, the shape of the curves is indicative of superlinear convergence. Moreover, this fast convergence does not only occur asymptotically, but over a large portion of the iterations. The first problem is solved much more smoothly and in less than half as many iterations while the second problem no longer fails to be solved.




Figure 7.2: Log-plot of the evolution of μ_k (dashed line) and $\|F(w_k) - \mu_k \tilde{e}\|$ (solid line) using Algorithm 6.3 on the same two instances of problem `dirichlet` as in Fig. 7.1.

Problem	IPOPT		EXTRAP		EXTRAP-LS		1-STEP-1.5		1-STEP-1.1	
	it	#f	it	#f	it	#f	it	#f	it	#f
bearing	19	20	20	42	20	42	23	42	27	28
camshape	-45	-46	-45	-46	-45	-46	-44	-45	-43	-44
catmix	18	19	23	24	23	24	19	20	28	29
dirichlet1	76	337	58	139	58	139	-546	-7695	30	33
dirichlet2	64	343	36	37	36	37	64	267	34	36
dirichlet3	-42	-43	35	36	35	36	-41	-42	36	39
gasoil	352	2855	213	1717	213	1717	225	1771	219	1658
glider	-379	-390	-379	-390	-379	-390	-1000	-1001	-1000	-1001
lane_emden1	25	72	24	61	24	61	-24	-334	27	36
lane_emden2	-325	-4572	-324	-4509	-314	-4349	28	30	39	41
lane_emden3	45	54	45	51	45	51	-57	-103	24	26
marine	15	16	15	16	15	16	15	16	29	30
methanol	15	19	19	23	19	23	10	11	30	31
minsurf	80	130	98	249	87	183	41	50	211	486
pinene	16	17	16	17	16	17	8	9	23	24
polygon	12	17	12	17	12	17	9	10	26	27
robot	38	134	38	134	38	134	17	18	26	27
rocket	53	61	58	66	58	66	542	543	89	313
steering	14	16	14	16	14	16	9	10	23	25
tetra_duct12	10	11	9	10	9	10	9	10	24	25
tetra_duct15	10	11	10	11	10	11	9	10	24	25
tetra_duct20	9	10	9	10	9	10	6	7	21	22
tetra_gear	9	10	9	10	9	10	7	8	23	24
tetra_hook	9	10	9	10	9	10	7	8	23	24
torsion	15	27	15	28	15	28	17	36	25	26
triangle_deer	7	8	7	8	7	8	7	8	23	24
triangle_pacman	7	8	7	8	7	8	7	8	23	24
triangle_turtle	9	10	9	10	9	10	7	8	24	30

Table 7.2: Results on the COPS 3.0 collection. A negative number specifies a failure with the given number of iterations and/or function evaluations. When failure occurs with less than 1000 iterations, it is due to CPU time being exceeded.

Table 7.2 compares the number of iterations and number of function evaluations across all variants. Algorithm IPOPT is the reference implementation using the updating rule

$$\mu_{j+1} = \max \left\{ \frac{\epsilon_{\text{tol}}}{11}, \min \{ \mu_j/5, \mu_j^{1.5} \} \right\},$$

EXTRAP is Algorithm 6.1, EXTRAP-LS is Algorithm 6.2, 1-STEP-1.5 is Algorithm 6.3 using (2.2) with $b = 1$ and $\gamma = 0.5$ and 1-STEP-1.1 is Algorithm 6.3 using (2.2) with $b = 1$ and $\gamma = 0.1$. Both EXTRAP and EXTRAP-LS use (2.2) with $b = 1$ and $\gamma = 0.5$ during extrapolation steps. During an extrapolation step in Algorithm 6.2 and in Algorithm 6.3, the merit function used was the exact merit function (7.1) with barrier parameter $\mu = \mu_k + d_k^\mu$.

All variants fail on problems `camshape`³ and `glider`⁴ while `minsurf` seems best solved with an aggressive decrease of μ . The performance of Algorithm 1-STEP-

³with the message “linesearch failure”

⁴because the maximum number of iterations was reached

1.1 on problems `steering` through `triangle_turtle` seems worse but it appeared in our tests that those problems are very efficiently solved with $\mu_0 = 10^{-9}$. We therefore attribute this decay to the initial value μ_0 rather than to the mechanism of Algorithm 6.3. Table 7.2 suggests that the function θ should, in general, not attempt to decrease μ too aggressively—rather, Newton’s method itself should take care of the fast convergence. Moreover, the 1-STEP-1.1 variant solves two more problems than the reference implementation: `dirichlet3` and `lane_emden2`.

A natural concern regarding Algorithm 6.3 is that it might not identify a local minimizer. Fortunately, in our tests, it always identified a point at which the objective function value is comparable to or lower than that obtained with the reference implementation. The only significant differences are on problems

- `dirichlet1` for the 1-STEP-1.5 variant, which fails. The variant 1-STEP-1.1 finishes successfully with a value 10% lower than the remaining three variants,
- `glider` on which all variants fail,
- `dirichlet3` with a relative difference of order 10^{-3} caused by variants IPOPT and 1-STEP-1.5 which fail. The largest relative difference across the remaining variants is of order 10^{-9} .
- `lane_emden3` with a relative difference of order 0.3% caused by the variant 1-STEP-1.5 which fails. The largest relative difference across the remaining variants is of order 10^{-9} .

On all other problems, the largest relative variation in final objective value is of order 10^{-6} . Although all variants fail on `camshape`, the largest relative difference is of order 10^{-8} , which leads us to believe that all algorithms are stuck in the same region.

8 Conclusions

We have presented a framework that attempts to give satisfactory theoretical grounding to so-called *adaptive*, or *dynamic* updating rules for the barrier parameter in interior-point methods for nonlinear programming. In the absence of a notion of duality and of a meaning to a duality gap, basing such dynamic rules on observations arising from linear or convex quadratic programming lacks a theoretical justification. We hope to have shown in this paper that instead of attempting to *design an updating rule*, the power of Newton’s method applied to an augmented primal-dual system provides the desired behaviour.

The present framework requires the choice of an updating function which determines the fastest allowed decrease of μ . The convexity of this function does not allow μ to increase.

It appears from our preliminary numerical tests that, in its more liberal form, this framework produces a much improved agreement between the barrier parameter and the barrier subproblem residual all along the iterations and a smaller number of iterations and function evaluations in many cases. There remains however much work to be done in order to devise globally-convergent algorithms based on the Newton/path-following scheme presented in this paper.

References

- [1] D. P. Bertsekas. *Constrained Optimization and Lagrange Multiplier Methods*. Academic Press, London, 1982.
- [2] R. H. Byrd, G. Liu, and J. Nocedal. On the local behavior of an interior point method for nonlinear programming. In D.F. Griffiths and D.J. Higham, editors, *Numerical Analysis 1997*, pages 37–56. Addison Wesley Longman, 1997. Proceedings of the Dundee 1997 Conference on Numerical Analysis.
- [3] E. D. Dolan, J. J. Moré, and T. S. Munson. Benchmarking optimization software with COPS 3.0. Technical Report ANL/MCS-273, Argonne National Laboratory, 2004.
- [4] J.-P. Dussault. Numerical stability and efficiency of penalty algorithms. *SIAM Journal on Numerical Analysis*, 32(1):296–317, 1995.
- [5] A. S. El-Bakry, R. A. Tapia, T. Tsuchiya, and Y. Zhang. On the formulation and theory of newton interior point methods for nonlinear programming. *Journal of Optimization Theory and Applications*, 89(3):507–541, 1996.
- [6] A. V. Fiacco and G. P. McCormick. *Nonlinear Programming: Sequential Unconstrained Minimization Techniques*. J. Wiley and Sons, Chichester, England, 1968. Reprinted as *Classics in Applied Mathematics*, SIAM, Philadelphia, USA, 1990.
- [7] D. M. Gay, M. L. Overton, and M. H. Wright. A primal-dual interior method for nonconvex nonlinear programming. In Y. Yuan, editor, *Advances in Nonlinear Programming*, pages 31–56, Dordrecht, The Netherlands, 1998. Kluwer Academic Publishers.
- [8] N. I. M. Gould, D. Orban, A. Sartenaer, and Ph. L. Toint. Superlinear Convergence of Primal-Dual Interior Point Algorithms for Nonlinear Programming. *SIAM Journal on Optimization*, 11(4):974–1002, 2001.
- [9] N. I. M. Gould, D. Orban, A. Sartenaer, and Ph. L. Toint. Componentwise fast convergence in the solution of full-rank systems of nonlinear equation. *Mathematical Programming, Series B*, 92(3):481–508, 2002.

- [10] C.T. Kelley. *Iterative Methods for Linear and Nonlinear Equations*. Frontiers in Applied Mathematics 16. SIAM, Belmont, 1995.
- [11] S. Mehrotra. On the implementation of a primal-dual interior point method. *SIAM Journal on Optimization*, 2(4):575–601, 1992.
- [12] J. Nocedal, A. Wächter, and R. A. Waltz. Adaptive barrier strategies for nonlinear interior methods. Technical Report, Northwestern University, IL, USA, 2005.
- [13] R. J. Vanderbei and D. F. Shanno. An interior point algorithm for nonconvex nonlinear programming. *Computational Optimization and Applications*, 13(2):231–252, 1999.
- [14] L. N. Vicente and S. J. Wright. Local convergence of a primal-dual method for degenerate nonlinear programming. *Computational Optimization and Applications*, 22(3):311–328, 2002.
- [15] A. Wächter and L. T. Biegler. On the implementation of an interior-point filter line-search algorithm for large-scale nonlinear programming. Research Report RC 23149, IBM T. J. Watson Research Center, NY, USA, 2004.
- [16] H. Yabe and H. Yamashita. Q-superlinear convergence of primal-dual interior point quasi-Newton methods for constrained optimization. *Journal of the Operations Research Society of Japan*, 40(3):415–436, 1997.
- [17] H. Yamashita and H. Yabe. Superlinear and quadratic convergence of some primal-dual interior point methods for constrained optimization. *Mathematical Programming, Series A*, 75(3):377–397, 1996.
- [18] H. Yamashita, H. Yabe, and T. Tanabe. A globally and superlinearly convergent primal-dual interior point trust region method for large scale constrained optimization. *Mathematical Programming, Series A*, 102(1):111–151, 2005.