

Dini Derivative and a Characterization for Lipschitz and Convex Functions on Riemannian Manifolds

O. P. Ferreira*

July 10, 2006

Abstract

Dini derivative on Riemannian manifold setting is studied in this paper. In addition, a characterization for Lipschitz and convex functions defined on Riemannian manifolds and sufficient optimality conditions for constraint optimization problems in terms of the Dini derivative are given.

Keywords: Dini derivative, Convex functions, Lipschitz functions, Riemannian manifolds.

1 Introduction

In the last few years, several important concepts of nonsmooth analysis have been extended from Euclidean space to Riemannian manifold setting, in order to go further in the study of optimization problems and related topics. Works dealing with this subject include the ones by D. Azagra, J. Ferrera and F. López-Mesas [1], D. Azagra, J. Ferrera [2], O. P. Ferreira [7] and Yu. S. Ledyev and Q. J. Zhu [11, 12, 13]. It is worthwhile to mention that extensions of concepts and techniques of optimization from Euclidean spaces to Riemannian manifolds have been extensively studied on several papers including [9], [15], [16], [17], and [20].

Lipschitz and convex functions play an important role in nonsmooth analysis on linear spaces and as well known the Dini derivative is a very useful tool in the analysis of that functions. Our aim in this paper is study some properties of Dini derivative on Riemannian manifolds context and provide a characterization for Lipschitz and convex functions in terms of this derivative. In addition, we obtain a sufficient optimality conditions for constraint optimization problems in that setting.

*Universidade Federal de Goiás, Instituto de Matemática e Estatística, Campus Samambaia, Caixa Postal 131, Goiânia, GO, 74001-970, Brazil (E-mail: orizon@mat.ufg.br). The author was supported in part by FUNAPE/UFG, PADCT-CNPq, PRONEX-Optimization(FAPERJ/CNPq) and PROCAD(CAPES).

A couple of papers have dealt with the issue of characterization for Lipschitz and convex functions in the context of linear spaces. F. H. Clarke, R. J. Stern and P. R. Wolenski [5] have given a characterization for Lipschitz functions in terms of both Dini derivative and proximal subgradient in the context of Hilbert space. R. A. Poliquin [18] provided a characterization for convex functions in terms of proximal subgradient on Euclidean Space \mathbb{R}^n and F. H. Clarke, R. J. Stern and P. R. Wolenski [5] using a novel approach extended this result to Hilbert space. R. Correa, A. Jofré and L. Thibault [6] have given a characterization for convex functions in terms of Clarke subdifferential in the context of the Banach space. Also, D. T. Luc and S. Swaminathan [14] have obtained a characterization for convex functions in terms of Dini derivative on real topological vector space.

The organization of the paper is as follows: In Subsection 1.1, we list some basic notations and auxiliary results used in this presentation. In Section 2 we introduce the concept of Dini derivative of functions defined on a Riemannian manifolds and obtain some results for it. In Section 3 we provide a characterization for Lipschitz functions in terms of Dini derivative on Riemannian manifold setting. In Section 4 we obtain a characterization for convex functions defined on Riemannian manifolds in terms of the Dini derivative. In Section 5 we give sufficient optimality conditions for constraint optimization problems in terms of the Dini derivative. We conclude this paper by making some remarks about extensions of our results.

1.1 Notation and auxiliary results

In this section we recall some notations, definitions and basic properties of Riemannian manifolds used throughout the paper. They can be found in many introductory books on Riemannian Geometry, for example in [19].

Throughout the paper, M is a smooth manifold and C^1 is the class of all continuously differentiable function on M . The space of vector fields on M is denoted by $\mathcal{X}(M)$, by $T_p M$ the tangent space of M at p and by $TM = \bigcup_{x \in M} T_x M$ the *tangent bundle* of M . Let M be endowed with a Riemannian metric $\langle \cdot, \cdot \rangle$, with corresponding norm denoted by $\| \cdot \|$, so that M is now a *Riemannian manifold*. Let us recall that the metric can be used to define the length of piecewise C^1 curve $c : [a, b] \rightarrow M$ joining p to q , i.e., such that $c(a) = p$ and $c(b) = q$, by $l(c) = \int_a^b \|c'(t)\| dt$. Minimizing this length functional over the set of all such curves we obtain a distance $d(p, q)$, which induces the original topology on M . Also, the metric induces a map $f \in C^1(M) \mapsto \text{grad } f \in \mathcal{X}(M)$, which associates to each f its *gradient* via the rule $\langle \text{grad } f, X \rangle = df(X)$, for all $X \in \mathcal{X}(M)$. The chain rule generalizes to this setting in the usual way: $(f \circ c)'(t) = \langle \text{grad } f(c(t)), c'(t) \rangle$, for all curve $c \in C^1$. Let c be a curve joining points p and q in M and let ∇ be a Levi-Civita connection associated to $(M, \langle \cdot, \cdot \rangle)$. For each $t \in [a, b]$, ∇ induces an isometry, relative to $\langle \cdot, \cdot \rangle$, $P(c)_t^a : T_{c(a)} M \rightarrow T_{c(t)} M$, the so-called *parallel translation* along c from $c(a)$ to $c(t)$. A vector field V along c is said to be *parallel* if $\nabla_{c'} V = 0$. If c' itself is parallel, then we say that c is a *geodesic*. The geodesic equation $\nabla_{\gamma'} \gamma' = 0$ is a second order nonlinear ordinary differential equation, so the geodesic γ is determined by its position and velocity at one point. It is easy to check that $\|\gamma'\|$ is constant. We say that γ is *normalized* if $\|\gamma'\| = 1$. A geodesic $\gamma : [a, b] \rightarrow M$ is said to be *minimal* if its length is equal the

distance of its end points, i.e. $l(\gamma) = d(\gamma(a), \gamma(b))$.

A finite dimensional Riemannian manifold is *complete* if its geodesics are defined for any values of t . The Hopf-Rinow's theorem asserts that if the Riemannian manifold M is complete, then any pair of points in M can be joined by a (not necessarily unique) minimal geodesic. Moreover, (M, d) is a complete metric space and its closed and bounded subsets are compact. *In this paper, we assume that all manifolds are complete and finite dimensional.*

The *exponential map* $\exp_p : T_p M \rightarrow M$ is defined by $\exp_p v = \gamma_v(1)$, where γ_v is the geodesic defined by it's position p and velocity v at p . We can prove that, $\gamma_v(t) = \exp_p tv$ for any values of t . For $p \in M$, let

$$i_p := \sup \left\{ r > 0 : \exp_p|_{B_r(o_p)} \text{ is diffeomorphism} \right\},$$

where o_p denotes the origin of $T_p M$ and $B_r(o_p) := \{v \in T_p M : \|v - o_p\| < r\}$. Note that if $0 < \delta < i_p$ then $\exp_p B_\delta(o_p) = B_\delta(p)$, where $B_\delta(p) := \{q \in M : d(p, q) < \delta\}$. The number i_p is called the *injetivity radius* of M at p . For any $p \in M$, the map $d^2(p, \cdot) \in C^\infty(B_{i_p}(p))$, where $B_{i_p}(p) = \{q \in M : d(p, q) < i_p\}$, and $\text{grad } d^2(p, q) = -2 \exp_q^{-1} p$, for all $q \in B_{i_p}(p)$. Throughout the paper M denotes a finite dimensional Riemannian manifold which is complete and for any $p \in M$, $\delta > 0$ we denote by $B_\delta(p)$ the open ball and by $B_\delta[p]$ the closed ball centered in p .

The set $C \subseteq M$ is said to be *convex*, if for any $p, q \in C$ the minimal geodesics joining p to q are contained in C .

Let U be an open subset of M . From now on, we denote by $\mathcal{F}(U)$ the class of all function $f : M \rightarrow (-\infty, +\infty]$ which are lower semicontinuous on U and $\text{dom}(f) \cap U \neq \emptyset$, where $\text{dom}(f) = \{x \in M : f(x) < +\infty\}$. If $U = M$ we denote \mathcal{F} for $\mathcal{F}(U)$.

2 Dini Derivative

In this section we study some properties of Dini derivative of locally Lipschitz functions. Our main result shows that Dini derivative does not depend on the curve, namely, it just depend on the direction.

Definition 1. A function $f \in \mathcal{F}(U)$ is said to be Lipschitz on V , of rank $L \geq 0$, if $V \subset \text{dom}(f)$ and there holds

$$|f(p) - f(q)| \leq Ld(p, q), \quad \forall p, q \in V,$$

where d is the Riemannian distance on M .

We denote the set of all Lipschitz function on V , of rank L , by $\text{Lip}_L(V)$.

Definition 2. A function $f \in \mathcal{F}(U)$ is said to be Lipschitz at p , of rank L_p , if there exists $\delta > 0$ such that $f \in \text{Lip}_{L_p}(B_\delta(p))$ and locally Lipschitz on V if it is Lipschitz at every $p \in V$.

Now, we are going to present three important examples of Lipschitz functions that emerge in the study of Riemannian manifold, see [19] and [10].

Example 1. Let M be a non-compact Riemannian manifold. A geodesic $\gamma : [0, +\infty) \rightarrow M$ parameterized by arc-length and emanating from p is called a ray emanating from p if $d(\gamma(t), \gamma(s)) = |t - s|$, for all $t, s > 0$. For a ray γ , the Busemann function $b_\gamma : M \rightarrow (-\infty, +\infty]$ is defined by

$$b_\gamma(q) = \lim_{t \rightarrow +\infty} (t - d(q, \gamma(t))),$$

is Lipschitz of rank 1, see [7].

Example 2. Let M be a Riemannian manifold and S a closed subset of M . The distance function $d_S(p) = \inf\{d(p, s) : s \in S\}$ is Lipschitz of rank 1, see [7].

Example 3. Let $f \in \mathcal{F}$ and $\lambda > 0$. Suppose that f is bounded below by the constant k . Then, it easy to show that the function $f_\lambda : M \rightarrow (-\infty, +\infty]$ defined by $f_\lambda(p) = \inf\{f(q) + \lambda d^2(p, q)\}$, is also bounded below by k . Moreover, following the same pattern used to prove the first part of the Theorem 5.1 on the page 44 of [3], we can prove that f_λ is locally Lipschitz on its domain.

We remark that, the Lipschitz properties depends on the Riemannian metric defined on M . In other words, if the metric on M is changed then the set of Lipschitz functions on M becomes different of previous one, see Example 4.4 in [7].

Definition 3. The lower Dini derivative of $f \in \mathcal{F}$ at $p \in \text{dom}(f)$ in the direction of $v \in T_p M$ is defined by

$$f'(p, v) = \liminf_{t \rightarrow 0^+} \frac{f(\gamma(t)) - f(p)}{t},$$

where $\gamma : \mathbb{R} \rightarrow M$ is the geodesic such that $\gamma(0) = p$ and $\gamma'(0) = v$. If $p \notin \text{dom}(f)$ define $f'(p, v) = -\infty$ for all $v \in T_p M$.

Note that if $f \in \mathcal{F}$ is a locally Lipschitz function on $\text{dom}(f)$, then $\text{dom}(f)$ is a open set and for all $p \in \text{dom}(f)$ and $v \in T_p M$ there exists the directional derivative and $f'(p, v) < +\infty$. Now, we are going to obtain some properties for lower Dini derivate. We begin with some preliminaries.

Let $p \in M$ and $c_1, c_2 : (-\varepsilon, \varepsilon) \rightarrow M$ be a two C^1 curves, such that $c_1(0) = c_2(0) = p$, $c_1'(0) = v$ and $c_2'(0) = w$. Assume that the image sets $c_1((-\varepsilon, \varepsilon))$, $c_2((-\varepsilon, \varepsilon))$ are in $B_{i_p}(p)$. Let $\alpha : [0, 1] \times (-\varepsilon, \varepsilon) \rightarrow M$ be a *variation of geodesics* given by

$$\alpha(t, s) = \exp_{c_1(s)}(t \exp_{c_1(s)}^{-1} c_2(s)). \quad (1)$$

Note that for each $s \in (-\varepsilon, \varepsilon)$ we have $\alpha(0, s) = c_1(s)$, $\alpha(1, s) = c_2(s)$ and the curve $\alpha_s : [0, 1] \rightarrow M$ given by $\alpha_s(t) = \alpha(t, s)$ is a geodesic. In particular, $\alpha_0(t) = \alpha(t, 0) = p$ is a constant geodesic. Now, consider the vector fields

$$T(\cdot, s) := \frac{\partial \alpha}{\partial t}(\cdot, s), \quad \text{and} \quad J(\cdot, s) = \frac{\partial \alpha}{\partial s}(\cdot, s).$$

Above definitions imply that $T(\cdot, s)$ is tangent to geodesic α_s and $J(\cdot, s)$, the *Jacobi vector field* through α_s , satisfies the following differential equation

$$\nabla_{\frac{\partial \alpha}{\partial t}} \nabla_{\frac{\partial \alpha}{\partial t}} J(t, s) + R(J(t, s), T(t, s))T(t, s) = 0, \quad (2)$$

where R is the curvature tensor, see [19].

Lemma 1. Let $p \in M$, c_1 and c_2 two C^1 curves in M such that $c_1(0) = c_2(0) = p$, $c_1'(0) = v$ and $c_2'(0) = w$. Then

$$\lim_{s \rightarrow 0^+} \frac{d(c_1(s), c_2(s))}{s} = \|w - v\|.$$

Proof. For simplifying the notations define $\psi(s) = d(c_1(s), c_2(s))$. Consider α the variation of geodesics defined by (1), then $\psi(s) = \|\alpha'_s(t)\| = \|T(t, s)\|$. Since $T(t, 0) = \alpha'_0(t) = 0$ the first derivative of ψ^2 at $s = 0$ is given by

$$\frac{d}{ds}(\psi^2(s))|_{s=0} = 2 \left\langle \nabla_{\frac{\partial \alpha}{\partial s}} T(t, 0), T(t, 0) \right\rangle = 0, \quad (3)$$

and the second derivative by

$$\begin{aligned} \frac{d^2}{ds^2}(\psi^2(s))|_{s=0} &= 2 \left\langle \nabla_{\frac{\partial \alpha}{\partial s}} \nabla_{\frac{\partial \alpha}{\partial s}} T(t, 0), T(t, 0) \right\rangle + 2 \left\langle \nabla_{\frac{\partial \alpha}{\partial s}} T(t, 0), \nabla_{\frac{\partial \alpha}{\partial s}} T(t, 0) \right\rangle \\ &= 2 \left\langle \nabla_{\frac{\partial \alpha}{\partial s}} T(t, 0), \nabla_{\frac{\partial \alpha}{\partial s}} T(t, 0) \right\rangle, \end{aligned} \quad (4)$$

in addition, the equation (2) becomes

$$\nabla_{\frac{\partial \alpha}{\partial t}} \nabla_{\frac{\partial \alpha}{\partial t}} J(t, 0) = 0.$$

Now, the latter equality together with the conditions $J(0, 0) = v$ and $J(1, 0) = w$ implies that $J(t, 0) = v + t(w - v)$. Using Symmetry's Lemma (see Lemma 2.2, pp. 35 in [19]) and this equality we have

$$\nabla_{\frac{\partial \alpha}{\partial s}} T(t, 0) = \nabla_{\frac{\partial \alpha}{\partial s}} \frac{\partial \alpha}{\partial t}(t, 0) = \nabla_{\frac{\partial \alpha}{\partial t}} \frac{\partial \alpha}{\partial s}(t, 0) = \nabla_{\frac{\partial \alpha}{\partial t}} J(t, 0) = w - v.$$

So, substituting last equality in (4) we obtain

$$\frac{d^2}{ds^2}(\psi^2(s))|_{s=0} = 2\|w - v\|^2.$$

Thus, as $\psi(0) = 0$ it follows from latter equality and (3) that $\psi^2(s) = \|w - v\|^2 s^2 + \mathcal{O}(s^2)$, with $\lim_{s \rightarrow 0^+} \mathcal{O}(s^2)/s^2 = 0$. Therefore, the result follows from definition of ψ . \square

Corollary 1. If $f \in \mathcal{F}$ is Lipschitz in p with constant L_p , then $f'(p, \cdot) \in \text{Lip}_{L_p}(T_p M)$.

Proof. Let v and w in $T_p M$. Let γ, η be the geodesics with $\gamma(0) = \eta(0) = p$, $\gamma'(0) = v$ and $\eta'(0) = w$. First note that

$$\left| \frac{f(\gamma(t)) - f(p)}{t} - \frac{f(\eta(t)) - f(p)}{t} \right| = \frac{|f(\gamma(t)) - f(\eta(t))|}{t}.$$

Since f is Lipschitz in p and $\gamma(0) = \eta(0) = p$ we have $|f(\gamma(t)) - f(\eta(t))| \leq L_p d(\gamma(t), \eta(t))$ for all $t \in [0, \varepsilon)$ and some $\varepsilon > 0$. This inequality together with the above equality imply that

$$\frac{f(\gamma(t)) - f(p)}{t} - L_p \frac{d(\gamma(t), \eta(t))}{t} \leq \frac{f(\eta(t)) - f(p)}{t} \leq \frac{f(\gamma(t)) - f(p)}{t} + L_p \frac{d(\gamma(t), \eta(t))}{t},$$

for all $t \in [0, \varepsilon)$. Now, as $\gamma'(0) = v$ and $\eta'(0) = w$ we obtain from Lemma 1 that

$$\lim_{t \rightarrow 0^+} \frac{d(\gamma(t), \eta(t))}{t} = \|v - w\|.$$

Therefore, taking \liminf in above inequality and considering the latter equality we obtain

$$f'(p, v) - L_p \|v - w\| \leq f'(p, w) \leq f'(p, v) + L_p \|v - w\|,$$

which implies that $f'(x, \cdot)$ is Lipschitz with constant L_p . □

Corollary 2. Let $f \in \mathcal{F}$ be a Lipschitz function in $p \in \text{dom}(f)$. For all C^1 curve $c : [0, \varepsilon) \rightarrow M$ satisfying $c(0) = p$ and $c'(0) = v$ there holds

$$f'(p, v) = \liminf_{t \rightarrow 0^+} \frac{f(c(t)) - f(p)}{t}.$$

Proof. Let γ be a geodesic with $\gamma(0) = p$ and $\gamma'(0) = v$. Since f is Lipschitz in p , a similar argument used in the proof of latter corollary implies

$$\frac{f(\gamma(t)) - f(p)}{t} - L_p \frac{d(\gamma(t), c(t))}{t} \leq \frac{f(c(t)) - f(p)}{t} \leq \frac{f(\gamma(t)) - f(p)}{t} + L_p \frac{d(\gamma(t), c(t))}{t},$$

for all $t \in [0, \bar{\varepsilon})$ and some $\bar{\varepsilon} \leq \varepsilon$, where L_p is a Lipschitz constant of f in p . Now, as $c'(0) = \gamma'(0) = v$ the Lemma 1 implies that $\lim_{t \rightarrow 0^+} d(\gamma(t), c(t))/t = 0$. So, taking \liminf in latter inequality we obtain the statement. □

Corollary 3. Let $f \in \mathcal{F}(U)$ be a locally Lipschitz function on U and $I \subset \mathbb{R}$ an open interval. If $c : I \rightarrow U$ be is a C^1 curve, then

$$(f \circ c)'(t, 1) = f'(c(t), c'(t)), \quad \forall t \in I.$$

Proof. It follows from Corollary 2. □

3 Characterization for Lipschitz Functions

In this section we present a characterization for Lipschitz functions defined on Riemannian manifolds. It is worth pointing out that our results in this section were obtained by adapting, for our context, the techniques introduced by F. H. Clarke, R. J. Stern and P. R. Wolenski [5] for characterizing Lipschitz functions in Hilbert spaces.

Proposition 1. Let U be an open convex subset of M and let $f \in \mathcal{F}(U)$. If f is locally Lipschitz on $\text{dom} f$ with constant L everywhere, then the following statements hold:

- i) $\text{dom} f = U$;
- ii) $f \in \text{Lip}_L(U)$.

Proof. For i). First note that $\text{dom}(f) \subset U$. It remains to show that $U \subset \text{dom}(f)$. For that, take $q \in U$. Now, take $p \in \text{dom}(f)$ and a minimal geodesic $\gamma : [0, 1] \rightarrow M$ such that $\gamma(0) = p$ and $\gamma(1) = q$. Since U is convex it follows that $\gamma([0, 1]) \subset U$. We claim that $q \in \text{dom}(f)$. Suppose not. Thus we have that

$$t_* := \sup \{t \in (0, 1] : f(\gamma(t)) < +\infty\} < 1,$$

and since $p \in \text{dom}(f)$ and f is locally Lipschitz on $\text{dom} f$ we also have $0 < t_*$. So, taking $t' \in (0, t_*)$ we have $\gamma([0, t']) \subset \text{dom} f$. Since $\gamma([0, t'])$ is compact and f is locally Lipschitz on $\text{dom} f$, we can take $0 = t_0 < t_1 < \dots < t_n = t'$ and positive numbers $\delta_0 \cdots \delta_{n-1}$ satisfying

$$\gamma([t_i, t_{i+1}]) \subset B_{\delta_i}(\gamma(t_i)) \subset U \quad \text{and} \quad f \in \text{Lip}_L(B_{\delta_i}(\gamma(t_i))),$$

for all $i = 0, \dots, n-1$. As the geodesic γ is minimal, using the local Lipschitz property we obtain

$$\begin{aligned} f(\gamma(t')) &= f(p) + \sum_{i=0}^{n-1} (f(\gamma(t_{i+1})) - f(\gamma(t_i))) \\ &\leq f(p) + \sum_{i=0}^{n-1} L d(\gamma(t_i), \gamma(t_{i+1})) \\ &\leq f(p) + L d(p, \gamma(t')). \end{aligned} \tag{5}$$

Since f is lower semicontinuous, letting t' goes to t_* in latter equation we conclude that

$$f(\gamma(t_*)) < +\infty,$$

which implies that $\gamma(t_*) \in \text{dom}(f)$. According to f is Lipschitz at $\gamma(t_*)$, definition of t_* is violated. Therefore, we obtain that f is finite on the entire $\gamma([0, 1])$. In particular, $q \in \text{dom}(f)$, so $U \subset \text{dom}(f)$ and the first statement follows.

For ii). Take $p, q \in U$ and a minimal geodesic $\gamma : [0, 1] \rightarrow M$ such that $\gamma(0) = p$ and $\gamma(1) = q$.

Since U is convex it follows from item i) that $\gamma([0, 1]) \subset \text{dom}(f)$. With an analogous argument used to obtain (5) we can show that

$$f(q) \leq f(p) + Ld(p, q).$$

Now, by reversing the roles of p and q , it easy to conclude that $f \in \text{Lip}_L(U)$, and the second statement is proved. \square

Theorem 1. Let $f \in \mathcal{F}$ and U be an open convex subset of M . Then f is Lipschitz on U of rank $L \geq 0$ if, and only if,

$$|f'(p, v)| \leq L\|v\|, \quad \forall p \in U, \quad \forall v \in T_p M.$$

Proof. First, suppose that f is Lipschitz on U of rank $L \geq 0$. Take $p \in U$, $v \in T_p M$. Let γ be the geodesic such that $\gamma(0) = p$ and $\gamma'(0) = v$. Since $\gamma(0) = p$ and f is Lipschitz on U of rank $L \geq 0$, there exists $\delta > 0$ such that

$$|f(\gamma(t)) - f(p)| \leq Ld(\gamma(t), p) \leq L\|v\|t, \quad \forall t \in [0, \delta].$$

So, using definition of lower Dini derivative, the above inequality implies that

$$|f'(p, v)| = \liminf_{t \rightarrow 0^+} \frac{|f(\gamma(t)) - f(p)|}{t} \leq L \liminf_{t \rightarrow 0^+} \frac{d(\gamma(t), p)}{t} = L\|v\|.$$

Reciprocally, suppose that $|f'(p, v)| \leq L\|v\|$, for all $p \in U$ and $v \in T_p M$. Let $p_0 \in \text{dom}(f)$. Since $f \in \mathcal{F}(U)$, take $0 < \delta < i_p$ such that f is bounded below on $B_{4\delta}(p_0) \subset U$. Let $K > L$ and $q \in B_\delta(p_0)$. Define $g : U \rightarrow [0, +\infty]$ as

$$g(p) = \begin{cases} Kd(p, q) & \text{if } p \in B_{2\delta}(q); \\ Kd(p, q) + \frac{(d(p, q) - 2\delta)^2}{3\delta - d(p, q)} & \text{if } p \in B_{3\delta}(q) \setminus B_{2\delta}(q); \\ +\infty & \text{otherwise.} \end{cases}$$

It is easy to see that $g \in \mathcal{F}(U)$. Moreover, as $g \in C^1(B_{3\delta}(q) \setminus \{q\})$ and $0 < \delta < i_p$ there holds

$$\text{grad } g(p) = \left[K + \frac{2(d(p, q) - 2\delta)(3\delta - d(p, q)) + d(p, q)(d(p, q) - 2\delta)}{(3\delta - d(p, q))^2} \right] (-\exp_p^{-1} q / d(p, q)),$$

for all $p \in B_{3\delta}(q) \setminus B_{2\delta}(q)$. Now, note that $f + g$ is in $\mathcal{F}(U)$, goes to $+\infty$ as p goes to the boundary of $B_{3\delta}(q)$ and is bounded below on $B_{3\delta}(q)$. Therefore, as M is a complete Riemannian manifolds of finite dimension, there exists $p_* \in B_{3\delta}(q)$ a minimizer for $f + g$. First we assume that $p_* \neq q$. Since $p_* \in B_{3\delta}(q) \setminus \{q\}$, $g \in C^1(B_{3\delta}(q) \setminus \{q\})$ and $f + g \in \mathcal{F}(U)$ we have

$$f'(p_*, v) + \langle \text{grad } g(p_*), v \rangle \geq 0 \quad \forall v \in T_{p_*} M. \quad (6)$$

For simplifying the notations set $v_* := \exp_{p_*}^{-1}q/d(p_*, q)$ and

$$\ell := \frac{2(d(p_*, q) - 2\delta)(3\delta - d(p_*, q)) + d(p_*, q)(d(p_*, q) - 2\delta)}{(3\delta - d(p_*, q))^2}.$$

So, $\text{grad } g(p_*) = -Kv_*$ if $p_* \in B_{2\delta}[q]$ and $\text{grad } g(p_*) = -[K + \ell]v_*$ if $p_* \in B_{3\delta}(q) \setminus B_{2\delta}(q)$. Thus, letting $v = v_*$ in (6) and taking in account that $\ell \geq 0$ for $p_* \in B_{3\delta}(q) \setminus B_{2\delta}(q)$ we obtain

$$f'(p_*, v_*) \geq K\|v_*\| > L\|v_*\|,$$

which contradicts our assumption. Consequently we must have $p_* = q$. Due the fact that $g(q) = 0$, the point q is a minimizer of $f + g$ and $B_\delta(p_0) \subset B_{2\delta}[q]$ we have

$$f(q) = (f + g)(q) \leq (f + g)(p) \leq f(p) + Kd(p, q), \quad \forall p \in B_\delta(p_0).$$

Since we can change the roles of q and p in the above argument the following inequality holds

$$|f(q) - f(p)| \leq Kd(p, q), \quad \forall p, q \in B_\delta(p_0).$$

Letting K goes to L , in the latter inequality, we conclude that for any $p_0 \in \text{dom } f$ there exists $\delta > 0$ such that $f \in \text{Lip}_L B_\delta(p_0)$. Thus we have shown that f is locally Lipschitz on $\text{dom}(f)$ with the same rank L everywhere. Therefore, for finishing the proof use the Proposition 1. \square

Example 4. Let S^n be the set of symmetric matrix endowed with the Frobenius metric defined by $\langle U, V \rangle = \text{tr}(VU)$ and let S_{++}^n be the set of positive definite symmetric matrix. Let $f : S_{++}^n \rightarrow \mathbb{R}$ be defined by $f(X) = \ln \det X$. It easy to see that the function f is not Lipschitz on S_{++}^n . For each $X \in S_{++}^n$ define a new inner product in S^n as

$$\langle\langle U, V \rangle\rangle = \langle X^{-1}UX^{-1}, V \rangle \quad \forall U, V \in S^n.$$

Endowing S_{++}^n with the Riemannian metric $\langle\langle \cdot, \cdot \rangle\rangle$ we obtain a complete Riemannian manifold. We denote by M this Riemannian manifold. Note that $f \in C^1$ on M . So,

$$f'(X, V) = \langle\langle \text{grad } f(X), V \rangle\rangle \quad \forall V \in T_X M.$$

Because the usual gradient of f on S_{++}^n is $\nabla f(X) = X^{-1}$, we have that the gradient of f on M is given by

$$\text{grad } f(X) = X\nabla f(X)X = X.$$

So, $\|\text{grad } f(X)\|^2 = \langle\langle \text{grad } f(X), \text{grad } f(X) \rangle\rangle = 1$. Thus,

$$|f'(X, V)| = |\langle\langle \text{grad } f(X), V \rangle\rangle| \leq \|\text{grad } f(X)\| \|V\| = \|V\|, \quad \forall V \in T_X M.$$

From Theorem 1 it follows that f is Lipschitz on M of rank $L = 1$.

Example 5. Let $\Omega = \{p = (p_1, p_2) \in \mathbb{R}^2 : p_2 > 0\}$ and let $f : \Omega \rightarrow \mathbb{R}$ be given by $f(p) = \ln(p_2)$. It is easy to see that f is not a Lipschitz function on Ω with respect to Euclidean metric $\langle \cdot, \cdot \rangle$. Let G be a 2×2 matrix defined by $G(p) = (g_{ij}(p))$, where

$$g_{11}(p) = g_{22}(p) = \frac{1}{p_2^2}, \quad g_{12}(p) = g_{21}(p) = 0.$$

Endowing Ω with the Riemannian metric $\langle \cdot, \cdot \rangle$ defined by $\langle u, v \rangle = \langle G(p)v, u \rangle$, we obtain a complete Riemannian manifold \mathbb{H}^2 , namely, the upper half-plane model of the Hyperbolic space. Note that $f \in C^1$ and the gradient of f in \mathbb{H}^2 is given by

$$\text{grad } f(p) = G(p)^{-1} \nabla f(p) = (0, p_2),$$

where ∇f is the usual gradient of f in Ω . It is simple to show that $\|\text{grad } f(p)\| = 1$ and so,

$$|f'(p, v)| = |\langle \text{grad } f(p), v \rangle| \leq \|v\|, \quad \forall v \in T_p \mathbb{H}^2.$$

Therefore, from Theorem 1 it follows that f is Lipschitz on \mathbb{H}^2 of rank $L = 1$.

4 Characterization for Convex Functions

In this section we obtain a characterization for convex functions defined on Riemannian manifolds. As usual, in the type of characterization we shall present, first we obtain a result like mean value theorem adapted to our setting.

Definition 4. A function $f : M \rightarrow (-\infty, +\infty]$ is said to be convex (respectively, strictly convex) if for all minimal geodesic $\gamma : [a, b] \rightarrow M$, the composition $f \circ \gamma : [a, b] \rightarrow (-\infty, +\infty]$ is convex (respectively, strictly convex).

It follows from above definition that if $f : M \rightarrow (-\infty, +\infty]$ is a convex function then $\text{dom}(f)$ and the sub-level sets $\{p \in M : f(p) \leq k\}$ are convex sets, for all $k \in \mathbb{R}$.

Example 6. Let $S^n = \{p \in \mathbb{R}^{n+1} : \|p\| = 1\}$ be a unitary sphere. Fix $\tilde{p} = (0, \dots, 0, 1) \in \mathbb{R}^{n+1}$. Setting $p = (p_1, \dots, p_n, p_{n+1})$, define $\varphi : S^n \rightarrow (-\infty, +\infty]$ as

$$\varphi(p) = \begin{cases} d(\tilde{p}, p), & \text{if } p_{n+1} \geq 0; \\ +\infty & \text{if } p_{n+1} < 0. \end{cases}$$

Note that $\varphi \in \mathcal{F}$, but φ is not a convex function and its $\text{dom}(\varphi) = \{p \in S^n : p_{n+1} \geq 0\}$ is not a convex set. Now, let $C \subset \{p \in S^n : p_{n+1} > 0\}$ a closed and convex set and define $\rho : S^n \rightarrow (-\infty, +\infty]$ as

$$\rho(p) = \begin{cases} d(\tilde{p}, p), & \text{if } p \in C; \\ +\infty & \text{if } p \notin C. \end{cases}$$

Note that ρ lower semicontinuous, convex and its domain C is closed. In general, for all $\tilde{p} \in M$ and $C \subset \{p \in M : d(\tilde{p}, p) < \pi/2\}$ a closed convex set, the function $\eta : S^n \rightarrow (-\infty, +\infty]$ defined as $\eta(p) = d(\tilde{p}, p)$, if $p \in C$ and $\eta(p) = +\infty$ if $p \notin C$, is lower semicontinuous and convex.

In above example $\text{dom}(\varphi) = \{p \in \mathbb{R}^n : p_{n+1} \geq 0\}$ is closed and its interior is convex. As $\text{dom}(\varphi)$ is not convex we conclude that, in general, the closure of a convex set is not convex. Note that φ is Lipschitz of rank $L = 1$ in $\text{int}(\text{dom}(\varphi))$, but not in $\text{dom}(\varphi)$.

Definition 5. A function $f \in \mathcal{F}$ is locally bounded in p if there exists $\delta > 0$ and $r > 0$ such that $f(q) \leq r$ for all $q \in B_\delta(p)$, and $f \in \mathcal{F}$ is locally bounded in $\text{dom}(f)$ if it is locally bounded in all point $p \in \text{dom}(f)$.

Note that if $f \in \mathcal{F}$ is locally bounded in $\text{dom}(f)$ then $\text{dom}(f)$ must be open.

Proposition 2. Let $f \in \mathcal{F}$ be locally bounded in $\text{dom}(f)$. If f is convex, then f is locally Lipschitz on $\text{dom}(f)$.

Proof. See Proposition 5.2 in [1]. □

We remark that in n -dimensional Euclidean spaces for proving that a convex function f is locally bounded in $\text{dom}(f)$ are used two important results, namely, for each $p \in \text{dom}(f)$ there exists a n -dimensional simplex $\Delta \subset \text{dom}(f)$ such that $p \in \text{int}(\Delta)$ and Jensen's inequality. However, as far as we know results like these in Riemannian context has not been studied yet.

From now on, we assume that $f \in \mathcal{F}$ and is locally Lipschitz on $\text{dom}(f)$ without explicitly mentioning them in the statements of our results. Note that in this case f is locally bounded in $\text{dom}(f)$ and $\text{dom}(f)$ is open.

Example 7. Let $S^n = \{p \in \mathbb{R}^{n+1} : \|p\| = 1\}$ be a unitary sphere. Fix $\tilde{p} = (0, \dots, 0, 1) \in \mathbb{R}^{n+1}$. Setting $p = (p_1, \dots, p_n, p_{n+1})$, define $\zeta : S^n \rightarrow (-\infty, +\infty]$ as

$$\zeta(p) = \begin{cases} -\ln(\pi/2 - d(\tilde{p}, p)), & \text{if } p_{n+1} > 0; \\ +\infty & \text{if } p_{n+1} < 0. \end{cases}$$

ζ is lower semicontinuous, convex and its domain is open. Moreover, ζ locally Lipschitz in $\text{dom}(\zeta)$.

Example 8. Let $M = \{p \in \mathbb{R}^n : \|p\| = 1\}$ be a unitary sphere. Setting $p = (p_1, \dots, p_n)$, define $\psi : M \rightarrow (-\infty, +\infty]$ as

$$\psi(p) = \begin{cases} -\sum_{i=1}^n \ln(p_i) & \text{if } p_1 > 0, \dots, p_n > 0, \\ +\infty & \text{otherwise.} \end{cases}$$

ψ is lower semicontinuous, convex and its domain is open. Moreover, ψ locally Lipschitz in $\text{dom}(\psi)$.

Proposition 3. If f is convex, then for all $p \in \text{dom}(f)$ and $v \in T_p M$ there holds

$$f'(p, v) = \lim_{t \rightarrow 0^+} \frac{f(\gamma(t)) - f(p)}{t} = \inf_{t > 0} \frac{f(\gamma(t)) - f(p)}{t},$$

where γ is a minimal geodesic such that $\gamma(0) = p$ and $\gamma'(0) = v$.

Proof. It follows from convexity of $f \circ \gamma$. □

Corollary 4. If f is convex and $\gamma : [a, b] \rightarrow \text{dom}(f)$ is a minimal geodesic, then there holds

$$(t - \tilde{t})f'(\gamma(\tilde{t}), \gamma'(\tilde{t})) + f(\gamma(\tilde{t})) \leq f(\gamma(t)), \quad \forall \tilde{t}, t \in [a, b].$$

Proof. Is an immediate consequence of Proposition 3. □

Definition 6. Let $f \in \mathcal{F}$ be a locally Lipschitz function on $\text{dom}(f)$. The lower Dini derivative $f' : TM \rightarrow [-\infty, +\infty]$ is said to be monotone (respectively, strictly monotone) if, for any minimal geodesic $\gamma : [a, b] \rightarrow M$ with its end points in $\text{dom}(f)$, the map $\varphi_\gamma : [a, b] \rightarrow [-\infty, +\infty]$ defined by $\varphi_\gamma(t) = f'(\gamma(t), \gamma'(t))$ is monotone non-decreasing (respectively, increasing).

Note that in above definition we not assume that $\text{dom}(f)$ is a convex set.

Lemma 2. If γ is a geodesic such that $\gamma(0) = p$, $\gamma(1) = q$ and $\gamma([0, 1]) \subset \text{dom}(f)$, then there exist $\tilde{t}, \hat{t} \in (0, 1)$ such that

$$f'(\gamma(\tilde{t}), \gamma'(\tilde{t})) \leq f(q) - f(p) \leq f'(\gamma(\hat{t}), \gamma'(\hat{t})).$$

Proof. Let $\psi : [0, 1] \rightarrow \mathbb{R}$ be defined by $\psi(s) = f(\gamma(s)) - s(f(q) - f(p))$. So, from Corollary 3 is sufficient to show that there exist $\tilde{t}, \hat{t} \in (0, 1)$ such that

$$\psi'(\tilde{t}, 1) \leq 0 \leq \psi'(\hat{t}, 1).$$

First we will prove that there exist $\tilde{t} \in (0, 1)$ such that $\psi'(\tilde{t}, 1) \leq 0$. Since ψ is continuous in $[0, 1]$ and $\psi(0) = \psi(1) = f(p)$, there exists $s_1 \in (0, 1]$ a global minimum of ψ . Now let $s_0 \in (0, s_1)$. We have two possibility: i) $\psi'(s_0, 1) \leq 0$ or ii) $\psi'(s_0, 1) > 0$. If i) occur take $\tilde{t} = s_0$, otherwise $\psi'(s_0, 1) > 0$ implies that there exists $\bar{s} \in (s_0, s_1)$ such that $\psi(s_0) < \psi(\bar{s})$. Hence, by the continuity of ψ there exists $\tilde{t} \in (s_0, s_1)$ such that $\psi(s) \leq \psi(\tilde{t})$ for all $s \in [s_0, s_1]$, which imply $\psi'(\tilde{t}, 1) \leq 0$. The second inequality can be shown by similar argument used in the first one. □

Corollary 5. If γ is a geodesic in M such that $t_2 > t_1$, $\gamma([t_1, t_2]) \subset \text{dom}(f)$, then there exist $\tilde{t}, \hat{t} \in (t_1, t_2)$ such that

$$f'(\gamma(\tilde{t}), \gamma'(\tilde{t})) \leq \frac{f(\gamma(t_2)) - f(\gamma(t_1))}{t_2 - t_1} \leq f'(\gamma(\hat{t}), \gamma'(\hat{t})).$$

Proof. Defining the geodesic $\alpha(s) = \gamma(st_2 + (1-s)t_1)$ we have $\alpha(0) = \gamma(t_1)$ and $\alpha(1) = \gamma(t_2)$. Now, note that $v \mapsto f'(p, v)$ is positively homogeneous, for all $p \in \text{dom}(f)$. So, applying Lemma 2 the statement follows. \square

Lemma 3. If f' is monotone, then $\text{dom}(f)$ is convex.

Proof. Take $p, q \in \text{dom}(f)$ and γ a minimal geodesic joining p to q . Assume without loss of generality that $\gamma(0) = p$ and $\gamma(1) = q$. Define

$$\hat{t} := \inf \{t \in [0, 1] : [\gamma(t), q] \in \text{dom}(f)\}, \quad \tilde{t} := \sup \{t \in [0, 1] : [p, \gamma(t)] \in \text{dom}(f)\}.$$

Since f is locally Lipschitz on $\text{dom}(f)$ we have that $\text{dom}(f)$ is open, so $0 \leq \hat{t} < 1$ and $0 < \tilde{t} \leq 1$. Assume by contraction that $\gamma([0, 1]) \not\subseteq \text{dom}(f)$ or equivalently that $0 < \hat{t}$ and $\tilde{t} < 1$. Now, take $0 < \lambda < \tilde{t} \leq \hat{t} < \tau < 1$. Definition of \hat{t} and \tilde{t} implies that the minimal geodesic segments joining $\gamma(\tau)$ to q and p to $\gamma(\lambda)$ are in $\text{dom}(f)$. Thus, from Corollary 5 there exist $t_1 \in (0, \lambda)$ and $t_2 \in (\tau, 1)$ such that

$$f(\gamma(\lambda)) - f(p) \leq \lambda f'(\gamma(t_1), \gamma'(t_1)), \quad (1 - \tau)f'(\gamma(t_2), \gamma'(t_2)) \leq f(q) - f(\gamma(\tau)).$$

Above inequalities together with monotonicity of the derivative f' imply that

$$\begin{aligned} \frac{f(\gamma(\lambda))}{\lambda} + \frac{f(\gamma(\tau))}{(1 - \tau)} &\leq \frac{f(p)}{\lambda} + f'(\gamma(t_1), \gamma'(t_1)) + \frac{f(q)}{(1 - \tau)} - f'(\gamma(t_2), \gamma'(t_2)) \\ &\leq \frac{f(p)}{\lambda} + \frac{f(q)}{(1 - \tau)}. \end{aligned}$$

Since $f \in \mathcal{F}$, $0 < \tilde{t}$ and $\hat{t} < 1$, letting λ goes to \tilde{t} and τ goes to \hat{t} in the above equation we obtain

$$\frac{f(\gamma(\tilde{t}))}{\tilde{t}} + \frac{f(\gamma(\hat{t}))}{(1 - \hat{t})} \leq \frac{f(p)}{\tilde{t}} + \frac{f(q)}{(1 - \hat{t})}.$$

So, $f(\gamma(\tilde{t})) < +\infty$ and $f(\gamma(\hat{t})) < +\infty$, i.e., $\gamma(\tilde{t}), \gamma(\hat{t}) \in \text{dom}(f)$. As $\text{dom}(f)$ is open we obtain a contradiction with the definitions of \hat{t} and \tilde{t} . Therefore, $\gamma([0, 1]) \subseteq \text{dom}(f)$ and the statement follows. \square

Theorem 2. f is convex if, and only if, f' is monotone.

Proof. Assume that f is convex. First note that $\text{dom}(f)$ is convex. Let $\gamma : [a, b] \rightarrow M$ be a minimal geodesic with its end points in $\text{dom}(f)$ and take $t_1, t_2 \in [a, b]$ such that $t_1 < t_2$. Since $\text{dom}(f)$ is convex we have that $\gamma([t_1, t_2]) \subseteq \text{dom}(f)$ and using Corollary 4 we obtain that

$$(t_2 - t_1)f'(\gamma(t_1), \gamma'(t_1)) + f(\gamma(t_1)) \leq f(\gamma(t_2)), \quad (t_1 - t_2)f'(\gamma(t_2), \gamma'(t_2)) + f(\gamma(t_2)) \leq f(\gamma(t_1)),$$

Therefore, as $t_1 < t_2$ it follows from latter two inequality that $f'(\gamma(t_1), \gamma'(t_1)) \leq f'(\gamma(t_2), \gamma'(t_2))$, hence f' is monotone.

Now, assume that f' is monotone. Let $\gamma : [a, b] \rightarrow M$ be a geodesic and let $t_1, t_2 \in [a, b]$ such that $t_1 < t_2$. If $\gamma(t_1) \notin \text{dom}(f)$ or $\gamma(t_2) \notin \text{dom}(f)$ we have

$$f \circ \gamma((1 - \lambda)t_1 + \lambda t_2) \leq (1 - \lambda)f(\gamma(t_1)) + \lambda f(\gamma(t_2)),$$

for all $\lambda \in [0, 1]$. Now, suppose that $\gamma(t_1), \gamma(t_2) \in \text{dom}(f)$. So, Lemma 3 implies that $\gamma([t_1, t_2]) \subseteq \text{dom}(f)$. Thus, for all $t \in (t_1, t_2)$ we obtain from Corollary 5 that there exist $s_1 \in (t_1, t)$ and $s_2 \in (t, t_2)$ such that

$$f(\gamma(t)) - f(\gamma(t_1)) \leq (t - t_1)f'(\gamma(s_1), \gamma'(s_1)), \quad (t_2 - t)f'(\gamma(s_2), \gamma'(s_2)) \leq (f(\gamma(t_2)) - f(\gamma(t))).$$

Thus, letting $t = (1 - \lambda)t_1 + \lambda t_2$ in above equation we obtain from monotonicity of f' that

$$\begin{aligned} (1 - \lambda)f(\gamma(t_1)) + \lambda f(\gamma(t_2)) - f(\gamma(t)) &= -(1 - \lambda)[f(\gamma(t)) - f(\gamma(t_1))] + \lambda[f(\gamma(t_2)) - f(\gamma(t))] \\ &\geq (1 - \lambda)\lambda(t_2 - t_1)[-f'(\gamma(s_1), \gamma'(s_1)) + f'(\gamma(s_2), \gamma'(s_2))] \\ &\geq 0, \end{aligned}$$

for all $\lambda \in [0, 1]$. Which implies that $f \circ \gamma((1 - \lambda)t_1 + \lambda t_2) \leq (1 - \lambda)f(\gamma(t_1)) + \lambda f(\gamma(t_2))$, for all $\lambda \in [0, 1]$. Therefore f is convex. \square

5 Sufficient Optimality Conditions for Optimization Problem

In this section we are going to obtain a sufficient optimality conditions for constraint optimization problems in terms of the Dini derivative.

Proposition 4. Let $C \subset M$ be a convex set. Assume that $f : M \rightarrow \mathbb{R}$ is convex in C . Consider the following nonlinear programming problem

$$(P) \quad \begin{cases} \min & f(p) \\ \text{s.t.} & p \in C. \end{cases}$$

Let $p^* \in C$. If for all $p \in C$ we have that $f'(p^*, \gamma'_{p^*p}(0)) \geq 0$, where γ_{p^*p} is a minimal geodesic from p^* to p with $\gamma'_{p^*p}(0) = p^*$ and $\gamma'_{p^*p}(1) = p$. Then $p^* \in C$ is a solution to (P) .

Proof. Given $p, p^* \in C$ and γ_{p^*p} a minimal geodesic from p^* to p with $\gamma'_{p^*p}(0) = p^*$ and $\gamma'_{p^*p}(1) = p$. Since C is convex we have that $\gamma_{p^*p}([0, 1]) \subset C$. Now, as f is convex we conclude from Corollary 4 that

$$f(p) \geq f(p^*) + f'(p^*, \gamma'_{p^*p}(0)).$$

Because $f'(p^*, \gamma'_{p^*p}(0)) \geq 0$, latter inequality implies that $f(p) \geq f(p^*)$. So, p^* is a solution to (P) . \square

Corollary 6. Let $f, g_i : M \rightarrow \mathbb{R}$ be convex, for $i = 1, \dots, m$. Consider the following nonlinear programming problem

$$(\tilde{P}) \quad \begin{cases} \min & f(p) \\ \text{s.t.} & g_i(p) \leq 0, \quad i = 1, \dots, m. \end{cases}$$

Let p^* be a feasible point to (\tilde{P}) . If for all p , a feasible point to (\tilde{P}) , there exist a vector $\mu = (\mu_1, \dots, \mu_m) \in \mathbb{R}^m$ such that

$$f'(x^*, \gamma'_{p^*p}(t_*)) + \sum_{i=1}^m \mu_i g'_i(x^*, \gamma'_{p^*p}(t_*)) \geq 0, \quad \mu \geq 0, \quad \text{and} \quad \sum_{i=1}^m \mu_i g_i(x^*) = 0, \quad (7)$$

where γ_{p^*p} is a minimal geodesic from p^* to p with $\gamma'_{p^*p}(0) = p^*$ and $\gamma'_{p^*p}(1) = p$. Then p^* is a solution to (\tilde{P}) .

Proof. Since $f, g_i : M \rightarrow \mathbb{R}$ are convex, for $i = 1, \dots, m$, and $\mu \geq 0$ we conclude that $C := \{p \in M : g_i(p) \leq 0, \quad i = 1, \dots, m\}$ is convex and $h : M \rightarrow \mathbb{R}$ defined by $h(p) = f(p) + \sum_{i=1}^m \mu_i g_i(p)$ is also convex. Moreover,

$$f(p) \geq h(p), \quad \text{for all } p \in C. \quad (8)$$

Take $p \in C$ and γ_{p^*p} a minimal geodesic from p^* to p with $\gamma'_{p^*p}(0) = p^*$ and $\gamma'_{p^*p}(1) = p$. From the first inequality in (7) we obtain that $h'(p^*, \gamma'_{p^*p}(0)) \geq 0$ and as $p^* \in C$ it follows from Proposition 4 that p^* satisfies $h(p) \geq h(p^*)$, for all $p \in C$. Thus, from (8) and the equality in (7) we obtain that

$$f(p) \geq h(p) \geq h(p^*) = f(p^*),$$

for all $p \in C$, and the proposition is proved. \square

6 Final Remarks

A complete, simply connected Riemannian manifold of nonpositive sectional curvature is called a *Hadamard manifold*, see Example 5 above. Hadamard-Cartan Theorem [19] asserts that the topological and differential structure of a Hadamard manifolds coincide with those of an Euclidean space of the same dimension. More precisely, at any point $p \in M$, the exponential map $\exp_p : T_p M \rightarrow M$ is a diffeomorphism. Furthermore, for any two points $p, q \in M$ there exists a unique geodesic joining p to q , which is minimal. As a consequence, all convex sets are strongly convex. So, the Definition 6 becomes

Definition 7. Let $f \in \mathcal{F}$ be a locally Lipschitz function on $\text{dom}(f)$. The lower Dini derivative $f' : TM \rightarrow [-\infty, +\infty]$ is said to be monotone if

$$f'(p, \exp_p^{-1}q) + f'(q, \exp_q^{-1}p) \leq 0,$$

for any two points $p, q \in M$.

Since we have obtained results only for locally Lipschitz functions defined on finite dimensional Riemannian manifolds, we expect that the results in this paper will be one more step toward a characterization for Lipschitz and convex functions on more general settings.

References

- [1] D. Azagra, J. Ferrera and F. López-Mesas, Nonsmooth analysis and Hamilton-Jacobi equations on Riemannian manifolds, *Journal of Functional Analysis* **220** (2), 304-361 (2005).
- [2] D. Azagra and J. Ferrera, Proximal Calculus on Riemannian Manifolds, *Mediterranean Journal of Mathematics*, Vol.**2**(4). pp. 437 - 450 (2005).
- [3] F. H. Clarke, Yu. S. Ledyaev, R. J. Stern and P.R. Wolenski, *Nonsmooth Analysis and Control Theory*, Springer-Verlag, (1998).
- [4] F. H. Clarke, Yu. S. Ledyaev and P.R. Wolenski, Proximal Analysis and Minimization Principles, *Journal of Mathematical Analysis and Applications* **196**, pp.722-735 (1995).
- [5] F. H. Clarke, R. J. Stern and P.R. Wolenski, Subgradient Criteria for Monotonicity, the Lipschitz Condition, and Convexity, *Canadian Journal of Mathematics*, Vol.**45**(6), pp. 1167-1183 (1993).
- [6] R. Correa, A. Jofré and L. Thibault, Characterization of Lower Semicontinuous Convex Functions, *Proceedings of the American Mathematical Society*, Vol.**116**(1), pp. 67-72 (1992).
- [7] O. P. Ferreira, Proximal subgradient and a characterization of Lipschitz Lipschitz function on Riemannian manifolds, *Journal of Mathematical Analysis and Applications* **313**, pp.587-597 (2006).
- [8] J.-B. Hiriart-Urruty and C. Lemaréchal, *Convex Analysis and Minimization Algorithms I and II*, Springer-Verlag, 1993.
- [9] D. Jiang, J. B. Moore and H. Ji, Self-Concordant functions for optimization on smooth manifolds, *43rd IEEE Conference on Decision and Control*, December 14-17, 2004 Atlantis, Paradise Island, Bahamas, (2004).
- [10] J. Jost, Convex functionals and generalized harmonic maps into spaces of non positive curvature, *Commentarii Mathematici Helvetici* Vol.**70**, pp. 659-673 (1995).
- [11] Yu. S. Ledyaev and Q. J. Zhu, Nonsmooth analysis on smooth manifold, to appear in *Transactions of the American Mathematical Society* (2006).

- [12] Yu. S. Ledyayev and Q. J. Zhu, Techniques for nonsmooth analysis on smooth manifolds. II. *Deformations and flows*, in *Optimal control, stabilization and nonsmooth analysis, Lecture Notes in Control and Inform. Sci.*, Vol. **301**, Springer, Berlin, pp. 299–311 (2004).
- [13] Yu. S. Ledyayev and Q. J. Zhu, Techniques for nonsmooth analysis on smooth manifolds. I. *Local problems*, *Optimal control, stabilization and nonsmooth analysis, Lecture Notes in Control and Inform. Sci.*, Vol. **301**, Springer, Berlin, pp. 283–297 (2004).
- [14] D. T. Luc and S. Swaminathan, A Characterization of Convex Functions, *Nonlinear Analysis: Theory, Methods & Applications*, Vol. **20**(6), pp. 697–701 (1993).
- [15] S. A. Miller and J. Malick, Newton methods for nonsmooth convex minimization: connections among -Lagrangian, Riemannian Newton and SQP methods *Mathematical Programming*, Vol. **104**(2-3), pp. 609–633 (2005).
- [16] T. Rapcsák, *Smooth Nonlinear Optimization in R^n* , Kluwer Academic Publishers, (1997).
- [17] S. T. Smith, *Optimization techniques on Riemannian Manifolds*, Vol. 3 of Fields Institute Communications, American Mathematical Society, Providence, R.I., pp. 113–146, (1994).
- [18] R. A. Poliquin, Subgradient Monotonicity and Convex Functions, *Nonlinear Analysis: Theory, Methods & Applications*, Vol. **14**(4), pp. 305–317 (1990).
- [19] T. Sakai, *Riemannian Geometry*, Translations of Mathematical Monographs, vol. **149**, American Mathematical Society, Providence, R.I. (1996).
- [20] C. Udriste, Convex Functions and Optimization Methods on Riemannian Manifolds, *Mathematics and its Applications*, Vol. **297**, Kluwer Academic Publishers (1994).