

Multivariate exponential integral approximations: a moment approach

Dimitris Bertsimas ^{*} Xuan Vinh Doan [†] Jean Lasserre [‡]

January 2007

Abstract

We propose a method to approximate a class of exponential multivariate integrals using moment relaxations. Using this approach, both lower and upper bounds of the integrals are obtained and we show that these bound values asymptotically converge to the real value of the integrals when the moment degree r increases. We further demonstrate the method by calculating both hypercubic and order statistic probabilities for multivariate normal distributions.

1 Introduction

Multivariate integrals arise in statistic, physics, engineering and finance applications among other areas. For example, these integrals are needed to calculate probabilities over compact sets for multivariate normal random variables. It is therefore important to compute or approximate multivariate integrals. Usual methods include Monte Carlo schemes (see Niederreiter [7] for details) and cubature formulae as shown in e.g. de la Harpe and Pache [2]. However, there are still many open problems currently and research on multivariate integrals is very much active due to its importance as well as its difficulties. For instance, most cubature formulas are restricted to special sets like boxes and simplices, and even in this particular context, determination of orthogonal polynomials used to construct a cubature, is not an easy task.

^{*}Boeing Professor of Operations Research, Sloan School of Management, co-director of the Operations Research Center, Massachusetts Institute of Technology, E40-147, Cambridge, MA 02139-4307, dbertsim@mit.edu.

[†]Operations Research Center, Massachusetts Institute of Technology, Cambridge, MA 02139-4307, vanxuan@mit.edu.

[‡]LAAS, 7 Avenue Du Colonel Roche, 31077 Toulouse Cédex 4, France, lasserre@laas.fr

Contributions and Paper Outline

In this paper, we attempt to approximate a class of exponential integrals, which in particular can be useful to calculate probabilities of multivariate normal random variables. Specifically, our contribution and structure of the paper are as follows:

- (1) In Section 2, we provide a general framework to calculate lower and upper bounds for the exponential integrals mentioned above. These bounds are calculated by solving specific semidefinite programming problems constructed from appropriate sequences of moments.
- (2) In Section 3, we prove that the two monotone sequences of lower and upper bounds generated by these semidefinite programming problems will asymptotically converge to the real value of the integral. The proof is due to some results from the problem of moments.
- (3) In Section 4, computational results are reported for probabilities of multivariate random variables and their order statistics over a hypercube. These results show that the proposed method is indeed applicable for this class of integrals.

2 General Framework

Before we provide a detailed analysis for general multivariate exponential integrals, and for clarity of exposition, we briefly describe the approach for simple integrals in two variables x and y on a box $[a, b] \times [c, d]$. Indeed, the multivariate case $n \geq 3$ essentially uses the same machinery but its detailed exposition is a little tedious as it requires unavoidable heavy notation, whereas the bivariate case on a box illustrates the approach with minimal effort.

2.1 Simple Integrals

Suppose that one wants to approximate:

$$\rho = \int_a^b \int_c^d g(x, y) e^{h(x, y)} dy dx, \quad (1)$$

where g, h are bivariate polynomials.

Define $\Omega = [a, b] \times [c, d] \in \mathbb{R}^2$ and let us consider the measure μ on \mathbb{R}^2 defined by

$$\mu(B) = \int_{\Omega \cap B} e^{h(x, y)} dy dx, \quad (2)$$

where $B \in \mathcal{B}(\mathbb{R}^2)$, and its sequence of moments $\mathbf{z} = \{z(\alpha, \beta)\}$:

$$z(\alpha, \beta) = \int_{\Omega} x^{\alpha} y^{\beta} e^{h(x,y)} dy dx \quad (3)$$

for all $(\alpha, \beta) \in \mathbb{N}^2$. Clearly, $\rho = \sum_{(\alpha, \beta) \in \mathbb{N}^2} g_{(\alpha, \beta)} z(\alpha, \beta) = \langle \mathbf{g}, \mathbf{z} \rangle$, where $g_{(\alpha, \beta)}$ is the coefficient of the monomial $x^{\alpha} y^{\beta}$ and \mathbf{g} is the sequence of those coefficients. Therefore, if we have all necessary moments $z(\alpha, \beta)$, we can compute ρ . Using integration by parts, we have:

$$z(\alpha, \beta) = \frac{1}{\beta + 1} \int_a^b x^{\alpha} \left[y^{\beta+1} e^{h(x,y)} \right]_{y=c}^{y=d} dx - \frac{1}{\beta + 1} \int_a^b \int_c^d x^{\alpha} y^{\beta+1} \frac{\partial h(x,y)}{\partial y} e^{h(x,y)} dy dx.$$

If $\{h_{(\gamma, \delta)}\}$ is the finite sequence of coefficients of the polynomial $h(x, y)$ then we have:

$$z(\alpha, \beta) = \frac{d^{\beta+1}}{\beta + 1} \int_a^b x^{\alpha} e^{h(x,d)} dx - \frac{c^{\beta+1}}{\beta + 1} \int_a^b x^{\alpha} e^{h(x,c)} dx - \sum_{(\gamma, \delta) \in \mathbb{N}^2} \frac{\delta h_{(\gamma, \delta)}}{\beta + 1} z(\alpha + \gamma, \beta + \delta).$$

If we define two measures ν and η on \mathbb{R} with their corresponding sequences of moments:

$$v(\alpha) = \int_a^b x^{\alpha} e^{h(x,d)} dx, \quad w(\alpha) = \int_a^b x^{\alpha} e^{h(x,c)} dx, \quad (4)$$

for all $\alpha \in \mathbb{N}$ respectively, then we have:

$$z(\alpha, \beta) = \frac{d^{\beta+1}}{\beta + 1} v(\alpha) - \frac{c^{\beta+1}}{\beta + 1} w(\alpha) - \sum_{(\gamma, \delta) \in \mathbb{N}^2} \frac{\delta h_{(\gamma, \delta)}}{\beta + 1} z(\alpha + \gamma, \beta + \delta). \quad (5)$$

Let us define $k(x) = h(x, d)$ and $l(x) = h(x, c)$. Clearly, $k(x)$ and $l(x)$ are univariate polynomials in x .

Using integration by parts again for $v(\alpha)$, we have:

$$v(\alpha) = \frac{1}{\alpha + 1} \left[x^{\alpha+1} e^{k(x)} \right]_{x=a}^{x=b} - \frac{1}{\alpha + 1} \int_a^b x^{\alpha+1} \frac{dk(x)}{dx} e^{k(x)} dx,$$

or

$$v(\alpha) = \frac{b^{\alpha+1} e^{k(b)}}{\alpha + 1} - \frac{a^{\alpha+1} e^{k(a)}}{\alpha + 1} - \sum_{\epsilon \in \mathbb{N}} \frac{\epsilon k_{\epsilon}}{\alpha + 1} v(\alpha + \epsilon), \quad (6)$$

where $\{k_{\epsilon}\}$ is the finite sequence of coefficients of the polynomial $x \mapsto k(x) = \sum_{\epsilon} k_{\epsilon} x^{\epsilon}$, of degree k_x .

Similarly, we have:

$$w(\alpha) = \frac{b^{\alpha+1} e^{l(b)}}{\alpha + 1} - \frac{a^{\alpha+1} e^{l(a)}}{\alpha + 1} - \sum_{\epsilon \in \mathbb{N}} \frac{\epsilon l_{\epsilon}}{\alpha + 1} w(\alpha + \epsilon), \quad (7)$$

where $\{l_{\epsilon}\}$ is the finite sequence of coefficients of the polynomial $x \mapsto l(x) = \sum_{\epsilon} l_{\epsilon} x^{\epsilon}$, of degree l_x .

Also it is clear that in view of (6) and (7), all $v(\alpha)$ and $w(\alpha)$ are affine functions of v_0, \dots, v_{k_x-1} , and w_0, \dots, w_{l_x-1} , respectively.

Methodology: Notice that in (1), the quantity ρ to approximate, is a linear combination $\langle \mathbf{g}, \mathbf{z} \rangle$ of moments of μ . Then, to compute upper and lower bounds on ρ , we build up *two* hierarchies of semidefinite programs (SDP) \mathcal{P}_r^u and \mathcal{P}_r^l , as follows. Consider the vectors of moments \mathbf{v}, \mathbf{w} and \mathbf{z} , up to order $2r$, $r \in \mathbb{N}$.

- The linear constraints of both \mathcal{P}_r^u and \mathcal{P}_r^l are coming from those equations (5), (6), and (7), that contain only moments of order up to $2r$.
- The LMI (linear matrix inequalities) constraints of both \mathcal{P}_r^u and \mathcal{P}_r^l (to be defined later), state necessary conditions on \mathbf{v}, \mathbf{w} and \mathbf{z} , to be *moments* of some measures supported on $[a, b]$, and $[a, b] \times [c, d]$, respectively.

Then the SDP \mathcal{P}_r^u (resp. \mathcal{P}_r^l) maximizes (resp. minimizes) the linear criterion $\langle \mathbf{g}, \mathbf{z} \rangle$ under the above constraints. Both are *SDP-relaxations* of the original problem (1), and so their respective optimal values provide upper and lower bounds on ρ . In addition, the quality of both bounds increases with r because more and more constraints (5), (6), and (7), are taken into account.

We will show in the general case $n > 2$ that the resulting monotone nonincreasing and nondecreasing sequences of upper and lower bounds both converge to the desired value ρ .

2.2 General Integrals

We now consider the class of more general multivariate exponential integrals:

$$\rho = \int_{\Omega} g(\mathbf{x}) e^{h(\mathbf{x})} d\mathbf{x}, \quad (8)$$

where $\mathbf{x} \in \mathbb{R}^n$, $g, h \in \mathbb{R}[\mathbf{x}]$, the ring of real polynomials and $\Omega \subset \mathbb{R}^n$ is a compact set defined as

$$\Omega = \{ \mathbf{x} \in \mathbb{R}^n : b_1^{(1)} \leq x_1 \leq b_1^{(2)}, b_i^{(1)}(\mathbf{x}[i-1]) \leq x_i \leq b_i^{(2)}(\mathbf{x}[i-1]) \quad \forall i = 2, \dots, n \} \quad (9)$$

where $\mathbf{x}[i] \in \mathbb{R}^i$ is the vector of first i elements of \mathbf{x} , $i = 1, \dots, n$, $b_i^{(1)}, b_i^{(2)} \in \mathbb{R}[\mathbf{x}[i-1]]$, $i = 2, \dots, n$, and $b_1^{(1)}, b_1^{(2)}$ are constants. Similar to the previous section, we can now analyze the general case as follows.

Define $\Omega_k = \{ \mathbf{x} \in \mathbb{R}^k : b_1^{(1)} \leq x_1 \leq b_1^{(2)}, b_i^{(1)}(\mathbf{x}[i-1]) \leq x_i \leq b_i^{(2)}(\mathbf{x}[i-1]) \quad \forall i = 2, \dots, k \}$, $k = 1, \dots, n$, we have: $\Omega_k \in \mathbb{R}^k$ and $\Omega_n = \Omega$. Clearly, these Ω_k sets are also compact in \mathbb{R}^k for all $k = 1, \dots, n$. Let us consider the measure $\mu_h^{(i)}$ on \mathbb{R}^i defined by

$$\mu_h^{(i)}(B) = \int_{\Omega_i \cap B} e^{h(\mathbf{x}[i])} d\mathbf{x}[i], \quad (10)$$

where $h \in \mathbb{R}[\mathbf{x}[i]]$, $B \in \mathcal{B}(\mathbb{R}^i)$, and its sequence of moments $\mathbf{z}_h^{(i)} = \{z_h^{(i)}(\boldsymbol{\alpha})\}$:

$$z_h^{(i)}(\boldsymbol{\alpha}) = \int_{\Omega_i} (\mathbf{x}[i])^\alpha e^{h(\mathbf{x}[i])} d\mathbf{x}[i], \quad (11)$$

for all $\boldsymbol{\alpha} \in \mathbb{N}^i$.

We then have $\rho = \sum_{\boldsymbol{\alpha} \in \mathbb{N}^n} g_\alpha z_h^{(n)}(\boldsymbol{\alpha}) = \langle \mathbf{g}, \mathbf{z}_h^{(n)} \rangle$, where g_α is the coefficient of monomial \mathbf{x}^α and \mathbf{g} is the sequence of those coefficients. Therefore, what we need to do is to calculate $z_h^{(n)}(\boldsymbol{\alpha})$ for all necessary $\boldsymbol{\alpha} \in \mathbb{N}^i$. Using integration by parts, we have:

$$z_h^{(n)}(\boldsymbol{\alpha}) = \frac{1}{\alpha_n + 1} (A_\alpha - B_\alpha),$$

where

$$A_\alpha = \int_{\Omega_{n-1}} (\mathbf{x}[n-1])^{\alpha[n-1]} \left[x_n^{\alpha_n+1} e^{h(\mathbf{x})} \right]_{b_n^{(1)}(\mathbf{x}[n-1])}^{b_n^{(2)}(\mathbf{x}[n-1])} d\mathbf{x}[n-1],$$

and

$$B_\alpha = \int_{\Omega} (\mathbf{x}[n-1])^{\alpha[n-1]} x_n^{\alpha_n+1} \frac{\partial h(\mathbf{x})}{\partial x_n} e^{h(\mathbf{x})} d\mathbf{x}.$$

We have:

$$\frac{\partial h(\mathbf{x})}{\partial x_n} = \sum_{\boldsymbol{\beta} \in \mathbb{N}^n} \beta_n h_\beta (\mathbf{x}[n-1])^{\beta[n-1]} x_n^{\beta_n-1}.$$

Therefore,

$$B_\alpha = \sum_{\boldsymbol{\beta} \in \mathbb{N}^n} \beta_n h_\beta \int_{\Omega} (\mathbf{x}[n-1])^{\alpha[n-1]} x_n^{\alpha_n+1} (\mathbf{x}[n-1])^{\beta[n-1]} x_n^{\beta_n-1} e^{h(\mathbf{x})} d\mathbf{x}$$

or

$$B_\alpha = \sum_{\boldsymbol{\beta} \in \mathbb{N}^n} \beta_n h_\beta \int_{\Omega} \mathbf{x}^{\alpha+\beta} e^{h(\mathbf{x})} d\mathbf{x} = \sum_{\boldsymbol{\beta} \in \mathbb{N}^n} \beta_n h_\beta z_h^{(n)}(\boldsymbol{\alpha} + \boldsymbol{\beta}).$$

Now consider A_α , we have: $A_\alpha = A_\alpha^2 - A_\alpha^1$ where

$$A_\alpha^i = \int_{\Omega_{n-1}} (\mathbf{x}[n-1])^{\alpha[n-1]} [b_n^{(i)}(\mathbf{x}[n-1])]^{\alpha_n+1} e^{h(\mathbf{x}[n-1], b_n^{(i)}(\mathbf{x}[n-1]))} d\mathbf{x}[n-1], \quad i = 1, 2.$$

Let

$$g_{i,\alpha}^{(n-1)}(\mathbf{x}) = \mathbf{x}^{\alpha[n-1]} [b_n^{(i)}(\mathbf{x})]^{\alpha_n+1}, \quad \mathbf{x} \in \mathbb{R}^{n-1}, \quad i = 1, 2,$$

and

$$h_i^{(n-1)}(\mathbf{x}) = h(\mathbf{x}, b_n^{(i)}(\mathbf{x})), \quad \mathbf{x} \in \mathbb{R}^{n-1}, \quad i = 1, 2.$$

All of these four functions are polynomials in $\mathbb{R}[\mathbf{x}[n-1]]$. Define two measures $\mu_1^{(n-1)} \equiv \mu_{h_1^{(n-1)}}^{(n-1)}$ and $\mu_2^{(n-1)} \equiv \mu_{h_2^{(n-1)}}^{(n-1)}$ over Ω_{n-1} on \mathbb{R}^{n-1} as defined in (10). With their two respective sequences of moments $\mathbf{z}_1^{(n-1)} \equiv \mathbf{z}_{h_1^{(n-1)}}^{(n-1)}$ and $\mathbf{z}_2^{(n-1)} \equiv \mathbf{z}_{h_2^{(n-1)}}^{(n-1)}$ as defined in (11), we have:

$$A_\alpha = \sum_{\boldsymbol{\beta} \in \mathbb{N}^{n-1}} \left(g_{2,\alpha}^{(n-1)} \right)_\beta z_2^{(n-1)}(\boldsymbol{\beta}) - \sum_{\boldsymbol{\beta} \in \mathbb{N}^{n-1}} \left(g_{1,\alpha}^{(n-1)} \right)_\beta z_1^{(n-1)}(\boldsymbol{\beta}).$$

Thus we have:

$$z_h^{(n)}(\boldsymbol{\alpha}) = \sum_{\boldsymbol{\beta} \in \mathbb{N}^{n-1}} \frac{\binom{g_{2,\boldsymbol{\alpha}}^{(n-1)}}{\boldsymbol{\beta}}}{\alpha_n + 1} z_2^{(n-1)}(\boldsymbol{\beta}) - \sum_{\boldsymbol{\beta} \in \mathbb{N}^{n-1}} \frac{\binom{g_{1,\boldsymbol{\alpha}}^{(n-1)}}{\boldsymbol{\beta}}}{\alpha_n + 1} z_1^{(n-1)}(\boldsymbol{\beta}) - \sum_{\boldsymbol{\beta} \in \mathbb{N}^n} \frac{\beta_n h_{\boldsymbol{\beta}}}{\alpha_n + 1} z_h^{(n)}(\boldsymbol{\alpha} + \boldsymbol{\beta}). \quad (12)$$

Equation (12) shows that in order to calculate $z_h^{(n)}$, we need to calculate $z_i^{(n-1)}$, $i = 1, 2$, which can then be calculated by some other moment sequences in lower dimensions. Let denote $h_1^{(n)}$ to be the function h and define

$$h_i^{(k)}(\mathbf{x}) = h_{\lfloor i/2 \rfloor}^{(k+1)}(\mathbf{x}, b_{\lfloor i/2 \rfloor + 1}^{(2 - \lfloor i/2 \rfloor + \lfloor i/2 \rfloor)}(\mathbf{x})), \quad \mathbf{x} \in \mathbb{R}^k, \quad k = 1, \dots, n-1, \quad i = 1, \dots, 2^{n-k}. \quad (13)$$

For each function $h_i^{(k)}$, a measure $\mu_i^{(k)}$ and its moments sequence $z_i^{(k)}$ are also defined over Ω_k in \mathbb{R}^k as in (10) and (11) respectively. We also need to define the function $g_{i,\boldsymbol{\alpha}}^{(k)}$. Let define

$$g_{i,\boldsymbol{\alpha}}^{(k)}(\mathbf{x}) = \mathbf{x}^{\boldsymbol{\alpha}[k]} [b_{k+1}^{(i)}(\mathbf{x})]^{\alpha_{k+1}+1}, \quad \mathbf{x} \in \mathbb{R}^k, \quad \boldsymbol{\alpha} \in \mathbb{N}^k, \quad i = 1, 2. \quad (14)$$

Then we have:

$$z_i^{(k)}(\boldsymbol{\alpha}) = \sum_{\boldsymbol{\beta} \in \mathbb{N}^{k-1}} \frac{\binom{g_{2,\boldsymbol{\alpha}}^{(k-1)}}{\boldsymbol{\beta}}}{\alpha_k + 1} z_{2i}^{(k-1)}(\boldsymbol{\beta}) - \sum_{\boldsymbol{\beta} \in \mathbb{N}^{k-1}} \frac{\binom{g_{1,\boldsymbol{\alpha}}^{(k-1)}}{\boldsymbol{\beta}}}{\alpha_k + 1} z_{2i-1}^{(k-1)}(\boldsymbol{\beta}) - \sum_{\boldsymbol{\beta} \in \mathbb{N}^k} \frac{\beta_k (h_i^{(k)})_{\boldsymbol{\beta}}}{\alpha_k + 1} z_i^{(k)}(\boldsymbol{\alpha} + \boldsymbol{\beta}) \quad (15)$$

or $z_i^{(k)}(\boldsymbol{\alpha}) = R_i^{(k)}(z_i^{(k)}, z_{2i}^{(k-1)}, z_{2i-1}^{(k-1)}, \boldsymbol{\alpha})$ for all $\boldsymbol{\alpha} \in \mathbb{R}^k$, $k = 2, \dots, n$, and $i = 1, \dots, 2^{n-k}$.

For $k = 1$ and $i = 1, \dots, 2^{n-1}$, we have:

$$z_i^{(1)}(\alpha) = \frac{(b_1^{(2)})^{\alpha+1}}{\alpha + 1} e^{h_i^{(1)}(b_1^{(2)})} - \frac{(b_1^{(1)})^{\alpha+1}}{\alpha + 1} e^{h_i^{(1)}(b_1^{(1)})} - \sum_{\beta \in \mathbb{N}} \frac{\beta (h_i^{(1)})_{\beta}}{\alpha + 1} z_i^{(1)}(\alpha + \beta) = R_i^{(1)}(z_i^{(1)}, \alpha). \quad (16)$$

With the general recursive formula presented in (15) and (16), the sequence of moments $z_h^{(n)}$ (or $z_1^{(n)}$) can now be calculated.

2.3 Moment Relaxation

Let consider the measure $\mu_i^{(k)}$ and its corresponding sequence of moments $z_i^{(k)}$. For any nonnegative integer r , the r -moment matrix associated with $\mu_i^{(k)}$ (or equivalently, with $z_i^{(k)}$) $M_r(\mu_i^{(k)}) \equiv M_r(z_i^{(k)})$ is a matrix of size $\binom{k+r}{r}$. Its rows and columns are indexed in the canonical basis $\{(\mathbf{x}[k])^{\boldsymbol{\alpha}}\}$ of $\mathbb{R}[\mathbf{x}[k]]$, and its elements are defined as follows:

$$M_r(z_i^{(k)})(\boldsymbol{\alpha}, \boldsymbol{\beta}) = z_i^{(k)}(\boldsymbol{\alpha} + \boldsymbol{\beta}), \quad \boldsymbol{\alpha}, \boldsymbol{\beta} \in \mathbb{N}^k, |\boldsymbol{\alpha}|, |\boldsymbol{\beta}| \leq r. \quad (17)$$

Similarly, given $\theta \in \mathbb{R}[\mathbf{x}[k]]$, the localizing matrix $M_r(\boldsymbol{\theta} \mathbf{z}_i^{(k)})$ associated with $\mathbf{z}_i^{(k)}$ and θ is defined by

$$M_r(\boldsymbol{\theta} \mathbf{z}_i^{(k)})(\boldsymbol{\alpha}, \boldsymbol{\beta}) := \sum_{\boldsymbol{\gamma} \in \mathbb{N}^k} \theta_{\boldsymbol{\gamma}} \mathbf{z}_i^{(k)}(\boldsymbol{\alpha} + \boldsymbol{\beta} + \boldsymbol{\gamma}), \quad \boldsymbol{\alpha}, \boldsymbol{\beta} \in \mathbb{N}^k, |\boldsymbol{\alpha}|, |\boldsymbol{\beta}| \leq r, \quad (18)$$

where $\boldsymbol{\theta} = \{\theta_{\boldsymbol{\gamma}}\}$ is the vector of coefficients of θ in the canonical basis $\{(\mathbf{x}[k])^{\boldsymbol{\alpha}}\}$.

If we define the matrix $M_r^{\boldsymbol{\gamma}}(\mathbf{z}_i^{(k)})$ with elements

$$M_r^{\boldsymbol{\gamma}}(\mathbf{z}_i^{(k)})(\boldsymbol{\alpha}, \boldsymbol{\beta}) = \mathbf{z}_i^{(k)}(\boldsymbol{\alpha} + \boldsymbol{\beta} + \boldsymbol{\gamma}), \quad \boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma} \in \mathbb{N}^k, |\boldsymbol{\alpha}|, |\boldsymbol{\beta}| \leq r,$$

then the localizing matrix can be expressed as $M_r(\boldsymbol{\theta} \mathbf{z}_i^{(k)}) = \sum_{\boldsymbol{\gamma} \in \mathbb{N}^k} \theta_{\boldsymbol{\gamma}} M_r^{\boldsymbol{\gamma}}(\mathbf{z}_i^{(k)})$.

Note that for every polynomial $f \in \mathbb{R}[\mathbf{x}[k]]$ of degree at most r with its vector of coefficients denoted by $\mathbf{f} = \{f_{\boldsymbol{\gamma}}\}$, we have:

$$\langle \mathbf{f}, M_r(\mathbf{z}_i^{(k)}) \mathbf{f} \rangle = \int f^2 d\mu_i^{(k)}, \quad \langle \mathbf{f}, M_r(\boldsymbol{\theta} \mathbf{z}_i^{(k)}) \mathbf{f} \rangle = \int \theta f^2 d\mu_i^{(k)}. \quad (19)$$

This property shows that necessarily, $M_r(\mathbf{z}_i^{(k)}) \succeq \mathbf{0}$ and $M_r(\boldsymbol{\theta} \mathbf{z}_i^{(k)}) \succeq \mathbf{0}$, whenever $\mu_i^{(k)}$ has its support contained in the level set $\{\mathbf{x} \in \mathbb{R}^k : \theta(\mathbf{x}) \geq 0\}$. If the sequence of moments is restricted to those moments used to construct the moment matrix $M_r(\mathbf{z}_i^{(k)})$ (up to moments of degree $2r$), then the second necessary condition is reduced to $M_{r-\lceil d/2 \rceil}(\boldsymbol{\theta} \mathbf{z}_i^{(k)}) \succeq \mathbf{0}$, where d is the degree of the polynomial θ . For more details on moment matrices, local matrices, and these necessary conditions, please refer to Laurent [6] and references therein.

Let us define

$$\theta_i^{(k)}(\mathbf{x}) = \left[b_i^{(2)}(\mathbf{x}[i-1]) - x_i \right] \left[x_i - b_i^{(1)}(\mathbf{x}[i-1]) \right], \quad \mathbf{x} \in \mathbb{R}^k, \quad \forall k \geq i, \quad (20)$$

for all $i = 2, \dots, n$. Similarly, define

$$\theta_1^{(k)}(\mathbf{x}) = \left[b_1^{(2)} - x_1 \right] \left[x_1 - b_1^{(1)} \right], \quad \mathbf{x} \in \mathbb{R}^k, \quad \forall k \geq 1. \quad (21)$$

For a fixed i ($i = 1, \dots, n$), the polynomials $\theta_i^{(k)}$ depends on the first i variables x_1, \dots, x_i exactly the same for all $k \geq i$. Thus they all have the same degree d_i for all $k \geq i$.

We also have $\Omega_k = \{\mathbf{x} \in \mathbb{R}^k : \theta_i^{(k)}(\mathbf{x}) \geq 0 \quad \forall i = 1, \dots, k\}$ for all $k = 1, \dots, n$. Thus necessary conditions for moment matrices of all measures $\mu_i^{(k)}$ can be written as follows:

$$M_r(\mathbf{z}_i^{(k)}) \succeq \mathbf{0}, \quad M_{r-\lceil d_j/2 \rceil}(\theta_j^{(k)} \mathbf{z}_i^{(k)}) \succeq \mathbf{0}, \quad \forall k = 1, \dots, n, \quad \forall i = 1, \dots, 2^{n-k}, \quad \forall j = 1, \dots, k. \quad (22)$$

Combining these necessary conditions and the recursive formulae for $\mathbf{z}_i^{(k)}$ in (15) and (16), we can find lower and upper bounds for ρ by solving the two following semidefinite programming problems:

$$\mathcal{P}_r^l \triangleq \left[\begin{array}{l} \inf \quad \langle \mathbf{g}, \mathbf{z}_1^{(n)} \rangle \\ \text{s.t.} \quad M_r(\mathbf{z}_i^{(k)}) \succeq 0 \quad \quad \quad k = 1, \dots, n, \quad i = 1, \dots, 2^{n-k}, \\ \quad \quad M_{r-\lceil d_j/2 \rceil}(\theta_j^{(k)} \mathbf{z}_i^{(k)}) \succeq 0 \quad \quad k = 1, \dots, n, \quad i = 1, \dots, 2^{n-k}, \quad j = 1, \dots, k, \\ \quad \quad z_i^{(1)}(\alpha) = R_i^{(1)}(\mathbf{z}_i^{(1)}, \alpha) \quad \quad i = 1, \dots, 2^{n-1}, \quad \alpha \in A_{i,r}^{(1)}, \\ \quad \quad z_i^{(k)}(\alpha) = R_i^{(k)}(\mathbf{z}_i^{(k)}, \mathbf{z}_{2i}^{(k-1)}, \mathbf{z}_{2i-1}^{(k-1)}, \alpha) \quad k = 2, \dots, n, \quad i = 1, \dots, 2^{n-k}, \quad \alpha \in A_{i,r}^{(k)}, \end{array} \right] \quad (23)$$

and

$$\mathcal{P}_r^u \triangleq \left[\begin{array}{l} \sup \quad \langle \mathbf{g}, \mathbf{z}_1^{(n)} \rangle \\ \text{s.t.} \quad M_r(\mathbf{z}_i^{(k)}) \succeq 0 \quad \quad \quad k = 1, \dots, n, \quad i = 1, \dots, 2^{n-k}, \\ \quad \quad M_{r-\lceil d_j/2 \rceil}(\theta_j^{(k)} \mathbf{z}_i^{(k)}) \succeq 0 \quad \quad k = 1, \dots, n, \quad i = 1, \dots, 2^{n-k}, \quad j = 1, \dots, k, \\ \quad \quad z_i^{(1)}(\alpha) = R_i^{(1)}(\mathbf{z}_i^{(1)}, \alpha) \quad \quad i = 1, \dots, 2^{n-1}, \quad \alpha \in A_{i,r}^{(1)}, \\ \quad \quad z_i^{(k)}(\alpha) = R_i^{(k)}(\mathbf{z}_i^{(k)}, \mathbf{z}_{2i}^{(k-1)}, \mathbf{z}_{2i-1}^{(k-1)}, \alpha) \quad k = 2, \dots, n, \quad i = 1, \dots, 2^{n-k}, \quad \alpha \in A_{i,r}^{(k)}, \end{array} \right] \quad (24)$$

where $A_{i,r}^{(k)}$ is the set of all $\alpha \in \mathbb{R}^k$ that the recursive formulae can be expressed by moments of degree up to $2r$ (used to construct the moment matrices M_r).

Clearly, $Z(\mathcal{P}_r^l) \leq \rho \leq Z(\mathcal{P}_r^u)$ and we will prove that these lower and upper bounds asymptotically converge to ρ when r tends to infinity in the next section.

3 Convergence

In order to prove the convergence of $Z(\mathcal{P}_r^l)$ and $Z(\mathcal{P}_r^u)$, we need to prove the recursive formulae in (15) and (16) define moment sequences for all measures $\mu_i^{(k)}$.

Lemma 1 *Let $h_i^{(k)}$ and $g_{i,\alpha}^{(k)}$ be defined as in (13) and (14) respectively, and let $\mathbf{z}_i^{(k)}$ be the moment sequences of some Borel measures $\psi_i^{(k)}$ on the compact sets Ω_k , which satisfy (15) and (16). Then $d\psi_i^{(k)} = d\mu_i^{(k)}$ for all $k = 1, \dots, n$ and $i = 1, \dots, 2^{n-k}$.*

Proof. We will prove the lemma by induction. Let consider the case $k = 1$, we have, according to (16):

$$z_i^{(1)}(\alpha) = \int x^\alpha d\psi_i^{(1)} = \left[\frac{x^{\alpha+1}}{\alpha+1} e^{h_i^{(1)}(x)} \right]_{b_1^{(1)}}^{b_1^{(2)}} - \int \frac{x^{\alpha+1}}{\alpha+1} (h_i^{(1)})'(x) d\psi_i^{(1)}. \quad (25)$$

We also have:

$$\int x^\alpha d\mu_i^{(1)} = \left[\frac{x^{\alpha+1}}{\alpha+1} e^{h_i^{(1)}(x)} \right]_{b_1^{(1)}}^{b_1^{(2)}} - \int \frac{x^{\alpha+1}}{\alpha+1} (h_i^{(1)})'(x) d\mu_i^{(1)}. \quad (26)$$

Let consider the *signed* measure $\phi_i^{(1)}$ on Ω_1 that satisfies $d\phi_i^{(1)} = d\psi_i^{(1)} - d\mu_i^{(1)}$, from (25) and (26), we have:

$$\int x^\alpha d\phi_i^{(1)} = - \int \frac{x^{\alpha+1}}{\alpha+1} (h_i^{(1)})'(x) d\phi_i^{(1)}. \quad (27)$$

Consider the polynomial $p(x) = \sum_{j=1}^d f_j x^j$, we have: $p'(x) = \sum_{j=0}^{d-1} f_{j+1} (j+1) x^j$. From Equation (27), we obtain the following equation:

$$\int \left[p'(x) + p(x) (h_i^{(1)})'(x) \right] d\phi_i^{(1)} = 0. \quad (28)$$

We now prove that Equation (28) is also true for all continuous function $f(x) = xg(x)$, where g is continuously differentiable on Ω_1 . We have, polynomials are dense in the space of continuously differentiable functions on Ω_1 under the sup-norm $\max\{\sup_{x \in \Omega_1} |f(x)|, \sup_{x \in \Omega_1} |f'(x)|\}$ (see e.g. Coattméc [1] and Hirsch [3]) Therefore for any $\epsilon > 0$, there exist $p_\epsilon \in \mathbb{R}[x]$ such that $\sup_{x \in \Omega_1} |g(x) - p_\epsilon(x)| \leq \epsilon$ and $\sup_{x \in \Omega_1} |g'(x) - p'_\epsilon(x)| \leq \epsilon$. Equation (28) is true for the polynomial $p(x) = xp_\epsilon(x)$, thus

$$\int \left[f'(x) + f(x) (h_i^{(1)})'(x) \right] d\phi_i^{(1)} = \int [x(g(x) - p_\epsilon(x))]' d\phi_i^{(1)} + \int x [g(x) - p_\epsilon(x)] (h_i^{(1)})'(x) d\phi_i^{(1)}.$$

We have: $[x(g(x) - p_\epsilon(x))]' = (g(x) - p_\epsilon(x)) + x(g'(x) - p'_\epsilon(x))$, thus

$$|[x(g(x) - p_\epsilon(x))]'| \leq (1 + \sup_{x \in \Omega_1} |x|) \epsilon \quad \forall x \in \Omega_1.$$

Similarly,

$$|x [g(x) - p_\epsilon(x)] (h_i^{(1)})'(x)| \leq \sup_{x \in \Omega_1} |x (h_i^{(1)})'(x)| \epsilon \quad \forall x \in \Omega_1.$$

So we have:

$$\left| \int \left[f'(x) + f(x) (h_i^{(1)})'(x) \right] d\phi_i^{(1)} \right| \leq \epsilon (1 + \sup_{x \in \Omega_1} |x| + \sup_{x \in \Omega_1} |x (h_i^{(1)})'(x)|) \int |d\phi_i^{(1)}|.$$

The constant term $M = (1 + \sup_{x \in \Omega_1} |x| + \sup_{x \in \Omega_1} |x (h_i^{(1)})'(x)|) \int |d\phi_i^{(1)}|$ is finite and the above equation is true for all $\epsilon > 0$, thus we have for all $f(x) = xg(x)$:

$$\int \left[f'(x) + f(x) (h_i^{(1)})'(x) \right] d\phi_i^{(1)} = 0. \quad (29)$$

For an arbitrary polynomial $g(x) = \sum_{j=0}^d g_j x^j$, define $G(x) = \sum_{j=0}^d \frac{g_j}{j+1} x^{j+1}$, we have $G'(x) = g(x)$. Let consider $f(x) = G(x) e^{-h_i^{(1)}(x)}$, we have: $\frac{f(x)}{x}$ is a continuously differentiable function and $f'(x) = g(x) e^{-h_i^{(1)}(x)} - f(x) (h_i^{(1)})'(x)$. Using Equation (29), we obtain the following equation

$$\int g(x) e^{-h_i^{(1)}(x)} d\phi_i^{(1)} = 0, \quad \forall g \in \mathbb{R}[x]. \quad (30)$$

If we define $d\nu_i^{(1)} = e^{-h_i^{(1)}(x)} d\phi_i^{(1)}$, we then have $\int f(x) d\nu_i^{(1)} = 0$ for all continuous function f in Ω_1 since polynomials are dense in the space of continuous functions. This implies that $\nu_i^{(1)}$ is a zero measure. In addition, $e^{-h_i^{(1)}(x)} > 0$ for all $x \in \mathbb{R}$, thus $\phi_i^{(1)}$ is also a zero measure or $d\psi_i^{(1)} = d\mu_i^{(1)}$ for all $i = 1, \dots, 2^{n-1}$.

Now assume that $d\psi_i^{(k-1)} = d\mu_i^{(k-1)}$, $2 \leq k \leq n$, for all $i = 1, \dots, 2^{n-k+1}$. We will prove that $d\psi_i^{(k)} = d\mu_i^{(k)}$ for all $i = 1, \dots, 2^{n-k}$. We have: $d\psi_{2i}^{(k-1)} = d\mu_{2i}^{(k-1)}$ and $d\psi_{2i-1}^{(k-1)} = d\mu_{2i-1}^{(k-1)}$ for $i : 1 \leq i \leq 2^{n-k}$. Thus $z_{2i}^{(k-1)}$ and $z_{2i-1}^{(k-1)}$ are moment sequences of two measures $\mu_{2i}^{(k-1)}$ and $\mu_{2i-1}^{(k-1)}$ respectively.

According to (15), we have:

$$\int \mathbf{x}^\alpha d\psi_i^{(k)} = \sum_{\beta \in \mathbb{N}^{k-1}} \frac{\binom{(k-1)}{g_{2,\alpha}}_\beta z_{2i}^{(k-1)}(\beta)}{\alpha_k + 1} - \sum_{\beta \in \mathbb{N}^{k-1}} \frac{\binom{(k-1)}{g_{1,\alpha}}_\beta z_{2i-1}^{(k-1)}(\beta)}{\alpha_k + 1} - \int \frac{\mathbf{x}^\alpha x_k}{\alpha_k + 1} \frac{\partial h_i^{(k)}(\mathbf{x})}{\partial x_k} d\psi_i^{(k)}. \quad (31)$$

Similarly,

$$\int \mathbf{x}^\alpha d\mu_i^{(k)} = \sum_{\beta \in \mathbb{N}^{k-1}} \frac{\binom{(k-1)}{g_{2,\alpha}}_\beta z_{2i}^{(k-1)}(\beta)}{\alpha_k + 1} - \sum_{\beta \in \mathbb{N}^{k-1}} \frac{\binom{(k-1)}{g_{1,\alpha}}_\beta z_{2i-1}^{(k-1)}(\beta)}{\alpha_k + 1} - \int \frac{\mathbf{x}^\alpha x_k}{\alpha_k + 1} \frac{\partial h_i^{(k)}(\mathbf{x})}{\partial x_k} d\mu_i^{(k)}. \quad (32)$$

Then if we consider the *signed* measure $\phi_i^{(k)}$ on Ω_k that satisfies $d\phi_i^{(k)} = d\psi_i^{(k)} - d\mu_i^{(k)}$, we have:

$$\int \mathbf{x}^\alpha d\phi_i^{(k)} = - \int \frac{\mathbf{x}^\alpha x_k}{\alpha_k + 1} \frac{\partial h_i^{(k)}(\mathbf{x})}{\partial x_k} d\phi_i^{(k)}. \quad (33)$$

Using similar arguments with simultaneous approximation of continuous functions, we obtain the following equation for all functions $f(\mathbf{x}) = x_k g(\mathbf{x})$, $\mathbf{x} \in \mathbb{R}^k$, where g and $\frac{\partial g}{\partial x_k}$ are continuous:

$$\int \left[\frac{\partial f(\mathbf{x})}{\partial x_k} + f(\mathbf{x}) \frac{\partial h_i^{(k)}(\mathbf{x})}{\partial x_k} \right] d\phi_i^{(k)} = 0. \quad (34)$$

For any polynomial $g \in \mathbb{R}[\mathbf{x}]$, define $G(\mathbf{x}) = x_k P(\mathbf{x})$, where $P(\mathbf{x}) \in \mathbb{R}[\mathbf{x}]$ and $\frac{\partial G(\mathbf{x})}{\partial x_k} = g(\mathbf{x})$, then with $f(\mathbf{x}) = G(\mathbf{x}) e^{-h_i^{(k)}(\mathbf{x})}$, we again obtain

$$\int g(\mathbf{x}) e^{-h_i^{(k)}(\mathbf{x})} d\phi_i^{(k)} = 0, \quad \forall g \in \mathbb{R}[\mathbf{x}]. \quad (35)$$

Similar arguments are again used and we have the measure $\phi_i^{(k)}$ is a zero measure or $d\psi_i^{(k)} = d\mu_i^{(k)}$ for all $i = 1, \dots, 2^{n-k}$. So the results are true for all $k = 1, \dots, n$. \square

With this lemma, we can now prove the following convergence theorem:

Theorem 1 *Let $g, h \in \mathbb{R}[\mathbf{x}]$, Ω be the compact set defined in (9), and consider the semidefinite programming problems \mathcal{P}_r^l and \mathcal{P}_r^u defined in (23) and (24) respectively. Then*

(i) Optimal values $Z(\mathcal{P}_r^l)$ and $Z(\mathcal{P}_r^u)$ are finite and in addition, both problems \mathcal{P}_r^l and \mathcal{P}_r^u are solvable for r large enough.

(ii) As $r \rightarrow \infty$, $Z(\mathcal{P}_r^l) \uparrow \rho$ and $Z(\mathcal{P}_r^u) \downarrow \rho$.

Proof. Clearly, the collection of truncated sequences of moments of $\mu_i^{(k)}$ is a feasible solution for both problems \mathcal{P}_r^l and \mathcal{P}_r^u . We have:

$$|z_i^{(k)}(\boldsymbol{\alpha})| \leq \int_{\Omega_k} |(\mathbf{x}[k])^\alpha e^{h_i^{(k)}(\mathbf{x}[k])}| d\mathbf{x}[k] \leq \sup_{\mathbf{x} \in \Omega_k} |\mathbf{x}^\alpha e^{h_i^{(k)}(\mathbf{x})}| \text{vol}(\Omega_k).$$

Ω_k is compact in \mathbb{R}^k ; therefore, we have $u_i^{(k)}(\boldsymbol{\alpha}) = \sup_{\mathbf{x} \in \Omega_k} |\mathbf{x}^\alpha e^{h_i^{(k)}(\mathbf{x})}| = \max_{\mathbf{x} \in \Omega_k} |\mathbf{x}^\alpha e^{h_i^{(k)}(\mathbf{x})}|$ is finite. Now consider the problems \mathcal{P}_r^l and \mathcal{P}_r^u with additional bound constraints $-u_i^{(k)}(\boldsymbol{\alpha}) - 1 \leq z_i^{(k)}(\boldsymbol{\alpha}) \leq u_i^{(k)}(\boldsymbol{\alpha}) + 1$ for all k, i and $\boldsymbol{\alpha}$, we have:

(i) The feasible sets of these two modified problems are bounded and closed. The objective functions are linear and both problems are feasible. Therefore, they are both solvable and their optimal values are finite.

(ii) Let $\{\mathbf{z}_{i,r}^{(k)}\}$ be the optimal solution of \mathcal{P}_r^l (with bound constraints), we extend these truncated sequences with zeros to make them become infinite sequences. According to Lasserre [5], there is a subsequence $\{r_m\}$ and infinite sequences $\{\mathbf{z}_{i,*}^{(k)}\}$ such that pointwise convergence holds with respect to the usual sup-norm. Therefore, we have:

$$M(\mathbf{z}_{i,*}^{(k)}) \succeq 0, \quad M(\theta_j^{(k)} \mathbf{z}_{i,*}^{(k)}) \succeq 0, \quad \forall k, i, j.$$

We also have Ω_k is compact in \mathbb{R}^k ; therefore, according to Putinar [8], there exist measures $\psi_i^{(k)}$ supported on Ω_k such that $\mathbf{z}_{i,*}^{(k)}$ are their moment sequences assuming the representation condition holds. In addition, $\{\mathbf{z}_{i,*}^{(k)}\}$ satisfy the recursive formulae (15) and (16) from the pointwise convergence. From Lemma 1, we obtain that $d\psi_i^{(k)} = d\mu_i^{(k)}$ for all $k = 1, \dots, n$ and $i = 1, \dots, 2^{n-k}$. Thus we have:

$$\lim_{m \rightarrow \infty} \langle \mathbf{g}, \mathbf{z}_{1,r_m}^{(n)} \rangle = \langle \mathbf{g}, \mathbf{z}_{1,*}^{(n)} \rangle = \rho.$$

Due to the construction of truncated moment matrices and localizing matrices as well as the sets $A_{i,r}^{(k)}$, clearly, a feasible solution of \mathcal{P}_{r+1}^l generates a feasible solution of \mathcal{P}_r^l . Thus with $r \geq \deg(g)$, then $\langle \mathbf{g}, \mathbf{z}_{1,r}^{(n)} \rangle \leq \langle \mathbf{g}, \mathbf{z}_{1,r+1}^{(n)} \rangle$ or we have:

$$\langle \mathbf{g}, \mathbf{z}_{1,r}^{(n)} \rangle \uparrow \rho.$$

Similar arguments can be applied for the modified problem \mathcal{P}_r^u .

So far, the results obtained are for problems \mathcal{P}_r^l and \mathcal{P}_r^u with additional bound constraints. However, at the limit with respect to the usual sup-norm, none of these bound constraints are tight. Thus, these constraints can be removed when r is large enough. In other words, the problems \mathcal{P}_r^l and \mathcal{P}_r^u have finite optimal solutions and solvable when r is large enough and

$$Z(\mathcal{P}_r^l) \uparrow \rho, \quad Z(\mathcal{P}_r^u) \downarrow \rho.$$

□

4 Computational Results

In order to demonstrate the method, we use two different kinds of Ω sets: hypercubes and order statistic integrals. The computational results are obtained for integrals over these sets of random multivariate normal distribution.

4.1 Integrals over Hypercubes

If Ω is a hypercube, then $b_k^{(i)}(\mathbf{x}) = b_k^{(i)}$ are constants for all $k = 2, \dots, n$. We have:

$$g_{i,\alpha}^{(k)}(\mathbf{x}) = [b_{k+1}^{(i)}]^{\alpha_{k+1}+1} \mathbf{x}^{\alpha^{[k]}}, \quad \mathbf{x} \in \mathbb{R}^k, \quad \alpha \in \mathbb{N}^k, \quad i = 1, 2. \quad (36)$$

The functions $h_i^{(k)}$ are still formulated as in (13):

$$h_i^{(k)}(\mathbf{x}) = h_{\lfloor i/2 \rfloor}^{(k+1)}(\mathbf{x}, b_{k+1}^{(2-\lfloor i/2 \rfloor + \lfloor i/2 \rfloor)}), \quad \mathbf{x} \in \mathbb{R}^k, \quad k = 1, \dots, n-1, \quad i = 1, \dots, 2^{n-k}.$$

The recursive formula simply becomes

$$z_i^{(k)}(\alpha) = \frac{[b_k^{(2)}]^{\alpha_k+1}}{\alpha_k+1} z_{2i}^{(k-1)}(\alpha[k-1]) - \frac{[b_k^{(1)}]^{\alpha_k+1}}{\alpha_k+1} z_{2i-1}^{(k-1)}(\alpha[k-1]) - \sum_{\beta \in \mathbb{N}^k} \frac{\beta(h_i^{(k)})_{\beta}}{\alpha_k+1} z_i^{(k)}(\alpha+\beta) \quad (37)$$

for all $k = 2, \dots, n$, $\alpha \in \mathbb{R}^k$, $i = 1, \dots, 2^{n-k}$ while for $k = 1$, it remains the same as in (16):

$$z_i^{(1)}(\alpha) = \frac{(b_1^{(2)})^{\alpha+1}}{\alpha+1} e^{h_i^{(1)}(b_1^{(2)})} - \frac{(b_1^{(1)})^{\alpha+1}}{\alpha+1} e^{h_i^{(1)}(b_1^{(1)})} - \sum_{\beta \in \mathbb{N}} \frac{\beta(h_i^{(1)})_{\beta}}{\alpha+1} z_i^{(1)}(\alpha+\beta), \quad \forall i = 1, \dots, 2^{n-1}.$$

The polynomials $\theta_i^{(k)}(\mathbf{x}) = [b_i^{(2)} - x_i] [x_i - b_i^{(1)}] = -b_i^{(1)}b_i^{(2)} + (b_i^{(1)} + b_i^{(2)})x_i - x_i^2$, $\mathbf{x} \in \mathbb{R}^k$ have the degree $d_i = 2$ for all $k \geq i$, $i = 1, \dots, n$ in this case.

4.2 Order Statistic Integrals

Order statistic integrals are calculated over the set $\Omega = \{\mathbf{x} \in \mathbb{R}^n : b^{(1)} \leq x_1 \leq \dots \leq x_n \leq b^{(2)}\}$. So we have: $b_k^{(2)} = b^{(2)}$ are constant for all $k = 1, \dots, n$ while $b_k^{(1)}(\mathbf{x}) = x_{k-1}$ for all $k = 2, \dots, n$ and $b_1^{(1)} = b^{(1)}$. The functions $g_{2,\alpha}^{(k)}$ are defined as in (36) while $g_{1,\alpha}^{(k)}$ are formulated as follows

$$g_{1,\alpha}^{(k)}(\mathbf{x}) = x_k^{\alpha_{k+1}+1} \mathbf{x}^{\alpha[k]} = \mathbf{x}^{\alpha[k+(\alpha_{k+1}+1)\mathbf{e}_k]}, \quad \mathbf{x} \in \mathbb{R}^k, \quad \alpha \in \mathbb{N}^k, \quad (38)$$

where \mathbf{e}_k is the k^{th} unit vector in \mathbb{R}^k .

The recursive formula is then

$$z_i^{(k)}(\alpha) = \frac{[b^{(2)}]^{\alpha_k+1}}{\alpha_k+1} z_{2i}^{(k-1)}(\alpha[k-1]) - \frac{1}{\alpha_k+1} z_{2i-1}^{(k-1)}(\alpha[k-1] + (\alpha_k+1)\mathbf{e}_k) - \sum_{\beta \in \mathbb{N}^k} \frac{\beta_k (h_i^{(k)})^\beta}{\alpha_k+1} z_i^{(k)}(\alpha + \beta) \quad (39)$$

for all $k = 2, \dots, n$, $\alpha \in \mathbb{R}^k$, $i = 1, \dots, 2^{n-k}$. For $k = 1$, the formula remains the same as in (16).

Finally, the polynomials $\theta_i^{(k)}(\mathbf{x}) = [b^{(2)} - x_i][x_i - x_{i-1}] = -b^{(2)}x_{i-1} + b^{(2)}x_i + x_{i-1}x_i - x_i^2$, $\mathbf{x} \in \mathbb{R}^k$ also have degree $d_i = 2$ for all $k \geq i$, $i = 2, \dots, n$. For $i = 1$, the polynomial $\theta_1^{(k)}(\mathbf{x})$ is the same as in the previous section with degree $d_1 = 2$.

4.3 Computational Results for Normal Distributions

Multivariate normal distributions are used to obtain computational results for integrals over hypercubes and order statistic integrals. The density function of a multivariate normal distribution with mean $\boldsymbol{\mu}$ and covariance matrix $\boldsymbol{\Sigma} = \mathbf{A}\mathbf{A}'$ is

$$f(\mathbf{x}) = \frac{1}{(2\pi)^{n/2} \det(\boldsymbol{\Sigma})^{1/2}} e^{-\frac{1}{2}(\mathbf{x}-\boldsymbol{\mu})'\boldsymbol{\Sigma}^{-1}(\mathbf{x}-\boldsymbol{\mu})}. \quad (40)$$

Thus, in this case, $g(\mathbf{x}) = [(2\pi)^{n/2} \det(\boldsymbol{\Sigma})^{1/2}]^{-1}$, a constant, and $h(\mathbf{x}) = -\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})'\boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu})$, a quadratic polynomials.

Algorithms for general g and h are implemented in Matlab for both hypercubic and order statistic integrals. Semidefinite programming problems are solved using SeDuMi routines. For the case of normal distributions, means and covariance matrices are created randomly using random vectors and matrices with elements within the range of $[-1, 1]$. The two Ω sets are defined on the hypercube $\{\mathbf{x} \in \mathbb{R}^n : -1 \leq x_i \leq 1 \quad \forall i = 1, \dots, n\}$.

The algorithms are run for different normal distributions in $n = 1, 2$, and 3-dimension space. The moment degree starts at $r = 2$ and increases until the tolerance is met. The tolerance $\epsilon =$

$\frac{1}{2} [Z(\mathcal{P}_r^u) - Z(\mathcal{P}_r^l)]$ is set to be 5×10^{-5} (4-digit accuracy). The results will be averaged from those different distributions in each case. All computations are done under a Windows environment on a Pentium III 1GHz with 256MB RAM.

Dimension n	Min. degree r_{\min}	Max. degree r_{\max}	Average degree \bar{r}	Computational time (seconds)			
				$r = 2$	$r = 3$	$r = 4$	$r = 5$
1	2	5	3.5	1.332	1.631	1.505	1.437
2	3	5	4.3	1.994	3.215	11.036	70.107

Table 1: Computational results for hypercubic integrals

Dimension n	Min. degree r_{\min}	Max. degree r_{\max}	Average degree \bar{r}	Computational time (seconds)			
				$r = 2$	$r = 3$	$r = 4$	$r = 5$
1	2	5	3.5	1.332	1.631	1.505	1.437
2	3	5	4.3	2.079	3.413	12.294	79.418

Table 2: Computational results for order statistic integrals

Table 1 and 2 show computational results for hypercubic and order statistic integrals respectively. Minimum degree r_{\min} is the smallest moment degree we need to use to meet the tolerance setting while r_{\max} is the largest moment degree. \bar{r} is the average value obtained from all distributions tested for each value of n . In the case $n = 1$, there is not much difference between computational times for different moment degrees and we need roughly $r = 3$ or $r = 4$ on average to obtain the desired results. These observations are for both hypercubic and order statistic integrals even though the latter need slightly more time to be calculated. For $n = 2$, the average moment degree is $\bar{r} = 4.3$ and computational times are significantly varied from 2 seconds for $r = 2$ up to approximately 80 seconds for $r = 5$. Due to the large amount of memory needed for SeDuMi routine, we cannot run the algorithms with $r = 5$ for the case $n = 3$ to meet the 4-digit accuracy requirement. Therefore, no result on required moment degrees for this accuracy setting is reported here for $n = 3$. However, we will report the real error gaps for this case up to $r = 4$ in Table 3. With respect to the computational time, it is not very favorable in the case $n = 3$ as we need more than 2000 seconds when $r = 4$ (as compared to approximately 5 and 100 seconds when $r = 2$ and $r = 3$ respectively).

Table 3 shows the real gaps between lower and upper bounds of integral values for some specific

Integral	Dimension n	Value $\bar{\rho}$	Error $\frac{1}{2} [Z(\mathcal{P}_r^u) - Z(\mathcal{P}_r^l)]$			
			$r = 2$	$r = 3$	$r = 4$	$r = 5$
Hypercubic	1	0.2665	1.4047E-04	8.1177E-07		
	2	0.1085	1.0376E+01	1.1200E-02	6.7765E-04	4.7855E-05
	3	0.0736	4.7420E-01	8.0400E-02	2.2000E-03	
Order statistic	2	0.0670	7.8969E+00	6.3000E-03	4.2279E-04	3.2933E-05
	3	0.0274	4.9390E-01	5.4000E-03	4.2253E-04	

Table 3: Real errors for specific normal distributions

normal distributions, both hypercubic and order statistic integrals. These values decrease when the moment order increases. For example, when $n = 3$, the gap between lower and upper bounds for $r = 4$ is already relatively small even though the accuracy requirement has not been satisfied in this case. The computational times for large n and r are yet very promising; however, we believe that this method is plausible to be implemented. One may also use the recursive formulae presented in Section 2, to eliminate some moments variables and substitute their expression in the remaining variables in the LMIs of the SDP-relaxations. This would substantially reduce the total number of variables. In addition, the framework we proposed could be further developed to calculate integrations not only with polynomials but also rational functions, which have been recently analyzed by Lasserre [4].

References

- [1] C. Coatmélec. Approximation et interpolation des fonctions différentiables de plusieurs variables. *Annales scientifiques de l'É.N.S 3^e série*, 83(4):271–341, 1966.
- [2] P. de la Harpe and C. Pache. Cubature formulas, geometrical designs, reproducing kernels, and Markov operators. In *International Conference on Group Theory*, Gaeta, June 2003.
- [3] M. W. Hirsch. *Differential topology*. Springer-Verlag, New York, 1976.
- [4] J. B. Lasserre. The K-moment problem with densities. Technical Report 05334, LAAS, June 2005.
- [5] J. B. Lasserre. A sum of squares approximation of nonnegative polynomials. *SIAM Journal on Optimization*, 16(2):610–628, 2005.
- [6] M. Laurent. Moment matrices and optimization over polynomials - a survey on selected topics. Preprint, September 2005.
- [7] H. Niederreiter. Random number generator and quasi-Monte Carlo method. *SIAM CBMS-NSF Regional Conference Series in Applied Mathematics*, 63, 1992.
- [8] M. Putinar. Positive polynomials on compact semi-algebraic sets. *Indiana University Mathematics Journal*, 42:969–984, 1993.