

Dedicated to Dr. Constantin Vârsan on the occasion of his 70th Birthday

Control problems with mixed constraints and application to an optimal investment problem

J.F. BONNANS and D. TIBA*

Abbreviated title: Control problems with mixed constraints

Abstract We discuss two optimal control problems of parabolic equations, with mixed state and control constraints, for which the standard qualification condition does not hold. Our first example is a bottleneck problem, and the second one is an optimal investment problem where a utility type function is to be minimized. By an adapted penalization technique, we derive optimality conditions from which useful information of the solution can be derived. In the case of a control entering linearly in the state equation and cost function, we obtain generalized bang-bang properties.

AMS 2000 Subject Classification: 49K20, 49M30, 35K05.

Key words: Parabolic control problems, mixed constraints, bottleneck problems, optimal investment, constraint qualification.

1 INTRODUCTION

Optimal control problems involving state constraints (and, in particular, mixed constraints) are well known for their intrinsic difficulty. There is a rich literature devoted to the optimality conditions and the regularity of the Lagrange multipliers for the case of parabolic control problems with mixed constraints:

*The work of the second author was supported by CNCSIS, Romania, grant ID-PCE 1192-09.

Arada and Raymond [1], Tröltzsch [17], De los Reyes and Tröltzsch [8], Rösch and Tröltzsch [15].

One special case very much discussed in the literature is the so-called "bottleneck problem", introduced by Bellman [3] further studied by Mirică [12], Bergounioux and Tiba [4], Bergounioux and Tröltzsch [5, 6]. We study in section 2 a variant of the bottleneck problem. We fix a "polynomial" cost functional and a linear parabolic state system, and we investigate the situation when the state is "dominated" by the control.

Section 3 is devoted to an optimal investment problem that in some sense is the opposite case of the bottleneck problem. Given a distribution of capital over space, we assume that one cannot invest more than a fraction of the capital, and that some diffusion of the capital occurs. We study the case of a small aversion to risk.

Our approach is based on the adapted penalization of the state equation, while the constraints are kept explicit. Elements of our technique have been previously used by Lions [11], Barbu and Precupanu [2], Bergounioux and Tiba [4]. The form of the optimality conditions that we obtain has the advantage of a certain symmetry: the control and the state play a similar role which is a natural characteristic for mixed constraints. They are also accessible for further analysis in order to obtain supplementary information like bang-bang or regularity properties for the unknowns, which is the main aim of this paper.

2 CONTROL DOMINATED PROBLEMS

We analyze the following model problem:

$$\text{Min } \left\{ \frac{1}{2} \int_Q (y - y_d)^2 dxdt + \frac{N}{q} \int_{Q_\omega} |Bu|^q dxdt \right\}, \quad (2.1)$$

$$y_t - \Delta y = Bu \text{ in } Q = \Omega \times]0, T[, \quad (2.2)$$

$$y(x, 0) = 0 \text{ in } \Omega, \quad y(x, t) = 0 \text{ on } \Sigma = \partial\Omega \times [0, T], \quad (2.3)$$

$$[y, u] \in D \subset C(0, T; H_0^1(\Omega)) \times U. \quad (2.4)$$

Above, U is the reflexive Banach space of controls, Ω is a bounded smooth domain in R^d , $d \geq 1$, ω is a measurable subset of Ω , and setting $Q_\omega = \omega \times]0, T[$, B is a linear continuous operator : $U \rightarrow L^q(Q_\omega)$, with extension by 0 to Q , D is a closed convex nonvoid subset of $C(0, T; H_0^1(\Omega)) \times U$, $q \geq 2$, $N \geq 0$, and $y_d \in L^2(Q)$ are given.

The cost functional is a direct generalization of the standard quadratic functional. More complex situations instead of (2.1)-(2.3) may be considered as well. Here, we concentrate on the treatment of the mixed constraint (2.4), which is formulated in a very general way.

Notice that the unique solution y of (2.2)-(2.3) belongs to $W^{2,1,q}(Q)$. If $q > \frac{1}{2}(n+2)$, then $y \in C(\overline{Q})$ by the Sobolev embedding theorem.

If the set of admissible pairs $[y, u] \in D$ and satisfying (2.2)-(2.3) is nonvoid and if $N > 0$, then it is well known that the problem (2.1)-(2.4) has a unique optimal pair $[y^*, u^*] \in W^{2,1,q}(Q) \times U$, by the coercivity and strict convexity of the cost functional, see e.g. Neittaanmäki and Tiba [13]. Since U is a reflexive space, existence may be obtained as well for $N = 0$ if the set of controls satisfying (2.4) is bounded in U . In the sequel, we assume:

$$\text{Problem (2.1)-(2.4) has at least one optimal pair } [y^*, u^*] \in D. \quad (2.5)$$

Examples for (2.4), that we have in mind are:

$$\frac{1}{2} \int_{\Omega} y(x, t)^2 dx \leq C(u)(t), \quad t \in [0, T], \quad (2.6)$$

$$u \in U_{ad}, \text{ with } U_{ad} \text{ closed convex subset of } U. \quad (2.7)$$

Here $C(\cdot)$ is a given operator $U \rightarrow L^1(0, T)$. For instance, if $C : U \rightarrow R$ is a positive constant, then (2.6)-(2.7) is a standard example of separate state and control constraints. If $U = L^q(0, T)$ and $B : U \rightarrow L^q(Q_\omega)$, $(Bu)(x, t) = f(x)u(t)$, $f \in L^\infty(\omega)$ and $C(u)(t) = u(t)$, $t \in [0, T]$, we obtain a variant of the bottleneck problem. The inequality (2.6) justifies the title of this section. In (2.7), under the above notations, one may take

$$U_{ad} = \{u \in L^q(0, T); \quad a(t) \leq u(t) \leq b(t) \quad \text{a.e. in } [0, T]\}$$

with a and b in $L^\infty(0, T)$. In this case, (2.5) holds even when $N = 0$. The adapted penalization method applied to problem (2.1)-(2.4) is based on the following approximation, for $\varepsilon > 0$:

$$\begin{aligned} \text{Min}_{[y, u]} \left\{ \frac{1}{2} \int_Q (y - y_d)^2 dx dt + \frac{N}{q} \int_{Q_\omega} |Bu|^q dx dt + \right. \\ \left. + |u - u^*|_U^2 + \frac{1}{q\varepsilon} \int_Q |y_t - \Delta y - Bu|^q dx dt \right\}, \quad (2.8) \end{aligned}$$

subject to:

$$y \in W^{2,1,q}(Q), \quad y(x, 0) = 0 \text{ in } \Omega, \quad y(x, t) = 0 \text{ in } \Sigma, \quad [y, u] \in D. \quad (2.9)$$

Due to the presence of the adapted term $|u - u^*|_U^2$, the minimization problem (2.8)-(2.9) has a unique minimal pair $[y_\varepsilon, u_\varepsilon]$. Moreover, since the $[y^*, u^*]$ satisfies (2.9), (2.2) and is feasible for (2.8), we have the inequality:

$$\begin{aligned} & \frac{1}{2} \int_Q (y_\varepsilon - y_d)^2 dxdt + \frac{N}{q} \int_{Q_\omega} |Bu_\varepsilon|^q dxdt + |u_\varepsilon - u^*|_U^2 + \\ & + \frac{1}{q\varepsilon} \int_Q |(y_\varepsilon)_t - \Delta y_\varepsilon - Bu_\varepsilon|^q dxdt \\ & \leq \frac{1}{2} \int_Q (y^* - y_d)^2 dxdt + \frac{N}{q} \int_{Q_\omega} |Bu^*|^q dxdt. \end{aligned} \quad (2.10)$$

Therefore $[y_\varepsilon, u_\varepsilon]$ is bounded in $W^{2,1,q}(Q) \times U$ and

$$(y_\varepsilon)_t - \Delta y_\varepsilon - Bu_\varepsilon \rightarrow 0 \text{ strongly in } L^q(Q). \quad (2.11)$$

Denote $r_\varepsilon = \varepsilon^{-1} [(y_\varepsilon)_t - \Delta y_\varepsilon - Bu_\varepsilon] \in L^q(Q)$. By (2.10), $\varepsilon^{(q-1)/q} r_\varepsilon$ is bounded in $L^q(Q)$. We may assume that (for some subsequence) $u_\varepsilon \rightarrow \hat{u}$ weakly in U , $y_\varepsilon \rightarrow \hat{y}$ weakly in $W^{2,1,q}(Q)$, and we get that $[\hat{y}, \hat{u}]$ satisfies (2.9) since D is weakly closed. Passing to the limit in (2.11), obtain

$$\hat{y}_t - \Delta \hat{y} - B\hat{u} = 0 \text{ in } Q,$$

i.e., the pair $[\hat{y}, \hat{u}]$ is feasible for problem (2.1)-(2.4). By (2.10) and the weak lower semicontinuity of the norm, we have:

$$\begin{aligned} & \frac{1}{2} \int_Q (\hat{y} - y_d)^2 dxdt + \frac{N}{q} \int_{Q_\omega} |B\hat{u}|^q dxdt + |\hat{u} - u^*|_U^2 \leq \\ & \leq \frac{1}{2} \int_Q (y^* - y_d)^2 dxdt + \frac{N}{q} \int_{Q_\omega} |Bu^*|^q dxdt. \end{aligned}$$

Therefore $[\hat{y}, \hat{u}]$ is optimal for the problem (2.1)-(2.4) and $\hat{u} = u^*$. Clearly, the weak convergences are in fact strong since $|u_\varepsilon - u^*|_U \rightarrow 0$. We have proved

Proposition 2.1. *The following holds:*

$$u_\varepsilon \rightarrow u^* \text{ strongly in } U, \quad (2.12)$$

$$y_\varepsilon \rightarrow y^* \text{ strongly in } W^{2,1,q}(Q), \quad (2.13)$$

$$\left\{ \varepsilon^{\frac{q-1}{q}} r_\varepsilon \right\} \text{ is bounded in } L^q(Q). \quad (2.14)$$

For a given pair $[y, u]$ satisfying (2.9), let us consider convex variations denoted $[y_s, u_s]$, with $y_s = y_\varepsilon + s(y - y_\varepsilon)$, $u_s = u_\varepsilon + s(u - u_\varepsilon)$, for s in $[0, 1]$. Obviously $[y_s, u_s]$ satisfies (2.9) and we can write the inequality

$$\begin{aligned}
& \frac{1}{2} \int_Q (y_\varepsilon - y_d)^2 dxdt + \frac{N}{q} \int_{Q_\omega} |Bu_\varepsilon|^q dxdt + |u_\varepsilon - u^*|_U^2 + \\
& + \frac{1}{q\varepsilon} \int_Q |(y_\varepsilon)_t - \Delta y_\varepsilon - Bu_\varepsilon|^q dxdt \leq \frac{1}{2} \int_Q (y_\varepsilon + s(y - y_\varepsilon) - y_d)^2 dxdt + \\
& + \frac{N}{q} \int_{Q_\omega} |Bu_\varepsilon + s(Bu - Bu_\varepsilon)|^q dxdt + |u_\varepsilon + s(u - u_\varepsilon) - u^*|_U^2 + \\
& + \frac{1}{q\varepsilon} \int_Q |(y_\varepsilon)_t + s(y - y_\varepsilon)_t - \Delta y_\varepsilon - s\Delta(y - y_\varepsilon) - Bu_\varepsilon - sB(u - u_\varepsilon)|^q dxdt.
\end{aligned} \tag{2.15}$$

Let us denote by $\text{sgn}(\cdot)$ the sign function, U^* the topological dual of U , and $F : U \rightarrow U^*$ the duality mapping. Standard computations in (2.15) allow to obtain the following result:

Proposition 2.2. *The pair $[y_\varepsilon, u_\varepsilon]$ satisfies the following necessary and sufficient first order optimality condition: for any $[y, u]$ for which (2.9) holds, we have that*

$$\begin{aligned}
0 \leq & \int_Q (y_\varepsilon - y_d)(y - y_\varepsilon) dxdt + N \int_{Q_\omega} |Bu_\varepsilon|^{q-1} \text{sgn}(Bu_\varepsilon)(Bu - Bu_\varepsilon) dxdt + \\
& + \langle F(u_\varepsilon - u^*), u - u_\varepsilon \rangle_{U^* \times U} + \int_Q \varepsilon^{q-2} |r_\varepsilon|^{q-1} \text{sgn}(r_\varepsilon)(y_t - \Delta y - Bu) dxdt.
\end{aligned} \tag{2.16}$$

Proof. As already mentioned, the necessity follows from (2.15), by dividing each side by $s > 0$ and taking the limit when $s \rightarrow 0$. The sufficiency of (2.16) is a consequence of the definition of the subdifferential since the right-hand side in (2.16) may be upper bounded by

$$\begin{aligned}
& \frac{1}{2} \int_Q (y - y_d)^2 dxdt - \frac{1}{2} \int_Q (y_\varepsilon - y_d)^2 dxdt + \frac{N}{q} \int_{Q_\omega} |Bu|^q dxdt - \\
& - \frac{N}{q} \int_{Q_\omega} |Bu_\varepsilon|^q dxdt + |u - u^*|_U^2 - |u_\varepsilon - u^*|_U^2 + \\
& \frac{1}{\varepsilon q} \int_Q |y_t - \Delta y - Bu|^2 dxdt - \frac{1}{q\varepsilon} \int_Q |(y_\varepsilon)_t - \Delta y_\varepsilon - Bu_\varepsilon|^q dxdt,
\end{aligned}$$

for any $[y, u]$ satisfying (2.9). The conclusion follows. \square

We consider now the main example of this section, the operator B having the form $(Bu)(x, t) = f(x)u(t)$, $u \in U = L^q(0, T)$ and (2.6)-(2.7) becomes

$$\frac{1}{2} \int_{\Omega} y(x, t)^2 dx dt \leq u(t), \quad \text{for a.a. } t \in [0, T]. \quad (2.17)$$

$$u \in L^q(0, T); \quad a(t) \leq u(t) \leq b(t) \quad \text{a.e. } [0, T], \quad (2.18)$$

and there exist constants $\alpha_a < 0$ and $\alpha_b > 0$ such that

$$a(t) \leq \alpha_a < 0 < \alpha_b \leq b(t), \quad \text{for a.a. } t \in (0, T). \quad (2.19)$$

Proposition 2.3. *If (2.17)-(2.19) hold, then $\{\varepsilon^{q-2} |r_\varepsilon|^{q-1}\}$ is bounded in $L^{q/(q-1)}(Q)$ (or equivalently, $\{\varepsilon^{(q-2)/(q-1)} r_\varepsilon\}$ is bounded in $L^q(Q)$).*

Proof. For $\lambda > 0$, let y^λ be the unique element of $W^{2,1,q}(Q)$ satisfying (2.3) and

$$y_t^\lambda - \Delta y^\lambda = f(x)\lambda \quad \text{in } Q.$$

That is, y^λ is the solution of (2.2)-(2.3) associated with $u^\lambda \equiv \lambda$. When $\lambda > 0$ is small enough, say $\lambda < \lambda_0$, with $\lambda_0 > 0$, then u^λ satisfies (2.18) and $\frac{\lambda}{2} \int_{\Omega} y^1(x, t)^2 dx \leq \frac{1}{2}$. Consequently

$$0 \leq \frac{1}{2} \int_{\Omega} y^\lambda(x, t)^2 dx = \frac{1}{2} \lambda^2 \int_{\Omega} y^1(x, t)^2 dx \leq \frac{1}{2} \lambda. \quad (2.20)$$

Given $\rho \in L^q(Q)$ with $|\rho|_{L^q(Q)} \leq 1$, define y_ρ as the solution of (2.3) and

$$(y_\rho)_t - \Delta y_\rho = \rho \quad \text{in } Q. \quad (2.21)$$

Then, $|y_\rho|_{C(\overline{Q})} \leq K$ (some positive constant) if $|\rho|_{L^q(Q)} \leq 1$. In view of (2.20)-(2.21), for any $\delta \in \mathbb{R}$, we have that

$$\begin{aligned} \frac{1}{2} \int_{\Omega} (y^\lambda + \delta y_\rho)^2 dx &= \frac{1}{2} \int_{\Omega} (y^\lambda)^2 dx + \frac{1}{2} \delta^2 \int_{\Omega} y_\rho^2 dx + \delta \int_{\Omega} y^\lambda y_\rho dx \\ &\leq \frac{1}{2} \lambda + \frac{1}{2} \delta^2 K^2 \text{mes}(\Omega) + |\delta| \lambda K \text{mes}(\Omega) |y^1|_{C(\overline{Q})}. \end{aligned} \quad (2.22)$$

Given $\lambda \in (0, \lambda_0)$, for small enough $\delta > 0$, (2.22) shows that the pair $(y^\lambda + \delta y_\rho, \lambda)$ belongs to D defined in (2.17)-(2.18). Use this pair in (2.16), we get

the inequality:

$$0 \leq \int_Q (y_\varepsilon - y_d)(y^\lambda + \delta y_\rho - y_\varepsilon) dxdt + N \int_{Q_\omega} |Bu_\varepsilon|^{q-1} \operatorname{sgn}(Bu_\varepsilon)(B\lambda - Bu_\varepsilon) dxdt + \langle F(u_\varepsilon - u^*), \lambda - u_\varepsilon \rangle_{U^* \times U} + \delta \int_Q \varepsilon^{q-2} |r_\varepsilon|^{q-1} \operatorname{sgn} r_\varepsilon \rho(x, t) dxdt.$$

Since all terms except the last remain uniformly bounded over $\varepsilon > 0$ (remember that here $\lambda > 0$ and $\delta > 0$ are fixed), the last integral is uniformly lower bounded. Since ρ is an arbitrary element of the closed unit ball, and the spaces $L^q(Q)$ and $L^{(q-1)/q}(Q)$ are dual to each other, the infimum of this integral over the unit ball is $-\varepsilon^{q-2} \|r_\varepsilon\|_{L^{(q-1)/q}(Q)}$. The conclusion follows. \square

Theorem 2.4. *If (2.17)-(2.19) hold, then the pair $[y^*, u^*] \in D$ is optimal for problem (2.1)-(2.4) iff there exists $r^* \in L^{\frac{q}{q-1}}(Q)$ such that, for any (y, u) satisfying (2.9):*

$$\begin{cases} 0 \leq \int_Q (y^* - y_d)(y - y^*) dxdt + N \int_{Q_\omega} |Bu^*|^{q-1} \operatorname{sgn}(Bu^*) \\ (Bu - Bu^*) dxdt + \int_Q r^*(y_t - \Delta y - Bu) dxdt. \end{cases} \quad (2.23)$$

Proof. Thanks to proposition 2.3, there exists a sequence $\varepsilon_k \downarrow 0$ such that $\{\varepsilon^{q-2} |r_\varepsilon|^{q-1}\}$ weakly converges in $L^{q/(q-1)}(Q)$ to r^* . Since spaces $L^{q/(q-1)}(Q)$ and $L^q(Q)$ are dual to each other, we may pass to the limit in (2.16), proving (2.23). Conversely, the sufficiency is obvious since, on admissible pairs $[y, u]$ satisfying (2.1)-(2.4), inequality (2.23) becomes

$$0 \leq \int_Q (y^* - y_d)(y - y^*) dxdt + N \int_{Q_\omega} |Bu^*|^{q-1} \operatorname{sgn}(Bu^*)(Bu - Bu^*) dxdt,$$

which immediately gives the optimality of $[y^*, u^*]$ by the definition of the subdifferential. \square

Remark 2.5. Notice the regularity (integrability) property of the Lagrange multiplier r^* .

Remark 2.6. Using (2.2), relation (2.23) may be rewritten as

$$0 \leq \int_Q (y^* - y_d)(y - y^*) dxdt + N \int_{Q_\omega} |Bu^*|^{q-1} \operatorname{sgn}(Bu^*)(Bu - Bu^*) dxdt + \int_Q r^*(y_t - \Delta y - Bu - y_t^* + \Delta y^* + Bu^*) dxdt.$$

When r^* is sufficiently smooth and $r^*(x, T) = 0$ in Ω , one can integrate by parts in the last integral. If I_D denotes the indicator function of the convex set D in $L^2(0, T; H_0^1(\Omega)) \times U$, then (2.23) may be rewritten as (B^* is the adjoint of B):

$$\left[y_d + y^* + r_t^* - \Delta r^*, -B^* r^* - NB^* \left(|Bu^*|^{q-1} \operatorname{sgn}(Bu^*) \right) \right] \in \partial I_D(y^*, u^*).$$

We denote by $\partial_1 I_D(y^*, u^*)$, $\partial_2 I_D(y^*, u^*)$ the two components of $\partial I_D(y^*, u^*)$ that occur above and we can write

$$\begin{aligned} r_t^* + \Delta r^* &\in y^* - y_D + \partial_1 I_D(y^*, u^*), \\ -B^* r^* &\in NB^* \left(|Bu^*|^{q-1} \operatorname{sgn}(Bu^*) \right) + \partial_2 I_D(y^*, u^*). \end{aligned}$$

This is the usual form of the optimality system, Barbu and Precupanu [2]. This formal interpretation may be made rigorous since r^* is the transposition solution of the above adjoint equation, Lions and Magenes [9, 10].

We next discuss the case when $N = 0$. In this case one typically expects that (a representative of) the optimal control u^* is piecewise continuous, i.e., continuous except for finitely many times (t_1, \dots, t_q) whose union is denoted \mathcal{T}^d . Reminding that $y^* \in W^{2,1,q}(Q) \subset C(\overline{Q})$ by the Sobolev embedding theorem, denote the *set of interior times* by

$$\mathcal{T} := \left\{ t \in [0, T] \setminus \mathcal{T}^d; \frac{1}{2} \int_{\Omega} (y^*(x, t))^2 dx < u^*(t) \right\}.$$

Then $Q_o := \Omega \times \mathcal{T}$ is an open subset. Since u^* is continuous over \mathcal{T} , for any $d \in \mathcal{D}(Q)$, with compact support in Q_o , and for $\delta \in R$ small enough, by the Weierstrass theorem, the pair $[y^* + \delta d, u^*]$ satisfies (2.9). Using theorem 2.4, we get

$$0 = \int_Q r^*(d_t - \Delta d) dx dt + \int_Q d(y^* - y_D) dx dt$$

and consequently

$$\begin{cases} r_t^* + \Delta r^* + j = y^* - y_D & \text{in } \mathcal{D}'(Q), \\ j \in \mathcal{D}'(Q) \text{ distribution with support in } \overline{Q} \setminus Q_o. \end{cases} \quad (2.24)$$

Note that

$$\overline{Q} \setminus Q_o = \left\{ (x, t) \in Q; t \in \mathcal{T}^d \text{ or } \frac{1}{2} \int_{\Omega} (y^*(x, t))^2 dx = u^*(t) \right\}.$$

Relation (2.24) is another well known form of the adjoint equation in the case when state or mixed constraints are present. Raymond and Arada [1], Rösch and Tröltzsch [15], De Los Reyes and Tröltzsch [8], studied the regularity properties of the multiplier j associated to the mixed constraint (2.4) under various interiority hypotheses.

Proposition 2.7. *Assume that u^* is piecewise continuous, the functions a and b are continuous, (2.17)-(2.19) hold, $N = 0$, and $y_d \in L^\infty(Q)$. Then, for all $t \in \mathcal{T}$, we have that*

$$\begin{cases} u^*(t) = a(t) & \text{in } \left\{ t \in \mathcal{T}; \int_{\Omega} f(x)r^*(x,t)dx < 0 \right\}, \\ u^*(t) = b(t) & \text{in } \left\{ t \in \mathcal{T}; \int_{\Omega} f(x)r^*(x,t)dx > 0 \right\}. \end{cases} \quad (2.25)$$

Proof. Let $t_o \in \mathcal{T}$ be a Lebesgue point of the function $t \rightarrow \int_{\Omega} f(x)r^*(x,t)dx$, such that $\int_{\Omega} f(x)r^*(x,t_o)dx < 0$. If $u^*(t_o) > a(t_o)$, since a and u^* are continuous at time t_o , define for $\eta > 0$ small enough

$$v^\eta(t) = \begin{cases} a(t_o) - u^*(t_o) & \text{if } |t - t_o| \leq \eta, \\ 0 & \text{otherwise.} \end{cases}$$

For small enough η , the pair $[y^*, u^* + v^\eta]$ belongs to D . By theorem 2.4, we have that

$$0 \leq (u^*(t_o) - a(t_o)) \int_{t_o-\eta}^{t_o+\eta} \left(\int_{\Omega} f(x)r^*(x,t)dx \right) dt. \quad (2.26)$$

Dividing (2.26) by η , and since t_o is a Lebesgue point of $\int_{\Omega} f(x)r^*(x,t)dx$, we obtain that $0 \leq \int_{\Omega} f(x)r^*(x,t_o)dx$, which is the desired contradiction. The second relation is proved in the same way. \square

3 OPTIMAL INVESTMENT AND STATE DOMINATED PROBLEMS

In this section, we discuss a variant of (2.1)-(2.4) corresponding in some sense to the "converse" of example (2.6):

$$\text{Min } \left\{ \int_Q F(x,t,y(x,t))dxdt + \int_Q (u + Nu^q)dxdt \right\}, \quad (3.1)$$

$$y_t - \Delta y + ay = u \quad \text{in } Q, \quad (3.2)$$

$$y(x, 0) = y_o(x) \text{ in } \Omega, \quad y(x, t) = 0 \text{ on } \Sigma, \quad (3.3)$$

$$0 \leq u(x, t) \leq cy(x, t) \quad \text{a.e. in } Q. \quad (3.4)$$

Here a and c are positive constants and $y_o \in W_o^{1,\infty}(\Omega) \cap W^{2,\infty}(\Omega)$, $y_o \geq 0$ a.e. in Ω and $y_o \not\equiv 0$ in Ω . The measurable function F is, for each (x, t) , convex and of class C^1 w.r.t. y , and such that $F(x, t, y(x, t))$ and $F_y(x, t, y(x, t))$ belong to $L^q(Q)$ for each continuous function y . A standard example is

$$F(x, t, y) = \mu(x, t)\pi(y), \quad (3.5)$$

where $\mu(x, t) > 0$ is an actualization coefficient, possibly depending on time, and $\pi : \mathbb{R}^+ \rightarrow \mathbb{R}$ is a *desutility function* (convex nonincreasing), and in that case the cost function can be interpreted as a compromise between the utility of y and the effort in resources u . The economic interpretation is as follows: $y(x, t) \geq 0$ is the capital at place x and time t . One cannot invest more than a fraction of the capital at every $(x, t) \in Q$. In addition there is a depreciation of the capital with constant rate a . Finally, the evolution of the capital depends also of what happens at neighbouring points, and this justifies the diffusion term. The cost function takes into account the preference for a certain type of evolution of the capital, and N can be viewed as a risk aversion coefficient (the preference for constant investment). Obviously there is a lot of freedom in the definition of the cost function. On the other hand, the problem has severe restrictions. If $u \in L^q(Q)$, then $y \in W^{2,1,q}(Q)$ and $y \in C(\bar{Q})$ if $q > \frac{1}{2}(n+2)$, $y \geq 0$ in Q if $u \in L^q(Q)_+$. The maximal state is obtained when taking $u = cy$, i.e., is solution of

$$\bar{y}_t - \Delta \bar{y} + (a - c)\bar{y} = 0 \quad \text{in } Q.$$

Therefore, if $c > a$, state decrease exponentially to zero, uniformly over the controls.

Remark 3.1. By the boundary conditions in (3.3), the constraint (3.4) excludes the standard "interiority" (Slater) assumptions used in the literature on control problems with state constraints. The interior of the set of feasible controls is also void, even in the $L^\infty(Q)$ topology. From this point of view, the constraint (3.4) is more difficult than (2.6).

Proposition 3.2. *The optimal control problem (3.1)-(3.4) has an optimal pair $[y^*, u^*] \in W^{2,1,q}(Q) \times L^\infty(Q)$.*

Proof. The control $u_o \equiv 0$ in Q , together with the corresponding solution of (3.1)-(3.3) is feasible. For any feasible pair $[y, u]$ in $W^{2,1,q}(Q) \times L^q(Q)$, we

have that $y \leq \bar{y}$, and hence $0 \leq u \leq \bar{u} := c\bar{y}$. So we have uniform bound on y and u in $W^{2,1,s}(Q)$ and $L^\infty(Q)$, resp. The usual passage to the limit in a minimizing sequence (using the fact that $L^\infty(Q)$ is the dual of the separable space $L^1(Q)$, and that an bounded sequence in the dual of a separable Banach space has a weakly * converging subsequence, and that the cost is l.s.c. for the weak * topology) allows to prove the existence of a solution to the problem (3.1)-(3.4). \square

The approximating problem is

$$\begin{aligned} \text{Min} \quad & \left\{ \int_Q F(x, t, y(x, t)) dxdt + \int_Q (u + Nu^q) dxdt + \frac{1}{q} \int_Q |u - u^*|^q dxdt + \right. \\ & \left. + \frac{1}{q\varepsilon} \int_Q |y_t - \Delta y + ay - u|^q dxdt \right\} \\ & \text{for all } [y, u] \in W^{2,1,q}(Q) \times L^q(Q), \text{ subject to (3.3)-(3.4).} \end{aligned} \quad (3.6)$$

This strongly convex problem has a unique solution $[y_\varepsilon, u_\varepsilon]$. Let $r_\varepsilon \in L^q(Q)$ be defined by

$$r_\varepsilon = \varepsilon^{-1}((y_\varepsilon)_t - \Delta y_\varepsilon + ay_\varepsilon - u_\varepsilon).$$

In the same way as in Section 2, we infer

Proposition 3.3. *The minimization problem (3.6) has a unique optimal pair $[y_\varepsilon, u_\varepsilon] \in W^{2,1,q}(Q) \times L^q(Q)$, $[y_\varepsilon, u_\varepsilon] \rightarrow (y^*, u^*)$ strongly in $W^{2,1,q}(Q) \times L^q(Q)$, and $\left\{ \varepsilon^{\frac{q-1}{q}} r_\varepsilon \right\}$ is bounded in $L^q(Q)$.*

Moreover, $(y_\varepsilon, u_\varepsilon)$ is characterized by the following relation: for any $(y, u) \in W^{2,1,q}(Q) \times L^q(Q)$ satisfying (3.3)-(3.4), we have that

$$\begin{aligned} 0 \leq & \int_Q F_y(x, t, y_\varepsilon(x, t))(y - y_\varepsilon) dxdt + \int_Q (1 + qNu_\varepsilon^{q-1})(u - u_\varepsilon) dxdt + \\ & + \int_Q |u_\varepsilon - u^*|^{q-1} \text{sgn}(u_\varepsilon - u^*)(u - u_\varepsilon) dxdt + \\ & + \int_Q \varepsilon^{q-2} |r_\varepsilon|^{q-1} \text{sgn}(r_\varepsilon)(y_t - \Delta y + ay - u) dxdt. \end{aligned} \quad (3.7)$$

Denote by $y^{oo} \in W^{2,1,q}(Q) \subset C(\bar{Q})$ the solution of (3.2)-(3.3) corresponding to $u_o \equiv 0$ in Q . We assume that y_o is non zero and that Ω is connected. It

follows that (see Protter and Weinberger [14]) that

$$y^{oo}(x, t) > 0, \quad \forall (x, t) \in Q. \quad (3.8)$$

Theorem 3.4. *Under hypothesis (3.8), the pair $(y^*, u^*) \in W^{2,1,q}(Q) \times L^q(Q)$ is optimal for problem (3.1)-(3.4) iff there exists $r^* \in M_{loc}(\overline{Q})$ such that*

$$\begin{aligned} 0 \leq & \int_Q F_y(x, t, y^*(x, t))(y - y^*) dxdt + \int_Q (1 + qN(u^*)^{q-1})(u - u^*) dxdt \\ & + \int_Q r^*(y_t - \Delta y + ay - u) dxdt, \end{aligned} \quad (3.9)$$

for any $(y, u) \in W^{2,1,q}(Q) \times L^q(Q)$ for which (3.3)-(3.4) hold, $y_t - \Delta y + ay - u \in L^\infty(Q)$, and there is $\mathcal{K} = \mathcal{K}_{y,u} \subset Q$ compact, such that

$$y_t - \Delta y + ay - u = 0 \quad \text{a.e. in } Q \setminus \mathcal{K}.$$

Remark 3.5. Here, $r^* \in M_{loc}(\overline{Q})$ means that for any $\mathcal{K} \subset Q$ compact, $r^* \in M(\mathcal{K})$, the dual of $L^\infty(\mathcal{K})$, i.e. $M_{loc}(\overline{Q}) = \cap \{L^\infty(\mathcal{K})^*, \mathcal{K} \subset Q, \text{ compact}\} \subset \mathcal{D}'(Q)$. Obviously, any admissible pair $[y, u]$ for (3.1)-(3.4) satisfies all the conditions on the test pairs in (3.9) since $y_t - \Delta y + ay - u = 0$ a.e. in Q by (3.2).

Proof. We show that $\varepsilon^{q-2} |r_\varepsilon|^{q-1}$ is bounded in $L^1_{loc}(Q)$. Let \mathcal{K} be a compact subset of Q , and let $\chi_{\mathcal{K}}$ denote its characteristic function. Take in (3.7) $\tilde{u} = \delta[\text{sgn } r_\varepsilon]_+ \chi_{\mathcal{K}}$ and the associated state denoted \tilde{y} , for small $\delta > 0$. The Weierstrass theorem and hypothesis (3.8) yields $y^{oo}|_{\mathcal{K}} \geq \alpha_{\mathcal{K}} > 0$. Then, the pair (y^{oo}, \tilde{u}) satisfies (3.3)-(3.4) and may be used in (3.7), if $\delta > 0$ is small enough.

By proposition 3.3 all terms except the last one in (3.7) are bounded independently of $\varepsilon > 0$ and we get

$$\delta \int_{\mathcal{K}} \varepsilon^{q-2} |r_\varepsilon|^{q-1} \text{sgn } r_\varepsilon [\text{sgn } r_\varepsilon]_+ dxdt \leq O(1), \quad \text{for all } \varepsilon > 0. \quad (3.10)$$

Take now $\hat{y} \in W^{2,1,q}(Q)$ to be the solution of (3.3) and

$$\hat{y}_t - \Delta \hat{y} + a\hat{y} = \chi_{\mathcal{K}} \quad \text{in } Q \quad (3.11)$$

and $\hat{u} \equiv 0$ in Q . Using the pair $[\hat{y}, \hat{u}]$ in (3.7), obtain

$$- \int_{\mathcal{K}} \varepsilon^{q-2} |r_\varepsilon|^{q-1} \text{sgn } r_\varepsilon dxdt \leq O(1), \quad \text{for all } \varepsilon > 0. \quad (3.12)$$

Multiplying (3.12) by $\delta > 0$ and adding (3.10) twice to it yields

$$\delta \int_{\mathcal{K}} \varepsilon^{q-2} |r_\varepsilon|^{q-1} |\operatorname{sgn} r_\varepsilon| dxdt \leq O(1), \quad \forall \varepsilon > 0,$$

where the $O(1)$ depend on \mathcal{K} . This proves that $\left\{ \varepsilon^{q-2} |r_\varepsilon|^{q-1} \right\}$ is bounded in $L^1_{loc}(Q)$. Next, for any compact subset \mathcal{K} of Q , we may define $r^*|_{L^\infty(\mathcal{K})} \in M(\mathcal{K})$ as the weak limit of $\varepsilon^{q-2} |r_\varepsilon|^{q-1}$ restricted to \mathcal{K} . Clearly, if $\hat{\mathcal{K}} \subset Q$ compact is such that $\mathcal{K} \subset \hat{\mathcal{K}}$, then the obtained limit extends the previous one as any element in $L^\infty(\mathcal{K})$ may be extended to $L^\infty(\hat{\mathcal{K}})$ by 0. In this way, we obtain $r^* \in M_{loc}(\overline{Q})$.

One can pass to the limit in (3.7) on any test pair $[y, u]$ satisfying the hypotheses of this theorem. This ends the proof of the necessity of (3.9). The sufficiency follows as in the previous section. \square

Corollary 3.6. *Assume that $N = 0$. Let Q_o be the interior of the set of points where u^* is continuous. Then*

$$\begin{cases} u^*(x, t) = cy^*(x, t) & \text{if } r^*(x, t) > 1, \text{ a.e. in } Q_o, \\ u^*(x, t) = 0 & \text{if } r^*(x, t) < 1, \text{ a.e. in } Q_o. \end{cases} \quad (3.13)$$

In addition, a.e. on Q_o , one of the three following statements hold:

$$\begin{cases} u^*(x, t) = cy^*(x, t) \text{ and } r^*(x, t) > 1, \\ u^*(x, t) = 0 \text{ and } r^*(x, t) < 1, \\ -a = F_y(x, t, y^*(x, t)). \end{cases} \quad (3.14)$$

Proof. Let $d \in \mathcal{D}(Q)$ have compact support in the open set

$$Q^* = \{(x, t) \in Q_o ; 0 \leq u^*(x, t) < cy^*(x, t)\}.$$

Then for λ close enough to 0, the pair $[y^* \pm \lambda d, u^*]$ may be taken in (3.9), and it follows by standard arguments that

$$r_t^* + \Delta r^* - ar^* + j = F_y(x, t, y^*(x, t)) \quad \text{in } \mathcal{D}'(Q) \quad (3.15)$$

where $j \in \mathcal{D}'(Q)$ is a distribution with support in $\overline{Q} \setminus Q^*$. Since $F_y(x, t, y^*)$ belongs to $L^q(Q)$, it follows that $r^* \in W_{loc}^{2,1,q}(Q^*) \subset C(Q^*)$. Take now $[y, u] \in D$ with $y = y^*$; since $N = 0$, (3.9) implies

$$0 \leq \int_Q (r^* - 1)(u^* - u) dxdt. \quad (3.16)$$

Let $(x_o, t_o) \in Q_o$ be such that $u^*(x_o, t_o) < cy^*(x_o, t_o)$. Then we may take $u = u^* + v$, with v nonnegative with small support near (x_o, t_o) , over which r^* is positive, and it follows with (3.16) that $\int_Q (r^* - 1)v \leq 0$, which gives the desired contradiction in the first relation of (3.13). The second one can be proved in the same way.

We next prove (3.14). By [7, p. 195], and in view of (3.15) we have that $F_y(x, t, y^*(x, t)) = -a$ a.e. over $\{(x, t) \in Q_o; r^* = 1\}$. Combining with (3.13), we obtain (3.14). \square

Remark 3.7. Note that relations (3.13) and (3.14) contain informations of different nature, and that neither of them implies the other one.

The previous result shows that, although the properties of r^* in Theorem 3.3 are very weak, (3.9) allows to obtain useful information on the optimal pair $[y^*, u^*]$.

Consider for instance the case when $\mu(x, t)$ has the constant value 1 and the desutility function is exponential, i.e. when $F(x, t, y) = e^{-y}$. Then $F_y(x, t, y^*) = -a$ iff $y^*(x, t) = -\log a$. Since $y^*(x, t)$ is positive over Q , this never occurs if $a \geq 1$. If $a \in (0, 1)$, since $y^* \in W_{loc}^{2,1,q}(Q_o)$, by the state equation and [7, p. 195], we obtain the additional information that $u^* = -a \log a$ a.e. on the set

$$\{(x, t) \in Q_o; y^*(x, t) = -\log a\}. \quad (3.17)$$

Therefore, in view (3.14), when $N = 0$, we have that

$$u^*(x, t) \in \{0, -a \log a, cy^*(x, t)\}, \quad \text{a.e. on } Q_o; \quad (3.18)$$

the same property holds on Q if u^* is continuous a.e.

Remark 3.8. Results such as the one in corollary 3.6 are called generalized bang-bang properties, see Tröltzsch [17], Bergounioux and Tiba [4]. Note that the sets where the constraints or relation $F_y(x, t, y^*(x, t)) = 0$ are active in Q need not be disjoint. In the work of Rösch and Tröltzsch [15], Hölder continuity properties are obtained for the optimal control in problems with mixed constraints, in a different setting.

REFERENCES

- [1] N. Arada and J.-P. Raymond. Optimal control problems with mixed control-state constraints. *SIAM J. Control Optim.*, 39(5):1391–1407 (electronic), 2000.

- [2] V. Barbu and Th. Precupanu. *Convexity and optimization in Banach spaces*, volume 10 of *Mathematics and its Applications (East European Series)*. D. Reidel Publishing Co., Dordrecht: D. Reidel, second edition, 1986.
- [3] R. Bellman. *Dynamic programming*. Princeton University Press, Princeton, 1961.
- [4] M. Bergounioux and D. Tiba. General optimality conditions for constrained convex control problems. *SIAM J. Control Optim.*, 34(2):698–711, 1996.
- [5] M. Bergounioux and F. Tröltzsch. Optimal control of linear bottleneck problems. *ESAIM Control Optim. Calc. Var.*, 3:235–250 (electronic), 1998.
- [6] M. Bergounioux and F. Tröltzsch. Optimal control of semilinear parabolic equations with state-constraints of bottleneck type. *ESAIM Control Optim. Calc. Var.*, 4:595–608 (electronic), 1999.
- [7] H. Brézis. *Analyse fonctionnelle*. Masson, Paris, 1983.
- [8] J.C. De los Reyes and F. Tröltzsch. Optimal control of the stationary Navier-Stokes equations with mixed control-state constraints. *SIAM J. Control Optim.*, 46(2):604–629 (electronic), 2007.
- [9] J.-L. Lions and E. Magenes. *Problèmes aux limites non homogènes et applications. Vol. 1*. Travaux et Recherches Mathématiques, No. 17. Dunod, Paris, 1968.
- [10] J.-L. Lions and E. Magenes. *Problèmes aux limites non homogènes et applications. Vol. 2*. Travaux et Recherches Mathématiques, No. 18. Dunod, Paris, 1968.
- [11] J.L. Lions. *Contrôle des systèmes distribués singuliers*. Dunod, Paris, 1983.
- [12] Ş. Mirică. Optimal feedback control for a class of bottleneck problems. *J. Math. Anal. Appl.*, 112(1):221–235, 1985.
- [13] P. Neittaanmäki and D. Tiba. *Optimal control of nonlinear parabolic systems*, volume 179 of *Monographs and Textbooks in Pure and Applied Mathematics*. Marcel Dekker Inc., New York, 1994. Theory, algorithms, and applications.

- [14] M.H. Protter and H.F. Weinberger. *Maximum principles in differential equations*. Springer-Verlag, New York, 1984. Corrected reprint of the 1967 original.
- [15] A. Rösch and F. Tröltzsch. On regularity of solutions and Lagrange multipliers of optimal control problems for semilinear equations with mixed pointwise control-state constraints. *SIAM J. Control Optim.*, 46(3):1098–1115 (electronic), 2007.
- [16] F. Tröltzsch. *Optimality conditions for parabolic control problems and applications*, volume 62 of *Teubner-Texte zur Mathematik [Teubner Texts in Mathematics]*. BSB B. G. Teubner Verlagsgesellschaft, Leipzig, 1984.
- [17] F. Tröltzsch. Regular Lagrange multipliers for control problems with mixed pointwise control-state constraints. *SIAM J. Optim.*, 15(2):616–634 (electronic), 2004/05.

*INRIA-Saclay, Centre de Mathématiques Appliquées,
Ecole Polytechnique, and
Laboratoire de Finance des Marchés d'Énergie, France
Frederic.Bonnans@inria.fr*

*Romanian Academy
Institute of Mathematics
P.O. Box 1-764, 014700 Bucharest
dan.tiba@imar.ro*