



COMPLEXITY BOUNDS FOR SECOND-ORDER OPTIMALITY  
IN UNCONSTRAINED OPTIMIZATION

by C. Cartis, N. I. M. Gould and Ph. L. Toint

Report NAXYS-11-2010

17 December 2010



University of Namur  
61, rue de Bruxelles, B5000 Namur (Belgium)

<http://www.fundp.ac.be/sciences/naxys>

# Complexity bounds for second-order optimality in unconstrained optimization

C. Cartis\*, N. I. M. Gould† and Ph. L. Toint‡

17 December 2010

## Abstract

This paper examines worst-case evaluation bounds for finding weak minimizers in unconstrained optimization. For the cubic regularization algorithm, Nesterov and Polyak (2006) and Cartis, Gould and Toint (2010a) show that at most  $O(\epsilon^{-3})$  iterations may have to be performed for finding an iterate which is within  $\epsilon$  of satisfying second-order optimality conditions. We first show that this bound can be derived for a version of the algorithm which only uses one-dimensional global optimization of the cubic model and that it is sharp. We next consider the standard trust-region method and show that a bound of the same type may also be derived for this method, and that it is also sharp in some cases. We conclude by showing that a comparison of the worst-case behaviour of the ARC and trust-region algorithms favours the first of these methods.

**Keywords:** evaluation complexity, worst-case analysis, nonconvex optimization, second-order optimality conditions.

## 1 Introduction

We consider algorithms for the solution of the unconstrained (possibly nonconvex) optimization problem

$$\min_x f(x) \tag{1.1}$$

where we assume that  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is smooth (in a sense to be specified later) and bounded below. All methods for the solution of (1.1) are iterative and, starting from some initial guess  $x_0$ , generate a sequence  $\{x_k\}$  of iterates approximating a critical point of  $f$ . Many such algorithms exist, and they are often classified according to their requirements in terms of computing derivatives of the objective function. In this paper, we focus on second-order methods, that is methods which evaluate the objective function  $f(x)$ , its gradient  $g(x)$  and its Hessian  $H(x)$  (or an approximation thereof) at every iteration. The advantage of these methods is that they can be expected to converge to solutions  $x_*$  satisfying the second-order optimality conditions

$$\nabla_x f(x_*) = 0, \text{ and } \lambda_{\min}(H(x_*)) \geq 0 \tag{1.2}$$

where  $\lambda_{\min}(A)$  is the smallest eigenvalue of the symmetric matrix  $A$ , rather than only satisfying first-order optimality (i.e., the first of these relations). In practice, however, a second-order algorithm is typically terminated as soon as an iterate  $x_k$  is found which is within  $\epsilon$  of satisfying (1.2), that is such that

$$\|\nabla_x f(x_*)\| \leq \epsilon_g \text{ and } \lambda_{\min}(H(x_*)) \geq -\epsilon_H, \tag{1.3}$$

---

\*School of Mathematics, University of Edinburgh, The King's Buildings, Edinburgh, EH9 3JZ, Scotland, UK. Email: coralia.cartis@ed.ac.uk

†Computational Science and Engineering Department, Rutherford Appleton Laboratory, Chilton, Oxfordshire, OX11 0QX, England, UK. Email: nick.gould@sftc.ac.uk

‡Namur Center for Complex Systems (NAXYS), FUNDP-University of Namur, 61, rue de Bruxelles, B-5000 Namur, Belgium. Email: philippe.toint@fundp.ac.be

for some user-specified tolerances  $\epsilon_g, \epsilon_H \in (0, 1)$ , where  $\|\cdot\|$  denotes the Euclidean norm. It is then of interest to bound the number of iterations which may be necessary to find an iterate satisfying (1.3) as a function of the thresholds  $\epsilon_g$  and  $\epsilon_H$ . It is the purpose of worst-case complexity analysis to derive such bounds. Many results are available in the literature for the case where the objective function  $f$  is convex (see, for instance, Nesterov 2004, 2008, Nemirovski, 1994, Agarwal, Bartlett, Ravikumar and Wainwright, 2009). The convergence to approximate first-order points in the nonconvex case has also been investigated for some time (see Vavasis 1992b, 1992a, 1993, Nesterov and Polyak, 2006, Gratton, Sartenaer and Toint, 2008, Cartis, Gould and Toint 2009a, 2010a, 2010b, 2010d, or Vicente, 2010).

Of particular interest here is the Adaptive Regularization with Cubics (ARC) algorithm independently proposed by Griewank (1981), Weiser, Deuffhard and Erdmann (2007) and Nesterov and Polyak (2006), whose worst-case complexity was shown in the last of these references to be of  $O(\epsilon_g^{-3/2})$  iterations for finding an iterate  $x_k$  satisfying the approximate first-order optimality conditions (the first relation in (1.3) only) and of  $O(\epsilon_H^{-3})$  iterations for finding an iterate  $x_k$  satisfying the whole of (1.3)<sup>(1)</sup>. These results were extended by Cartis et al. (2010a) to an algorithm no longer requiring the computation of exact second-derivatives (but merely of a suitably accurate approximation), nor an (also possibly approximate) knowledge of the objective function's Hessian's Lipschitz constant. More importantly, these authors showed that the  $O(\epsilon_g^{-3/2})$  complexity bound for convergence to first-order critical points can be achieved without requiring multi-dimensional global optimization of the cubic model (see Cartis, Gould and Toint, 2008). However, such a global minimization on nested Krylov subspaces of increasing dimensions was still required to obtain the  $O(\epsilon_H^{-3})$  convergence to second-order critical points.

The present paper focusses on worst-case complexity bounds for convergence to second-order critical points and shows that, as in the first-order case, multi-dimensional global minimization of the cubic model is unnecessary for obtaining the mentioned  $O(\epsilon_H^{-3})$  bound for the ARC algorithm. This latter bound is also shown to be sharp. We also prove that a bound of the same type holds for the standard trust-region method. Moreover, we show that it is also sharp for a range of relative values of  $\epsilon_g$  and  $\epsilon_H$ . We finally compare the known bounds for the ARC and trust-region algorithms and show that the ARC algorithm is always as good or better from this point of view.

The ARC algorithm is recalled in Section 2 and the associated complexity bounds are derived without multidimensional global minimization. Section 3 then discusses an example showing that the bound on convergence of the ARC algorithm to approximate second-order critical points is sharp. A bound of this type is derived in Section 4 for the trust-region methods, its sharpness for suitable values of  $\epsilon_g$  and  $\epsilon_H$  is demonstrated, and the comparison with the ARC algorithm discussed. Conclusions and perspectives are finally presented in Section 5.

## 2 The ARC algorithm and its worst-case complexity

The Adaptive Regularization with Cubics (ARC) algorithm is based on the approximate minimization, at iteration  $k$ , of the (possibly nonconvex) cubic model

$$m_k(s) = \langle g_k, s \rangle + \frac{1}{2} \langle s, B_k s \rangle + \frac{1}{3} \sigma_k \|s\|^3, \quad (2.1)$$

where  $\langle \cdot, \cdot \rangle$  denotes the Euclidean inner product. Here  $B_k$  is a symmetric  $n \times n$  approximation of  $H(x_k) \stackrel{\text{def}}{=} H_k$ ,  $\sigma_k > 0$  is a regularization weight and  $g_k = \nabla_x m_k(0) = \nabla_x f(x_k)$ . By ‘‘approximate minimization’’, we mean that a step  $s_k$  is computed to ensure the following conditions.

We first require that the step satisfies the conditions

$$\langle g_k, s_k \rangle + \langle s_k, B_k s_k \rangle + \sigma_k \|s_k\|^3 = 0 \quad (2.2)$$

and

$$\langle s_k, B_k s_k \rangle + \sigma_k \|s_k\|^3 \geq 0, \quad (2.3)$$

As noted in Cartis et al. (2010a), these conditions must hold if  $s_k$  is a global minimizer of  $m_k$  along the direction  $s_k$  (see Lemma 3.2 in Cartis et al., 2009a). In order to guarantee convergence to first-order

---

<sup>(1)</sup>It appears that this latter result is the first worst-case complexity bound for convergence to approximate second-order critical points ever proved.

critical points, we also require the familiar ‘‘Cauchy condition’’

$$m_k(s_k) \leq m_k(s_k^C) \quad (2.4)$$

with

$$s_k^C = -\alpha_k^C g_k \quad \text{and} \quad \alpha_k^C = \arg \min_{\alpha \geq 0} m_k(-\alpha g_k). \quad (2.5)$$

Because we are, in addition, interested in convergence to second-order critical points, we also require the following variant of the ‘‘eigen condition’’ whenever  $B_k$  is not positive semi-definite (see Section 6.6.1 in Conn, Gould and Toint, 2000): we require in that case that

$$m_k(s_k) \leq m_k(s_k^E), \quad (2.6)$$

where

$$s_k^E = \alpha_k^E u_k \quad \text{and} \quad \alpha_k^E = \arg \min_{\alpha} m_k(\alpha u_k), \quad (2.7)$$

with  $u_k$  being an approximate eigenvector of  $B_k$  associated with its smallest eigenvalue  $\lambda_{\min}(H_k) \stackrel{\text{def}}{=} \tau_k$ , in the sense that

$$\langle g_k, u_k \rangle \leq 0 \quad \text{and} \quad \langle u_k, B_k u_k \rangle \leq \kappa_{\text{snc}} \tau_k \|u_k\|^2 \quad (2.8)$$

for some constant  $\kappa_{\text{snc}} \in (0, 1]$ . The knowledge of  $\tau_k$  and  $u_k$  may be obtained, for instance, by applying the power method to  $B_k$ . Note that we require the minimization in (2.5) and (2.7) to be global, which means that (2.2) and (2.3) also hold with  $s_k$  replaced by  $s_k^C$  and  $s_k^E$ . Finally, we may also optionally require that

$$\|\nabla_x m_k(s_k)\| = \|g_k + B_k s_k + \sigma_k \|s_k\| s_k\| \leq \kappa_{\theta} \min[1, \|s_k\|] \|g_k\|, \quad (2.9)$$

for some given constant  $\kappa_{\theta} \in (0, 1)$  if we wish to accelerate the convergence to first-order critical points.

Remarkably, conditions (2.2)-(2.9) can all be ensured algorithmically and hold, in particular, if  $s_k$  is a (computable) global minimizer of  $m_k$  (see Griewank, 1981, Nesterov and Polyak, 2006, see also Cartis, Gould and Toint, 2009b). We also note that, if  $s_k$  is computed as the global minimizer of  $m_k$  in a subspace  $\mathcal{L}_k$  containing the gradient and satisfies (2.9), then all the above conditions also hold with  $u_k = Q_k w_k$ , where  $\tau_k$  and  $w_k$  are respectively the most negative eigenvalue of  $Q_k^T B_k Q_k$  and its corresponding eigenvector, and  $Q_k$  is an orthonormal basis of  $\mathcal{L}_k$ . We also note that they require global minimization of the cubic model along  $-g_k$ , (possibly)  $u_k$  and  $s_k$ , but that global minimization in subspaces of dimension larger than one is not necessary.

The ARC algorithm may then be stated as presented on the following page. In this description, we assume that the constants satisfy  $\gamma_2 \geq \gamma_1 > 1$ ,  $1 > \eta_2 \geq \eta_1 > 0$  and  $\sigma_0 > 0$ .

Let  $\mathcal{S}$  denote the index set of all successful or very successful iterations in the sense of (2.11), and define

$$\mathcal{S}_j = \{k \in \mathcal{S} \mid k \leq j\} \quad \text{and} \quad \mathcal{U}_j = \{0, \dots, j\} \setminus \mathcal{S}_j, \quad (2.12)$$

the sets of successful and unsuccessful iterations up to iteration  $j$ .

We now recall the main complexity results for this method, as well as the assumptions under which these hold. We first restate our assumptions.

**A.1:** The objective function  $f$  is twice continuously differentiable on  $\mathbb{R}^n$  and its gradient and Hessian are Lipschitz continuous on the path of iterates with Lipschitz constants  $L_g$  and  $L_H$ , i.e., for all  $k \geq 0$  and all  $\alpha \in [0, 1]$ ,

$$\|\nabla_x f(x_k) - \nabla_x f(x_k + \alpha s_k)\| \leq L_g \alpha \|s_k\| \quad (2.13)$$

and

$$\|\nabla_{xx} f(x_k) - \nabla_{xx} f(x_k + \alpha s_k)\| \leq L_H \alpha \|s_k\|. \quad (2.14)$$

**A.2:** The objective function  $f$  is bounded below, i.e. there exists a constant  $f_{\text{low}}$  such that, for all  $x \in \mathbb{R}^n$ ,

$$f(x) \geq f_{\text{low}}.$$

**Algorithm 2.1: ARC**

**Step 0:** A starting point  $x_0$ , an initial regularization parameter  $\sigma_0$ , and user-defined accuracy thresholds  $\epsilon_g, \epsilon_H \in (0, 1)$  are given. Set  $k = 0$ .

**Step 1:** If conditions (1.3) hold, terminate with approximate solution  $x_k$ .

**Step 2:** Compute a Hessian approximation  $B_k$  and a step  $s_k$  satisfying (2.2)–(2.9).

**Step 3:** Compute  $f(x_k + s_k)$  and

$$\rho_k = \frac{f(x_k) - f(x_k + s_k)}{-m_k(s_k)}. \quad (2.10)$$

Set  $x_{k+1} = x_k + s_k$  if  $\rho_k \geq \eta_1$ , or  $x_{k+1} = x_k$  otherwise.

**Step 4:** Set

$$\sigma_{k+1} \in \begin{cases} (0, \sigma_k] & \text{if } \rho_k > \eta_2, & \text{[very successful iteration]} \\ [\sigma_k, \gamma_1 \sigma_k] & \text{if } \eta_1 \leq \rho_k \leq \eta_2, & \text{[successful iteration]} \\ [\gamma_1 \sigma_k, \gamma_2 \sigma_k] & \text{otherwise.} & \text{[unsuccessful iteration]} \end{cases} \quad (2.11)$$

Increment  $k$  by one and return to Step 1.

**A.3:** For all  $k \geq 0$ , the Hessian approximation  $B_k$  satisfies

$$\|B_k\| \leq \kappa_B \quad (2.15)$$

and

$$\|(\nabla_{xx} f(x_k) - B_k)s_k\| \leq \kappa_{BH} \|s_k\|^2 \quad (2.16)$$

for some constants  $\kappa_B > 1$  and  $\kappa_{BH} > 0$ .

We start by noting that the form of the cubic model (2.1) and (2.2)–(2.3) ensure a remarkable bound on the the step norm and model decrease.

**Lemma 2.1** [Lemma 4.2 in Cartis et al. (2010a)] *We have that*

$$m_k(s_k) \leq -\frac{1}{6} \sigma_k \|s_k\|^3. \quad (2.17)$$

For our purposes it is also useful to consider the following bounds on the value of the regularization parameter.

**Lemma 2.2** *Suppose that (2.13) and (2.15) hold. Then there exists a constant  $\kappa_\sigma > 0$  independent of  $n$  such that, for all  $k \geq 0$*

$$\sigma_k \leq \max \left[ \sigma_0, \frac{\kappa_\sigma}{\epsilon_g} \right]. \quad (2.18)$$

*If, in addition, (2.14) and (2.16) also hold, then there exists a constant  $\sigma_{\max} > 0$  independent of  $n$ ,  $\epsilon_g$  and  $\epsilon_H$  such that, for all  $k \geq 0$ ,*

$$\sigma_k \leq \sigma_{\max}. \quad (2.19)$$

**Proof.** See Lemmas 3.2 and 3.3 in Cartis et al. (2010a) for the proof of (2.18) and Lemma 5.2 in Cartis et al. (2009a) for that of (2.19).  $\square$

A first complexity bound can then be derived.

**Lemma 2.3** [Corollary 3.4 in Cartis et al. (2010a)] Assume that (2.13), **A.2** and (2.15) hold. Then there exists a constant  $\kappa_{\text{ARC},S}^0 > 0$  independent of  $n$  such that  $N_{\text{ARC},S}^1$ , the total number of successful and very successful iterations of the ARC algorithm with  $\|g_k\| \geq \epsilon_g$ , is bounded above by  $\lceil \kappa_{\text{ARC},S}^0 \epsilon_g^{-2} \rceil$ .

If we are ready to strengthen our assumption by assuming (2.14) and to impose (2.9), then, crucially, the step  $s_k$  can then be proved to be sufficiently long compared to the gradient's norm at iteration  $k+1$ .

**Lemma 2.4** [Lemma 5.2 in Cartis et al. (2010a)] Suppose that **A.1**, **A.3** and (2.9) hold. Then, for all  $k \geq 0$ , one has that, for some  $\kappa_g > 0$  independent of  $n$ ,

$$\|s_k\| \geq \kappa_g \sqrt{\|\nabla_x f(x_k + s_k)\|}. \quad (2.20)$$

Combining (2.17) with this last result, it is then not difficult to show the second complexity result.

**Lemma 2.5** [Corollary 5.3 in Cartis et al. (2010a)] Suppose that **A.1–A.3** and (2.9) hold. Suppose also that

$$\sigma \geq \sigma_{\min} \quad (2.21)$$

for some  $\sigma_{\min} > 0$ . Then there exists a constant  $\kappa_{\text{ARC},S}^1 > 0$  independent of  $n$  such that  $N_{\text{ARC},S}^1$  is bounded above by  $\lceil \kappa_{\text{ARC},S}^1 \epsilon_g^{-3/2} \rceil$ .

The final important observation in the first-order analysis is that the total number of iterations required by the ARC algorithm to terminate may be bounded in terms of the number of successful iterations needed.

**Lemma 2.6** [Theorem 2.1 in Cartis et al. (2010a)] Suppose that (2.21) holds and, for any fixed  $j \geq 0$ , let  $\mathcal{S}_j$  and  $\mathcal{U}_j$  be defined in (2.12). Then one has that

$$|\mathcal{U}_j| \leq \left\lceil (|\mathcal{S}_j| + 1) \frac{1}{\log \gamma_1} \log \left( \frac{\sigma_{\max}}{\sigma_{\min}} \right) \right\rceil. \quad (2.22)$$

We may now use this last result with Lemmas 2.3 and 2.5, and deduce the following worst-case bounds.

**Theorem 2.7** [See Corollary 5.5 in Cartis et al. (2010a)] Suppose that (2.13), **A.2** and (2.15) hold. Suppose also that there exist a  $\sigma_{\min} > 0$  such that (2.21) hold for all  $k \geq 0$ . Then, the ARC algorithm produces an iterate  $x_k$  satisfying the first part of (1.3) after at most

$$\lceil \kappa_{\text{ARC},S}^{\text{1st}} \epsilon_g^{-2} \rceil \quad (2.23)$$

successful iterations and at most

$$\lceil \kappa_{\text{ARC}}^{\text{1st}} \epsilon_g^{-2} \rceil \quad (2.24)$$

iterations in total, where  $\kappa_{\text{ARC},S}^{\text{1st}}$  and  $\kappa_{\text{ARC}}^{\text{1st}}$  are positive constants independent of  $n$ . Moreover, if (2.14) and (2.9) also hold, then the bounds (2.23) and (2.24) respectively become

$$\lceil \kappa_{\text{ARC},S}^{\text{1st}} \epsilon_g^{-3/2} \rceil \quad \text{and} \quad \lceil \kappa_{\text{ARC}}^{\text{1st}} \epsilon_g^{-3/2} \rceil. \quad (2.25)$$

The bounds (2.25) are known to be qualitatively<sup>(2)</sup> tight and optimal for a wide class of second-order methods (see Cartis et al. 2010b, 2010d).

After reviewing the complexity of convergence to first-order critical points, we now turn to the analysis of the number of iterations necessary to ensure the second part of (1.3) under our present assumptions (which do not require multi-dimensional global model minimization).

**Lemma 2.8** Suppose that **A.1–A.3** hold. Then there exists a constant  $\kappa_S^2 > 0$  independent of  $n$  such that  $N_{\text{ARC},S}^2$ , the total number of successful and very successful iterations of the ARC algorithm with  $\tau_k < -\epsilon_H$ , is bounded above by  $\lceil \kappa_{\text{ARC},S}^2 \epsilon_H^{-3} \rceil$ .

<sup>(2)</sup>The constants may not be optimal.

**Proof.** We first note that, when  $\tau_k < 0$ ,  $s_k^E$  gives a minimizer of the model in the direction  $u_k$  by (2.7), from which we derive, using (2.3) for  $s_k^E$ , that, for all  $k \geq 0$ ,

$$\sigma_k \|s_k^E\| \geq -\frac{\langle s_k^E, B_k s_k^E \rangle}{\|s_k^E\|^2} \geq \kappa_{\text{snc}} |\tau_k|, \quad (2.26)$$

where we have used (2.7) to derive the last inequality. Combining this bound with (2.17) applied for  $s_k^E$ , and (2.19), we then obtain that

$$-m_k(s_k) \geq -m_k(s_k^E) \geq \frac{\kappa_{\text{snc}} |\tau_k|^3}{6\sigma_k^2} \geq \frac{\kappa_{\text{snc}}}{6\sigma_{\max}^2} |\tau_k|^3 \geq \frac{\kappa_{\text{snc}} \epsilon_H^3}{6\sigma_{\max}^2} \quad (2.27)$$

for all  $k$  such that the second part of (1.3) fails. If we now restrict our attention to the subset of those iterations which are successful or very successful, we obtain, using **A.2** and the monotonically decreasing nature of the sequence  $\{f(x_k)\}$ , that

$$f(x_0) - f_{\text{low}} \geq \sum_{k=0, k \in \mathcal{S}} (f(x_k) - f(x_{k+1})) \geq N_{\text{ARC}, \mathcal{S}}^2 \frac{\eta_1 \kappa_{\text{snc}} \epsilon_H^3}{6\sigma_{\max}^2}.$$

We therefore obtain the desired result with  $\kappa_{\text{ARC}, \mathcal{S}}^2 \stackrel{\text{def}}{=} 6\sigma_{\max}^2 (f(x_0) - f_{\text{low}}) / \kappa_{\text{snc}} \eta_1$ .  $\square$

As was the case for convergence to first-order critical points, we may now combine Lemmas 2.3 and 2.6 with our last result to obtain worst-case complexity bounds for convergence of the ARC algorithm to approximate second-order critical points.

**Theorem 2.9** *Suppose that **A.1–A.3** hold. Suppose also that there exist a  $\sigma_{\min} > 0$  such that (2.21) hold for all  $k \geq 0$ . Then, the ARC algorithm produces an iterate  $x_k$  satisfying (1.3) (and thus terminates) after at most*

$$\lceil \kappa_{\text{ARC}, \mathcal{S}}^{2\text{nd}} \max[\epsilon_g^{-2}, \epsilon_H^{-3}] \rceil \quad (2.28)$$

*successful or very successful iterations and at most*

$$\lceil \kappa_{\text{ARC}}^{2\text{nd}} \max[\epsilon_g^{-2}, \epsilon_H^{-3}] \rceil \quad (2.29)$$

*iterations in total, where  $\kappa_{\text{ARC}, \mathcal{S}}^{2\text{nd}}$  and  $\kappa_{\text{ARC}}^{2\text{nd}}$  are positive constants independent of  $n$ . Moreover, if (2.9) also holds, then the bounds (2.28) and (2.29) respectively become*

$$\lceil \kappa_{\text{ARC}, \mathcal{S}}^{2\text{nd}} \max[\epsilon_g^{-3/2}, \epsilon_H^{-3}] \rceil \quad \text{and} \quad \lceil \kappa_{\text{ARC}}^{2\text{nd}} \max[\epsilon_g^{-3/2}, \epsilon_H^{-3}] \rceil. \quad (2.30)$$

**Proof.** Lemmas 2.3 and 2.8 yield that the total number of successful iterations such that the first or the second part of (1.3) is violated cannot exceed

$$\kappa_{\text{ARC}, \mathcal{S}}^0 \epsilon_g^{-2} + \kappa_{\text{ARC}, \mathcal{S}}^2 \epsilon_H^{-3}.$$

We thus immediately deduce (2.28) with  $\kappa_{\text{ARC}, \mathcal{S}}^{2\text{nd}} \stackrel{\text{def}}{=} \kappa_{\text{ARC}, \mathcal{S}}^0 + \kappa_{\text{ARC}, \mathcal{S}}^2$ . The bound (2.29) follows by applying Lemma 2.6, while (2.30) directly is obtained by using Lemma 2.5 instead of Lemma 2.3 in this reasoning.  $\square$

### 3 An example of slow convergence of ARC

We now show by an example that the bounds (2.28) and (2.29) cannot be improved. Our example is unidimensional and is inspired by the technique used in Cartis et al. (2010b) and Cartis et al. (2010d).

We first choose the starting point and sequences of gradient and Hessian values and steps to be, for all  $k \geq 0$ ,

$$x_0 = 0, \quad g_k = 0, \quad s_k = \left( \frac{1}{k+1} \right)^{\frac{1}{3} + \delta} \quad \text{and} \quad B_k = H_k = \tau_k = - \left( \frac{1}{k+1} \right)^{\frac{1}{3} + \delta} \quad (3.1)$$

where  $\delta \in (0, 1)$  is a (small) positive constant. Because it is straightforward to verify that the conditions (2.2)–(2.9) hold with this choice and  $\sigma_k = 1$  for all  $k$ , we may consider these values as produced by the  $k$ -th iteration of the ARC algorithm at iterate  $x_k = x_0 + \sum_{j=0}^{k-1} s_j$ . We also define  $f_k \stackrel{\text{def}}{=} f(x_k)$  for all  $k$  by the relations

$$f_0 = \zeta(1 + 3\delta) \quad \text{and} \quad f_{k+1} = f_k - \left( \frac{1}{k+1} \right)^{1+3\delta}, \quad (3.2)$$

where  $\zeta(t) \stackrel{\text{def}}{=} \sum_{k=1}^{\infty} k^{-t}$  is the Riemann zeta function, which is finite for all  $t > 1$  (and thus for  $t = 1 + 3\delta$ ). Observe that, since (2.2) and (2.3) both hold as equalities, we have that

$$-m_k(s_k) = \frac{1}{6} \|s_k\|^3 = \frac{1}{6} \left( \frac{1}{k+1} \right)^{1+3\delta}$$

and (3.2) therefore implies that all iterations are very successful, allowing us to keep  $\sigma_k$  fixed to 1.

We now use Hermite interpolation to construct the objective function  $f$  on the successive intervals  $[x_k, x_{k+1}]$ , and define

$$f(x) = p_k(x - x_k) + f_{k+1} \quad \text{for } x \in [x_k, x_{k+1}] \quad \text{and } k \geq 0, \quad (3.3)$$

where  $p_k$  is the polynomial

$$p_k(s) = c_{0,k} + c_{1,k}s + c_{2,k}s^2 + c_{3,k}s^3 + c_{4,k}s^4 + c_{5,k}s^5,$$

with coefficients defined by the interpolation conditions

$$\begin{aligned} p_k(0) &= f_k - f_{k+1}, & p_k(s_k) &= 0; \\ p'_k(0) &= g_k, & p'_k(s_k) &= g_{k+1}; \\ p''_k(0) &= H_k, & p''_k(s_k) &= H_{k+1}. \end{aligned} \quad (3.4)$$

These conditions yield the following values for the first three coefficients

$$c_{0,k} = f_k - f_{k+1}, \quad c_{1,k} = g_k = 0, \quad c_{2,k} = \frac{1}{2}H_k;$$

and the remaining coefficients satisfy

$$\begin{pmatrix} s_k^3 & s_k^4 & s_k^5 \\ 3s_k^2 & 4s_k^3 & 5s_k^4 \\ 6s_k & 12s_k^2 & 20s_k^3 \end{pmatrix} \begin{pmatrix} c_{3,k} \\ c_{4,k} \\ c_{5,k} \end{pmatrix} = \begin{pmatrix} \Delta f_k - g_k s_k - \frac{1}{2} s_k H_k s_k \\ \Delta g_k - H_k s_k \\ \Delta H_k \end{pmatrix},$$

where

$$\Delta f_k = f_{k+1} - f_k, \quad \Delta g_k = g_{k+1} - g_k \quad \text{and} \quad \Delta H_k = H_{k+1} - H_k.$$

Hence we obtain, also from (3.1) and  $\Delta g_k = 0$ , that

$$\begin{aligned} c_{3,k} &= 10 \frac{\Delta f_k}{s_k^3} - 4 \frac{\Delta g_k}{s_k^2} + \frac{\Delta H_k}{2s_k} - 10 \frac{g_k}{s_k^2} - \frac{H_k}{s_k} = 10 \frac{\Delta f_k}{s_k^3} + \frac{\Delta H_k}{2s_k} - \frac{H_k}{s_k}; \\ c_{4,k} &= -15 \frac{\Delta f_k}{s_k^4} + 7 \frac{\Delta g_k}{s_k^3} - \frac{\Delta H_k}{s_k^2} + 15 \frac{g_k}{s_k^3} + \frac{H_k}{2s_k^2} = -15 \frac{\Delta f_k}{s_k^4} - \frac{\Delta H_k}{s_k^2} + \frac{H_k}{2s_k^2}; \\ c_{5,k} &= 6 \frac{\Delta f_k}{s_k^5} - 3 \frac{\Delta g_k}{s_k^4} + \frac{\Delta H_k}{2s_k^3} - 6 \frac{g_k}{s_k^4} = 6 \frac{\Delta f_k}{s_k^5} + \frac{\Delta H_k}{2s_k^3}. \end{aligned} \quad (3.5)$$

It remains to show that the constructed  $f$  satisfies **A.1–A.3**. One easily sees from its construction that  $f$  is twice continuously differentiable. Moreover its third derivative exists everywhere and, on the  $k$ -th interval, satisfies the bound

$$|f'''(x_k + s)| = |p_k'''(s)| \leq 6|c_{3,k}| + 24|c_{4,k}|s_k + 60|c_{5,k}|s_k^2, \quad s \in [0, s_k].$$

But (3.1), (3.5) and the resulting inequality  $|\Delta H_k| \leq |H_k|$  imply that  $|c_{3,k}|$ ,  $|c_{4,k}|s_k$  and  $|c_{5,k}|s_k^2$  are uniformly bounded, and thus so is  $f'''$ . As a consequence,  $f$  has a Lipschitz continuous Hessian and

**A.1** holds. The definition (3.2) and the definition of the Riemann function together imply that **A.2** holds with  $f_{\text{low}} = 0^{(3)}$  and **A.3** directly results from (3.1). Figure 3.1 shows plots of  $f$  and its first three derivatives for  $\delta = 0.0001$  and for  $k = 0, \dots, 15$ . The figure reveals the objective functions's nonconvexity and monotonically decreasing nature.




Figure 3.1: The function  $f$  and its first three derivatives (from top to bottom and left to right) on the first 16 intervals

We have thus verified that the ARC algorithm applied on  $f$  (which satisfies **A.1–A.3**) starting from  $x_0 = 0$  and  $\sigma_0 = 1$  produces iterates such that

$$\lambda_{\min}(H_k) = - \left( \frac{1}{k+1} \right)^{\frac{1}{3} + \delta}$$

and for which the second part of (1.3) fails for exactly

$$\left\lceil \frac{1}{\epsilon_H^{3(1+\delta)}} \right\rceil - 1$$

iterations, for any  $\epsilon_H, \epsilon_g$  and  $\delta$  in  $(0, 1)$ . Since we know from Cartis et al. (2010b) that the bound in  $O(\epsilon_g^{-3/2})$  is sharp for obtaining a mere first-order approximate critical point, we deduce that the bound (2.30) cannot be improved. As a consequence it is sharp as far as the ARC algorithm is concerned.

## 4 Second-order complexity for the trust-region method

We may wonder if the worst-case complexity for convergence to approximate second-order points is better or worse for the standard trust-region method than for ARC. We show in this section that they

<sup>(3)</sup>Note that we have shown that  $f(x)$  is bounded below for  $x \geq 0$ , which is the domain of interest since  $x_k \geq 0$ ; we may extend  $f$  by continuity for  $x < 0$ .

are (qualitatively) identical. Our first step is to establish an upper bound on this complexity for the trust-region method, which requires revisiting some of its convergence theory. For the sake of completeness, we briefly recall the basic formulation of this method, as based on Section 6.1 of Conn et al. (2000). The main idea of the trust-region method is similar to that of the ARC algorithm: at iteration  $k$ , a quadratic model

$$m_k(s) \stackrel{\text{def}}{=} \langle g_k, s \rangle + \frac{1}{2} \langle s, B_k s \rangle \quad (4.1)$$

is minimized in the “trust region” defined by

$$\mathcal{B}_k \stackrel{\text{def}}{=} \{s \in \mathbb{R}^n \mid \|s\| \leq \Delta_k\}, \quad (4.2)$$

where  $\Delta_k$  is the (dynamically updated) trust-region radius. The other conditions on the step  $s_k$  are again similar to what happens for the ARC method: one typically requires  $s_k$  to satisfy (2.4)–(2.8) where the model  $m_k(s)$  is now defined by (4.1) instead of (2.1) and where minimization in (2.4) and (2.6) is restricted to the trust-region. Note that, in this context, global optimization of the model along  $s_k^C$  or  $s_k^E$  within the trust region no longer implies (2.2) and (2.3). In practice, the condition (2.9) is often replaced by

$$\|\nabla_x m_k(s_k)\| = \|g_k + B_k s_k\| \leq \kappa_\theta \min[1, \|g_k\|^\alpha] \|g_k\|, \quad (4.3)$$

for some given constant  $\kappa_\theta \in (0, 1)$  and some exponent  $\alpha > 0$ , but this is irrelevant for the complexity analysis developed below. Global optimization of the model along  $s_k$  within the trust region is not necessary.

The basic trust-region algorithm may then be stated as follows.

#### Algorithm 4.1: Trust-region algorithm

**Step 0:** A starting point  $x_0$ , an initial radius  $\Delta_0 > 0$  and user-defined accuracy thresholds  $\epsilon_g, \epsilon_H \in (0, 1)$  are given. Set  $k = 0$ .

**Step 1:** If conditions (1.3) holds, terminate with approximate solution  $x_k$ .

**Step 2:** Compute a Hessian approximation  $B_k$  and a step  $s_k \in \mathcal{B}_k$  satisfying (2.4)–(2.8) and (optionally) (4.3).

**Step 3:** Compute  $f(x_k + s_k)$  and  $\rho_k$  given by (2.10). Set  $x_{k+1} = x_k + s_k$  if  $\rho_k \geq \eta_1$ , or  $x_{k+1} = x_k$  otherwise.

**Step 4:** Set

$$\Delta_{k+1} \in \begin{cases} [\Delta_k, \gamma_3 \Delta_k] & \text{if } \rho_k > \eta_2, & \text{[very successful iteration]} \\ [\gamma_2 \Delta_k, \Delta_k] & \text{if } \eta_1 \leq \rho_k \leq \eta_2, & \text{[successful iteration]} \\ [\gamma_1 \Delta_k, \gamma_2 \Delta_k] & \text{otherwise.} & \text{[unsuccessful iteration]} \end{cases} \quad (4.4)$$

Increment  $k$  by one and return to Step 1.

In this algorithm, we have assumed that the constants satisfy the inequalities

$$\Delta_0 \leq \Delta_{\max}, \quad 0 < \eta_1 \leq \eta_2 < 1 \quad \text{and} \quad 0 < \gamma_1 \leq \gamma_2 < 1 \leq \gamma_3 \quad (4.5)$$

and we define the sets of very successful, successful and unsuccessful iterations just as in (2.12).

In order to establish the desired complexity bound, we start by re-examining the size of the discrepancy between the model and the objective function in the case where Lipschitz continuity of the Hessian is assumed (an assumption never made in Chapter 6 of Conn et al., 2000).

**Lemma 4.1** *Suppose that A.1 and A.3 hold. Then, for each  $k \geq 0$ ,*

$$|f(x_k + s_k) - m_k(s_k)| \leq \kappa_{\text{fm}} \Delta_k^3. \quad (4.6)$$

for some  $\kappa_{\text{fm}} > 0$ .

**Proof.** (See the proof of Lemma 6.4.1 in Conn et al., 2000.) Using A.1, we may apply the mean-value theorem on the objective function and obtain that

$$f(x_k + s_k) = f(x_k) + \langle g_k, s_k \rangle + \frac{1}{2} \langle s_k, H(\xi_k) s_k \rangle$$

for some  $\xi_k$  in the segment  $[x_k, x_k + s_k]$ . Subtracting (4.1), taking absolute values and using A.1, A.3, the inequality  $\|\xi_k - x_k\| \leq \|s_k\|$  and the Cauchy-Schwarz inequality yields that

$$\begin{aligned} |f(x_k + s_k) - m_k(s_k)| &= \frac{1}{2} |\langle s_k, H(\xi_k) s_k \rangle - \langle s_k, B_k s_k \rangle| \\ &\leq \frac{1}{2} |\langle s_k, [H(\xi_k) - H(x_k) + H(x_k) - B_k] s_k \rangle| \\ &\leq \frac{1}{2} [L_H \|s_k\|^3 + \kappa_{\text{BH}} \|s_k\|^3] \end{aligned}$$

and (4.6) with  $\kappa_{\text{fm}} = \frac{1}{2}(L_H + \kappa_{\text{BH}})$  then follows from the inequality  $\|s_k\| \leq \Delta_k$ .  $\square$

We then recall a standard result on the model decrease in the presence of significant gradient or negative curvature.

**Lemma 4.2** [Theorems 6.3.1 and 6.6.1 in Conn et al. (2000)] *Suppose that  $m_k$  is given by (4.1). Then, if  $\|g_k\| > 0$ , we have that*

$$-m_k(s_k) \geq -m_k(s_k^C) \geq \frac{1}{2} \|g_k\| \min \left[ \frac{\|g_k\|}{\kappa_{\text{B}}}, \Delta_k \right] \quad (4.7)$$

while, if  $\tau_k < 0$ , then

$$-m_k(s_k) \geq -m_k(s_k^E) \geq \frac{1}{2} \kappa_{\text{snc}} |\tau_k| \Delta_k^2. \quad (4.8)$$

From this result, we may deduce the following crucial lemma.

**Lemma 4.3** *Suppose that A.1 and A.2 hold and that  $m_k$  is given by (4.1). Suppose furthermore that  $\tau_k < 0$  and that*

$$\Delta_k \leq \frac{(1 - \eta_2) \kappa_{\text{snc}} |\tau_k|}{2 \kappa_{\text{fm}}}. \quad (4.9)$$

Then iteration  $k$  of the trust-region algorithm is very successful and  $\Delta_{k+1} \geq \Delta_k$ .

**Proof.** Suppose that (4.9) holds. We obtain from (4.6) and (4.8) that

$$|\rho_k - 1| = \left| \frac{f(x_k + s_k) - m_k(s_k)}{-m_k(s_k)} \right| \leq \frac{\kappa_{\text{fm}}}{\frac{1}{2} \kappa_{\text{snc}} |\tau_k|} \Delta_k \leq 1 - \eta_2,$$

where we used (4.9) to deduce the last inequality. Thus  $\rho_k \geq \eta_2$  and the mechanism of the trust-region algorithm then ensures that iteration  $k$  is very successful and, by (4.4), that  $\Delta_{k+1} \geq \Delta_k$ .  $\square$

We may then use this result to show that, as long as second-order optimality is not reached in the sense of (1.3), then the trust-region radius is bounded away from zero. To make our result more precise we first observe that

$$\text{either } \|g_k\| \geq \epsilon_g \quad \text{or} \quad -\min(0, \tau_k) \geq \epsilon_H \quad (4.10)$$

as long as the trust-region algorithm does not terminate.

**Lemma 4.4** *Suppose that A.1 and A.2 hold and that  $m_k$  is given by (4.1). Then, there exists a constant  $\kappa_{\Delta} \in (0, 1)$  independent of  $n$  such that, if the trust-region algorithm does not terminate at iteration  $k$ ,*

$$\Delta_k \geq \kappa_{\Delta} \min[\epsilon_g, \epsilon_H]. \quad (4.11)$$

**Proof.** Assume, for the purpose of deriving a contradiction, that iteration  $k$  is the first such that

$$\Delta_{k+1} \leq \gamma_1 \min \left[ \frac{1}{2\kappa_B}, \frac{\kappa_{\text{snc}}}{2\kappa_{\text{fm}}} \right] (1 - \eta_2) \min [\epsilon_g, \epsilon_H]. \quad (4.12)$$

Then we have from (4.4) that, either

$$\Delta_k \leq \min \left[ \frac{1}{2\kappa_B}, \frac{\kappa_{\text{snc}}}{2\kappa_{\text{fm}}} \right] (1 - \eta_2) \min [\epsilon_g, \epsilon_H] \leq \frac{(1 - \eta_2)}{2\kappa_B} \epsilon_g \leq \frac{(1 - \eta_2)}{2\kappa_B} \|g_k\|$$

if the first part of (4.10) holds, or

$$\Delta_k \leq \min \left[ \frac{1}{2\kappa_B}, \frac{\kappa_{\text{snc}}}{2\kappa_{\text{fm}}} \right] (1 - \eta_2) \min [\epsilon_g, \epsilon_H] \leq \frac{(1 - \eta_2)\kappa_{\text{snc}}}{2\kappa_{\text{fm}}} \epsilon_H \leq \frac{(1 - \eta_2)\kappa_{\text{snc}}|\tau_k|}{2\kappa_{\text{fm}}}.$$

if the second part of (4.10) holds. In the first case, Theorem 6.4.3 in Conn et al. (2000) implies that iteration  $k$  is very successful and  $\Delta_{k+1} \geq \Delta_k$ . In the second case, the same conclusion follows from Lemma 4.3. Thus  $\Delta_{k+1} \geq \Delta_k$  in both cases and our assumption that iteration  $k$  is the first such that (4.12) holds must be false. As a consequence, there cannot be any iteration such that inequality (4.12) holds as long as the algorithm does not terminate, and we obtain the desired conclusion with

$$\kappa_\Delta \stackrel{\text{def}}{=} \gamma_1 \min \left[ \frac{1}{2\kappa_B}, \frac{\kappa_{\text{snc}}}{2\kappa_{\text{fm}}} \right] (1 - \eta_2) < 1, \quad (4.13)$$

the last inequality following from the bound  $\kappa_B \geq 1$  and (4.5).  $\square$

We may now compute an upper bound on the number of successful or very successful iterations such that (1.3) does not hold.

**Lemma 4.5** *Suppose that A.1 and A.2 hold and that  $m_k$  is given by (4.1). Then there exists a constant  $\kappa_{\text{TR,S}}^{2\text{nd}} > 0$  independent of  $n$  such that  $N_{\text{TR,S}}^{2\text{nd}}$ , the number of successful or very successful iterations of the trust-region method before (1.3) holds, is bounded above by  $\lceil \kappa_{\text{TR,S}}^{2\text{nd}} \max [\epsilon_g^{-2} \epsilon_H^{-1}, \epsilon_H^{-3}] \rceil$ .*

**Proof.** Consider an iteration  $k$  of the trust-region algorithm (before it terminates). Then either  $\|g_k\| > \epsilon_g$  or  $\tau_k < -\epsilon_H$ . In the first of these cases, (4.7), (4.11) and (4.13) yield that

$$-m_k(s_k) \geq \frac{1}{2} \epsilon_g \min \left[ \frac{\epsilon_g}{\kappa_B}, \kappa_\Delta \min [\epsilon_g, \epsilon_H] \right] = \frac{1}{2} \kappa_\Delta \epsilon_g \min [\epsilon_g, \epsilon_H],$$

while we obtain, in the second case, that

$$-m_k(s_k) \geq \frac{1}{2} \kappa_{\text{snc}} |\tau_k| \Delta_k^2 \geq \frac{1}{2} \kappa_{\text{snc}} \kappa_\Delta^2 \epsilon_H \min [\epsilon_g, \epsilon_H]^2$$

from (4.8) and (4.11). We thus obtain, using A.2 and the monotonically decreasing nature of the sequence  $\{f(x_k)\}$ , that

$$\begin{aligned} f(x_0) - f_{\text{low}} &\geq \sum_{k=0}^{\infty} [f(x_k) - f(x_{k+1})] \\ &\geq \sum_{k=0, k \in \mathcal{S}} [f(x_k) - f(x_{k+1})] \\ &\geq \frac{1}{2} \eta_1 \sum_{k=0, k \in \mathcal{S}} \min \left[ \kappa_\Delta \epsilon_g \min [\epsilon_g, \epsilon_H], \kappa_{\text{snc}} \kappa_\Delta^2 \epsilon_H \min [\epsilon_g, \epsilon_H]^2 \right] \\ &= \frac{1}{2} \eta_1 \kappa_{\text{snc}} \kappa_\Delta^2 \sum_{k=0, k \in \mathcal{S}} \min [\epsilon_g, \epsilon_H] \min [\epsilon_g, \epsilon_H \epsilon_g, \epsilon_H^2] \\ &= \frac{1}{2} N_{\text{TR,S}}^{2\text{nd}} \eta_1 \kappa_{\text{snc}} \kappa_\Delta^2 \epsilon_H \min [\epsilon_g, \epsilon_H]^2 \end{aligned} \quad (4.14)$$

where  $N_{\text{TR,S}}^{2\text{nd}}$  is the total number of successful or very successful iterations such that (1.3) fails, and where we used the inequalities  $\kappa_\Delta < 1$ ,  $\kappa_{\text{snc}} \leq 1$  and  $\max[\epsilon_g, \epsilon_H] < 1$ . The desired conclusion follows from this last inequality with

$$\kappa_{\text{TR,S}}^{2\text{nd}} \stackrel{\text{def}}{=} \frac{2(f(x_0) - f_{\text{low}})}{\eta_1 \kappa_{\text{snc}} \kappa_\Delta^2}.$$

□

Before concluding, we still need an analog of Lemma 2.6 for the trust-region algorithm. Such a result is also described in Gratton et al. (2008), but we formalize it for the sake of clarity.

**Lemma 4.6** *Suppose that A.1 and A.3 hold and, for any fixed  $j \geq 0$ , let  $\mathcal{S}_j$  and  $\mathcal{U}_j$  be defined in (2.12). Then one has that*

$$|\mathcal{U}_j| \leq \left\lceil \frac{\log \gamma_3}{|\log \gamma_2|} |\mathcal{S}_j| + \frac{1}{|\log \gamma_2|} \log \left( \frac{\Delta_0}{\kappa_\Delta \min[\epsilon_g, \epsilon_H]} \right) \right\rceil. \quad (4.15)$$

**Proof.** It follows from the mechanism of the trust-region algorithm that

$$\Delta_{k+1} \leq \gamma_3 \Delta_k \quad \text{for all } k \in \mathcal{S}_j \quad \text{and} \quad \Delta_{k+1} \leq \gamma_2 \Delta_k \quad \text{for all } k \in \mathcal{U}_j.$$

Thus we obtain that

$$\Delta_j \leq \Delta_0 \gamma_2^{|\mathcal{U}_j|} \gamma_3^{|\mathcal{S}_j|}.$$

But Lemma 4.4 gives that, as long as the trust-region algorithm has not terminated, (4.11) must hold. Therefore, we obtain that

$$|\mathcal{S}_j| \log \gamma_3 + |\mathcal{U}_j| \log \gamma_2 \geq \log \left( \frac{\kappa_\Delta \min[\epsilon_g, \epsilon_H]}{\Delta_0} \right).$$

Reorganizing this inequality using  $\gamma_2 < 1$  and taking into account that  $|\mathcal{U}_j|$  is an integer then yields (4.15). □

We may now state the final worst-case complexity bound for convergence of the trust-region algorithm to approximate second-order critical points.

**Theorem 4.7** *Suppose that A.1–A.3 hold. Then, the trust-region algorithm produces an iterate  $x_k$  satisfying (1.3) (and thus terminates) after at most*

$$\left\lceil \kappa_{\text{TR,S}}^{2\text{nd}} \max[\epsilon_g^{-2} \epsilon_H^{-1}, \epsilon_H^{-3}] \right\rceil \quad (4.16)$$

*successful iterations and at most*

$$\left\lceil \kappa_{\text{TR}}^{2\text{nd}} \max[\epsilon_g^{-2} \epsilon_H^{-1}, \epsilon_H^{-3}] \right\rceil \quad (4.17)$$

*iterations in total, where  $\kappa_{\text{TR,S}}^{2\text{nd}}$  and  $\kappa_{\text{TR}}^{2\text{nd}}$  are positive constants independent of  $n$ .*

**Proof.** The first part of the theorem immediately results from Lemma 4.5. The second bound follows by applying Lemma 4.6 and noting that the term in  $\log(1/\epsilon)$  arising from the second term on the left-hand side of (4.15) is dominated by the first as, obviously,  $\log(1/\epsilon) = O(\epsilon^{-3})$  for  $\epsilon \in (0, 1)$ . □

As for the ARC algorithm, we now show that the bound stated in Theorem 4.7 cannot be improved. Again this is achieved by exhibiting a unidimensional example where this bound is attained. The example is itself a modification of that introduced in Section 3 and uses the definitions of  $x_0$ ,  $g_k$  and  $B_k = H_k = \tau_k$  given by (3.1). We now define

$$s_k = \Delta_k = \left( \frac{1}{k+1} \right)^{\frac{1}{3} + \delta} \quad (4.18)$$

(which gives the same steps as in Section 3) and

$$f(x_0) = \frac{1}{4}(\eta_1 + \eta_2)\zeta(1 + 3\delta) \quad \text{and} \quad f_{k+1} = f_k - \frac{1}{4}(\eta_1 + \eta_2) \left( \frac{1}{k+1} \right)^{1+3\delta} \quad (4.19)$$

and therefore the sequence  $\{f_k\}$  is bounded below by zero. It is also clear from the derivation of the example in Section 3 that we may use Hermite interpolation to define the objective function  $f$  on  $\mathbb{R}$  such that it is twice continuously differentiable and has a Lipschitz continuous Hessian. It therefore satisfies both **A.1** and **A.2**. In order to verify that these functions and step values may be generated by a trust-region algorithm, we first note, using (3.1), (4.18) and (4.19), that

$$f_{k+1} = f_k - \frac{1}{4}(\eta_1 + \eta_2)|\tau_k|\Delta_k^2.$$

Hence we obtain<sup>(4)</sup> from (2.10) that

$$\rho_k = \frac{\frac{1}{4}(\eta_1 + \eta_2)|\tau_k|\Delta_k^2}{\frac{1}{2}|\tau_k|\Delta_k^2} = \frac{1}{2}(\eta_1 + \eta_2),$$

for each  $k \geq 0$ . Every iteration is therefore successful (but not very successful). According to (4.4), we may then choose  $\Delta_{k+1}$  in the range  $[\gamma_2\Delta, \Delta_k]$  and our choice

$$\Delta_{k+1} = \left(\frac{k+1}{k+2}\right)^{\frac{1}{3}+\delta} \Delta_k$$

is thus acceptable assuming, without loss of generality, that  $\gamma_2 \leq (\frac{1}{2})^{\frac{1}{3}+\delta}$ .

As in Section 3, we have constructed an objective function  $f$  satisfying **A.1-A.2** on which the trust-region algorithm will need, for any  $\epsilon_g, \epsilon_H$  and  $\delta$  in  $(0, 1)$ , at least of the order of  $O(\epsilon_H^{-3/(1+\delta)})$  successful iterations to achieve approximate second-order optimality. The bounds given by (4.16) and (4.17) are therefore sharp when  $\epsilon_g \geq \epsilon_H$ . We have not been able to show that these bounds are sharp whenever  $\epsilon_g \leq \epsilon_H$ .

We conclude this paper by comparing the bounds for achieving (1.3) given for the ARC algorithm by (2.30) in Theorem 2.9 and for the trust-region algorithm by (4.16)-(4.17) in Theorem 4.7.

- If one assumes that  $\epsilon_H \leq \epsilon_g$ , then the two sets of bounds are qualitatively identical<sup>(5)</sup>, and we have seen that both are sharp.
- If  $\epsilon_g < \epsilon_H < \epsilon_g^{1/2}$ , then the worst-case bound for the trust-region method is  $O(\epsilon_g^{-2}\epsilon_H^{-1}) = O(\epsilon_H^{-\theta})$  iterations at most for some  $\theta \in (3, 5)$ , while the corresponding (sharp) bound for the ARC algorithm remains  $O(\epsilon_H^{-3})$ , which is more favourable.
- Finally, if  $\epsilon_g^{1/2} \leq \epsilon_H$ , the worst-case bound for the trust-region method is now  $O(\epsilon_g^{-2}\epsilon_H^{-1}) = O(\epsilon_g^{-5/2})$  iterations at most, but Cartis et al. (2010b) show that it is also at least  $O(\epsilon_g^{-2})$ . By comparison, the worst-case bound for the ARC algorithm is shown to be no worse than  $O(\epsilon_g^2)$ , while if (2.9) holds this improves to  $O(\epsilon_g^{-3/2})$ , which, according to Cartis et al. (2010b) is sharp. The choice of  $\epsilon_H$  of the order of the square root of  $\epsilon_g$  (which falls at the limit between this third case and the second) makes sense if one wishes to ensure independence of the stopping rule (1.3) from the effect of linear transformations of the problem's variables, and we note that such a choice is also implied by the definition of the measure of local optimality in Nesterov and Polyak (2006).

We therefore see that the ARC algorithm has equal or better worst-case bounds than the trust-region algorithm in all cases, and that the difference is largest for the most practically relevant choice of the relative sizes of the first- and second-order stopping tolerances.

We conclude this section by observing that both presented examples are independent of the value of  $\epsilon_g$  relative to  $\epsilon_H$ , disentangling the interaction between the first- and second-order optimality measures. In particular, this is notable for the trust-region case, where Lemma 4.4 implies a strong interaction between the measures, reflected in Theorem 4.7. Note however, that in both Lemma 4.4 and in Theorem 4.7, if  $\|g_k\| \leq \epsilon_g$  for all  $k$  (which is the case of our example), then it must be that  $\tau_k < -\epsilon_H$  for all  $k$  until termination. Furthermore, then (4.11) becomes  $\Delta_k \geq \kappa_\Delta \epsilon_H$ , only the second-order model decrease applies in the proof of Lemma 4.5 and depends entirely on  $\epsilon_H$ , yielding an upper bound of order  $\epsilon_H^{-3}$

<sup>(4)</sup>Note that  $\kappa_{\text{src}} = 1$  because our example is unidimensional.

<sup>(5)</sup>The constants differ.

for the evaluation complexity of trust-region. Thus, for the particular case when only the curvature condition needs to be satisfied, this upper bound is sharp for the trust-region algorithm. (Similarly, when only the size of the gradient needs to be decreased, Theorem 4.7 yields an upper bound of order  $\epsilon_g^{-2}$ , which was shown in (Cartis et al. 2010b) to be sharp for trust-region.) These remarks illustrate that it is not just the relationship between  $\epsilon_g$  and  $\epsilon_H$  which matters for the worst-case bounds, but also how “close”  $\|g_k\|$  and  $|\tau_k|$  are to these thresholds.

## 5 Summary and perspectives

We have considered the worst-case complexity of achieving approximate second-order optimality for the ARC and trust-region algorithms. We have started by showing that the known bound of  $O(\epsilon_H^{-3})$  ARC iterations can be derived for a variant of the algorithm not requiring multi-dimensional global optimization, and have then shown that the obtained bound is sharp. In addition, we have proved that a bound of the same type also holds for the standard trust-region algorithm, and that this second bound is also sharp whenever  $\epsilon_H = O(\epsilon_g)$ . We also showed that the worst-case behaviour of the ARC algorithm is always as good or preferable to that of the trust-region method.

An obvious next step is to extend the worst-case analysis for second-order optimality to finite-difference and derivative-free schemes, in the spirit of Cartis, Gould and Toint (2010c), and to constrained problems, possibly working along the lines of Cartis et al. (2008). It is also interesting to verify if the optimality properties of the ARC algorithm for convergence to approximate first-order point (Cartis et al., 2010d) can be extended to the ARC algorithm for the second-order case.

### Acknowledgements

The work of the second author is funded by EPSRC Grant EP/E053351/1. All three authors are grateful to the Royal Society for its support through the International Joint Project 14265.

## References

- A. Agarwal, P. L. Bartlett, P. Ravikumar, and M. J. Wainwright. Information-theoretic lower bounds on the oracle complexity of convex optimization. *in* ‘Proceedings of the 23rd Annual Conference on Neural Information Processing Systems’, 2009.
- C. Cartis, N. I. M. Gould, and Ph. L. Toint. An adaptive cubic regularisation algorithm for nonconvex optimization with convex constraints and its function-evaluation complexity. Technical Report 08/05, Department of Mathematics, FUNDP - University of Namur, Namur, Belgium, 2008.
- C. Cartis, N. I. M. Gould, and Ph. L. Toint. Adaptive cubic overestimation methods for unconstrained optimization. Part I: motivation, convergence and numerical results. *Mathematical Programming, Series A*, 2009a. DOI: 10.1007/s10107-009-0286-5, 51 pages.
- C. Cartis, N. I. M. Gould, and Ph. L. Toint. Trust-region and other regularisation of linear least-squares problems. *BIT*, **49**(1), 21–53, 2009b.
- C. Cartis, N. I. M. Gould, and Ph. L. Toint. Adaptive cubic overestimation methods for unconstrained optimization. Part II: worst-case function-evaluation complexity. *Mathematical Programming, Series A*, 2010a. DOI: 10.1007/s10107-009-0337-y.
- C. Cartis, N. I. M. Gould, and Ph. L. Toint. On the complexity of steepest descent, Newton’s and regularized Newton’s methods for nonconvex unconstrained optimization. *SIAM Journal on Optimization*, **20**(6), 2833–2852, 2010b.
- C. Cartis, N. I. M. Gould, and Ph. L. Toint. On the oracle complexity of first-order and derivative-free algorithms for smooth nonconvex minimization. Technical Report NAXYS-03-2010, Department of Mathematics, FUNDP - University of Namur, Namur, Belgium, 2010c.

- C. Cartis, N. I. M. Gould, and Ph. L. Toint. Worst-case complexity of second-order methods for unconstrained nonconvex optimization. Technical Report (in preparation), Department of Mathematics, FUNDP - University of Namur, Namur, Belgium, 2010*d*.
- A. R. Conn, N. I. M. Gould, and Ph. L. Toint. *Trust-Region Methods*. Number 01 in ‘MPS-SIAM Series on Optimization’. SIAM, Philadelphia, USA, 2000.
- S. Gratton, A. Sartenaer, and Ph. L. Toint. Recursive trust-region methods for multiscale nonlinear optimization. *SIAM Journal on Optimization*, **19**(1), 414–444, 2008.
- A. Griewank. The modification of Newton’s method for unconstrained optimization by bounding cubic terms. Technical Report NA/12, Department of Applied Mathematics and Theoretical Physics, University of Cambridge, Cambridge, United Kingdom, 1981.
- A. S. Nemirovski. Efficient methods in convex programming. Lectures notes (online) available on <http://www2.isye.gatech.edu/~nemirovs/OPTILectureNotes.pdf>, 1994.
- Yu. Nesterov. *Introductory Lectures on Convex Optimization*. Applied Optimization. Kluwer Academic Publishers, Dordrecht, The Netherlands, 2004.
- Yu. Nesterov. Accelerating the cubic regularization of Newton’s method on convex problems. *Mathematical Programming, Series A*, **112**(1), 159–181, 2008.
- Yu. Nesterov and B. T. Polyak. Cubic regularization of Newton method and its global performance. *Mathematical Programming, Series A*, **108**(1), 177–205, 2006.
- S. A. Vavasis. Approximation algorithms for indefinite quadratic programming. *Mathematical Programming*, **57**(2), 279–311, 1992*a*.
- S. A. Vavasis. *Nonlinear Optimization: Complexity Issues*. International Series of Monographs on Computer Science. Oxford University Press, Oxford, England, 1992*b*.
- S. A. Vavasis. Black-box complexity of local minimization. *SIAM Journal on Optimization*, **3**(1), 60–80, 1993.
- L. N. Vicente. Worst case complexity of direct search. Technical report, Department of Mathematics, University of Coimbra, Coimbra, Portugal, May 2010.
- M. Weiser, P. Deuffhard, and B. Erdmann. Affine conjugate adaptive Newton methods for nonlinear elastomechanics. *Optimization Methods and Software*, **22**(3), 413–431, 2007.