

Convexity Conditions of Kantorovich Function and Related Semi-infinite Linear Matrix Inequalities

Yun-Bin Zhao *

Abstract. The Kantorovich function $(x^T Ax)(x^T A^{-1}x)$, where A is a positive definite matrix, is not convex in general. From a matrix or convex analysis point of view, it is interesting to address the question: When is this function convex? In this paper, we prove that the 2-dimensional Kantorovich function is convex if and only if the condition number of its matrix is less than or equal to $3 + 2\sqrt{2}$. Thus the convexity of the function with two variables can be completely characterized by the condition number. The upper bound ' $3 + 2\sqrt{2}$ ' is turned out to be a necessary condition for the convexity of the Kantorovich function in any finite-dimensional spaces. We also point out that when the condition number of the matrix (which can be any dimensional) is less than or equal to $\sqrt{5 + 2\sqrt{6}}$, the Kantorovich function is convex. Furthermore, we prove that this general sufficient convexity condition can be improved to $2 + \sqrt{3}$ in 3-dimensional space. Our analysis shows that the convexity of the function is closely related to some modern optimization topics such as the semi-infinite linear matrix inequality or 'robust positive semi-definiteness' of symmetric matrices. In fact, our main result for 3-dimensional cases has been proved by finding an explicit solution range to some semi-infinite linear matrix inequalities.

Keywords. Matrix analysis, condition number, Kantorovich function, convex analysis, positive definite matrix.

*School of Mathematics, University of Birmingham, Edgbaston, Birmingham B15 2TT, United Kingdom (zhaoyy@maths.bham.ac.uk).

1 Introduction

Denote by

$$K(x) = (x^T Ax)(x^T A^{-1}x)$$

where $x \in R^n$ and A is a given $n \times n$ symmetric, positive definite real matrix with eigenvalues $0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$. Then, we have the following Kantorovich inequality (see e.g. [18, 22, 23, 25, 35]):

$$\|x\|_2^4 \geq \frac{4\lambda_1\lambda_n}{\lambda_1^2 + \lambda_n^2} (x^T Ax)(x^T A^{-1}x) \text{ for any } x \in R^n.$$

This inequality and its variants have many applications in matrix analysis, statistics, numerical algebra, and optimization (see e.g. [8, 12, 16, 17, 23, 24, 26, 27, 34, 36, 37, 38]). In this paper, $K(x)$ is referred to as the ‘Kantorovich function’. While $K(x)$ has been widely studied and generalized to different forms in the literature, from a matrix or convex analysis point of view some fundamental questions associated with this function remain open or are not fully addressed. For instance, when is this function convex? Let A be a 2×2 diagonal matrix with diagonal entries 1 and 6. Then A^{-1} is diagonal with diagonal entries 1 and $1/6$, and it is easy to verify that for this example the function $K(x)$ is not convex (since its Hessian matrix is not positive semi-definite at $(1,1)$). Thus, the Kantorovich function is not convex in general.

The investigation of the convexity of $K(x)$ is also inspired by the study of the so-called multiplicative programming problems, which have been widely studied in the field of global optimization (see for example, [28, 29, 3]). In these problems, the product of several convex functions is to be minimized under certain constraints. Matsui [30] showed that multiplicative programming problems are NP-hard, even for the product of linear functions. The existing computational methods for this class of problems are largely global optimization techniques. To find efficient algorithms for multiplicative programming problems, the convexity and/or special structure of the product function should be exploited. Unfortunately, the question of when the product of convex functions is convex remains open, even for the convexity condition for a special product function like $K(x)$ remains open (while $K(x)$ has been widely studied and used in linear algebra and optimization). Until recently, some efforts in this aspect have been made by Zhao [39] and Hiriart-Urruty [20]. The development of convexity conditions for product functions can identify a polynomially solvable subclass of the NP-hard multiplicative programming problems, so that some high-performance computational methods (such as interior-point algorithms and other modern convex optimization methods) can be used to solve such a subclass of problems. Thus, in this paper, we focus on this special product function, $K(x)$, and develop some sufficient and/or necessary convexity conditions for this function.

The main contribution of this paper is as follows. We prove that the Kantorovich function in 2-dimensional space is convex if and only if the condition number of its matrix is less than or equal to $3 + 2\sqrt{2}$. Therefore, the convexity of this function can be characterized completely by the condition number of its matrix, and thus the aforementioned question is fully answered in

2-dimensional space. For higher dimensional cases, we prove that the upper bound ‘ $3 + 2\sqrt{2}$ ’ of the condition number is a necessary condition for the convexity of $K(x)$, i.e., if $K(x)$ is convex, the condition number of A must not exceed this constant. On the other hand, we show that if the condition number of the matrix is less than or equal to $\sqrt{5 + 2\sqrt{6}}$, then $K(x)$ must be convex. An immediate question is how tight this bound is. Can such a general sufficient condition be improved? A remarkable progress in this direction can be achieved for 3-dimensional Kantorovich functions, for which we prove that the bound $\sqrt{5 + 2\sqrt{6}}$ can be improved to $2 + \sqrt{3}$. The proof of such a result is far more than straightforward. It is worth mentioning that we do not know at present whether the result in 2-dimensional space remains valid in higher dimensional spaces, i.e., it is not clear whether or not there is a constant γ such that the following result holds: $K(x)$ is convex if and only if the condition number is less than or equal to γ . By our result, we may conclude that if such a constant γ exists, then $\sqrt{5 + 2\sqrt{6}} \leq \gamma \leq 3 + 2\sqrt{2}$.

The analysis of this paper will yield some new challenging questions and links to certain topics of modern optimization and matrix analysis as well. First, the analysis of this paper indicates that the convexity issue of $K(x)$ is directly related to the so-called (semi-infinite) linear matrix inequality problem which is one of the central topics of modern convex optimization and has found broad applications in control theory, continuous and discrete optimization, geometric distance problems, and so on (see e.g. [9, 11, 12, 15, 19, 32]). In fact, the convexity condition of $K(x)$ can be formulated as a semi-infinite linear matrix inequality problem. In 3-dimensional space, we will show that how such a semi-infinite matrix inequality is explicitly solved in order to develop a convexity condition for $K(x)$.

The convexity issue of $K(x)$ can be also viewed as the so-called ‘*robust positive semi-definiteness*’ of certain symmetric matrices, arising naturally from the analysis to the Hessian matrix of $K(x)$. The typical robust problem on the positive semi-definiteness can be stated as follows: Let the entries of a matrix be multi-variable functions, and some of the variables can take any values in some intervals. The question is what range of values the other variables should take such that the matrix is positive semi-definite. Clearly, this question is also referred to as a robust optimization problem or robust feasibility/stability problem [2, 4, 5, 6, 7, 13, 14, 31]. Our analysis shows that the study of the convexity of $K(x)$ is closely related to these topics. Finally, the convexity issue of $K(x)$ may stimulate the study of more general functions than $K(x)$. Denote by $q_A(x) = \frac{1}{2}x^T A x$. Notice that $q_A^*(x) = \frac{1}{2}x^T A^{-1}x = q_{A^{-1}}(x)$ is the Legendre-Fenchel transform of $q_A(x)$ (see e.g. [1, 10, 21, 33]). The Kantorovich function can be rewritten as $K(x) = 4q_A(x)q_A^*(x)$. Thus, $K(x)$ can be viewed as the product of the quadratic form $q_A(x)$ and its Legendre-Fenchel transform $q_A^*(x)$. Thus, one of the generalization of $K(x)$ is the product function $\widetilde{K}(x) = h(x)h^*(x)$, where h is a convex function and $h^*(x)$ is the Legendre-Fenchel transform of h . However, to our knowledge, the function like $\widetilde{K}(x)$ has not been discussed in the literature.

This paper is organized as follows. In Section 2, we establish some general sufficient, necessary conditions for the convexity of $K(x)$, and point out that the convexity issue of $K(x)$ can

be reformulated as a semi-infinite linear matrix inequality or robust positive semi-definiteness of matrices. In Section 3, we characterize the convexity of the 2-dimensional $K(x)$ by the condition number of its matrix. In the Section 4, we prove an improved sufficient convexity condition for the function $K(x)$ in 3-dimensional space by finding an explicit solution range to a class of semi-infinite linear matrix inequalities. Conclusions are given in the last section.

2 Sufficient, necessary conditions for the convexity of $K(x)$

Throughout this paper, we use $A \succ 0$ ($\succeq 0$) to denote the positive definite (positive semi-definite) matrix. $\kappa(A)$ denotes the condition number of A , i.e., the ratio of its largest and smallest eigenvalues: $\kappa(A) = \lambda_{\max}(A)/\lambda_{\min}(A)$. $q_A(x)$ denotes the quadratic form $(1/2)x^T Ax$.

First of all, we note that a sufficient convexity condition for $K(x)$ can be obtained by Theorem 3.2 in [39] which claims that the following result holds for the product of any two positive definite quadratic forms: *Let A, B be two $n \times n$ matrices and $A, B \succ 0$. If $\kappa(B^{-1/2}AB^{-1/2}) \leq 5 + 2\sqrt{6}$, then the function $\left(\frac{1}{2}x^T Ax\right)\left(\frac{1}{2}x^T Bx\right)$ is convex.* By setting $B = A^{-1}$ and noting that in this case $\kappa(B^{-1/2}AB^{-1/2}) = \kappa(A^2)$, we have the following result.

Theorem 2.1. *Let A be any $n \times n$ matrix and $A \succ 0$. If $\kappa(A) \leq \sqrt{5 + 2\sqrt{6}}$, then the Kantorovich function $K(x)$ is convex.*

Thus, an immediate question arises: What is a necessary condition for the convexity of $K(x)$? The remainder of this section is devoted to answering this question. Let A be an $n \times n$ positive definite matrix. Denote by

$$f(x) := q_A(x)q_{A^{-1}}(x) = \left(\frac{1}{2}x^T Ax\right)\left(\frac{1}{2}x^T A^{-1}x\right) = \frac{1}{4}K(x). \quad (1)$$

Clearly, the convexity of $K(x)$ is exactly the same as that of $f(x)$. Since f is twice continuously differentiable in R^n , the function f is convex if and only if its Hessian matrix $\nabla^2 f$ is positive semi-definite at any point in R^n . It is easy to verify that the Hessian matrix of f is given by

$$\nabla^2 f(x) = q_A(x)A^{-1} + q_{A^{-1}}(x)A + Axx^T A^{-1} + A^{-1}xx^T A. \quad (2)$$

Since A is positive definite, there exists an orthogonal matrix U (i.e., $U^T U = I$) such that

$$A = U^T \Lambda U, \quad A^{-1} = U^T \Lambda^{-1} U, \quad (3)$$

where Λ is a diagonal matrix whose diagonal entries are eigenvalues of A and arranged in non-decreasing order, i.e.,

$$\Lambda = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_n), \quad 0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n.$$

By setting $y = Ux$, we have

$$q_A(x) = \frac{1}{2}x^T(U^T\Lambda U)x = q_\Lambda(y), \quad q_{A^{-1}}(x) = \frac{1}{2}x^T(U^T\Lambda^{-1}U)x = q_{\Lambda^{-1}}(y).$$

Notice that

$$q_\Lambda(y) = \frac{1}{2}y^T\Lambda y = \frac{1}{2}\sum_{i=1}^n \lambda_i y_i^2, \quad q_{\Lambda^{-1}}(y) = \frac{1}{2}y^T\Lambda^{-1}y = \frac{1}{2}\sum_{i=1}^n \frac{1}{\lambda_i} y_i^2.$$

Thus, we have

$$\begin{aligned} & q_\Lambda(y)\Lambda^{-1} + q_{\Lambda^{-1}}(y)\Lambda \\ &= \begin{pmatrix} \frac{1}{2}\sum_{i=1}^n \left(\frac{\lambda_i}{\lambda_1} + \frac{\lambda_1}{\lambda_i}\right) y_i^2 & & & & \\ & \frac{1}{2}\sum_{i=1}^n \left(\frac{\lambda_i}{\lambda_2} + \frac{\lambda_2}{\lambda_i}\right) y_i^2 & & & \\ & & \ddots & & \\ & & & \frac{1}{2}\sum_{i=1}^n \left(\frac{\lambda_i}{\lambda_n} + \frac{\lambda_n}{\lambda_i}\right) y_i^2 & \\ & y_1^2 + \frac{1}{2}\sum_{i=2}^n \Delta_{1i} y_i^2 & & & \\ & & y_2^2 + \frac{1}{2}\sum_{i=1, i \neq 2}^n \Delta_{2i} y_i^2 & & \\ & & & \ddots & \\ & & & & y_n^2 + \frac{1}{2}\sum_{i=1, i \neq n}^n \Delta_{ni} y_i^2 \end{pmatrix} \end{aligned} \quad (4)$$

where

$$\Delta_{ij} = \frac{\lambda_j}{\lambda_i} + \frac{\lambda_i}{\lambda_j}, \quad i, j = 1, \dots, n, \quad i \neq j. \quad (5)$$

It is evident that

$$\Delta_{ij} = \Delta_{ji} \text{ and } \Delta_{ij} \geq 2 \text{ for any } i \neq j. \quad (6)$$

Thus, throughout the paper we do not distinguish Δ_{ij} and Δ_{ji} , and we denote by $\Delta = (\Delta_{ij})$, the $\left(\frac{1}{2}(n-1)n\right)$ -dimensional vector whose components are Δ_{ij} 's where $1 \leq i < j \leq n$. On the other hand, it is easy to verify that

$$\Lambda^{-1}yy^T\Lambda + \Lambda yy^T\Lambda^{-1} = \begin{pmatrix} 2y_1^2 & \Delta_{12}y_1y_2 & \cdots & \Delta_{1n}y_1y_n \\ \Delta_{21}y_2y_1 & 2y_2^2 & \cdots & \Delta_{2n}y_2y_n \\ \vdots & \vdots & \ddots & \vdots \\ \Delta_{n1}y_ny_1 & \Delta_{n2}y_ny_2 & \cdots & 2y_n^2 \end{pmatrix} \quad (7)$$

where Δ_{ij} is given by (5). Therefore, by (2), (3), (4) and (7), we have

$$\begin{aligned} \nabla^2 f(x) &= U^T \left(q_\Lambda(y)\Lambda^{-1} + q_{\Lambda^{-1}}(y)\Lambda + \Lambda yy^T\Lambda^{-1} + \Lambda^{-1}yy^T\Lambda \right) U \\ &= U^T H_n(\Delta, y) U, \end{aligned} \quad (8)$$

where

$$H_n(\Delta, y)$$

where all stated entries are positive numbers. Thus, $H_n(\Delta, \tilde{y}) \succeq 0$ implies that its principle submatrix

$$\begin{pmatrix} 3 + \frac{1}{2}\Delta_{ij} & \Delta_{ij} \\ \Delta_{ji} & 3 + \frac{1}{2}\Delta_{ji} \end{pmatrix}$$

is positive semi-definite. Since the diagonal entries of this submatrix are positive, the submatrix is positive semi-definite if and only if its determinant is nonnegative, i.e.,

$$\left(3 + \frac{1}{2}\Delta_{ij}\right)\left(3 + \frac{1}{2}\Delta_{ji}\right) - \Delta_{ij}\Delta_{ji} \geq 0,$$

which, by (6), can be written as $9 + 3\Delta_{ij} - \frac{3}{4}\Delta_{ij}^2 \geq 0$. Thus, $\Delta_{ij} \leq 6$. Since e_i and e_j can be any columns of the identity matrix, the inequality $\Delta_{ij} \leq 6$ holds for any $i, j = 1, \dots, n$ and $i \neq j$. Lemma 2.2 implies that $\Delta_{1n} = \max_{1 \leq i, j \leq n, i \neq j} \Delta_{ij}$, and hence

$$\kappa(A) + \frac{1}{\kappa(A)} = \frac{\lambda_n}{\lambda_1} + \frac{\lambda_1}{\lambda_n} = \Delta_{1n} = \max_{1 \leq i, j \leq n, i \neq k} \Delta_{ij} \leq 6,$$

which is equivalent to $\kappa(A) \leq 3 + 2\sqrt{2}$. \square

Notice that for each fixed y , $H_n(\Delta, y) \succeq 0$ is a linear matrix inequality (LMI) in Δ . Since y is any vector in R^n , the system (10) is actually a semi-infinite system of LMIs. Recall that for a given function $g(u, v) : R^p \times R^q \rightarrow R$, the semi-infinite inequality (in u) is defined as $g(u, v) \geq 0$ for any $v \in S$, where $S \subseteq R^q$ is a set containing infinite many points. Such inequalities have been widely used in robust control [9, 15] and so-called semi-infinite programming problems [19], and they can be also interpreted as robust feasibility or stability problems when the value of v is uncertain or cannot be given precisely, in which case S means all the possible values of v . Recently, robust problems have wide applications in such areas as mathematical programming, structure design, dynamic system, and financial optimization [2, 4, 5, 7, 13, 14, 31]. Since Δ should be in certain range such that $H_n(\Delta, y) \succeq 0$ for any $y \in R^n$, (10) can be also called ‘robust positive semi-definiteness’ of the matrix H_n .

Any sufficient convexity condition of $K(x)$ can provide some explicit solution range to the system (10). In fact, by Theorem 2.1, we immediately have the next result.

Corollary 2.4. *Consider the semi-infinite linear matrix inequality (in Δ):*

$$\begin{bmatrix} 3y_1^2 + \sum_{i=2}^n \frac{1}{2}\Delta_{1i}y_i^2 & \Delta_{12}y_1y_2 & \cdots & \Delta_{1n}y_1y_n \\ \Delta_{21}y_2y_1 & 3y_2^2 + \sum_{i=1, i \neq 2}^n \frac{1}{2}\Delta_{2i}y_i^2 & \cdots & \Delta_{2n}y_2y_n \\ \vdots & \vdots & \ddots & \vdots \\ \Delta_{n1}y_ny_1 & \Delta_{n2}y_ny_2 & \cdots & 3y_n^2 + \sum_{i=1, i \neq n}^n \frac{1}{2}\Delta_{ni}y_i^2 \end{bmatrix} \succeq 0$$

for any $y \in R^n$, where Δ_{ij} 's satisfy (6). Then the solution set Δ^* of the above system is nonempty, and any vector $\Delta = (\Delta_{ij})$ where $\Delta_{ij} \in \left[2, \sqrt{5 + 2\sqrt{6}}\right]$ is in Δ^* .

Conversely, any range of the feasible solution Δ to the semi-infinite linear matrix inequality (10) can provide a sufficient condition for the convexity of $K(x)$. This idea is used to prove

an improved sufficient convexity condition for $K(x)$ in 3-dimensional space (see Section 4 for details).

Combining Theorems 2.1 and 2.3 leads to the following corollary.

Corollary 2.5. *Assume that there exists a constant, denoted by γ^* , such that the following statement is true: $K(x)$ is convex if and only if $\kappa(A) \leq \gamma^*$. Then such a constant must satisfy that $\sqrt{5 + 2\sqrt{6}} \leq \gamma^* \leq 3 + 2\sqrt{2}$.*

However, the question is: Does such a constant exist? If the answer is ‘yes’, we obtain a complete characterization of the convexity of $K(x)$ by merely the condition number. In the next section, we prove that this question can be fully addressed for 2-dimensional Kantorovich functions, to which the constant is given by $\gamma^* = 3 + 2\sqrt{2}$ (as a result, the bound given in Theorem 2.3 is tight). For higher dimensional cases, the answer to this question is not clear at present. However, for 3-dimensional cases we can prove that the lower bound $\sqrt{5 + 2\sqrt{6}}$ can be significantly improved (see Section 4 for details).

3 Convexity characterization for $K(x)$ in 2-dimensional space

We now consider the Kantorovich function with two variables and prove that the necessary condition in Theorem 2.3 is also sufficient for this case.

Theorem 3.1. *Let A be any 2×2 positive definite matrix. Then $K(x)$ is convex if and only if $\kappa(A) \leq 3 + 2\sqrt{2}$.*

Proof. Notice that there exists an orthogonal matrix U (i.e., $U^T U = I$) such that

$$A = U^T \begin{bmatrix} \beta_1 & 0 \\ 0 & \beta_2 \end{bmatrix} U, \quad A^{-1} = U^T \begin{bmatrix} \frac{1}{\beta_1} & 0 \\ 0 & \frac{1}{\beta_2} \end{bmatrix} U,$$

where β_1, β_2 are eigenvalues of A . By using the same notation in Section 2, and setting $y = Ux$ and $n = 2$ in (9), we have

$$H_2(\Delta, y) = \begin{bmatrix} 3y_1^2 + \frac{1}{2}\Delta_{12}y_2^2 & \Delta_{12}y_1y_2 \\ \Delta_{21}y_2y_1 & 3y_2^2 + \frac{1}{2}\Delta_{21}y_1^2 \end{bmatrix}. \quad (11)$$

For 2-dimensional cases, the vector Δ is reduced to the scalar $\Delta = \Delta_{12}$. If $f(x)$ given by (1) is convex in R^n , it follows from (8) that $H_2(\Delta, y) \succeq 0$ for any $y \in R^2$. In particular, it must be positive semi-definite at $y = e = (1, 1)$, thus

$$H_2(\Delta, e) = \begin{bmatrix} 3 + \frac{1}{2}\Delta_{12} & \Delta_{12} \\ \Delta_{21} & 3 + \frac{1}{2}\Delta_{21} \end{bmatrix} \succeq 0.$$

Since $\Delta_{12} = \Delta_{21}$, it follows that $\det H_2(\Delta, e) = \left(3 + \frac{1}{2}\Delta_{12}\right)^2 - \Delta_{12}^2 \geq 0$, i.e.

$$9 + 3\Delta_{12} - \frac{3}{4}\Delta_{12}^2 \geq 0. \quad (12)$$

Conversely, if (12) holds, we can prove that f is convex. Indeed, we see that the diagonal entries of $H_2(\Delta, y)$ are positive, and that

$$\begin{aligned} \det H_2(\Delta, y) &= \left(3y_1^2 + \frac{1}{2}\Delta_{12}y_2^2\right) \left(3y_2^2 + \frac{1}{2}\Delta_{12}y_1^2\right) - \Delta_{12}^2y_1^2y_2^2 \\ &= 9y_1^2y_2^2 + \frac{3}{2}\Delta_{12}(y_1^4 + y_2^4) - \frac{3}{4}\Delta_{12}^2y_1^2y_2^2 \\ &\geq 9y_1^2y_2^2 + 3\Delta_{12}y_1^2y_2^2 - \frac{3}{4}\Delta_{12}^2y_1^2y_2^2 \\ &= \left(9 + 3\Delta_{12} - \frac{3}{4}\Delta_{12}^2\right)y_1^2y_2^2 \geq 0. \end{aligned}$$

The first inequality above follows from the fact $y_1^4 + y_2^4 \geq 2y_1^2y_2^2$ and the second inequality follows from (12). Thus, $H_2(\Delta, y) \succeq 0$ for any $y \in R^2$, which implies that f is convex.

Therefore, $f(x)$ is convex (i.e., $K(x)$ is convex) if and only if (12) holds. Notice that the roots of the quadratic function $9 + 3t - \frac{3}{4}t^2 = 0$ are $t_1^* = -2$ and $t_2^* = 6$, and notice that $\Delta_{12} \geq 2$ (by(6)). We conclude that (12) holds if and only if $\Delta_{12} \leq 6$. By the definition of Δ_{12} , we have

$$\Delta_{12} = \frac{\beta_1}{\beta_2} + \frac{\beta_2}{\beta_1} = \kappa(A) + \frac{1}{\kappa(A)},$$

Thus the inequality $\Delta_{12} \leq 6$ can be written as $\kappa(A)^2 - 6\kappa(A) + 1 \leq 0$, which is equivalent to $\kappa(A) \leq 3 + 2\sqrt{2}$. \square

It is worth stressing that the above result can be also obtained by solving the semi-infinite linear matrix inequality (10). In fact, by Theorem 2.3, it suffices to prove that $\kappa(A) \leq 3 + 2\sqrt{2}$ is sufficient for the convexity of $K(x)$ in 2-dimensional space. Suppose that $\kappa(A) \leq 3 + 2\sqrt{2}$, which is equivalent to $\Delta_{12} = \frac{\beta_1}{\beta_2} + \frac{\beta_2}{\beta_1} \leq 6$. Thus, $\Delta = \Delta_{12} \in [2, 6]$. We now prove that $K(x)$ is convex. Define $v(\Delta_{12}, y) := \det H_2(\Delta, y)$. It is easy to verify that for any given $y \in R^2$ the function $v(\Delta_{12}, y)$ is concave with respect to Δ_{12} , and hence $v(\Delta_{12}, y) \geq \min\{v(2, y), v(6, y)\}$. Notice that for any $y \in R^2$ we have

$$v(2, y) = \det \begin{bmatrix} 3y_1^2 + y_2^2 & 2y_1y_2 \\ 2y_2y_1 & 3y_2^2 + y_1^2 \end{bmatrix} \geq 0, \quad v(6, y) = \det \begin{bmatrix} 3y_1^2 + 3y_2^2 & 6y_1y_2 \\ 6y_2y_1 & 3y_2^2 + 3y_1^2 \end{bmatrix} \geq 0.$$

Therefore, $v(\Delta_{12}, y) \geq 0$ for any $\Delta_{12} \in [2, 6]$ and $y \in R^2$. This implies that when $\Delta_{12} \in [2, 6]$ the matrix $H_2(\Delta, y) \succeq 0$ for any $y \in R^2$, and hence $K(x)$ is convex.

4 An improved convexity condition for $K(x)$ in 3-dimensional space

In this section, we prove that for $K(x)$ in 3-dimensional space, the upper bound of the condition number given in Theorem 2.1 can be improved to $2 + \sqrt{3}$. To prove this result, we try to find an explicit solution range to a class of semi-infinite linear matrix inequalities, which will immediately yield an improved sufficient convexity condition for $K(x)$ in 3-dimensional space. First, we give some useful inequalities.

Lemma 4.1. *If $2 \leq \delta_1, \delta_2, \delta_3 \leq 4$, then all the functions below are nonnegative:*

$$\chi_1(\delta_1, \delta_2, \delta_3) := 6\delta_3 + \delta_1\delta_2 - \frac{1}{2}\delta_3\delta_2^2 \geq 0, \quad (13)$$

$$\chi_2(\delta_1, \delta_2, \delta_3) := 6\delta_1 + \delta_3\delta_2 - \frac{1}{2}\delta_1\delta_2^2 \geq 0, \quad (14)$$

$$\chi_3(\delta_1, \delta_2, \delta_3) := 6\delta_2 + \delta_1\delta_3 - \frac{1}{2}\delta_2\delta_1^2 \geq 0, \quad (15)$$

$$\chi_4(\delta_1, \delta_2, \delta_3) := 6\delta_3 + \delta_1\delta_2 - \frac{1}{2}\delta_3\delta_1^2 \geq 0, \quad (16)$$

$$\psi(\delta_1, \delta_2, \delta_3) := 12 + \delta_1\delta_2\delta_3 - \delta_1^2 - \delta_2^2 - \delta_3^2 \geq 0. \quad (17)$$

Proof. For any given $\delta_1, \delta_3 \in [2, 4]$, we consider the quadratic function (in t): $\rho(t) = 6\delta_3 + \delta_1 t - \frac{1}{2}\delta_3 t^2$ which is concave in t . Let t_2^* be the largest root of $\rho(t) = 0$. Then

$$t_2^* = \frac{\delta_1 + \sqrt{\delta_1^2 + 12\delta_3^2}}{\delta_3} = \frac{\delta_1}{\delta_3} + \sqrt{\left(\frac{\delta_1}{\delta_3}\right)^2 + 12} \geq \frac{2}{4} + \sqrt{\left(\frac{2}{4}\right)^2 + 12} = 4,$$

where the inequality follows from the fact that $\frac{\delta_1}{\delta_3} \geq \frac{2}{4}$. It is easy to check that the least root of $\rho(t) = 0$ is non-positive, i.e., $t_1^* \leq 0$. Thus, the interval $[2, 4] \subset [t_1^*, t_2^*]$ in which the quadratic function $\rho(t) \geq 0$. Since $2 \leq \delta_2 \leq 4$, we conclude that

$$\chi_1(\delta_1, \delta_2, \delta_3) = \rho(\delta_2) = 6\delta_3 + \delta_1\delta_2 - \frac{1}{2}\delta_3\delta_2^2 \geq 0.$$

Thus (13) holds. All inequalities (14)-(16) can be proved by the same way, or simply by exchanging the role of δ_1, δ_2 and δ_3 in (13).

We now prove (17). It is easy to see that ψ is concave with respect to its every variable. The minimum value of ψ with respect to δ_1 attains at the boundary of the interval $[2, 4]$. Thus, we have

$$\psi(\delta_1, \delta_2, \delta_3) \geq \min\{\psi(2, \delta_2, \delta_3), \psi(4, \delta_2, \delta_3)\} \quad \text{for any } \delta_2, \delta_3 \in [2, 4]. \quad (18)$$

Notice that $\psi(2, \delta_2, \delta_3)$ and $\psi(4, \delta_2, \delta_3)$ are concave with respect to $\delta_2 \in [2, 4]$. Thus we have that

$$\psi(2, \delta_2, \delta_3) \geq \min\{\psi(2, 2, \delta_3), \psi(2, 4, \delta_3)\} \quad \text{for any } \delta_2, \delta_3 \in [2, 4], \quad (19)$$

$$\psi(4, \delta_2, \delta_3) \geq \min\{\psi(4, 2, \delta_3), \psi(4, 4, \delta_3)\} \quad \text{for any } \delta_2, \delta_3 \in [2, 4]. \quad (20)$$

Similarly, $\psi(2, 2, \delta_3)$, $\psi(2, 4, \delta_3)$, $\psi(4, 2, \delta_3)$ and $\psi(4, 4, \delta_3)$ are concave in δ_3 . Thus,

$$\begin{aligned}\psi(2, 2, \delta_3) &\geq \min\{\psi(2, 2, 2), \psi(2, 2, 4)\} = \min\{8, 4\} > 0 \text{ for any } \delta_3 \in [2, 4], \\ \psi(2, 4, \delta_3) &\geq \min\{\psi(2, 4, 2), \psi(2, 4, 4)\} = \min\{4, 8\} > 0 \text{ for any } \delta_3 \in [2, 4], \\ \psi(4, 2, \delta_3) &\geq \min\{\psi(4, 2, 2), \psi(4, 2, 4)\} = \min\{4, 8\} > 0 \text{ for any } \delta_3 \in [2, 4], \\ \psi(4, 4, \delta_3) &\geq \min\{\psi(4, 4, 2), \psi(4, 4, 4)\} = \min\{8, 28\} > 0 \text{ for any } \delta_3 \in [2, 4].\end{aligned}$$

Thus, combining (18)-(20) and the last four inequalities above yields (17). \square

We now focus on developing explicit solution range to certain semi-infinite linear matrix inequalities which will be used later to establish an improved sufficient convexity condition for $K(x)$ in 3-dimensional space.

4.1 The solution range to semi-infinite linear matrix inequalities

Consider the following 3×3 matrix whose entries are the functions in (ω, α, β) :

$$M(\omega, \alpha, \beta) = \begin{bmatrix} 3 + \frac{1}{2}\omega_1\alpha^2 + \frac{1}{2}\omega_2\beta^2 & \omega_1\alpha & \omega_2\beta \\ \omega_1\alpha & \frac{1}{2}\omega_1 + 3\alpha^2 + \frac{1}{2}\omega_3\beta^2 & \omega_3\alpha\beta \\ \omega_2\beta & \omega_3\alpha\beta & \frac{1}{2}\omega_2 + \frac{1}{2}\omega_3\alpha^2 + 3\beta^2 \end{bmatrix} \quad (21)$$

where $\omega = (\omega_1, \omega_2, \omega_3) \in R^3$, $\omega_j \in [2, 6]$ for $j = 1, 2, 3$, and $\alpha, \beta \in [-1, 1]$. We are interested in finding the range of ω_1, ω_2 , and ω_3 in the interval $[2, 6]$ such that

$$M(\omega, \alpha, \beta) \succeq 0 \text{ for any } \alpha, \beta \in [-1, 1],$$

which is a semi-infinite linear matrix inequality. To this end, we seek the condition for ω , under which all the principle minors of M are nonnegative for any $\alpha, \beta \in [-1, 1]$.

First of all, we see from (21) that all diagonal entries (which are the first order principle minors) of M are positive in the intervals considered. Secondly, since $\omega_j \in [2, 6]$ ($j = 1, 2, 3$), and $\alpha \in [-1, 1]$, it is easy to see that

$$\begin{aligned}3 + \frac{1}{2}\omega_1\alpha^2 &= \frac{1}{2}(6 + \omega_1\alpha^2) \geq \frac{1}{2}(\omega_1 + \omega_1\alpha^2) \geq \omega_1|\alpha|, \\ \frac{1}{2}\omega_1 + 3\alpha^2 &= \frac{1}{2}(\omega_1 + 6\alpha^2) \geq \frac{1}{2}(\omega_1 + \omega_1\alpha^2) \geq \omega_1|\alpha|.\end{aligned}$$

Therefore, the second order principle submatrix of (21)

$$\begin{bmatrix} 3 + \frac{1}{2}\omega_1\alpha^2 + \frac{1}{2}\omega_2\beta^2 & \omega_1\alpha \\ \omega_1\alpha & \frac{1}{2}\omega_1 + 3\alpha^2 + \frac{1}{2}\omega_3\beta^2 \end{bmatrix}$$

is diagonally dominant for any $\omega_j \in [2, 6]$ ($j = 1, 2, 3$) and $\alpha, \beta \in [-1, 1]$. Similarly, the second order principle submatrices

$$\begin{bmatrix} 3 + \frac{1}{2}\omega_1\alpha^2 + \frac{1}{2}\omega_2\beta^2 & \omega_2\beta \\ \omega_2\beta & \frac{1}{2}\omega_2 + \frac{1}{2}\omega_3\alpha^2 + 3\beta^2 \end{bmatrix},$$

$$\begin{bmatrix} \frac{1}{2}\omega_1 + 3\alpha^2 + \frac{1}{2}\omega_3\beta^2 & \omega_3\alpha\beta \\ \omega_3\alpha\beta & \frac{1}{2}\omega_2 + \frac{1}{2}\omega_3\alpha^2 + 3\beta^2 \end{bmatrix}$$

are also diagonally dominant for any $\omega_j \in [2, 6]$ ($j = 1, 2, 3$) and $\alpha, \beta \in [-1, 1]$. Thus, all second order principle minors of M are nonnegative in the intervals considered. It is sufficient to find the range of $\omega = (\omega_1, \omega_2, \omega_3)$ such that the third order principle minor is nonnegative, i.e.,

$$\det M(\omega, \alpha, \beta) \geq 0 \quad \text{for any } \alpha, \beta \in [-1, 1].$$

However, this is not straightforward. In order to find such a range, we calculate up to the sixth order partial derivative of $\det M(\omega, \alpha, \beta)$ with respect to α . The first order partial derivative is given as

$$\frac{\partial \det M(\omega, \alpha, \beta)}{\partial \alpha} = \det M_1(\omega, \alpha, \beta) + \det M_2(\omega, \alpha, \beta) + \det M_3(\omega, \alpha, \beta), \quad (22)$$

where $M_i(\omega, \alpha, \beta)$ is the matrix that coincides with the matrix $M(\omega, \alpha, \beta)$ except that every entry in the i th row is differentiated with respect to α , i.e.,

$$M_1(\omega, \alpha, \beta) = \begin{bmatrix} \omega_1\alpha & \omega_1 & 0 \\ \omega_1\alpha & \frac{1}{2}\omega_1 + 3\alpha^2 + \frac{1}{2}\omega_3\beta^2 & \omega_3\alpha\beta \\ \omega_2\beta & \omega_3\alpha\beta & \frac{1}{2}\omega_2 + \frac{1}{2}\omega_3\alpha^2 + 3\beta^2 \end{bmatrix}, \quad (23)$$

$$M_2(\omega, \alpha, \beta) = \begin{bmatrix} 3 + \frac{1}{2}\omega_1\alpha^2 + \frac{1}{2}\omega_2\beta^2 & \omega_1\alpha & \omega_2\beta \\ \omega_1 & 6\alpha & \omega_3\beta \\ \omega_2\beta & \omega_3\alpha\beta & \frac{1}{2}\omega_2 + \frac{1}{2}\omega_3\alpha^2 + 3\beta^2 \end{bmatrix}, \quad (24)$$

$$M_3(\omega, \alpha, \beta) = \begin{bmatrix} 3 + \frac{1}{2}\omega_1\alpha^2 + \frac{1}{2}\omega_2\beta^2 & \omega_1\alpha & \omega_2\beta \\ \omega_1\alpha & \frac{1}{2}\omega_1 + 3\alpha^2 + \frac{1}{2}\omega_3\beta^2 & \omega_3\alpha\beta \\ 0 & \omega_3\beta & \omega_3\alpha \end{bmatrix}. \quad (25)$$

Similarly, the notation $M_{ij}(\omega, \alpha, \beta)$ means the resulting matrix by differentiating, with respect to α , every entry of the i th and j th rows of $M(\omega, \alpha, \beta)$, respectively; $M_{ijk}(\omega, \alpha, \beta)$ means the matrix obtained by differentiating every entry of the i th, j th and k th rows of $M(\omega, \alpha, \beta)$, respectively. All other matrices $M_{ijkl\dots}(\omega, \alpha, \beta)$ are understood this way. In particular, the matrix such as $M_{iij}(\omega, \alpha, \beta)$ with some identical indices means the matrix obtained by differentiating every entry of the i th row of $M(\omega, \alpha, \beta)$ twice and differentiating every entry of its j th row once. Notice that every entry of $M(\omega, \alpha, \beta)$ is at most quadratic in α . Thus if we differentiate a row three times or more, the resulting matrix contains a row with all entries zero, and hence its determinant is equal to zero. For example, $\det M_{1113}(\omega, \alpha, \beta) = 0$.

Clearly, the second order partial derivative is given as follows:

$$\begin{aligned} & \frac{\partial^2 \det M(\omega, \alpha, \beta)}{\partial^2 \alpha} \\ &= \frac{\partial \det M_1(\omega, \alpha, \beta)}{\partial \alpha} + \frac{\partial \det M_2(\omega, \alpha, \beta)}{\partial \alpha} + \frac{\partial \det M_3(\omega, \alpha, \beta)}{\partial \alpha}, \\ &= \sum_{j=1}^3 \det M_{1j}(\omega, \alpha, \beta) + \sum_{j=1}^3 \det M_{2j}(\omega, \alpha, \beta) + \sum_{j=1}^3 \det M_{3j}(\omega, \alpha, \beta) \end{aligned}$$

$$\begin{aligned}
&= \sum_{i=1}^3 \sum_{j=1}^3 \det M_{ij}(\omega, \alpha, \beta) \\
&= \det M_{11}(\omega, \alpha, \beta) + \det M_{22}(\omega, \alpha, \beta) + \det M_{33}(\omega, \alpha, \beta) \\
&\quad + 2 \det M_{12}(\omega, \alpha, \beta) + 2 \det M_{13}(\omega, \alpha, \beta) + 2 \det M_{23}(\omega, \alpha, \beta), \tag{26}
\end{aligned}$$

where the last equality follows from the fact $M_{ij}(\omega, \alpha, \beta) = M_{ji}(\omega, \alpha, \beta)$. By differentiating (26) and noting that $\det M_{111} = \det M_{222} = \det M_{333} = 0$, we have

$$\begin{aligned}
\frac{\partial^3 \det M(\omega, \alpha, \beta)}{\partial^3 \alpha} &= \sum_{i=1}^3 \sum_{j=1}^3 \sum_{k=1}^3 \det M_{ijk}(\omega, \alpha, \beta) \\
&= 3 \det M_{112}(\omega, \alpha, \beta) + 3 \det M_{113}(\omega, \alpha, \beta) + 3 \det M_{122}(\omega, \alpha, \beta) \\
&\quad + 6 \det M_{123}(\omega, \alpha, \beta) + 3 \det M_{133}(\omega, \alpha, \beta) + 3 \det M_{223}(\omega, \alpha, \beta) \\
&\quad + 3 \det M_{233}(\omega, \alpha, \beta). \tag{27}
\end{aligned}$$

By differentiating it again and noting that $\det M_{iiij} = 0$, we have

$$\begin{aligned}
\frac{\partial^4 \det M(\omega, \alpha, \beta)}{\partial^4 \alpha} &= \sum_{i=1}^3 \sum_{j=1}^3 \sum_{k=1}^3 \sum_{l=1}^3 \det M_{ijkl}(\omega, \alpha, \beta), \\
&= 6 \det M_{1122}(\omega, \alpha, \beta) + 6 \det M_{1133}(\omega, \alpha, \beta) + 6 \det M_{2233}(\omega, \alpha, \beta) \\
&\quad + 12 \det M_{1123}(\omega, \alpha, \beta) + 12 \det M_{1223}(\omega, \alpha, \beta) + 12 \det M_{1233}(\omega, \alpha, \beta). \tag{28}
\end{aligned}$$

Finally,

$$\frac{\partial^5 \det M(\omega, \alpha, \beta)}{\partial^5 \alpha} = \sum_{i=1}^3 \sum_{j=1}^3 \sum_{k=1}^3 \sum_{l=1}^3 \sum_{p=1}^3 \det M_{ijklp}(\omega, \alpha, \beta), \tag{29}$$

$$\frac{\partial^6 \det M(\omega, \alpha, \beta)}{\partial^6 \alpha} = \sum_{i=1}^3 \sum_{j=1}^3 \sum_{k=1}^3 \sum_{l=1}^3 \sum_{p=1}^3 \sum_{q=1}^3 \det M_{ijklpq}(\omega, \alpha, \beta). \tag{30}$$

Our first technical result is given as follows.

Lemma 4.2. *Let $\omega_1, \omega_2, \omega_3 \in [2, 4]$ and $\alpha, \beta \in [-1, 1]$. Then the function $\frac{\partial^4 \det M(\omega, \alpha, \beta)}{\partial^4 \alpha}$ is convex with respect to α , and $\frac{\partial^4 \det M(\omega, \alpha, \beta)}{\partial^4 \alpha} \geq 0$.*

Proof. Let $\beta \in [-1, 1]$ and $\omega_1, \omega_3, \omega_2 \in [2, 4]$ be arbitrarily given. Define $g(\alpha) = \frac{\partial^4 \det M(\omega, \alpha, \beta)}{\partial^4 \alpha}$, which is a function in α . As we mentioned earlier, if we differentiate a row of $M(\omega, \alpha, \beta)$ three times or more, the determinant of the resulting matrix is equal to zero. Thus, the nonzero terms on the right-hand side of (30) are only those matrices obtained by differentiating every row of $M(\omega, \alpha, \beta)$ exactly twice, i.e, the terms

$$\det M_{ijklpq}(\omega, \alpha, \beta) = \det M_{112233}(\omega, \alpha, \beta) = \det \begin{bmatrix} \omega_1 & 0 & 0 \\ 0 & 6 & 0 \\ 0 & 0 & \omega_3 \end{bmatrix} = 6\omega_1\omega_3 > 0,$$

and hence $g''(\alpha) = \frac{\partial^6 \det M(\omega, \alpha, \beta)}{\partial^6 \alpha} > 0$. This implies that $g(\alpha) = \frac{\partial^4 \det M(\omega, \alpha, \beta)}{\partial^4 \alpha}$ is convex with respect to $\alpha \in [-1, 1]$. Notice that $g'(\alpha) = \frac{\partial^5 \det M(\omega, \alpha, \beta)}{\partial^5 \alpha}$. We now prove that

$$g'(0) = \frac{\partial^5 \det M(\omega, \alpha, \beta)}{\partial^5 \alpha} \Big|_{\alpha=0} = 0.$$

When the matrix M is differentiated 5 times with respect to α , there are only three possible cases in which we have nonzero determinants for the resulting matrices.

Case 1: Rows 1, 2 are differentiated twice, and row 3 once. In this case we have

$$\det M_{11223}(\omega, \alpha, \beta) = \det \begin{bmatrix} \omega_1 & 0 & 0 \\ 0 & 6 & 0 \\ 0 & \omega_3 \beta & \omega_3 \alpha \end{bmatrix} = 6\omega_1 \omega_3 \alpha.$$

Case 2: Rows 1, 3 are differentiated twice, and row 2 once. We have

$$\det M_{11233}(\omega, \alpha, \beta) = \det \begin{bmatrix} \omega_1 & 0 & 0 \\ \omega_1 & 6\alpha & \omega_3 \beta \\ 0 & 0 & \omega_3 \end{bmatrix} = 6\omega_1 \omega_3 \alpha.$$

Case 3: Rows 2, 3 are done twice, and row 1 once. Then

$$\det M_{12233}(\omega, \alpha, \beta) = \det \begin{bmatrix} \omega_1 \alpha & \omega_1 & 0 \\ 0 & 6 & 0 \\ 0 & 0 & \omega_3 \end{bmatrix} = 6\omega_1 \omega_3 \alpha.$$

Clearly,

$$\begin{aligned} g'(\alpha) &= m_1 \det M_{11223}(\omega, \alpha, \beta) + m_2 \det M_{11233}(\omega, \alpha, \beta) + m_3 \det M_{12233}(\omega, \alpha, \beta) \\ &= 6(m_1 + m_2 + m_3)\omega_1 \omega_3 \alpha, \end{aligned}$$

where m_1, m_2, m_3 are positive integers due to the duplication of the terms in (29), such as $M_{11223}(\omega, \alpha, \beta) = M_{12123}(\omega, \alpha, \beta) = M_{22131}(\omega, \alpha, \beta)$. Therefore, $g'(0) = 0$. By the convexity of $g(\alpha)$, the minimum value of g attains at $\alpha = 0$. We now prove that this minimum value is nonnegative, and hence $g(\alpha) \geq 0$. Indeed, it is easy to see that

$$M_{1223}(\omega, \alpha, \beta) = \begin{bmatrix} \omega_1 \alpha & \omega_1 & 0 \\ 0 & 6 & 0 \\ 0 & \omega_3 \beta & \omega_3 \alpha \end{bmatrix}, \quad M_{1123}(\omega, \alpha, \beta) = \begin{bmatrix} \omega_1 & 0 & 0 \\ \omega_1 & 6\alpha & \omega_3 \beta \\ 0 & \omega_3 \beta & \omega_3 \alpha \end{bmatrix}$$

$$M_{1233}(\omega, \alpha, \beta) = \begin{bmatrix} \omega_1 \alpha & \omega_1 & 0 \\ \omega_1 & 6\alpha & \omega_3 \beta \\ 0 & 0 & \omega_3 \end{bmatrix},$$

$$M_{1133}(\omega, \alpha, \beta) = \begin{bmatrix} \omega_1 & 0 & 0 \\ \omega_1 \alpha & \frac{1}{2}\omega_1 + 3\alpha^2 + \frac{1}{2}\omega_3 \beta^2 & \omega_3 \alpha \beta \\ 0 & 0 & \omega_3 \end{bmatrix},$$

$$M_{1122}(\omega, \alpha, \beta) = \begin{bmatrix} \omega_1 & 0 & 0 \\ 0 & 6 & 0 \\ \omega_2\beta & \omega_3\alpha\beta & \frac{1}{2}\omega_2 + \frac{1}{2}\omega_3\alpha^2 + 3\beta^2 \end{bmatrix},$$

$$M_{2233}(\omega, \alpha, \beta) = \begin{bmatrix} 3 + \frac{1}{2}\omega_1\alpha^2 + \frac{1}{2}\omega_2\beta^2 & \omega_1\alpha & \omega_2\beta \\ 0 & 6 & 0 \\ 0 & 0 & \omega_3 \end{bmatrix}.$$

Therefore, by (28) we have

$$\begin{aligned} g(0) &= \frac{\partial^4 \det M(\omega, \alpha, \beta)}{\partial^4 \alpha} \Big|_{\alpha=0} \\ &= 12 [\det M_{1123}(\omega, \alpha, \beta)|_{\alpha=0} + \det M_{1223}(\omega, \alpha, \beta)|_{\alpha=0} + \det M_{1233}(\omega, \alpha, \beta)|_{\alpha=0}] \\ &\quad + 6 [\det M_{1122}(\omega, \alpha, \beta)|_{\alpha=0} + \det M_{1133}(\omega, \alpha, \beta)|_{\alpha=0} + \det M_{2233}(\omega, \alpha, \beta)|_{\alpha=0}] \\ &= 18 \left(6\omega_1 + \omega_2\omega_3 - \frac{1}{2}\omega_1\omega_3^2 \right) \beta^2 + 18 \left(6\omega_3 + \omega_1\omega_2 - \frac{1}{2}\omega_1^2\omega_3 \right). \end{aligned}$$

By Lemma 4.1, when $\omega_1, \omega_2, \omega_3 \in [2, 4]$, we have

$$\begin{aligned} 6\omega_1 + \omega_2\omega_3 - \frac{1}{2}\omega_1\omega_3^2 &= \chi_2(\omega_1, \omega_3, \omega_2) \geq 0, \\ 6\omega_3 + \omega_1\omega_2 - \frac{1}{2}\omega_1^2\omega_3 &= \chi_3(\omega_1, \omega_3, \omega_2) (= \chi_4(\omega_1, \omega_2, \omega_3)) \geq 0. \end{aligned}$$

Therefore, $g(\alpha) \geq g(0) \geq 0$ for any $\alpha \in [-1, 1]$. Since $\beta \in [-1, 1]$, and $\omega_1, \omega_3, \omega_2 \in [2, 4]$ are arbitrarily given points, the desired results follows. \square

Lemma 4.3. *Let $\omega_1, \omega_2, \omega_3 \in [2, 4]$ and $\alpha, \beta \in [-1, 1]$. Then $\frac{\partial^2 \det M(\omega, \alpha, \beta)}{\partial^2 \alpha}$ is convex with respect to α , and $\alpha = 0$ is a minimizer of it, i.e.*

$$\frac{\partial^2 \det M(\omega, \alpha, \beta)}{\partial^2 \alpha} \geq \frac{\partial^2 \det M(\omega, \alpha, \beta)}{\partial^2 \alpha} \Big|_{\alpha=0} \quad (31)$$

for any α, β, ω in the above-mentioned intervals.

Proof. The convexity of $\frac{\partial^2 \det M(\omega, \alpha, \beta)}{\partial^2 \alpha}$ with respect to α is an immediate consequence of Lemma 4.2 since $\frac{\partial^4 \det M(\omega, \alpha, \beta)}{\partial^4 \alpha}$, the second order derivative of $\frac{\partial^2 \det M(\omega, \alpha, \beta)}{\partial^2 \alpha}$, is nonnegative. We now prove (31). To this end, we first show that

$$\frac{\partial^3 \det M(\omega, \alpha, \beta)}{\partial^3 \alpha} \Big|_{\alpha=0} = 0, \quad (32)$$

which implies that $\alpha = 0$ is a minimizer of the second order partial derivative. In fact, by (27), to calculate the third order partial derivative with respect to α we need to calculate the determinant of $M_{ijk}(\omega, \alpha, \beta)$. Clearly,

$$M_{123}(\omega, \alpha, \beta) = \begin{bmatrix} \omega_1\alpha & \omega_1 & 0 \\ \omega_1 & 6\alpha & \omega_3\beta \\ 0 & \omega_3\beta & \omega_3\alpha \end{bmatrix}.$$

and thus $\det M_{123}(\omega, \alpha, \beta)|_{\alpha=0} = 0$. Similarly, we have

$$\begin{aligned} M_{112}(\omega, \alpha, \beta) &= \begin{bmatrix} \omega_1 & 0 & 0 \\ \omega_1 & 6\alpha & \omega_3\beta \\ \omega_2\beta & \omega_3\alpha\beta & \frac{1}{2}\omega_2 + \frac{1}{2}\omega_3\alpha^2 + 3\beta^2 \end{bmatrix}, \\ M_{113}(\omega, \alpha, \beta) &= \begin{bmatrix} \omega_1 & 0 & 0 \\ \omega_1\alpha & \frac{1}{2}\omega_1 + 3\alpha^2 + \frac{1}{2}\omega_3\beta^2 & \omega_3\alpha\beta \\ 0 & \omega_3\beta & \omega_3\alpha \end{bmatrix}, \\ M_{221}(\omega, \alpha, \beta) &= \begin{bmatrix} \omega_1\alpha & \omega_1 & 0 \\ 0 & 6 & 0 \\ \omega_2\beta & \omega_3\alpha\beta & \frac{1}{2}\omega_2 + \frac{1}{2}\omega_3\alpha^2 + 3\beta^2 \end{bmatrix}, \\ M_{223}(\omega, \alpha, \beta) &= \begin{bmatrix} 3 + \frac{1}{2}\omega_1\alpha^2 + \frac{1}{2}\omega_2\beta^2 & \omega_1\alpha & \omega_2\beta \\ 0 & 6 & 0 \\ 0 & \omega_3\beta & \omega_3\alpha \end{bmatrix}, \end{aligned}$$

which imply that $\det M_{112}(\omega, \alpha, \beta)|_{\alpha=0} = \det M_{113}(\omega, \alpha, \beta)|_{\alpha=0} = \det M_{221}(\omega, \alpha, \beta)|_{\alpha=0} = \det M_{223}(\omega, \alpha, \beta)|_{\alpha=0} = 0$. Finally, we have

$$\begin{aligned} M_{331}(\omega, \alpha, \beta) &= \begin{bmatrix} \omega_1\alpha & \omega_1 & 0 \\ \omega_1\alpha & \frac{1}{2}\omega_1 + 3\alpha^2 + \frac{1}{2}\omega_3\beta^2 & \omega_3\alpha\beta \\ 0 & 0 & \omega_3 \end{bmatrix}, \\ M_{332}(\omega, \alpha, \beta) &= \begin{bmatrix} 3 + \frac{1}{2}\omega_1\alpha^2 + \frac{1}{2}\omega_2\beta^2 & \omega_1\alpha & \omega_2\beta \\ \omega_1 & 6\alpha & \omega_3\beta \\ 0 & 0 & \omega_3 \end{bmatrix}. \end{aligned}$$

Clearly, we also have that $\det M_{331}(\omega, \alpha, \beta)|_{\alpha=0} = \det M_{332}(\omega, \alpha, \beta)|_{\alpha=0} = 0$. From the above calculation, by (27) we see that (32) holds. This means that $\alpha = 0$ is the minimizer of $\frac{\partial^2 M(\omega, \alpha, \beta)}{\partial^2 \alpha}$ for any given $\omega_1, \omega_2, \omega_3 \in [2, 4]$ and $\beta \in [-1, 1]$, and thus (31) holds. \square

Based on the above fact, we may further show that the second order partial derivative is positive in the underlying intervals.

Lemma 4.4. *If $\omega_1, \omega_3, \omega_2 \in [2, 4]$ and $\alpha, \beta \in [-1, 1]$, then $\frac{\partial^2 M(\omega, \alpha, \beta)}{\partial^2 \alpha} \geq 0$.*

Proof. By Lemma 4.3, $\alpha = 0$ is a minimizer of $\frac{\partial^2 M(\omega, \alpha, \beta)}{\partial^2 \alpha}$. Thus, it is sufficient to show that at $\alpha = 0$ the second order partial derivative is nonnegative for any given $\beta \in [-1, 1]$ and $\omega_1, \omega_3, \omega_2 \in [2, 4]$. Indeed,

$$\begin{aligned} &\frac{\partial^2 \det M(\omega, \alpha, \beta)}{\partial^2 \alpha} \\ &= \det M_{11}(\omega, \alpha, \beta) + 2[\det M_{12}(\omega, \alpha, \beta) + \det M_{13}(\omega, \alpha, \beta) + \det M_{23}(\omega, \alpha, \beta)] \\ &\quad + \det M_{22}(\omega, \alpha, \beta) + \det M_{33}(\omega, \alpha, \beta) \end{aligned}$$

$$\begin{aligned}
&= \det \begin{bmatrix} \omega_1 & 0 & 0 \\ \omega_1\alpha & \frac{1}{2}\omega_1 + 3\alpha^2 + \frac{1}{2}\omega_3\beta^2 & \omega_3\alpha\beta \\ \omega_2\beta & \omega_3\alpha\beta & \frac{1}{2}\omega_2 + \frac{1}{2}\omega_3\alpha^2 + 3\beta^2 \end{bmatrix} \\
&+ 2 \det \begin{bmatrix} \omega_1\alpha & \omega_1 & 0 \\ \omega_1 & 6\alpha & \omega_3\beta \\ \omega_2\beta & \omega_3\alpha\beta & \frac{1}{2}\omega_2 + \frac{1}{2}\omega_3\alpha^2 + 3\beta^2 \end{bmatrix} \\
&+ 2 \det \begin{bmatrix} \omega_1\alpha & \omega_1 & 0 \\ \omega_1\alpha & \frac{1}{2}\omega_1 + 3\alpha^2 + \frac{1}{2}\omega_3\beta^2 & \omega_3\alpha\beta \\ 0 & \omega_3\beta & \omega_3\alpha \end{bmatrix} \\
&+ 2 \det \begin{bmatrix} 3 + \frac{1}{2}\omega_1\alpha^2 + \frac{1}{2}\omega_2\beta^2 & \omega_1\alpha & \omega_2\beta \\ \omega_1 & 6\alpha & \omega_3\beta \\ 0 & \omega_3\beta & \omega_3\alpha \end{bmatrix} \\
&+ \det \begin{bmatrix} 3 + \frac{1}{2}\omega_1\alpha^2 + \frac{1}{2}\omega_2\beta^2 & \omega_1\alpha & \omega_2\beta \\ 0 & 6 & 0 \\ \omega_2\beta & \omega_3\alpha\beta & \frac{1}{2}\omega_2 + \frac{1}{2}\omega_3\alpha^2 + 3\beta^2 \end{bmatrix} \\
&+ \det \begin{bmatrix} 3 + \frac{1}{2}\omega_1\alpha^2 + \frac{1}{2}\omega_2\beta^2 & \omega_1\alpha & \omega_2\beta \\ \omega_1\alpha & \frac{1}{2}\omega_1 + 3\alpha^2 + \frac{1}{2}\omega_3\beta^2 & \omega_3\alpha\beta \\ 0 & 0 & \omega_3 \end{bmatrix},
\end{aligned}$$

and hence, by setting $\alpha = 0$ and by a simple calculation we have

$$\begin{aligned}
\frac{\partial^2 \det M(\omega, \alpha, \beta)}{\partial^2 \alpha} \Big|_{\alpha=0} &= 9\omega_2 + \frac{3}{2}\omega_1\omega_3 - \frac{3}{4}\omega_1^2\omega_2 + \frac{9}{2} \left(12 + \omega_1\omega_2\omega_3 - \omega_1^2 - \omega_3^2 - \omega_2^2 \right) \beta^2 \\
&+ \left(9\omega_2 + \frac{3}{2}\omega_1\omega_3 - \frac{3}{4}\omega_2\omega_3^2 \right) \beta^4.
\end{aligned}$$

By Lemma 4.1, for any $\omega_1, \omega_2, \omega_3 \in [2, 4]$, we have the following inequalities:

$$\begin{aligned}
9\omega_2 + \frac{3}{2}\omega_1\omega_3 - \frac{3}{4}\omega_1^2\omega_2 &= \frac{3}{2}\chi_4(\omega_1, \omega_3, \omega_2) \geq 0, \\
9\omega_2 + \frac{3}{2}\omega_1\omega_3 - \frac{3}{4}\omega_2\omega_3^2 &= \frac{3}{2}\chi_1(\omega_1, \omega_3, \omega_2) \geq 0, \\
12 + \omega_1\omega_2\omega_3 - \omega_1^2 - \omega_3^2 - \omega_2^2 &= \psi(\omega_1, \omega_3, \omega_2) \geq 0.
\end{aligned}$$

Thus, the minimum value of $\frac{\partial^2 \det M(\alpha, \beta)}{\partial^2 \alpha}$ is nonnegative. \square

We now prove the main result of this subsection.

Theorem 4.5. *If $\omega_1, \omega_3, \omega_2 \in [2, 4]$, then $M(\omega, \alpha, \beta) \succeq 0$ for any $\alpha, \beta \in [-1, 1]$.*

Proof. By Lemma 4.4, $\det M(\omega, \alpha, \beta)$ is convex with respect to α since its second partial derivative is nonnegative. We now prove the $\alpha = 0$ is a minimizer of $\det M(\omega, \alpha, \beta)$ for an arbitrarily given $\beta \in [-1, 1]$ and ω_{ij} in $[2, 4]$. Indeed, from (23), (24) and (25), it is easy to see that

$$\det M_1(\omega, \alpha, \beta)|_{\alpha=0} = \det M_2(\omega, \alpha, \beta)|_{\alpha=0} = \det M_3(\omega, \alpha, \beta)|_{\alpha=0} = 0,$$

and hence by (22), we have

$$\frac{\partial \det M(\omega, \alpha, \beta)}{\partial \alpha} \Big|_{\alpha=0} = 0,$$

which together with the convexity of $\det M(\omega, \alpha, \beta)$ implies that the minimum value of $\det M(\omega, \alpha, \beta)$ attains at $\alpha = 0$. Substituting $\alpha = 0$ into $\det M(\omega, \alpha, \beta)$ we have

$$\begin{aligned} \det M(\omega, \alpha, \beta) \Big|_{\alpha=0} &= \det \begin{bmatrix} 3 + \frac{1}{2}\omega_2\beta^2 & 0 & \omega_2\beta \\ 0 & \frac{1}{2}\omega_1 + \frac{1}{2}\omega_3\beta^2 & 0 \\ \omega_2\beta & 0 & \frac{1}{2}\omega_2 + 3\beta^2 \end{bmatrix} \\ &= \frac{3}{2} (\omega_1 + \omega_3\beta^2) \left(\frac{1}{2}\omega_2 + \left(3 - \frac{1}{4}\omega_2^2\right) \beta^2 + \frac{1}{2}\omega_2\beta^4 \right). \end{aligned} \quad (33)$$

Notice that the quadratic function $9 - \frac{5}{2}t + \frac{1}{16}t^2 \leq 0$ for any $t \in [4, 36]$. Since $\omega_2 \in [2, 4]$, we have that $\omega_2^2 \in [4, 16] \subset [4, 36]$ we conclude that

$$\left(3 - \frac{1}{4}\omega_2^2\right)^2 - 4 \left(\frac{1}{2}\omega_2\right) \left(\frac{1}{2}\omega_2\right) = 9 - \frac{5}{2}\omega_2^2 + \frac{1}{16}\omega_2^4 \leq 0,$$

which means the determinant of the quadratic function $\frac{1}{2}\omega_2 + \left(3 - \frac{1}{4}\omega_2^2\right)t + \frac{1}{2}\omega_2 t^2$ (in t) is non-positive, and thus

$$\frac{1}{2}\omega_2 + \left(3 - \frac{1}{4}\omega_2^2\right)\beta^2 + \frac{1}{2}\omega_2\beta^4 \geq 0,$$

which together with (33) implies that

$$\det M(\omega, \alpha, \beta) \geq \det M(\omega, \alpha, \beta) \Big|_{\alpha=0} \geq 0.$$

Thus, the third order principle minor of $M(\omega, \alpha, \beta)$ is nonnegative. As we mentioned at the beginning of this subsection, under our conditions the diagonal entries of M are positive, and all second order principle minors of M are also nonnegative. Thus, $M(\omega, \alpha, \beta) \succeq 0$ under our conditions. \square

By symmetry, we have the following result:

Theorem 4.6. *If $\omega_1, \omega_3, \omega_2 \in [2, 4]$, then $P(\omega, \alpha, \beta) \succeq 0$ and $Q(\omega, \alpha, \beta) \succeq 0$ for any $\alpha, \beta \in [-1, 1]$, where*

$$\begin{aligned} P(\omega, \alpha, \beta) &= \begin{bmatrix} 3\alpha^2 + \frac{1}{2}\omega_1 + \frac{1}{2}\omega_2\beta^2 & \omega_1\alpha & \omega_2\alpha\beta \\ \omega_1\alpha & \frac{1}{2}\omega_1\alpha^2 + 3 + \frac{1}{2}\omega_3\beta^2 & \omega_3\beta \\ \omega_2\alpha\beta & \omega_3\beta & \frac{1}{2}\omega_2\alpha^2 + \frac{1}{2}\omega_3 + 3\beta^2 \end{bmatrix}, \\ Q(\omega, \alpha, \beta) &= \begin{bmatrix} 3\alpha^2 + \frac{1}{2}\omega_1\beta^2 + \frac{1}{2}\omega_2 & \omega_1\alpha\beta & \omega_2\alpha \\ \omega_1\alpha\beta & \frac{1}{2}\omega_1\alpha^2 + 3\beta^2 + \frac{1}{2}\omega_3 & \omega_3\beta \\ \omega_2\alpha & \omega_3\beta & \frac{1}{2}\omega_2\alpha^2 + \frac{1}{2}\omega_3\beta^2 + 3 \end{bmatrix}. \end{aligned}$$

Proof. This result can be proved by the same way of Theorem 4.5. However, repeating the whole similar proof is too tedious. We may prove the result by symmetry. In fact, notice

that all the analysis and results in this Subsection depend only on the following conditions: $\omega_1, \omega_2, \omega_3 \in [2, 4]$ and $\alpha, \beta \in [-1, 1]$. By permuting the rows 1 and 2, and columns 1 and 2 of $P(\omega, \alpha, \beta)$, and by setting the substitutions $\omega_1 = \tilde{\omega}_1$, $\omega_2 = \tilde{\omega}_3$ and $\omega_3 = \tilde{\omega}_2$, then $P(\omega, \alpha, \beta)$ is transformed to $M(\tilde{\omega}, \alpha, \beta)$ where $\tilde{\omega} = (\tilde{\omega}_1, \tilde{\omega}_2, \tilde{\omega}_3)$. Clearly, we have $\tilde{\omega}_j \in [2, 4]$ ($j = 1, 2, 3$). Thus, the positive semi-definiteness of P immediately follows from that of M . Similarly, by swapping rows 2 and 3 and swapping columns 2 and 3 of Q , and setting the substitutions $\omega_1 = \tilde{\omega}_2$, $\omega_2 = \tilde{\omega}_1$, and $\omega_3 = \tilde{\omega}_3$, it is easy to see that $Q(\omega, \alpha, \beta)$ can be transformed into $P(\tilde{\omega}, \alpha, \beta)$. Therefore, the result follows from Theorem 4.5 immediately. \square

4.2 The improved sufficient convexity condition

For general Kantorovich functions, Theorem 2.3 claims that when $K(x)$ is convex, it must satisfy that $\kappa(A) \leq 3 + 2\sqrt{2}$ which is the necessary condition for $K(x)$ to be convex. This necessary condition is equivalent to $2 \leq \Delta_{ij} \leq 6$ for any $i \neq j$. Thus any sufficient condition for the convexity of $K(x)$ must fall into this range. Theorem 2.2 shows that if $\kappa(A) \leq \sqrt{5 + 2\sqrt{6}} \approx 3.14626$ (which is equivalent to $2 \leq \Delta_{ij} \leq \frac{6+2\sqrt{6}}{\sqrt{5+2\sqrt{6}}} \approx 3.4641$ for any $i \neq j$), then $K(x)$ is convex. Based on the result of Subsection 4.1, we now prove that this sufficient convexity condition can be improved in 3-dimensional space.

Notice that in 3-dimensional space, by (8) and (9), the positive semi-definiteness of the Hessian matrix of $K(x)$ is equivalent to that of

$$H_3(y) = \begin{bmatrix} 3y_1^2 + \frac{1}{2}\Delta_{12}y_2^2 + \frac{1}{2}\Delta_{13}y_3^2 & \Delta_{12}y_1y_2 & \Delta_{13}y_1y_3 \\ \Delta_{21}y_2y_1 & \frac{1}{2}\Delta_{21}y_1^2 + 3y_2^2 + \frac{1}{2}\Delta_{23}y_3^2 & \Delta_{23}y_2y_3 \\ \Delta_{31}y_3y_1 & \Delta_{32}y_3y_2 & \frac{1}{2}\Delta_{31}y_1^2 + \frac{1}{2}\Delta_{32}y_2^2 + 3y_3^2 \end{bmatrix}. \quad (34)$$

The main result of this section is given below.

Theorem 4.7. *If $\kappa(A) \leq 2 + \sqrt{3}$, then $H_3(y) \succeq 0$ for any $y \in R^3$, and hence the function $K(x) = x^T A x x^T A^{-1} x$ is convex.*

Proof. Let $H_3(y)$ be given by (34). Clearly, $H_3(0) \succeq 0$. In what follows we assume that $y = (y_1, y_2, y_3) \neq 0$. There are three possible cases:

Case 1. y_1 is the component with the largest absolute value: $|y_1| \geq \max\{|y_2|, |y_3|\}$. Denote by $\alpha = \frac{y_2}{y_1}$, $\beta = \frac{y_3}{y_1}$. Notice that $\Delta_{ij} = \Delta_{ji}$ for any $i \neq j$ (see (6)). By setting $\omega = \Delta = (\Delta_{12}, \Delta_{13}, \Delta_{23})$, it is easy to see from (34) that

$$\begin{aligned} H_3(y) &= y_1^2 \begin{bmatrix} 3 + \frac{1}{2}\Delta_{12}\alpha^2 + \frac{1}{2}\Delta_{13}\beta^2 & \Delta_{12}\alpha & \Delta_{13}\beta \\ \Delta_{21}\alpha & \frac{1}{2}\Delta_{21} + 3\alpha^2 + \frac{1}{2}\Delta_{23}\beta^2 & \Delta_{23}\alpha\beta \\ \Delta_{31}\beta & \Delta_{32}\alpha\beta & \frac{1}{2}\Delta_{31} + \frac{1}{2}\Delta_{32}\alpha^2 + 3\beta^2 \end{bmatrix} \\ &= y_1^2 M(\Delta, \alpha, \beta) \end{aligned}$$

where $M(\cdot, \cdot, \cdot)$ is defined by (21). Thus if $M(\Delta, \alpha, \beta) \succeq 0$ for any $\alpha, \beta \in [-1, 1]$, then $H_3(y) \succeq 0$ for any y such that $|y_1| \geq \max\{|y_2|, |y_3|\}$.

Case 2. y_2 is the component with the largest absolute value: $|y_2| \geq \max\{|y_1|, |y_3|\}$. Denote by $\alpha = \frac{y_1}{y_2}$, $\beta = \frac{y_3}{y_2}$. Then, we have

$$\begin{aligned} H_3(y) &= y_2^2 \begin{bmatrix} 3\alpha^2 + \frac{1}{2}\Delta_{12} + \frac{1}{2}\Delta_{13}\beta^2 & \Delta_{12}\alpha & \Delta_{13}\alpha\beta \\ \Delta_{21}\alpha & \frac{1}{2}\Delta_{21}\alpha^2 + 3 + \frac{1}{2}\Delta_{23}\beta^2 & \Delta_{23}\beta \\ \Delta_{31}\alpha\beta & \Delta_{32}\beta & \frac{1}{2}\Delta_{31}\alpha^2 + \frac{1}{2}\Delta_{32} + 3\beta^2 \end{bmatrix} \\ &= y_2^2 P(\Delta, \alpha, \beta) \end{aligned}$$

where $P(\cdot, \cdot, \cdot)$ is defined as in Theorem 4.6. Thus if $P(\Delta, \alpha, \beta) \succeq 0$ for any $\alpha, \beta \in [-1, 1]$, then $H_3(y) \succeq 0$ for any y such that $|y_2| \geq \max\{|y_1|, |y_3|\}$.

Case 3. y_3 is the component with the largest absolute value: $|y_3| \geq \max\{|y_1|, |y_2|\}$. Denote by $\alpha = \frac{y_1}{y_3}$, $\beta = \frac{y_2}{y_3}$. Then,

$$\begin{aligned} H_3(y) &= y_3^2 \begin{bmatrix} 3\alpha^2 + \frac{1}{2}\Delta_{12}\beta^2 + \frac{1}{2}\Delta_{13} & \Delta_{12}\alpha\beta & \Delta_{13}\alpha \\ \Delta_{21}\alpha\beta & \frac{1}{2}\Delta_{21}\alpha^2 + 3\beta^2 + \frac{1}{2}\Delta_{23} & \Delta_{23}\beta \\ \Delta_{31}\alpha & \Delta_{32}\beta & \frac{1}{2}\Delta_{31}\alpha^2 + \frac{1}{2}\Delta_{32}\beta^2 + 3 \end{bmatrix} \\ &= y_3^2 Q(\Delta, \alpha, \beta) \end{aligned}$$

where $Q(\cdot, \cdot, \cdot)$ is defined as in Theorem 4.6. Therefore, if $Q(\Delta, \alpha, \beta) \succeq 0$ for any $\alpha, \beta \in [-1, 1]$, then $H_3(y) \succeq 0$ for any y such that $|y_3| \geq \max\{|y_1|, |y_2|\}$.

If $\kappa(A) \leq 2 + \sqrt{3}$ which is equivalent to $\Delta_{ij} \in [2, 4]$ where $i, j = 1, 2, 3$ and $i \neq j$, by setting $\omega = \Delta$, i.e., $(\omega_1, \omega_2, \omega_3) = (\Delta_{12}, \Delta_{13}, \Delta_{23})$, from Theorems 4.5 and 4.6 we have that

$$M(\Delta, \alpha, \beta) \succeq 0, \quad P(\Delta, \alpha, \beta) \succeq 0, \quad Q(\Delta, \alpha, \beta) \succeq 0 \text{ for any } \alpha, \beta \in [-1, 1].$$

Thus, $H_3(y) \succeq 0$ for any $y \in R^n$, and hence $K(x)$ is convex. \square

The above theorem shows that the upper bound of the condition number in Theorem 2.2 is improved to $2 + \sqrt{3}$ (≈ 3.73205) in 3-dimensional space. As a result, Corollary 2.5 can be improved accordingly in 3-dimensional space.

Corollary 4.8. *Let $n = 3$. Assume that there exists a constant γ^* such that the following statement is true: $\kappa(A) \leq \gamma^*$ if and only if $K(x)$ is convex. Then the constant γ^* must satisfy that $2 + \sqrt{3} \leq \gamma^* \leq 3 + 2\sqrt{2}$.*

5 Conclusions

We have shown that if the Kantorovich function is convex, the condition number of its matrix must be less than or equal to $3 + 2\sqrt{2}$. It turns out that such a necessary condition is also sufficient for any Kantorovich functions in 2-dimensional space. For higher dimensional cases ($n \geq 3$), we point out that a sufficient condition for $K(x)$ to be convex is that the condition number is less than or equal to $\sqrt{5 + 2\sqrt{6}}$. Via certain semi-infinite linear matrix inequalities, we have proved that this general sufficient convexity condition can be improved to $2 + \sqrt{3}$ in

3-dimensional space. Our analysis shows that the convexity issue of the Kantorovich function is closely related to a class of semi-infinite linear matrix inequality problems. Some interesting and challenging questions on $K(x)$ are worthwhile for the future work. For instance, *can we obtain a complete convexity characterization for $n \geq 3$? In another word, for $n \geq 3$, does there exist a constant γ^* such that $\kappa(A) \leq \gamma^*$ if and only if $K(x)$ is convex?* We conjecture that γ^* exists and $\gamma^* = 3 + 2\sqrt{2}$.

References

- [1] J.P. Aubin, *Optima and Equilibria: An Introduction to Nonlinear Analysis*, Springer-Verlag, 1993.
- [2] I. Averbakh and Y.B. Zhao, Explicit reformulations of robust optimization problems with general uncertainty sets, *SIAM J. Optim.*, 18 (2008), pp. 1436-1466.
- [3] H.P. Benson and G.M. Boger, Multiplicative programming problems: analysis and efficient point search heuristic, *J. Optim. Theory Appl.*, 94 (1997), pp. 487–510.
- [4] A. Ben-Tal and A. Nemirovski, Robust convex optimization, *Math. Oper. Res.*, 23 (1998), pp. 769-805.
- [5] A. Ben-Tal, L. El-Ghaoui and A. Nemirovski, *Robust Optimization*, Princeton University Press, 2009.
- [6] D. Bertsimas and M. Sim, The price of robustness, *Oper. Res.*, 52 (2004), pp. 35-53.
- [7] D. Bertsimas and M. Sim, Tractable approximations to robust conic optimization problems, *Math Program., Ser. B*, 107 (2006), pp. 5-36.
- [8] P. Bloomfield and G.S. Watson, The inefficiency of least squares, *Biometrika*, 62 (1975), pp. 121-128.
- [9] S. Boyd, L. El Ghaoui, E. Feron and V. Balakrishnan, *Linear Matrix Inequalities in System and Control Theory*, SIAM, 1994.
- [10] S. Boyd and L. Vandenberghe, *Convex Optimization*, Cambridge University Press, 2004.
- [11] J. Dattorro, *Convex Optimization and Euclidean Distance Geometry*, Meboo, 2009.
- [12] S.S. Dragomir, New inequalities of the Kantorovich type for bounded linear operators in Hilbert spaces, *Linear Algebra and its Applications*, 428 (2008), pp. 2750-2760.
- [13] L. El Ghaoui and H. Lebret, Robust solutions to least-square problems with uncertain data matrices, *SIAM J. Matrix Anal. Appl.*, 18 (1997), pp. 1035-1064.

- [14] L. El Ghaoui, F. Oustry and H. Lebret, Robust solutions to uncertain semi-definite programs, *SIAM J. Optim.*, 9 (1998), pp. 33-52.
- [15] L. El Ghaoui and S.-I. Niculescu (editors), *Advances in Linear Matrix Inequality Methods in Control*, Advances in Control and Design, SIAM, Philadelphia, PA, 2000.
- [16] A.A. Goldstein, A modified Kantorovich inequality for the convergence of Newton's method, *Contemp. Math.*, Vol. 114, Amer. Math. Soc., Providence, RI, 1990.
- [17] A.A. Goldstein, A global Newton method II: Analytic centers, *Math. Program. Ser. B*, 62 (1993), pp. 223-237.
- [18] W. Greub and W. Rheinboldt, On a generalization of an inequality of L.V. Kantorovich, *Proc. American Math Soc.*, 10 (1959), pp. 407-415.
- [19] R. Hettich and K.O. Kortanek, Semi-infinite programming: theory, methods, and applications, *SIAM review*, 35 (1993), pp. 380-429.
- [20] J.B. Hiriart-Urruty, *A note on the Legendre-Fenchel transform of convex composite functions*, in Nonsmooth Mechanics and Analysis (Book Series of Advances in Mathematics and Mechanics, Springer), Vol. 12, 2006, pp. 34-56.
- [21] J.B. Hiriart-Urruty and C. Lemaréchal, *Fundamentals of Convex Analysis*, Grundlehren Text Ed., Springer-Verlag, Berlin, 2001.
- [22] R.A. Horn and C.R. Johnson, *Matrix Analysis*, Cambridge University Press, 1986.
- [23] A.S. Householder, The Kantorovich and some related inequalities, *SIAM Rev.*, 7 (1965), pp. 463-473.
- [24] J. Huang and J. Zhou, A direct proof and a generalization for a Kantorovich type inequality, *Linear Algebra Appl.*, 397 (2005), pp. 185-192.
- [25] L.V. Kantorovich, Functional analysis and applied mathematics (Russian), *Uspekhi Mat Nauk (NS)*, 3 (1948), pp. 89-185.
- [26] C.G. Khatri and C.B. Rao, Some extensions of the Kantorovich inequality and statistical applications, *Journal of Multivariate Analysis*, 11(1981), pp. 498-505.
- [27] M. Knott, On the minimum efficiency of least squares, *Biometrika*, 62 (1975), pp.129-132.
- [28] H. Konno and T. Kuno, Linear multiplicative programming, *Math. Program.*, 56 (1992), pp. 51-64
- [29] H. Konno and T. Kuno, *Multiplicative programming problems*, In: R. Horst and P.M.Pardalos (eds.), *Handbook of Global Optimization*, Kluwer Academic Publishers, Netherlands, 1995, pp. 369-405.

- [30] T. Matsui, NP-hardness of linear multiplicative programming and related problemsn, *J. Global Optim.*, 9 (1996), pp. 113–119.
- [31] M. Mönnigmann, W. Marquardt, C. Bischof, T. Beelitz, B. Lang and P. Willems, A hybrid approach for efficient robust design of dynamic systems, *SIAM Review*, 49 (2007), pp. 236-254.
- [32] Y. Nesterov and A. Nemirovsky, *Interior-point polynomial methods in convex programming*, SIAM, Philadelphia, PA, 1994.
- [33] R.T. Rockafellar, *Convex Analysis*, Princeton University Press, 1970.
- [34] J.C. Spall, The Kantorovich inequality for error analysis of the Kalman filter with unknown noise distributions, *Automatica J. IFAC*, 31 (1995), pp. 1513-1517.
- [35] W.G. Strang, On the Kantorovich inequality, *Proc. American Math Soc.*, 11 (1959), pp. 468.
- [36] G.S. Watson, A method for discovering Kantorovich-type inequalities and a probabilistic interpretation, *Linear Algebra Appl.*, 97 (1987), pp. 211-217.
- [37] G.S. Watson, G. Alpargu and G.P.H. Styan, Some comments on six inequalities associated with the inefficiency of ordinary least squares with one regressor, *Linear Algebra Appl.*, 264 (1997), pp. 13-54.
- [38] T. Yamazaki, An extension of Kantorovich inequality to n -operators via the geometric mean by Ando-Li-Mathias, *Linear Algebra Appl.*, 416 (2006), pp. 688-695.
- [39] Y.B. Zhao, The Legendre-Fenchel conjugate of the product of two positive definite quadratic forms, *SIAM Journal on Matrix Analysis and Applications*, 31(2010), no.4, pp. 1792-1811.