

About error bounds in metric spaces

Marian J. Fabian, René Henrion, Alexander Y. Kruger and Jiří V. Outrata

Abstract The paper presents a general primal space classification scheme of necessary and sufficient criteria for the error bound property incorporating the existing conditions. Several primal space derivative-like objects – slopes – are used to characterize the error bound property of extended-real-valued functions on metric spaces.

1 Introduction

In this paper f is an extended-real-valued function on a metric space X , $f(\bar{x}) = 0$, $S_f := \{x \in X \mid f(x) \leq 0\}$, and $f_+(x) := \max(f(x), 0)$. We are looking for characterizations of the *error bound property*.

Definition 1. f has a local error bound at \bar{x} if there exists a $c > 0$ such that

$$d(x, S_f) \leq cf_+(x) \quad \text{for all } x \text{ near } \bar{x}. \quad (1)$$

For the summary of the theory of error bounds and its various applications, the reader is referred to the survey papers [2, 5, 9, 10], as well as the book [1]. Recent extensions to vector-valued functions can be found in [3].

M. Fabian

Mathematical Institute, Academy of Sciences of the Czech Republic, Žitná 25, 11567 Prague 1, Czech Republic; e-mail: fabian@math.cas.cz

R. Henrion

Weierstrass Institute for Applied Analysis and Stochastics, 10117 Berlin, Germany; e-mail: henrion@wias-berlin.de

A. Kruger

School of Information Technology and Mathematical Sciences, Centre for Informatics and Applied Optimization, University of Ballarat, POB 663, Ballarat, Vic, 3350, Australia; e-mail: a.kruger@ballarat.edu.au

J. Outrata

Institute of Information Theory and Automation, Academy of Sciences of the Czech Republic, 18208 Prague, Czech Republic; e-mail: outrata@utia.cas.cz

Property (1) can be equivalently defined in terms of the *error bound modulus* [5]:

$$\text{Er}(\bar{x}) := \liminf_{\substack{x \rightarrow \bar{x} \\ f(x) > 0}} \frac{f(x)}{d(x, S_f)}, \quad (2)$$

namely, f has a local error bound at \bar{x} if and only if $\text{Er} f(\bar{x}) > 0$. Constant (2) provides a quantitative characterization of this property.

2 Slopes

Primal space characterizations of error bounds can be formulated in terms of slopes.

Recall that the (strong) *slope* [4] of f at x ($|f(x)| < \infty$) is defined as

$$|\nabla f|(x) := \limsup_{u \rightarrow x} \frac{(f(x) - f(u))_+}{d(u, x)}. \quad (3)$$

The following modifications of (3) can be convenient for characterizing the error bound property:

$$|\nabla f|^0(\bar{x}) = \liminf_{x \rightarrow \bar{x}} \frac{f(x) - f(\bar{x})}{\|x - \bar{x}\|}, \quad (4)$$

$$\overline{|\nabla f|}(\bar{x}) = \liminf_{x \rightarrow \bar{x}, f(x) \rightarrow f(\bar{x})} |\nabla f|(x), \quad (5)$$

$$\overline{|\nabla f|}^>(\bar{x}) = \liminf_{x \rightarrow \bar{x}, f(x) \downarrow f(\bar{x})} |\nabla f|(x), \quad (6)$$

$$\overline{|\nabla f|}^\circ(\bar{x}) := \liminf_{x \rightarrow \bar{x}, f(x) \downarrow f(\bar{x})} \sup_{u \neq x} \frac{(f(x) - f_+(u))_+}{d(u, x)}. \quad (7)$$

Constants (4)–(7) are called the *internal slope*, *strict slope*, *strict outer slope*, and *uniform strict slope* of f at \bar{x} respectively.

The relationships between the constants are straightforward.

- Proposition 1.** (i) $\overline{|\nabla f|}(\bar{x}) \leq \overline{|\nabla f|}^>(\bar{x}) \leq \overline{|\nabla f|}^\circ(\bar{x})$.
(ii) $|\nabla f|(\bar{x}) = (-|\nabla f|^0(\bar{x}))_+$.
(iii) $|\nabla f|^0(\bar{x}) \leq \overline{|\nabla f|}^\circ(\bar{x})$.
(iv) If $|\nabla f|^0(\bar{x}) > 0$ then $|\nabla f|^0(\bar{x}) = \text{Er} f(\bar{x})$.

The inequalities in Proposition 1 can be strict.

Example 1. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined as follows:

$$f(x) = \begin{cases} 0 & \text{if } x < 0, \\ x & \text{if } x \geq 0. \end{cases}$$

Obviously $\overline{|\nabla f|}(0) = |\nabla f|(0) = |\nabla f|^0(0) = 0$. At the same time, $\overline{|\nabla f|}^>(0) = \overline{|\nabla f|}^\circ(0) = 1$.

The next example is a modification of the corresponding one in [6].

Example 2. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined as follows:

$$f(x) = \begin{cases} -x & \text{if } x \leq 0, \\ \frac{1}{i} & \text{if } \frac{1}{i+1} < x \leq \frac{1}{i}, i = 1, 2, \dots, \\ x & \text{if } x > 1. \end{cases}$$

Obviously $\overline{|\nabla f|}^>(0) = |\nabla f|(0) = 0$. At the same time, $\overline{|\nabla f|}^\circ(0) = |\nabla f|^0(0) = 1$.

The function in the above example is discontinuous. However, the second inequality in Proposition 1 (i) can be strict for continuous and even Lipschitz continuous functions. The function in the next example is piecewise linear and Clarke regular at 0 (that is, directionally differentiable, and its Clarke generalized directional derivative coincides with the usual one).

Example 3. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined as follows:

$$f(x) = \begin{cases} -x & \text{if } x \leq 0, \\ x \left(1 + \frac{1}{i}\right) - \frac{1}{i(i+1)} & \text{if } \frac{1}{i+1} < x \leq \frac{1}{i+1} + \frac{1}{(i+1)^2}, i = 1, 2, \dots, \\ \frac{1}{i} & \text{if } \frac{1}{i+1} + \frac{1}{(i+1)^2} < x \leq \frac{1}{i}, i = 1, 2, \dots, \\ x & \text{if } x > 1. \end{cases}$$

f is everywhere Fréchet differentiable except for a countable number of points. One can find a point $x > 0$ arbitrarily close to 0 with $|\nabla f|(x) = 0$ (on a horizontal part of the graph). The slopes of non-horizontal parts of the graph decrease monotonously to 1 as $x \downarrow 0$. It is not difficult to check that $\overline{|\nabla f|}^>(0) = |\nabla f|(0) = 0$ while $\overline{|\nabla f|}^\circ(0) = |\nabla f|^0(0) = 1$.

If f is convex then the second inequality in Proposition 1 (i) holds as equality.

For the function f in Example 1, it holds $|\nabla f|(0) < \overline{|\nabla f|}^>(0)$. In the nonconvex case one can also have the opposite inequality.

Example 4. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined as follows:

$$f(x) = \begin{cases} x & \text{if } x < 0, \\ x^2 & \text{if } x \geq 0. \end{cases}$$

Obviously $|\nabla f|(0) = 1$ while $\overline{|\nabla f|}^>(0) = 0$. Note that despite slope $|\nabla f|(0)$ being positive, the function in this example does not have a local error bound at 0. Hence, condition $|\nabla f|(\bar{x}) > 0$ is not in general sufficient for the error bound property to hold at \bar{x} .

3 Error bound criteria

The next theorem generalizes and strengthens [5, Theorem 2].

Theorem 1. (i) $\text{Er} f(\bar{x}) \leq \overline{|\nabla f|}^\circ(\bar{x})$.

(ii) If X is complete and f_+ is lower semicontinuous near \bar{x} , then $\text{Er} f(\bar{x}) = \overline{|\nabla f|}^\circ(\bar{x})$.

Proof. (i) If $\text{Er} f(\bar{x}) = 0$ or $\overline{|\nabla f|}^\circ(\bar{x}) = \infty$, the conclusion is trivial. Let $0 < \gamma < \text{Er} f(\bar{x})$ and $\overline{|\nabla f|}^\circ(\bar{x}) < \infty$. We are going to show that $\overline{|\nabla f|}^\circ(\bar{x}) \geq \gamma$. By (2), there is a $\delta > 0$ such that

$$\frac{f(x)}{d(x, S_f)} > \gamma. \quad (8)$$

for any $x \in B_\delta(\bar{x})$ with $f(x) \geq f(\bar{x})$. Take any $x \in B_\delta(\bar{x})$ with $f(\bar{x}) < f(x) \leq f(\bar{x}) + \delta$ (Such points x exist since $\overline{|\nabla f|}^\circ(\bar{x}) < \infty$.) By (8), one can find a $w \in S_f$ such that

$$\frac{f(x)}{d(x, w)} > \gamma.$$

It follows that $\overline{|\nabla f|}^\circ(\bar{x}) \geq \gamma$.

(ii) Let X be complete and f_+ be lower semicontinuous near \bar{x} . Thanks to (i), we only need to prove that $\text{Er} f(\bar{x}) \geq \overline{|\nabla f|}^\circ(\bar{x})$. If $\text{Er} f(\bar{x}) = \infty$, the inequality is trivial. Let $\text{Er} f(\bar{x}) < \gamma < \infty$. Chose a $\delta > 0$ such that f_+ is lower semicontinuous on $B_{(\gamma^{-1}+1)\delta}(\bar{x})$. Then by (2), there is an $x \in B_{\delta \min(1/2, \gamma^{-1})}(\bar{x})$ such that

$$0 < f(x) < \gamma d(x, S_f).$$

Put $\varepsilon = f(x)$. Then $f_+(x) \leq \inf f_+ + \varepsilon$. Applying to f_+ the Ekeland variational principle with an arbitrary $\lambda \in (\gamma^{-1}\varepsilon, d(x, S_f))$, one can find a w such that $f(w) \leq f(x)$, $d(w, x) \leq \lambda$ and

$$f_+(u) + (\varepsilon/\lambda)d(u, w) \geq f_+(w), \quad \forall u \in B_{(\gamma^{-1}+1)\delta}(\bar{x}). \quad (9)$$

Obviously,

$$\begin{aligned} d(w, x) &< d(x, S_f) \leq d(x, \bar{x}), \\ d(w, \bar{x}) &\leq d(w, x) + d(x, \bar{x}) < 2d(x, \bar{x}) \leq \delta, \\ f(w) &\leq f(x) < \gamma d(x, \bar{x}) \leq \delta. \end{aligned} \quad (10)$$

Besides, $f(w) > 0$ due to the first inequality in (10). It follows from (9) that

$$f(w) \leq f_+(u) + (\varepsilon/\lambda)d(u, w) \leq \gamma d(u, w)$$

for all $u \in B_{(\gamma^{-1}+1)\delta}(\bar{x})$. If $u \notin B_{(\gamma^{-1}+1)\delta}(\bar{x})$, then $d(u, w) > (\gamma^{-1} + 1)\delta - d(w, \bar{x}) > \gamma^{-1}\delta$, and consequently

$$f(w) < \delta < \gamma d(u, w)$$

Thus, in both cases

$$\sup_{u \neq w} \frac{f(w) - f_+(u)}{d(u, w)} \leq \gamma.$$

This implies the inequality $\overline{|\nabla f|}^\diamond(\bar{x}) \leq \text{Er } f(\bar{x})$. \square

Without lower semicontinuity, the inequality in Theorem 1 (i) can be strict.

Example 5. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined as follows:

$$f(x) = \begin{cases} -3x & \text{if } x \leq 0, \\ 3x - \frac{1}{2^i} & \text{if } \frac{1}{2^{i+1}} < x \leq \frac{1}{2^i}, i = 0, 1, \dots, \\ 2x & \text{if } x > 1. \end{cases}$$

Obviously, $\text{Er } f(0) = 1$ while $\overline{|\nabla f|}^\diamond(0) = 3$.

Example 6. Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined as follows:

$$f(x_1, x_2) = \begin{cases} x_1 + x_2 & \text{if } x_1 > 0, x_2 > 0, \\ -x_1 & \text{if } x_1 > 0, x_2 \leq 0, \\ -x_2 & \text{if } x_2 > 0, x_1 \leq 0, \\ 0 & \text{otherwise,} \end{cases}$$

and let \mathbb{R}^2 be equipped with the Euclidean norm. The function is discontinuous on the set $\{(t, 0) \in \mathbb{R}^2 : t > 0\} \cup \{(0, t) \in \mathbb{R}^2 : t > 0\}$. Then $\text{Er } f(0) = 2$ and $\overline{|\nabla f|}^\diamond(0) = 3$.

In view of Theorem 1, inequality $\overline{|\nabla f|}^\diamond(\bar{x}) > 0$ provides a necessary and sufficient error bound criterion for lower semicontinuous functions on complete metric spaces. In a slightly different form, a similar condition for the calmness [q] property of level set maps first appeared in [8, Proposition 3.4]; see also [7, Corollary 4.3].

Taking into account Proposition 1, inequalities


$$|\nabla f|^0(\bar{x}) > 0, \quad \overline{|\nabla f|}(\bar{x}) > 0 \quad \text{and} \quad \overline{|\nabla f|}^>(\bar{x}) > 0$$

provide sufficient error bound criteria

The relationships among the primal space error bound criteria are illustrated in Fig. 1 (X is complete and f_+ is lower semicontinuous near \bar{x}).

In Banach spaces, it is possible to formulate corresponding dual space error bound criteria in terms of subdifferential slopes [5].

Acknowledgements The research was partially supported by the Institutional Research Plan of the Academy of Sciences of Czech Republic, AVOZ 101 905 03; the Grant Agency of the Czech Republic, Projects 201/09/1957 and 201/11/0345; the DFG Research Center MATHEON “Mathematics for key technologies” in Berlin; and the Australian Research Council, Project DP110102011.

Fig. 1 Primal space criteria

References

1. Auslender, A., Teboulle, M.: Asymptotic Cones and Functions in Optimization and Variational Inequalities. Springer Monographs in Mathematics. Springer-Verlag, New York (2003)
2. Azé, D.: A survey on error bounds for lower semicontinuous functions. In: Proceedings of 2003 MODE-SMAI Conference, *ESAIM Proc.*, vol. 13, pp. 1–17. EDP Sci., Les Ulis (2003)
3. Bednarczuk, E.M., Kruger, A.Y.: Error bounds for vector-valued functions: necessary and sufficient conditions. *Nonlinear Anal.* (2011). DOI 10.1016/j.na.2011.05.098
4. De Giorgi, E., Marino, A., Tosques, M.: Problems of evolution in metric spaces and maximal decreasing curve. *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur.* (8) **68**(3), 180–187 (1980)
5. Fabian, M.J., Henrion, R., Kruger, A.Y., Outrata, J.V.: Error bounds: necessary and sufficient conditions. *Set-Valued Var. Anal.* **18**(2), 121–149 (2010).
6. Ioffe, A.D., Outrata, J.V.: On metric and calmness qualification conditions in subdifferential calculus. *Set-Valued Anal.* **16**(2–3), 199–227 (2008)
7. Klatte, D., Kruger, A.Y., Kummer, B.: From convergence principles to stability and optimality conditions. *J. Convex Anal.* To be published
8. Kummer, B.: Inclusions in general spaces: Hoelder stability, solution schemes and Ekeland’s principle. *J. Math. Anal. Appl.* **358**(2), 327–344 (2009).
9. Lewis, A.S., Pang, J.S.: Error bounds for convex inequality systems. In: Generalized Convexity, Generalized Monotonicity: Recent Results (Luminy, 1996), *Nonconvex Optim. Appl.*, vol. 27, pp. 75–110. Kluwer Acad. Publ., Dordrecht (1998)
10. Pang, J.S.: Error bounds in mathematical programming. *Math. Programming, Ser. B* **79**(1-3), 299–332 (1997). Lectures on Mathematical Programming (ISMP97) (Lausanne, 1997)