

Approximate Maximum Principle for Discrete Approximations of Optimal Control Systems with Nonsmooth Objectives and Endpoint Constraints

BORIS S. MORDUKHOVICH¹

Department of Mathematics

Wayne State University

Detroit, MI 48202

Email: boris@math.wayne.edu

ILYA SHVARTSMAN

Department of Mathematics and Computer Science

Penn State University - Harrisburg

Middletown, PA 17110

Email: ius13@psu.edu

Abstract. The paper studies discrete/finite-difference approximations of optimal control problems governed by continuous-time dynamical systems with endpoint constraints. Finite-difference systems, considered as parametric control problems with the decreasing step of discretization, occupy an intermediate position between continuous-time and discrete-time (with fixed steps) control processes and play a significant role in both qualitative and numerical aspects of optimal control. In this paper we derive an enhanced version of the Approximate Maximum Principle for finite-difference control systems, which is new even for problems with smooth endpoint constraints on trajectories and occurs to be the first result in the literature that holds for nonsmooth objectives and endpoint constraints. The results obtained establish necessary optimality conditions for constrained nonconvex finite-difference control systems and justify stability of the Pontryagin Maximum Principle for continuous-time systems under discrete approximations.

Key words. discrete and continuous control systems, discrete approximations, constrained optimal control, maximum principles

AMS subject classification. 49K15, 49M25, 49J52, 49J53, 93C55

Abbreviated title. Approximate Maximum Principle

¹Research of this author was partially supported by the USA National Science Foundation under grant DMS-1007132, by the Australian Research Council under grant DP-12092508, by the European Regional Development Fund (FEDER), and by the following Portuguese agencies: Foundation for Science and Technology, Operational Program for Competitiveness Factors, and Strategic Reference Framework under grant PTDC/MAT/111809/2009.

1 Introduction

The classical result of optimal control theory is the *Pontryagin Maximum Principle* (PMP), which provides necessary optimality conditions for control systems governed by differential equations under various constraints; see the seminal book [8] and also, e.g., [4, 11] with the references therein for more recent developments and applications. For the basic optimal control problem written as

$$(P) \left\{ \begin{array}{l} \text{minimize } \varphi_0(x(t_1)) \\ \text{subject to} \\ \dot{x}(t) = f(x(t), u(t), t) \text{ a.e. } t \in T := [t_0, t_1], \\ x(t_0) = x_0 \in \mathbb{R}^n, \\ u(t) \in U(t) \text{ a.e. } t \in [t_0, t_1], \\ \varphi_i(x(t_1)) \leq 0, \quad i = 1, \dots, l, \\ \varphi_i(x(t_1)) = 0, \quad i = l + 1, \dots, l + q, \end{array} \right.$$

the PMP asserts that every optimal control $\bar{u}(t)$ satisfies the maximum condition

$$H(\bar{x}(t), p(t), \bar{u}(t), t) = \max_{u \in U(t)} H(\bar{x}(t), p(t), u, t) \text{ a.e. } t \in T, \quad (1.1)$$

along the corresponding trajectory $\bar{x}(t)$ of the primal system in (P) and the solution $p(t)$ of the adjoint system

$$\dot{p}(t) = -\nabla_x H(\bar{x}(t), p(t), \bar{u}(t), t) \text{ a.e. } t \in T, \quad (1.2)$$

when f is smooth in x , with an appropriate transversality condition on $p(t_1)$ via gradients or subgradients of smooth or nonsmooth functions φ_i at $\bar{x}(t_1)$. The *Hamilton-Pontryagin function*, or the unmaximized Hamiltonian, in (1.1) and (1.2) is given by

$$H(x, p, u, t) := \langle p, f(x, u, t) \rangle, \quad p \in \mathbb{R}^n. \quad (1.3)$$

This paper is devoted to the study of *discrete approximations* of the continuous-time control problem (P) that are formalized as follows:

$$(P_N) \left\{ \begin{array}{l} \text{minimize } \varphi_0(x_N(t_1)) \\ \text{subject to} \\ x_N(t + h_N) = x_N(t) + h_N f(x_N(t), u_N(t), t), \\ x_N(t_0) = x_0 \in \mathbb{R}^n, \\ u_N(t) \in U(t), \quad t \in T_N := \{t_0, t_0 + h_N, \dots, t_1 - h_N\}, \\ \varphi_i(x_N(t_1)) \leq 0, \quad i = 1, \dots, l, \\ |\varphi_i(x_N(t_1))| \leq \delta > 0, \quad i = l + 1, \dots, l + q, \\ h_N := \frac{t_1 - t_0}{N}, \quad N \in \mathbb{N} := \{1, 2, \dots\}. \end{array} \right.$$

We treat (P_N) as a *sequence* of discrete-time optimal control problems depending on the natural parameter $N = 1, 2, \dots$. This sequence clearly arises from the Euler finite-difference replacement of the derivative in the differential equation of (P) :

$$\dot{x}(t) \approx \frac{x(t+h) - x(t)}{h} \quad \text{as } h \rightarrow 0. \quad (1.4)$$

It is worth noting that we use here the uniform Euler scheme (1.4) just for simplicity; the approach and results obtained below can be carried over to other (including higher-order) approximation schemes.

Apart from the inevitable usage of discrete approximations in computer simulations and calculations of control systems with continuous time, there are deep interrelations between *qualitative aspects* of optimal control for continuous-time systems and their finite-difference approximations. On one hand, the method of discrete approximations has been very instrumental in deriving new necessary optimality conditions for various kinds of continuous-time control systems governed by differential equations and inclusions; see, e.g., [3, 4, 10] and the references therein. On the other hand, it is well known that the *Discrete Maximum Principle* (DMP), as an analog of the PMP with the *exact* maximum condition (1.1), does not generally hold for the discrete-time problems (P_N) with *fixed* steps h_N if the admissible velocity sets $f(x, U(t), t)$ are not convex; see [4, Sections 6.4.1 and 6.5.23] for more details, examples, and historical background. This convexity assumption is rather restrictive, while the possibility to avoid it for continuous-time systems is due to a certain property of “hidden convexity” that is inherent in such systems and plays a crucial role (in one way or another) in any known proof of the PMP.

The failure of the DMP may create a potential instability of the PMP in modeling and numerical calculations of nonconvex continuous-time systems when using discrete approximations. Observe, however, that such *stability* would be justified by showing that merely an *approximate* counterpart of the PMP holds, with a certain perturbation of the maximum condition (1.1) diminishing together with the discretization step $h_N \downarrow 0$ as $N \rightarrow \infty$.

For smooth optimal control problems with no endpoint constraints, the first result of this type was obtained by Gabasov and Kirillova [1] under the name of “quasi-maximum principle” by using an analytic device exploiting the *smooth* and *unconstrained* nature of the family of parametric discrete problems considered therein.

Considering *constrained* control problems of type (P_N) with smooth data required a completely different approach invoking the geometry of endpoint constraints under multineedle variations of optimal controls. It has been initiated by Mordukhovich [2], where a certain finite-difference counterpart of the hidden convexity property for sequences of constrained discrete-time systems was revealed and the first version of the *Approximate Maximum Principle* (AMP) for *smooth* control problems (P_N) was established; the reader can find all the details and discussions in [4, Section 6.4] and commentaries therein.

The version of the AMP given in [2, 4] for smooth constrained problems (P_N) can be formulated as follows. Let the pair (\bar{u}_N, \bar{x}_N) be optimal to (P_N) for each $N \in \mathbb{N}$ such that the

control sequence $\{\bar{u}_N\}$ is *proper* (see [2, 4] and Section 3 below for the exact definition and discussions). Then for any $\varepsilon > 0$ there exist multipliers λ_{iN} as $i = 0, \dots, l + q$ normalized to

$$\lambda_{0N}^2 + \dots + \lambda_{l+qN}^2 = 1 \quad (1.5)$$

satisfying the *sign* and *perturbed complementary slackness* conditions

$$\lambda_{iN} \geq 0 \text{ for } i = 0, \dots, l \text{ and } |\lambda_{iN} \varphi_i(\bar{x}_N(t_1))| < \varepsilon \text{ for } i = 1, \dots, l, \quad (1.6)$$

and such that the *approximate maximum condition*

$$H(\bar{x}_N(t), p_N(t + h_N), \bar{u}_N(t), t) \geq \max_{u \in U(t)} H(\bar{x}_N(t), p_N(t + h_N), u, t) - \varepsilon, \quad t \in T_N, \quad (1.7)$$

holds for all N sufficiently large along the corresponding trajectory of the *adjoint system*

$$p_N(t) = p_N(t + h_N) + h_N \frac{\partial H}{\partial x}(\bar{x}_N(t), p_N(t + h_N), \bar{u}_N(t), t), \quad t \in T_N, \quad (1.8)$$

with the *endpoint/transversality condition*

$$-p_N(t_1) = \sum_{i=0}^{l+q} \lambda_{iN} \nabla \varphi_i(\bar{x}_N(t_1)). \quad (1.9)$$

The presented version of the AMP from [2, 4] plays actually the same role as the DMP for studying and solving discrete-time smooth optimal control problems with sufficiently small stepsizes without imposing the restrictive convexity assumption on the sets $f(\bar{x}_N(t), U(t), t)$, $t \in T_N$. We refer the reader to, e.g., [4, 7] and the bibliographies therein for more discussions and applications of the AMP to chemical engineering, periodic control, biotechnological and ecological processes, etc.

By analogy with the continuous-time problem (P), it would be natural to expect that an extension of the AMP in form (1.5)–(1.9) could be obtained for problems (P_N) with *nonsmooth* cost and constraint functions φ_i via well-recognized subdifferentials of convex and nonsmooth analysis. But it has been surprisingly shown in [5] that this is *not true* even in the case of the simplest *convex nondifferentiable* cost function $\varphi_0(x) = |x - r|$ in the absence of any endpoint constraints on trajectories of linear one-dimensional control systems in (P_N). This reveals a potential *instability* of the PMP under discrete approximations of problem (P) with such a simple and standard type of nonsmoothness when the AMP in form (1.5)–(1.9) is applied. Note that nonsmooth control problems are not only highly challenging theoretically, but are important for various practical applications to engineering design, finance, environmental science, economics, etc.; see, e.g., [4, 10, 11] and the references therein.

The main goal of this paper is to develop a *new version* of the AMP that holds for problems (P_N) with *nonsmooth objectives* and *endpoint inequality constraints* represented as *sums* of *convex* and *smooth* (in fact, strictly differentiable) functions, while equality constraint functions remain smooth. In this version of the AMP, which is new even for problems

(P_N) with all functions φ_i being smooth, we keep the approximate maximum condition (1.7) together with the adjoint system (1.8), as well as the nontriviality and sign conditions on λ_{iN} in (1.5) and (1.6), respectively. What we modify is the perturbed complementarity slackness condition in (1.6), which we replace by its *exact* counterpart

$$\lambda_{iN}\varphi_i(\bar{x}) = 0 \quad \text{for all } i = 1, \dots, l$$

and we also replace the transversality condition (1.9) by its *subdifferential* analog

$$-p_N(t_1) \in \sum_{i=0}^l \lambda_{iN} \partial\varphi_i(\bar{x}) + \sum_{i=l+1}^{l+q} \lambda_{iN} \nabla\varphi_i(\bar{x}), \quad (1.10)$$

where \bar{x} in (1.10) is a *limiting point* of the sequence $\{\bar{x}_N(t_1)\}$, and where $\partial\varphi_i(\bar{x})$ stands for the sum of the subdifferential of convex analysis and the classical gradient at \bar{x} of the corresponding terms in the aforementioned representations of φ_i as $i = 0, \dots, l$.

The rest of the paper is organized as follows. In Section 2 we recall some definitions and preliminaries needed for formulations and proofs of the main results. Section 3 contains the exact formulations and discussions of the assumptions and the statement of the new version of the AMP developed in the paper for the sequence of problems (P_N) . We also discuss the corresponding results for discrete approximation problems of type (P_N) with relaxations of endpoint constraints. Section 4 presents the proof of the main theorem split into several lemmas, which are of their own interest. In the concluding Section 5 we discuss endpoint constraints of the equality type and give an example showing that the new version of the AMP does not hold for such problems without equality constraint relaxations.

Throughout the paper we use the standard notation from variational analysis, optimization, and optimal control; see, e.g., the books [4, 11].

2 Preliminaries

For the reader's convenience, in this section we present several definitions and well-known facts widely used in what follows.

Recall that $\varphi: \mathbb{R}^n \rightarrow \mathbb{R}$ is *strictly differentiable* at \bar{x} if there is a vector $\nabla\varphi(\bar{x})$ such that

$$\lim_{x_1, x_2 \rightarrow \bar{x}} \frac{\varphi(x_1) - \varphi(x_2) - \langle \nabla\varphi(\bar{x}), x_1 - x_2 \rangle}{\|x_1 - x_2\|} = 0.$$

Strict differentiability at a point is a stronger property than just (Fréchet) differentiability, but weaker than continuous differentiability (i.e., C^1) around \bar{x} . A classical example of a function that is differentiable while not strictly differentiable at $\bar{x} = 0$ is

$$\varphi(x) := \begin{cases} x^2 \sin \frac{1}{x}, & x \neq 0, \\ 0, & x = 0. \end{cases}$$

Note that just differentiability of the cost and constraint functions is needed for the PMP to hold in the continuous-time problem (P) (see [4, Theorem 6.37]) in contrast to the corresponding *strict* differentiability assumptions required for the validity of the AMP in the previous and new forms; cf. Theorem 6.59 and Example 6.55 in [4] (the latter is taken from [5, Example 2.3]) and Section 3 below.

If $\varphi: \mathbb{R}^n \rightarrow \overline{\mathbb{R}} := \mathbb{R} \cup \{\infty\}$ is a *convex* function finite at \bar{x} , its *subdifferential* at this point in the sense of convex analysis is defined by

$$\partial\varphi(\bar{x}) := \{v \in \mathbb{R}^n \mid \langle v, x - \bar{x} \rangle \leq \varphi(x) - \varphi(\bar{x}) \text{ for all } x \in \mathbb{R}^n\}. \quad (2.1)$$

It is well known (see, e.g., [9]) that $\partial\varphi(\bar{x}) \neq \emptyset$ for any $\bar{x} \in \text{dom } \varphi := \{x \in \mathbb{R}^n \mid \varphi(x) < \infty\}$, that φ is locally Lipschitz continuous on the interior of $\text{dom } \varphi$ and that

$$\|v\| \leq \ell \text{ for all } v \in \partial\varphi(\bar{x})$$

where $\ell > 0$ is the local Lipschitz constant of φ at \bar{x} . Note also that the subdifferential $\partial\varphi(\bar{x})$ can be equivalently represented as

$$\partial\varphi(\bar{x}) = \{v \in \mathbb{R}^n \mid \langle v, z \rangle \leq \varphi'(\bar{x}; z) \text{ for all } z \in \mathbb{R}^n\}$$

via the *directional derivative* $\varphi'(\bar{x}; z)$ of φ at \bar{x} in the direction z defined by

$$\varphi'(\bar{x}; z) := \lim_{\alpha \downarrow 0} \frac{\varphi(\bar{x} + \alpha z) - \varphi(\bar{x})}{\alpha}. \quad (2.2)$$

This implies, in turn, the directional derivative representation

$$\varphi'(\bar{x}; z) = \sup_{v \in \partial\varphi(\bar{x})} \langle v, z \rangle. \quad (2.3)$$

Given a convex set $\Omega \subset \mathbb{R}^n$ and a point $\bar{x} \in \Omega$, the *normal cone* to Ω at \bar{x} is defined by

$$\mathcal{N}(\bar{x}; \Omega) := \{v \in \mathbb{R}^n \mid \langle v, x - \bar{x} \rangle \leq 0 \text{ for all } x \in \Omega\}. \quad (2.4)$$

The following results of convex analysis are widely used in the paper; see [9] for more details.

Subdifferential Sum Rule (Moreau-Rockafellar Theorem). Let φ, ψ be convex functions on \mathbb{R}^n , and let $\bar{x} \in \text{int}(\text{dom } \psi) \cap \text{int}(\text{dom } \varphi)$. Then we have

$$\partial(\psi + \varphi)(\bar{x}) = \partial\psi(\bar{x}) + \partial\varphi(\bar{x}). \quad (2.5)$$

In particular, assuming in addition that ψ is differentiable at \bar{x} gives us

$$\partial(\psi + \varphi)(\bar{x}) = \nabla\psi(\bar{x}) + \partial\varphi(\bar{x}).$$

Carathéodory Theorem. If a point $x \in \mathbb{R}^n$ lies in the convex hull of a set P , there is a subset $\tilde{P} \subset P$ consisting of at most $n + 1$ points such that x belongs to the convex hull of \tilde{P} .

Separation Theorem. Let $C, D \subset \mathbb{R}^n$ be convex sets such that $\text{int } C \neq \emptyset$ and $\text{int } C \cap D = \emptyset$. Then there exists a nonzero vector $v \in \mathbb{R}^n$ such that

$$\langle v, x_1 \rangle \leq \langle v, x_2 \rangle \text{ for all } x_1 \in C, x_2 \in D.$$

3 The Main Result and Discussions

It what follows we assume that the control sets $U(t)$ in (P_N) are compact subsets of a metric space (\mathcal{U}, d) and that the set-valued mapping $U: [t_0, t_1] \rightrightarrows \mathcal{U}$ is continuous with respect to the Hausdorff distance. Following [2, 4] we say that a sequence of discrete-time control $\{u_n\}$ in (P_N) is *proper* if for every sequence of mesh points

$\tau_{\theta(N)} := t_0 + \theta(N)h_N \in T_N$ with $\theta(N) = 0, \dots, N-1$ and $\tau_{\theta(N)} \rightarrow t \in [t_0, t_1]$ as $N \rightarrow \infty$

we have the following relationships:

$$\text{either } d(u_N(\tau_{\theta(N)}), u_N(\tau_{\theta(N)+s})) \rightarrow 0 \text{ or } d(u_N(\tau_{\theta(N)}), u_N(\tau_{\theta(N)-s})) \rightarrow 0,$$

or both, hold as $N \rightarrow \infty$ with any natural constant s .

Note that this property postulates at least one-sided continuity in the limiting sense of the control sequence in (P_N) at every point and is a discrete-time analog of Lebesgue points in continuous time, which comprise a set of full measure. Thus is not required for continuous-time control systems; see [4] for more details. However, this property occurs to be essential for the validity of the AMP even for one-dimensional linear systems with linear cost and constraint functions; see [5, Example 2.1] and [4, Theorem 6.60].

Theorem 3.1 (Approximate Maximum Principle for Nonsmooth Control Problems). *Let the pair (\bar{u}_N, \bar{x}_N) be optimal to problems (P_N) as $N \in \mathbb{N}$, the sequence $\{\bar{x}_N(t_1)\}$ be uniformly bounded, and let \bar{x} be a limiting point of this sequence. Assume that:*

(a) *The function f is continuous with respect to its variables and continuously differentiable with respect to x in a tube containing the optimal trajectories $\bar{x}_N(t)$ for all large N .*

(b) *The cost function φ_0 and the inequality constraint functions φ_i admit the representations*

$$\varphi_i(x) = \psi_i(x) + \vartheta_i(x) \text{ for } i = 0, \dots, l,$$

where each ψ_i is strictly differentiable at \bar{x} , and where each ϑ_i is convex.

(c) *The equality constraint functions φ_i as $i = l+1, \dots, l+q$ are strictly differentiable at \bar{x} .*

(d) *The sequence of optimal controls $\{\bar{u}_N\}$ is proper.*

Then for any $\varepsilon > 0$ there exist a natural number N_ε and multipliers $\lambda_{iN} \in \mathbb{R}$ as $i = 0, \dots, l+q$ such that for all $N \geq N_\varepsilon$ the following hold:

- *the SIGN and NONTRIVIALITY CONDITIONS*

$$\lambda_{iN} \geq 0 \text{ as } i = 0, \dots, l \text{ and } \sum_{i=0}^l \lambda_{iN}^2 = 1; \quad (3.1)$$

- *the COMPLEMENTARY SLACKNESS CONDITIONS*

$$\lambda_{iN} \varphi_i(\bar{x}) = 0 \text{ as } i = 1, \dots, l; \quad (3.2)$$

- the APPROXIMATE MAXIMUM CONDITION (1.7), where $p_N(t)$ as $t \in T_N \cup \{t_1\}$ is the solution of the adjoint system (1.8) satisfying
- the ENDPOINT/TRANSVERSALITY CONDITION

$$-p_N(t_1) \in \sum_{i=0}^l \lambda_{iN} (\nabla \psi_i(\bar{x}) + \partial \vartheta_i(\bar{x})) + \sum_{i=l+1}^{l+q} \lambda_{iN} \nabla \varphi_i(\bar{x}). \quad (3.3)$$

Note that the new AMP of Theorem 3.1 is an appropriate finite-difference counterpart of the corresponding nonsmooth PMP for continuous-time system (P) ; see [4, 11]. Also, we would like to emphasize that the new result differs from the previous version of the AMP for smooth problems (P_N) discussed in Section 1 in the following two major aspects:

- (i) In contrast to (1.6), the complementary slackness conditions (3.2) are *exact* and are evaluated at the *limiting point* \bar{x} rather than at $\bar{x}_N(t_1)$.
- (ii) In contrast to (1.9), the right-hand side of the transversality condition (3.3) is evaluated at the *limiting point* \bar{x} rather than at $\bar{x}_N(t_1)$.

Note that the result of Theorem 3.1 was announced in [6] for the case of convex cost and endpoint constraint functions φ_i , $i = 0, \dots, l$.

Remark 3.2 (AMP under constraint perturbations). It can be seen from the proof of Theorem 3.1 given in the next section that this theorem is valid without any changes in the assertions when the inequality constraints

$$\varphi_i(x_N(t_1)) \leq 0, \quad i = 1, \dots, l,$$

in (P_N) are perturbed to

$$\varphi_i(x_N(t_1)) \leq \gamma_{iN} \quad \text{with} \quad \gamma_{iN} \rightarrow 0 \quad \text{as} \quad N \rightarrow \infty, \quad i = 1, \dots, l,$$

with *arbitrary* vanishing sequences $\{\gamma_{iN}\}$, $i = 1, \dots, l$. Perturbations of this type were considered in [2, 4], where the AMP was derived in form (1.5)–(1.9) in the case of smooth cost and constraint functions. Note that the previous form of the AMP was also obtained in [2, 4] for perturbations of the *equality constraints*

$$|\varphi_{iN}(x_N(t_1))| \leq \delta_{iN} \quad \text{with} \quad \limsup_{N \rightarrow \infty} \frac{h_N}{\delta_{iN}} = 0 \quad \text{for all} \quad i = l+1, \dots, l+q. \quad (3.4)$$

The question about the validity of the AMP in the new form of Theorem 3.1 under the general *consistent* equality constraint perturbations as in (3.4) remains open. On the other hand, in Section 5 we present a counterexample to this statement for $\delta_{iN} \equiv 0$ as well as for the case of too quickly vanishing (inconsistent) perturbations δ_{iN} in (3.4).

4 Proof of the Theorem

Assume first that Theorem 3.1 is valid in the case $q = 0$, that is, in the absence of the relaxed equality constraints $|\varphi_i(x_N(t_1))| \leq \delta$, $i = l + 1, \dots, l + q$. Let us show that Theorem 3.1 is then valid when $q > 0$.

Indeed, each of these constraints can be written as the two inequalities:

$$\varphi_i(x_N(t_1)) \leq \delta \quad \text{and} \quad -\varphi_i(x_N(t_1)) \leq \delta \quad \text{for } i = l + 1, \dots, l + q. \quad (4.1)$$

Fix an index $i \in \{l + 1, \dots, l + q\}$ in (4.1). Due to our assumption on validity of Theorem 3.1 in the absence of relaxed equality constraints, we can find $\lambda_{iN}^+ \geq 0$ and $\lambda_{iN}^- \geq 0$ for which the complementary slackness conditions yield

$$\lambda_{iN}^+(\varphi_i(\bar{x}) - \delta) = 0 \quad \text{and} \quad \lambda_{iN}^-(\varphi_i(\bar{x}) + \delta) = 0. \quad (4.2)$$

Since both $\varphi_i(\bar{x}) - \delta$ and $\varphi_i(\bar{x}) + \delta$ cannot be equal to zero at the same time, we have $\lambda_{iN}^+ \lambda_{iN}^- = 0$. Denoting finally $\lambda_{iN} := \lambda_{iN}^+ - \lambda_{iN}^-$ for the chosen index $i \in \{l + 1, \dots, l + q\}$ in the last term on (3.3), we arrive at all the conclusions of Theorem 3.1, as formulated.

Therefore, in what follows, *we prove Theorem 3.1 in the case $q = 0$* , breaking down the proof into a series of lemmas.

The first lemma presents a simple subdifferential property of convex continuous functions needed in what follows.

Lemma 4.1 (subdifferential property of convex functions). *Let $\varphi: \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex continuous function, and let $y_k \rightarrow 0$ as $k \rightarrow \infty$ for some sequence $\{y_k\} \subset \mathbb{R}^n$. Then given $\bar{x} \in \mathbb{R}^n$, there exists a subgradient $v \in \partial\varphi(\bar{x})$ such that*

$$\varphi(\bar{x} + y_k) - \varphi(\bar{x}) = \langle v, y_k \rangle + o(\|y_k\|) \quad \text{as } k \rightarrow \infty$$

along a subsequence of $\{y_k\}$ (without relabeling).

Proof. Since the assertion of the lemma is obvious when $y_k \equiv 0$, we suppose without loss of generality that $\{y_k\}$ contains a nonzero subsequence. Let $y_k/\|y_k\| \rightarrow z$ along such a subsequence. Then we have the representation

$$\varphi(\bar{x} + y_k) - \varphi(\bar{x}) = \varphi'(\bar{x}; z)\|y_k\| + o(\|y_k\|)$$

via the directional derivative (2.2) of φ at \bar{x} in the direction z . Taking into account the directional derivative representation (2.3) as well as the nonemptiness and compactness of the subdifferential of convex and continuous (hence Lipschitz continuous) functions, we find a subgradient $v \in \partial\varphi(\bar{x})$ such that

$$\varphi'(\bar{x}; z) = \langle v, z \rangle.$$

This implies, therefore, the relationships

$$\varphi(\bar{x} + y_k) - \varphi(\bar{x}) = \langle v, z \rangle \|y_k\| + o(\|y_k\|) = \left\langle v, \frac{y_k}{\|y_k\|} \right\rangle \|y_k\| + o(\|y_k\|) = \langle v, y_k \rangle + o(\|y_k\|),$$

which complete the proof of the lemma. \square

Next we prove a technical lemma that plays a significant role in the proof of the main result.

Lemma 4.2 (limiting property of convex functions). *Let $\varphi: \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex continuous function, $\{x_k\}, \{y_k\} \subset \mathbb{R}^n$ and $\{\tilde{c}_k\} \subset \mathbb{R}$ be such sequences that $x_k \rightarrow \bar{x}$, $y_k \rightarrow 0$, and $\tilde{c}_k = o(\|y_k\|)$ as $k \rightarrow \infty$, and let*

$$\varphi(x_k + y_k) \geq \varphi(x_k) + \tilde{c}_k \quad \text{for all } k \in \mathbb{N}. \quad (4.3)$$

Then there exists a sequence $\{c_k\} \subset \mathbb{R}$ such that $c_k = o(\|y_k\|)$ as $k \rightarrow \infty$ and

$$\varphi(\bar{x} + y_k) \geq \varphi(\bar{x}) + c_k \quad \text{for all } k \in \mathbb{N}. \quad (4.4)$$

Proof. Note first that by modifying \tilde{c}_k , if necessary, (4.3) can be reduced to a strict inequality, that is,

$$\varphi(x_k + y_k) > \varphi(x_k) + \tilde{c}_k \quad \text{for all } k \in \mathbb{N},$$

which can be restated as $(x_k + y_k, \varphi(x_k) + \tilde{c}_k) \notin \text{epi } \varphi$ or, equivalently, as $(x_k, \varphi(x_k)) + (y_k, \tilde{c}_k) \notin \text{epi } \varphi$. Denote

$$\alpha_k := \max \{ \alpha \geq 0 \mid (x_k, \varphi(x_k)) + \alpha(y_k, \tilde{c}_k) \in \text{epi } \varphi \} \quad (4.5)$$

observing that the maximum is reached in (4.5) for some $\alpha_k \in [0, 1)$ due to closedness and convexity of the epigraph $\text{epi } \varphi$ of φ .

Suppose without loss of generality that $\{y_k\}$ contains a nonzero subsequence, and let A be the set of the limiting points of $\{y_k/\|y_k\|\}$, i.e.,

$$A := \left\{ z \in \mathbb{R}^n \mid z = \lim_{k \rightarrow \infty} \frac{y_k}{\|y_k\|} \text{ along a subsequence of } k \in \mathbb{N} \right\}.$$

Considering further the union set

$$B := \bigcup_{\beta > 0} (\bar{x} + \beta A),$$

we claim the relationship

$$(B, f(\bar{x})) \cap \text{int}(\text{epi } f) = \emptyset. \quad (4.6)$$

Indeed, assume otherwise that there exist $\tilde{\beta} > 0$ and $z \in A$ such that

$$(\bar{x} + \tilde{\beta}z, \varphi(\bar{x})) \in \text{int}(\text{epi } \varphi).$$

This implies that for any sequence converging to $(\bar{x} + \tilde{\beta}z, \varphi(\bar{x}))$ all its terms belong to the epigraph of φ starting with a certain number. In particular, there exists a subsequence of y_k (without relabeling) such that

$$\left(x_k + \alpha_k y_k + \tilde{\beta} \frac{y_k}{\|y_k\|}, \varphi(x_k) + \alpha_k \tilde{c}_k + \tilde{\beta} \frac{\tilde{c}_k}{\|y_k\|}\right) \in \text{epi } \varphi$$

for sufficiently large $k \in \mathbb{N}$; this can be written as

$$\left(x_k + \left(\alpha_k + \frac{\tilde{\beta}}{\|y_k\|}\right) y_k, \varphi(x_k) + \left(\alpha_k + \frac{\tilde{\beta}}{\|y_k\|}\right) \tilde{c}_k\right) \in \text{epi } \varphi.$$

It contradicts the definition of α_k in (4.5) and thus justifies the empty intersection in (4.6).

Furthermore, we have from the construction of the set A that

$$\bar{x} + y_k = \bar{x} + \|y_k\| \left(\frac{y_k}{\|y_k\|}\right) \in \bar{x} + \|y_k\|A + z_k$$

with some sequence $\{z_k\} \subset \mathbb{R}^n$ of order $o(\|y_k\|)$. This implies that

$$(\bar{x} + y_k - z_k, \varphi(\bar{x})) \in (\bar{x} + \|y_k\|A, \varphi(\bar{x})) \subset (B, f(\bar{x}))$$

and hence $(\bar{x} + y_k - z_k, \varphi(\bar{x})) \notin \text{int}(\text{epi } \varphi)$ due to (4.6), which means that

$$\varphi(\bar{x} + y_k - z_k) \geq \varphi(\bar{x}). \quad (4.7)$$

Due to Lipschitz continuity of φ at \bar{x} with some constant ℓ we get

$$\varphi(\bar{x} + y_k - z_k) \leq \varphi(\bar{x} + y_k) + \ell \|z_k\|. \quad (4.8)$$

Setting finally $c_k := -\ell \|z_k\|$ and combining (4.7) with (4.8) ensures the validity of (4.4) and thus completes the proof of the lemma. \square

Note that the assertion of Lemma 4.2 does *not* generally hold for *nonconvex* nonsmooth functions. A simple example is provided by $\varphi(x) = -|x|$ with the sequences $x_k = -1/k$, $y_k = 1/k$, and $\tilde{c}_k = 0$. Then we have (4.3) along these sequences while (4.4) reduces to $-1/k \geq 0 + c_k$, which does not hold whenever $c_k = o(1/k)$ as $k \rightarrow \infty$.

Having in hand the technical lemmas established above, we are now ready to proceed with the proof of the AMP in Theorem 3.1. First we recall a significant property of finite-difference control systems related to needle variations of optimal controls.

Let (\bar{x}_N, \bar{u}_N) be optimal process to problems (P_N) for $N \in \mathbb{N}$. Fix a natural number $r \in \{1, \dots, N-1\}$, mesh points $\tau_j(N) \in T_N$, and control impulses $v_j(N) \in U(\tau_j(N))$ and then define an *r-needle variation* of the optimal control \bar{u}_N by

$$u_N^r(t) := \begin{cases} v_j(N), & t = \tau_j(N), \\ \bar{u}_N(t), & t \in T_N, t \neq \tau_j(N), j = 1, \dots, r. \end{cases} \quad (4.9)$$

The case of $r = 1$ in (4.9) defines a *single needle variation* of \bar{u} , and we omit the index $r = 1$ in this case. The corresponding trajectory increments are denoted by $\Delta^r x_N(t) := x_N^r(t) - \bar{x}_N(t)$ and $\Delta x_N(t) := x_N(t) - \bar{x}_N(t)$ when $r = 1$. Next we pick natural numbers p and m_j as $j = 1, \dots, p$ independent of N and consider the *integer combination*

$$\Delta_N(p, m_j) := \sum_{j=1}^p m_j \Delta_j x_N(t_1) \quad (4.10)$$

of the endpoint trajectory increments $\Delta_j x_N(t_1)$ generated by p *single needle* control variations $u_{Nj}(t)$. The following lemma taken from [4, Lemma 6.62] shows that any single needle integer combination of type (4.10) can be approximated up to a *small quantity* of order $o(h_N)$ by an endpoint trajectory increment generated by a *multineedle* control variation.

Lemma 4.3 (needle variations). *Let the reference sequence of optimal controls $\{\bar{u}_N\}$ to (P_N) be proper, and let $\Delta_N(p, m_j)$ be the integer combination (4.10) generated by single needle control variations $u_{Nj}(t)$, $j = 1, \dots, p$. Then there exists an r -needle control variation $u_N^r(t)$ of type (4.9) with $r := \sum_{j=1}^p m_j$ and a vector quantity of order $o(h_N)$ such that*

$$\Delta_N(p, m_j) = \Delta^r x_N(t_1) + o(h_N) \quad \text{as } N \rightarrow \infty$$

for the corresponding endpoint trajectory increments.

To proceed further, assume that the first $s \leq l$ constraints in (P_N) are *asymptotically active* and the remaining $l - s$ constraints are *asymptotically inactive*, i.e.,

$$\varphi_i(\bar{x}) = 0 \quad \text{for } i = 1, \dots, s; \quad \varphi_i(\bar{x}) < 0 \quad \text{for } i = s + 1, \dots, l.$$

Recall the forms of the cost and inequality constraint functions given in assumption in (b) of Theorem 3.1 and on this basis introduce the following vector mappings from \mathbb{R}^n to \mathbb{R}^{s+1} involving only asymptotically active inequality constraints and defined by

$$\begin{aligned} \Phi(x) &:= (\varphi_0(x), \dots, \varphi_s(x)), & \Psi(x) &:= (\psi_0(x), \dots, \psi_s(x)), & \Upsilon(x) &:= (\vartheta_0(x), \dots, \vartheta_s(x)), \\ L(x) &:= \Psi(\bar{x}) + \Psi'(\bar{x})(x - \bar{x}) + \Upsilon(x). \end{aligned} \quad (4.11)$$

It is obvious that all the components L_i of the mapping L from (4.11) are convex and represented as sums of convex and linear functions. Thus

$$L_i(\bar{x}) = \varphi_i(\bar{x}) \quad \text{and} \quad \partial L_i(x) = \nabla \psi_i(\bar{x}) + \partial \vartheta_i(\bar{x}) \quad \text{for all } i = 0, \dots, s$$

by the subdifferential sum rule mentioned in Section 2.

The next lemma presents an asymptotic consequence of optimality in (P_N) by using multi-needle variations of optimal controls.

Lemma 4.4 (asymptotic consequence of optimality.) *Fix a natural number r independent of N and consider a sequence $\{\Delta^r x_N(t_1)\}$, $N \in \mathbb{N}$, of endpoint trajectory increments generated by r -needle variations of optimal controls. Then there exists an index $i_0 \in \{0, \dots, s\}$ independent of N such that*

$$\liminf_{h_N \rightarrow 0} \frac{L_{i_0}(\bar{x} + \Delta^r x_N(t_1)) - L_{i_0}(\bar{x})}{h_N} \geq 0 \quad (4.12)$$

for the corresponding component of the mapping L defined in (4.11).

Proof. Select a subsequence of $\{N\}$ (without relabeling) such that $\bar{x}_N(t_1) \rightarrow \bar{x}$. Due to optimality of the sequence $\{\bar{x}_N\}$, there is an index $i_0 = i_0(N) \in \{0, \dots, s\}$ such that

$$\varphi_{i_0}(\bar{x}_N(t_1) + \Delta^r x_N(t_1)) \geq \varphi_{i_0}(\bar{x}_N(t_1)). \quad (4.13)$$

(Indeed, inequalities $\varphi_i(\bar{x}_N(t_1) + \Delta^r x_N(t_1)) < \varphi_i(\bar{x}_N(t_1))$, for all $i = 0, \dots, s$ contradict optimality of \bar{x}_N .) By selecting a further subsequence, if necessary, we can assume that i_0 does not depend on N . It follows from (4.13), by the strict differentiability of ψ at \bar{x} , that

$$\begin{aligned} \vartheta_{i_0}(\bar{x}_N(t_1) + \Delta^r x_N(t_1)) - \vartheta_{i_0}(\bar{x}_N(t_1)) &\geq -(\psi_{i_0}(\bar{x}_N(t_1) + \Delta^r x_N(t_1)) - \psi_{i_0}(\bar{x}_N(t_1))) \\ &= -\langle \nabla \psi_{i_0}(\bar{x}), \Delta^r x_N(t_1) \rangle + \tilde{c}_N, \end{aligned}$$

where $\tilde{c}_N = o(\|\Delta^r x_N(t_1)\|) = o(h_N)$ due to $\|\Delta^r x_N(t_1)\| = O(h_N)$ as shown in the proof of [4, Lemma 6.62]. We conclude from the relation above that

$$\begin{aligned} \vartheta_{i_0}(\bar{x}_N(t_1) + \Delta^r x_N(t_1)) + \psi_{i_0}(\bar{x}) + \langle \nabla \psi_{i_0}(\bar{x}), \bar{x}_N(t_1) + \Delta^r x_N(t_1) - \bar{x} \rangle \\ \geq \vartheta_{i_0}(\bar{x}_N(t_1)) + \psi_{i_0}(\bar{x}) + \langle \nabla \psi_{i_0}(\bar{x}), \bar{x}_N(t_1) - \bar{x} \rangle + \tilde{c}_N \end{aligned}$$

which can be written, in the notation of (4.11), as

$$L_{i_0}(\bar{x}_N(t_1) + \Delta^r x_N(t_1)) \geq L_{i_0}(\bar{x}_N(t_1)) + \tilde{c}_N. \quad (4.14)$$

Applying now Lemma 4.2, we conclude that there exists a sequence $\{c_N\} \subset \mathbb{R}$ of order $c_N = o(h_N)$ as $N \rightarrow \infty$ such that

$$L_{i_0}(\bar{x} + \Delta^r x_N(t_1)) - L_{i_0}(\bar{x}) \geq c_N,$$

which implies in turn that

$$\liminf_{h_N \rightarrow 0} h_N^{-1} \left(L_{i_0}(\bar{x} + \Delta^r x_N(t_1)) - L_{i_0}(\bar{x}) \right) \geq 0$$

and thus completes the proof of this lemma. \square

Next we form the set E by

$$E := \left\{ (x, \nu_0, \dots, \nu_s) \in \mathbb{R}^{n+s+1} \mid L_i(x) = \psi_i(\bar{x}) + \langle \nabla \psi_i(\bar{x}), x - \bar{x} \rangle + \vartheta_i(x) \leq \nu_i, \quad i = 0, \dots, s \right\}, \quad (4.15)$$

which is a combination of the epigraphs of the asymptotically active components of the mapping L in (4.11). Observe that the set E is convex due to the convexity of the functions ϑ_i as $i = 0, \dots, s$ and also that $\text{int } E \neq \emptyset$.

Lemma 4.5 (representation of the normal cone to E). *If*

$$(\xi, -\lambda_0, \dots, -\lambda_s) \in \mathcal{N}((\bar{x}, L(\bar{x})), E), \quad (4.16)$$

then we have the relationships

$$\begin{cases} \lambda_i \geq 0 \text{ for } i = 0, \dots, s \text{ and} \\ \xi \in \sum_{i=0}^s \lambda_i \partial L_i(\bar{x}) = \sum_{i=0}^s \lambda_i (\nabla \psi_i(\bar{x}) + \partial \vartheta_i(\bar{x})). \end{cases}$$

Proof. From the definition (2.4) of the normal cone and (4.16) we conclude that

$$\langle \xi, x - \bar{x} \rangle - \sum_{i=0}^s \lambda_i (\nu_i - L_i(\bar{x})) \leq 0$$

for all x and $\nu_i \geq L_i(x)$, $i = 0, \dots, s$. Setting $x = \bar{x}$ we can see that this equality is possible only if $\lambda_i \geq 0$, $i = 0, \dots, s$. Setting $\nu_i = L_i(x)$, $i = 0, \dots, s$ we conclude that

$$\langle \xi, x - \bar{x} \rangle \leq \sum_{i=0}^s \lambda_i L_i(x) - \sum_{i=0}^s \lambda_i L_i(\bar{x}),$$

which implies via (2.1) that

$$\xi \in \partial \left(\sum_{i=0}^s \lambda_i L_i \right) (\bar{x}) = \sum_{i=0}^s \lambda_i \partial L_i(\bar{x}),$$

where the last equality is due to the subdifferential sum rule (2.5). □

Define further the convex hull

$$\Omega_N := \text{co} \{ \Delta x_N(t_1) \} \quad (4.17)$$

of the endpoint increments of the optimal trajectory \bar{x}_N to (P_N) generated by *all single needle* variations (4.9) with $r = 1$ of the optimal control \bar{u}_N for each $N \in \mathbb{N}$. The following crucial lemma asserts that the convex set $(\bar{x} + \Omega_N, L(\bar{x}))$ can be shifted by a vector c_N of order $o(h_N)$ so that it does not intersect with the interior of the set E . We can treat this result as a sequence of primal optimality conditions for (P_N) as $N \rightarrow \infty$.

Lemma 4.6 (primal optimality conditions for discrete approximations). *There exists a sequence $\{c_N\} \subset \mathbb{R}^{s+1}$ of order $o(h_N)$ as $N \rightarrow \infty$ such that*

$$(\bar{x} + \Omega_N, L(\bar{x}) + c_N) \cap \text{int } E = \emptyset, \quad N \in \mathbb{N}. \quad (4.18)$$

Proof. Consider the sequence of real numbers

$$\sigma_N := \min_y \max_{i \in \{0, \dots, s\}} (L_i(\bar{x} + y) - L_i(\bar{x})), \quad (4.19)$$

where the minimum is taken over the set of $y \in \Omega_N$ with $(\bar{x} + y, L(\bar{x})) \in E$; observe that the minimum is reached since this set is compact and nonempty containing $y = 0$. From the construction of σ_N in (4.19) it follows that for any $y_N \in \Omega_N$ with $(\bar{x} + y_N, L(\bar{x})) \in E$ there exists an index $i_1 = i_1(N) \in \{0, \dots, s\}$ such that

$$\sigma_N \leq L_{i_1}(\bar{x} + y_N) - L_{i_1}(\bar{x}). \quad (4.20)$$

Define $c_N := (\sigma_N, \dots, \sigma_N) \in \mathbb{R}^{s+1}$ and observe that

$$(\bar{x} + y_N, L(\bar{x}) + c_N) \notin \text{int } E, \quad (4.21)$$

since the contrary would mean that $L_i(\bar{x} + y_N) < L_i(\bar{x}) + \sigma_N$ for all i and thus contradict (4.20). It follows from (4.19) with $y = 0$ that $\sigma_N \leq 0$. The assertion of the lemma will now follow from (4.21) if we show that $\sigma_N = o(h_N)$ as $N \rightarrow \infty$.

Assume to the contrary that $\sigma_N \neq o(h_N)$ and find a sequence of $y_N \in \Omega_N$ and negative constants β_i as $i = 0, \dots, s$ such that

$$\liminf_{h_N \rightarrow 0} h_N^{-1} \left(L_i(\bar{x} + y_N) - L_i(\bar{x}) \right) \leq \beta_i < 0.$$

Recalling the definition (4.17) of the set Ω_N and invoking the Carathéodory Theorem, we represent any element $y_N \in \Omega_N$ by

$$y_N = \sum_{j=1}^p \alpha_j(N) \Delta_j x_N(t_1) \quad \text{with } p = n + 1,$$

where $\Delta_j x_N(t_1)$ are optimal trajectory increments generated by single needle variations of the optimal control \bar{u}_N with the needle variation parameters $(\tau_j(N), v_j(N))$ for each $j = 1, \dots, p$ and $N \in \mathbb{N}$, and where $\alpha_j(N) \geq 0$ with $\sum_{j=1}^p \alpha_j(N) = 1$. Therefore

$$\liminf_{h_N \rightarrow 0} h_N^{-1} \left(L_i(\bar{x} + \sum_{j=1}^p \alpha_j(N) \Delta_j x_N) - L_i(\bar{x}) \right) \leq \beta_i < 0, \quad i = 0, \dots, s. \quad (4.22)$$

For a given number $\eta > 0$ to be specified later, define the quantities

$$\gamma_j(N) := [\alpha_j(N)/\eta], \quad j = 1, \dots, p,$$

where $[a]$ stands, as usual, for the integer part of the positive number a . Along a subsequence (without relabeling) we have $\gamma_j(N) \rightarrow \gamma_j^0$ as $N \rightarrow \infty$ for $j = 1, \dots, p$. Due to Lemma 4.1,

there exist subgradients $\xi_i \in \partial L_i(\bar{x})$ as $i = 0, \dots, s$ such that

$$\begin{aligned} L_i(\bar{x} + \sum_{j=1}^p \gamma_j^0 \Delta_j x_N(t_1)) - L_i(\bar{x}) &= \left\langle \xi_i, \sum_{j=1}^p \gamma_j(N) \Delta_j x_N(t_1) \right\rangle + o(h_N) \\ &= \frac{1}{\eta} \left\langle \xi_i, \sum_{j=1}^p \alpha_j(N) \Delta_j x_N(t_1) \right\rangle + \left\langle \xi_i, \sum_{j=1}^p \left(\gamma_j - \frac{\alpha_j(N)}{\eta} \right) \Delta_j x_N(t_1) \right\rangle + o(h_N) \end{aligned} \quad (4.23)$$

along a subsequence of $N \rightarrow \infty$. Let M and M_1 be positive constants such that that $\|\Delta_j x_N(t_1)\| \leq M h_N$ and $\|\xi_i\| \leq M_1$ for als i and j . Since $\langle \xi_i, y \rangle \leq L_i(\bar{x} + y) - L_i(\bar{x})$ for all y we get from (4.23) that

$$\begin{aligned} L_i(\bar{x} + \sum_{j=1}^p \gamma_j^0 \Delta_j x_N(t_1)) - L_i(\bar{x}) &\leq \frac{1}{\eta} \left(L_i(\bar{x} + \sum_{j=1}^p \alpha_j(N) \Delta_j x_N(t_1)) - L_i(\bar{x}) \right) \\ &\quad + p M M_1 h_N + o(h_N). \end{aligned} \quad (4.24)$$

Now select an index $i_2 \in \{0, \dots, s\}$ so that $|\beta_{i_2}| = \max \{ |\beta_i| \mid 0 \leq i \leq s \}$ and set η above as

$$\eta = \frac{\beta_{i_2}}{\beta_{i_2} - p M M_1} \in (0, 1).$$

Then it follows from (4.24) and (4.22) that

$$\begin{aligned} \liminf_{h_N \rightarrow 0} h_N^{-1} \left(L_i(\bar{x} + \sum_{j=1}^p \gamma_j^0 \Delta_j x_N) - L_i(\bar{x}) \right) &\leq \frac{\beta_i}{\eta} + p M M_1 \\ &= \frac{\beta_i (\beta_{i_2} - p M M_1)}{\beta_{i_2}} + p M M_1 = \frac{p M M_1 (-\beta_i + \beta_{i_2}) + \beta_i \beta_{i_2}}{\beta_{i_2}} \leq \beta_i < 0 \end{aligned} \quad (4.25)$$

for all $i = 1, \dots, s$. Thus inequality (4.22) for convex combinations of $\Delta_j x_N$ as $j = 1, \dots, p$ implies the same type of inequality (4.25) for their integer combinations. Due to Lemma 4.3 there exists an r -needle variation of the optimal control \bar{u}_N with $r = \sum_{j=1}^p \gamma_j^0$ such that

$$\sum_{j=1}^p \gamma_j^0 \Delta_j x_N(t_1) = \Delta^r x_N(t_1) + o(h_N),$$

which ensures by (4.25) the inequalities

$$\liminf_{h_N \rightarrow 0} h_N^{-1} \left(L_i(\bar{x} + \Delta^r x_N(t_1)) - L_i(\bar{x}) \right) \leq \beta_i < 0 \quad \text{for all } i = 0, \dots, s.$$

The latter inequalities contradict (4.12) and thus complete the proof of the lemma. \square

The next lemma based on convex separation transforms the primal optimality conditions (4.18) of Lemma 4.6 into dual conditions via appropriate Lagrange multipliers.

Lemma 4.7 (Lagrange multipliers for discrete approximation problems). *Given a limiting point \bar{x} for the endpoint sequence $\{\bar{x}_N(t_1)\}$ of optimal trajectories to (P_N) , there exist sequences of Lagrange multipliers $\lambda_N = (\lambda_{0N}, \dots, \lambda_{lN}) \in \mathbb{R}^{l+1}$ and dual vectors $\xi_N \in \mathbb{R}^n$ satisfying the conditions*

$$\begin{cases} \lambda_{iN} \geq 0 \text{ for } i = 0, \dots, l, & \sum_{i=0}^l \lambda_{iN}^2 = 1, \\ \lambda_{iN} \varphi_i(\bar{x}) = 0 \text{ for } i = 1, \dots, l, & \text{and} \\ \xi_N \in \sum_{i=0}^l \lambda_{iN} \partial L_i(\bar{x}) = \sum_{i=0}^l \lambda_{iN} (\nabla \psi_i(\bar{x}) + \partial \vartheta_i(\bar{x})). \end{cases} \quad (4.26)$$

Furthermore, for an arbitrary sequence of endpoint increments $\Delta x_N(t_1)$ generated by single needle variations of optimal controls we have

$$\langle \xi_N, \Delta x_N(t_1) \rangle \geq o(h_N) \text{ as } N \rightarrow \infty. \quad (4.27)$$

Proof. Applying the classical Separation Theorem to the convex sets $(\bar{x} + \Omega_N, L(\bar{x}) + c_N)$ and E with empty intersection (4.18), we find dual elements $\tilde{\lambda}_N = (\tilde{\lambda}_{0N}, \dots, \tilde{\lambda}_{sN}) \in \mathbb{R}^{s+1}$ and $\tilde{\xi}_N \in \mathbb{R}^n$ such that

$$\|\tilde{\xi}_N\|^2 + \sum_{i=0}^s \tilde{\lambda}_{iN}^2 = 1$$

and that for any $\Delta x_N(t_1) \in \Omega_N$ and $(x, y) \in E$ the inequality

$$\langle \tilde{\xi}_N, \bar{x} + \Delta x_N(t_1) \rangle - \langle \tilde{\lambda}_N, L(\bar{x}) + c_N \rangle \geq \langle \tilde{\xi}_N, x \rangle - \langle \tilde{\lambda}_N, y \rangle,$$

holds, which can be equivalently written as

$$\langle \tilde{\xi}_N, \Delta x_N(t_1) \rangle - \langle \tilde{\lambda}_N, c_N \rangle \geq \langle \tilde{\xi}_N, x - \bar{x} \rangle - \langle \tilde{\lambda}_N, y - L(\bar{x}) \rangle. \quad (4.28)$$

Setting $\Delta x_N(t_1) = 0$ in (4.28) gives us

$$\langle \tilde{\xi}_N, x - \bar{x} \rangle - \langle \tilde{\lambda}_N, y - L(\bar{x}) \rangle \leq -\langle \tilde{\lambda}_N, c_N \rangle, \quad (4.29)$$

while by setting $x = \bar{x}$ and $y = L(\bar{x})$ in (4.28) we obtain

$$\langle \tilde{\xi}_N, \Delta x_N(t_1) \rangle \geq \langle \tilde{\lambda}_N, c_N \rangle = o(h_N) \text{ for any } \Delta x_N(t_1) \in \Omega_N. \quad (4.30)$$

We can see from (4.29) and the definition of the normal cone (2.4) that the vector $(\tilde{\xi}_N, -\tilde{\lambda}_N)$ “almost belongs” to $\mathcal{N}(((\bar{x}, L(\bar{x})), E))$, or, more precisely,

$$\text{dist} \left((\tilde{\xi}_N, -\tilde{\lambda}_N), \mathcal{N}(((\bar{x}, L(\bar{x})), E)) \right) \rightarrow 0 \text{ as } N \rightarrow \infty.$$

Denote by $(\xi_N, -\lambda_N) = (\xi_N, -\lambda_{0N}, \dots, -\lambda_{sN}) \in \mathcal{N}((\bar{x}, L(\bar{x})); E)$ the (unique) element of the normal cone to the convex set (4.15) at $(\bar{x}, L(\bar{x}))$ that is nearest to $(\tilde{\xi}_N, -\tilde{\lambda}_N)$. As we noticed above, we have

$$\|(\xi_N, \lambda_N) - (\tilde{\xi}_N, \tilde{\lambda}_N)\| \rightarrow 0 \quad \text{as } N \rightarrow \infty, \quad (4.31)$$

and, therefore,

$$\|\xi_N\|^2 + \sum_{i=0}^s \lambda_{iN}^2 \rightarrow 1 \quad \text{as } N \rightarrow \infty. \quad (4.32)$$

Invoking Lemma 4.5 we conclude that

$$\begin{cases} \lambda_{iN} \geq 0 \quad \text{for } i = 0, \dots, s, \\ \xi_N \in \sum_{i=0}^s \lambda_{iN} \partial L_i(\bar{x}) = \sum_{i=0}^s \lambda_{iN} (\nabla \psi_i(\bar{x}) + \partial \vartheta_i(\bar{x})). \end{cases}$$

Relation (4.32) implies that the multipliers λ_{iN} as $i = 0, \dots, s$ can be renormalized to satisfy $\sum_{i=0}^s \lambda_{iN}^2 = 1$. Setting now $\lambda_{iN} = 0$, $i = s+1, \dots, l$ for the Lagrange multipliers corresponding to the asymptotically inactive constraints, we get all the conditions in (4.26). Finally, it follows from (4.30) and (4.31) that condition (4.27) also holds, which thus concludes the proof of the lemma. \square

To complete the proof of Theorem 3.1 it remains to show that the obtained estimate (4.27) implies the approximate maximum condition (1.7) along the adjoint trajectory $p_N(t)$ of system (1.8) satisfying the endpoint/transversality condition

$$-p_N(t_1) \in \sum_{i=0}^l \lambda_{iN} (\nabla \psi_i(\bar{x}) + \partial \vartheta_i(\bar{x})). \quad (4.33)$$

Completing the proof of Theorem 3.1. Using Lemma 4.7, we find vectors $\xi_N \in \mathbb{R}^n$ and $\lambda_N = (\lambda_{0N}, \dots, \lambda_{lN}) \in \mathbb{R}^{l+1}$ satisfying conditions (4.26) and (4.27) for all $N \in \mathbb{N}$. Define now $p_N(t_1) := -\xi_N$ and observe that conditions (4.26) reduce to those in (3.1)–(3.3) of Theorem 3.1 for problems (P_N) with $q = 0$, while condition (4.27) can be rewritten in the form

$$\langle p_N(t_1), \Delta x_N(t_1) \rangle \leq o(h_N). \quad (4.34)$$

Recall that $\Delta x_N(t)$ are optimal trajectory increments generated by the following single needle variations of optimal control $\bar{u}_N(t)$:

$$u_N(t) = \begin{cases} v(N), & t = \tau(N), \\ \bar{u}_N(t), & t \in T_N, t \neq \tau(N). \end{cases}$$

For any function $p_N(t)$, $t \in T_N \cup \{t_1\}$, we have the identity

$$\begin{aligned} \langle p_N(t_1), \Delta x_N(t_1) \rangle &= \sum_{t=t_0}^{t_1-h_N} \langle p_N(t+h_N) - p_N(t), \Delta x_N(t) \rangle \\ &+ \sum_{t=t_0}^{t_1-h_N} \langle p_N(t+h_N), \Delta x_N(t+h_N) - \Delta x_N(t) \rangle. \end{aligned} \quad (4.35)$$

Using further the notation

$$\begin{aligned} \Delta f(t) &:= f(x_N(t), u_N(t), t) - f(\bar{x}_N(t), \bar{u}_N(t), t), \\ \Delta_u f(t) &:= f(\bar{x}_N(t), u_N(t), t) - f(\bar{x}_N(t), \bar{u}_N(t), t), \\ \Delta_u H(t) &:= \langle p_N(t+h_N), \Delta_u f(t) \rangle \end{aligned}$$

we obtain the representation

$$\Delta x_N(t+h_N) - \Delta x_N(t) = h_N \Delta f(t) = h_N \left[\Delta_u f(t) + \frac{\partial f}{\partial x}(\bar{x}_N(t), \bar{u}_N(t), t) \Delta x_N(t) + \eta_N(t) \right],$$

where the remainder $\eta_N(t)$ is defined by

$$\eta_N(t) := \left(\frac{\partial f}{\partial x}(\bar{x}_N(t), u_N(t), t) - \frac{\partial f}{\partial x}(\bar{x}_N(t), \bar{u}_N(t), t) \right) \Delta x_N(t) + o(\|\Delta x_N(t)\|).$$

Since $\Delta x_N(t) = 0$ for $t \leq \tau = \tau(N)$ and $(\frac{\partial f}{\partial x}(\bar{x}_N(t), u_N(t), t) - \frac{\partial f}{\partial x}(\bar{x}_N(t), \bar{u}_N(t), t)) = 0$ for all $t \in T_N$ except $t = \tau$, the first term in the formula for $\eta_N(t)$ vanishes, and therefore we arrive at the relationships $\eta_N(t) = o(\|\Delta x_N(t)\|) = o(h_N)$.

This allows us to represent the second sum on the right-hand side of (4.35) as

$$\begin{aligned} &\sum_{t=t_0}^{t_1-h_N} \langle p_N(t+h_N), \Delta x_N(t+h_N) - \Delta x_N(t) \rangle \\ &= h_N \sum_{t=t_0}^{t_1-h_N} \Delta_u H(t) + h_N \sum_{t=t_0}^{t_1-h_N} \left\langle p_N(t+h_N), \frac{\partial f}{\partial x}(\bar{x}_N(t), \bar{u}_N(t), t) \Delta x_N(t) + \eta_N(t) \right\rangle \quad (4.36) \\ &= h_N \Delta_u H(\tau) + h_N \sum_{t=t_0}^{t_1-h_N} \left\langle p_N(t+h_N), \frac{\partial f}{\partial x}(\bar{x}_N(t), \bar{u}_N(t), t) \Delta x_N(t) \right\rangle + o(h_N). \end{aligned}$$

If $p_N(t)$ satisfies the adjoint system (1.8), we can rewrite the first sum on the right-hand side of (4.35) as

$$\begin{aligned} &\sum_{t=t_0}^{t_1-h_N} \langle p_N(t+h_N) - p_N(t), \Delta x_N(t) \rangle = -h_N \sum_{t=t_0}^{t_1-h_N} \left\langle \frac{\partial H}{\partial x}(p_N(t+h_N) \bar{x}_N(t), \bar{u}_N(t), t), \Delta x_N(t) \right\rangle \\ &= -h_N \sum_{t=t_0}^{t_1-h_N} \left\langle p_N(t+h_N), \frac{\partial f}{\partial x}(\bar{x}_N(t), \bar{u}_N(t), t) \Delta x_N(t) \right\rangle. \end{aligned} \quad (4.37)$$

It follows from (4.34)–(4.37) that

$$o(h_N) \geq \langle p_N(t_1), \Delta x_N(t_1) \rangle = h_N \Delta_u H(\tau) + o(h_N),$$

which can be written in the form

$$h_N \Delta_u H(\tau) \leq o(h_N).$$

This implies the approximate maximum condition (1.7) at the point $t = \tau(N)$. Since $\tau(N) \in T_N$ was chosen arbitrarily, we justified the approximate maximum condition for all $t \in T_N$ and thus completed the proof of the theorem. \square

5 Problems with Equality Constraints

In the concluding section of the paper we show that the Approximate Maximum Principle of Theorem 3.1 does not hold if $\delta = 0$ in problems (P_N) , i.e., in the case of unrelaxed equality constraints. The following simple example illustrates this phenomenon.

Example 5.1 (violation of the AMP in discrete approximation problems with unrelaxed equality constraints). Consider a two-dimensional continuous-time problem

$$(P) \left\{ \begin{array}{l} \text{minimize } x(1) \text{ subject to} \\ \dot{x} = u \text{ a.e. } t \in T := [0, 1], \\ \dot{y} = v \text{ a.e. } t \in T, \\ x(0) = y(0) = 0, \\ (u(t), v(t)) \in U := \{(0, 0), (-1, \sqrt{2}), (-1, -1)\} \text{ a.e. } t \in T, \\ y(1) = 0. \end{array} \right.$$

The corresponding discrete approximation problems (P_N) defined in Section 1 with $\delta = 0$ are given by

$$(P_N) \left\{ \begin{array}{l} \text{minimize } x_N(1) \text{ subject to} \\ x_N(t + h_N) = x_N(t) + h_N u_N(t), \quad t \in T_N := \{0, h_N, \dots, 1 - h_N\}, \\ y_N(t + h_N) = y_N(t) + h_N v_N(t), \quad t \in T_N, \\ x_N(0) = y_N(0) = 0, \\ (u_N(t), v_N(t)) \in U, \quad t \in T_N, \\ y_N(1) = 0, \\ h_N = \frac{1}{N}, \quad N = 1, 2, \dots \end{array} \right.$$

It is easy to see that the optimal control to problem (P) is

$$(\bar{u}(t), \bar{v}(t)) = \begin{cases} (-1, \sqrt{2}) & \text{for } 0 \leq t \leq 1/(1 + \sqrt{2}), \\ (-1, -1) & \text{for } 1/(1 + \sqrt{2}) < t \leq 1. \end{cases}$$

We can also observe that, due to the incommensurability of the radical $\sqrt{2}$ and the discretization step $h_N = 1/N$, the only feasible control for each discrete approximation problem (P_N) satisfying the constraint $y_N(1) = 0$ is $(u_N(t), v_N(t)) \equiv (0, 0)$. It remains to show that this control does not satisfy the necessary optimality conditions of Theorem 3.1.

To proceed, note that the Hamilton-Pontryagin function (1.3) reduces in the this case to

$$H(p_{1N}, p_{2N}, u_N, v_N) = p_{1N}u_N + p_{2N}v_N \text{ for all } t \in T_N, \quad (5.1)$$

where both p_{1N} and p_{2N} are constants due to the adjoint system (1.6). The endpoint condition (3.3) is written now as

$$(p_{1N}(1), p_{2N}(1)) = -\nabla_{(x,y)}(\lambda_{0N}x + \lambda_{1N}y)|_{(x,y)=(x_N(1), y_N(1))} = (-\lambda_{0N}, -\lambda_{1N})$$

with $(\lambda_{0N}, \lambda_{1N})$ satisfying the sign and nontriviality conditions

$$\lambda_{0N} \geq 0 \text{ and } \lambda_{0N}^2 + \lambda_{1N}^2 = 1 \quad (5.2)$$

by (3.1). This allows us to rewrite function (5.1) in the form

$$H(\lambda_{0N}, \lambda_{1N}, u_N, v_N) = -\lambda_{0N}u_N - \lambda_{1N}v_N.$$

A simple analysis shows that for any λ_{0N} and λ_{1N} satisfying (5.2) there exists a constant $\alpha > 0$ such that, depending on the sign of λ_{1N} , we have

$$\begin{cases} \text{either } H(\lambda_{0N}, \lambda_{1N}, -1, \sqrt{2}) > H(\lambda_{0N}, \lambda_{1N}, 0, 0) + \alpha \\ \text{or } H(\lambda_{0N}, \lambda_{1N}, -1, -1) > H(\lambda_{0N}, \lambda_{1N}, 0, 0) + \alpha, \end{cases}$$

which contradicts the approximate maximum condition (1.7).

We wish to note that this example can be slightly modified (we are not presenting the details) to illustrate the failure of the AMP if the equality constraint $y_N(1) = 0$ in problems (P_N) of Example 5.1 is relaxed to $|y_N(1)| \leq \delta_N$, where $\delta_N \downarrow 0$ too fast as $N \rightarrow \infty$; cf. the discussions in Remark 3.2.

References

- [1] R. Gabasov and F.M. Kirillova, On the extension of the maximum principle by L.S. Pontryagin to discrete systems, *Autom. Remote Control* **27** (1966), 1878–1882.
- [2] B.S. Mordukhovich, Approximate maximum principle for finite-difference control systems, *Comput. Maths. Math. Phys.* **28** (1988), 106–114.
- [3] B.S. Mordukhovich, Discrete approximations and refined Euler-Lagrange conditions for nonconvex differential inclusions, *SIAM J. Control Optim.* **33** (1995), 882–915.

- [4] B.S. Mordukhovich, *Variational Analysis and Generalized Differentiation, I: Basic Theory, II: Applications*, Springer, Berlin, 2006.
- [5] B.S. Mordukhovich and I. Shvartsman, The approximate maximum principle in constrained optimal control, *SIAM J. Control Optim.* **43** (2004), 1037–1062.
- [6] B.S. Mordukhovich and I. Shvartsman, Nonsmooth approximate maximum principle in optimal control, *Proc. 50th IEEE Conf. Dec. Cont.*, Orlando, FL, 2011.
- [7] K. Nitka-Styczen, Approximate discrete maximum principle for the discrete approximation of optimal periodic control problems, *Int. J. Control* **50** (1989), 1863–1871.
- [8] L.C. Pontryagin, V.G. Boltyanskii, R.V. Gamkrelidze and E.F. Mishchenko, *The Mathematical Theory of Optimal Processes*, Wiley, New York, 1962.
- [9] R.T. Rockafellar, *Convex Analysis*, Princeton University Press, Princeton, NJ, 1973.
- [10] G.V. Smirnov, *Introduction to the Theory of Differential Inclusions*, American Mathematical Society, Providence, RI, 2002.
- [11] R.B. Vinter, *Optimal Control*, Birkhäuser, Boston, 2000.