

Global convergence of trust-region algorithms for constrained minimization without derivatives

P.D. Conejo[§] E.W. Karas[¶] A.A. Ribeiro[¶] L.G. Pedroso[¶] M. Sachine[¶]

September 27, 2012

Abstract

In this work we propose a trust-region algorithm for the problem of minimizing a function within a convex closed domain. We assume that the objective function is differentiable but no derivatives are available. The algorithm has a very simple structure and allows a great deal of freedom in the choice of the models. Under reasonable assumptions for derivative-free schemes, we prove global convergence for the algorithm, that is to say, that all accumulation points of the sequence generated by the algorithm are stationary.

1 Introduction

In this work we will discuss global convergence of a general derivative-free trust-region algorithm for solving the nonlinear programming problem

$$\begin{aligned} & \text{minimize} && f(x) \\ & \text{subject to} && x \in \Omega, \end{aligned} \tag{1}$$

where $\Omega \subset \mathbb{R}^n$ is a nonempty closed convex set and $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is a differentiable function. Although the objective function is smooth, we assume that its derivatives are not available. This situation is common in a wide range of applications [8], particularly when the objective function is provided by a simulation package or a black box. Such practical situations have been motivating research on derivative-free optimization in the recent years [8, 10].

There are many derivative-free methods with global convergence results for handling the problem (1). When the feasible set is defined by linear constraints, the problem can be solved by GSS method (*Generating Set Search*) presented in [18]. This method encompasses many algorithms, including Generalized Pattern Search [19]. In [2], the authors proposed an inexact-restoration scheme where the GSS algorithm is used in the optimality phase, therefore avoiding the evaluation of the gradient of the objective function. In [13, 17], the authors present Augmented Lagrangian methods for generally constrained problems. In [17], the subproblems are linearly constrained and solved by GSS algorithm. In [13], any class of constraints can be considered on the subproblems, since one can provide a suitable derivative-free algorithm for handling them.

Here we consider the class of derivative-free trust-region methods, which has been pioneered by Winfield [30] and exhaustively studied for unconstrained and box-constrained problems by Powell [21, 22, 23, 24, 26], Conn and Toint [11], Conn, Scheinberg and Toint [7], Conn, Scheinberg and Vicente [10], Fasano, Morales and Nocedal [14], Gratton, Toint and Tröltzsch [16], and for linearly constrained problems by Powell [25]. Global convergence results for the unconstrained case are presented in [10, 16, 26]. In these works

[§]Ph.D. Program in Numerical Methods in Engineering, Federal University of Paraná. Partially supported by Fundação Araucária. State University of Paraná, Dep. of Mathematics, 85819-110, Cascavel, PR, Brazil. E-mail: paulo.conejo@unioeste.br

[¶]Department of Mathematics, Federal University of Paraná. Cx. Postal 19081, 81531-980 Curitiba, PR, Brazil. Partially supported by CNPq, Grants 472313/2011-8 and 307714/2011-0. E-mail: ewkaras@ufpr.br, lucaspedroso@ufpr.br, ademir.ribeiro@ufpr.br, mael@ufpr.br.

the models are based on polynomial interpolation. Essentially, the difference between them consists in the interpolation set updating.

With regard to derivative based trust-region methods, for both constrained and unconstrained problems, well-established algorithms with global convergence results can be found in the literature, for example in [4, 5, 6, 20, 27]. When the derivatives of the objective function are not available, there are also trust-region methods with good practical performance [1, 24, 28, 29]. However, until now, theoretical results have not been established for the constrained case. As far as we know, this work is the first one to present global convergence results for a class of derivative-free trust-region methods for constrained optimization.

The proposed algorithm considers quadratic models that approximate the objective function based only on zero-order information. Nevertheless, the gradient of the models must represent adequately the gradient of the objective function at the current point. This property can be achieved by many derivative-free techniques, most of them based in polynomial interpolation [3, 9, 10, 12].

At each iteration, the algorithm considers the Euclidean projection of the gradient of the model at the current point onto the feasible set Ω . The model is minimized subject to Ω and to the trust region, in the sense that an approximate solution of this subproblem must satisfy a Cauchy-type condition. This solution will be accepted or rejected as the new iterate according to the ratio between the actual and predicted reductions, a classical trust-region procedure. In particular, the proposed algorithm can be applied for solving unconstrained problems. In this case, the considered projection is reduced to the norm of the gradient of the model. Furthermore, the classical condition on the Cauchy step can be used as an acceptance criterion for the solution of the subproblem.

The paper is organized as follows. In Section 2, we propose a derivative-free trust-region algorithm. In Section 3, we present its global convergence analysis. Conclusions are stated in Section 4. Throughout the paper, the symbol $\|\cdot\|$ denotes the Euclidean norm.

2 The algorithm

In this section we present a general trust-region algorithm for solving the problem (1) that generates a sequence of approximate minimizers of quadratic constrained subproblems. The algorithm allows a great deal of freedom in the construction and resolution of the subproblems.

At each iteration $k \in \mathbb{N}$, we consider the current iterate $x^k \in \Omega$ and the quadratic model

$$Q_k(d) = f(x^k) + (g^k)^T d + \frac{1}{2} d^T G_k d,$$

where $g^k \in \mathbb{R}^n$ and $G_k \in \mathbb{R}^{n \times n}$ is a symmetric matrix. Any quadratic model in this form can be used, as long as it provides a sufficiently accurate approximation of the objective function, in the sense that $g^k = \nabla Q_k(0)$ and G_k satisfy Hypotheses H3 and H4 discussed ahead. We assume little about the Hessian of the models, just symmetry and uniform boundedness, allowing that even linear models may be used by setting $G_k = 0$. We do not use Taylor models because we are interested in the case where, although derivatives exist, they are not available.

We consider the stationarity measure at x^k for the problem of minimizing Q_k over the set Ω defined by

$$\pi_k = \|P_\Omega(x^k - g^k) - x^k\|,$$

where P_Ω denotes the orthogonal projection onto Ω , which exists because it is a closed convex set. We say that a point $x^* \in \Omega$ is stationary for the original problem (1) when $\|P_\Omega(x^* - \nabla f(x^*)) - x^*\| = 0$. This definition is based on the fact that, since Ω is convex, a point satisfies the Karush-Kuhn-Tucker (KKT) conditions if and only if the norm of the projection onto Ω of the gradient of the objective function vanishes at this point [6, 27].

For proving convergence to stationary points, we assume that an approximate solution $d^k \in \mathbb{R}^n$ of the trust-region subproblem

$$\begin{aligned} & \text{minimize} && Q_k(d) \\ & \text{subject to} && x^k + d \in \Omega \\ & && \|d\| \leq \Delta_k \end{aligned} \tag{2}$$

satisfies the efficiency condition given by

$$Q_k(0) - Q_k(d^k) \geq c_1 \pi_k \min \left\{ \frac{\pi_k}{1 + \|G_k\|}, \Delta_k, 1 \right\}, \quad (3)$$

where $c_1 > 0$ is a constant independent of k .

Conditions of this kind are well-known in trust-region approaches and were used by several authors in different situations. In the unconstrained case, in which $\Omega = \mathbb{R}^n$, the stationarity measure π_k is simply $\|g^k\|$ and the classical Cauchy step satisfies a similar condition, as proved in [20, Lemma 4.5] and [10, Theorem 10.1] with and without derivatives of the objective function, respectively. Conditions of this type also appear throughout [6], under different contexts. In [15], the authors prove the global convergence of a filter method for nonlinear programming by assuming that an approximate solution of the subproblems satisfies a condition analogous to (3). For bound-constrained nonlinear optimization without derivatives, Tröltzsch [29] also assumes such a condition.

Once computed an approximate solution of (2), we analyse whether or not it produces a satisfactory decrease in the model. As usual in trust-region methods, the trial step is assessed by means of the ratio

$$\gamma_k = \frac{f(x^k) - f(x^k + d^k)}{Q_k(0) - Q_k(d^k)}. \quad (4)$$

We present now a general derivative-free trust-region algorithm with no specification on the model update and on the internal solver for the subproblems. Later we will state assumptions to prove that any accumulation point of the sequence generated by the algorithm is stationary.

Algorithm 1. *General Algorithm*

Data: $x^0 \in \Omega$, $\alpha > 0$, $\Delta_0 > 0$, $\eta \in (0, \frac{3}{4})$.

Set $k = 0$.

REPEAT

 Construct the model Q_k .

 IF $\Delta_k > \alpha \pi_k$, then

$$\Delta_{k+1} = \frac{\Delta_k}{2}, d^k = 0 \text{ and } x^{k+1} = x^k.$$

 ELSE

 Find an approximate solution d^k of (2).

 IF $\gamma_k \geq \eta$, then

$$x^{k+1} = x^k + d^k \text{ and } \Delta_{k+1} = \Delta_k.$$

 ELSE

$$x^{k+1} = x^k \text{ and } \Delta_{k+1} = \frac{\Delta_k}{2}.$$

$k = k + 1$.

Note that the model is updated whenever a new point is computed. Otherwise, the trust-region radius is halved. We point out that any other factor could be used to perform the radius reduction. Observe that the sequence of trust-region radii is nonincreasing. Moreover, we will prove in the next section that $\Delta_k \rightarrow 0$ as $k \rightarrow \infty$, what will be important in the proofs of the convergence results. This also suggests that, given a tolerance $\varepsilon > 0$, the condition $\Delta_k \leq \varepsilon$ can be used as a stopping criterion on implementations of the algorithm. When π_k is small, the iterate is probably close to a solution of the problem of minimizing the model within the feasible set Ω . On the other hand, if Δ_k is large, we cannot guarantee that the model represents properly the objective function. Hence, when $\Delta_k > \alpha \pi_k$, the trust-region radius is reduced in the attempt of finding a more accurate model. Although we could always set $\alpha = 1$, this parameter might be used to balance the magnitude of π_k and Δ_k , according to the problem.

3 Convergence

In this section we prove, under reasonable assumptions, that every accumulation point of the sequence generated by the algorithm is stationary. From now on we assume that the algorithm generates an infinite sequence $(x^k) \subset \Omega$.

For proving the convergence we consider the following hypotheses.

H1. The function f is differentiable and its gradient ∇f is Lipschitz continuous with constant $L > 0$, in Ω .

H2. The function f is bounded below in Ω .

H3. The matrices G_k are uniformly bounded, that is, there exists a constant $\beta \geq 1$ such that $\|G_k\| \leq \beta - 1$ for all $k \geq 0$.

H4. There exists a constant $c_2 > 0$ such that

$$\|g^k - \nabla f(x^k)\| \leq c_2 \Delta_k$$

for all $k \in \mathbb{N}$.

Hypotheses H1 and H2 impose conditions on the objective function, whereas H3 and H4 describe properties that the interpolation models must satisfy. The first three hypotheses are usual in convergence analysis for both, derivative-free and derivative-based trust-region algorithms. Hypothesis H4 states that the model has to properly represent the objective function near the current point. There are algorithms able to find models with such properties without computing $\nabla f(x^k)$, for instance [10, Chapter 6]. The proposed algorithm allows the usage of any technique to fulfill Hypothesis H4, although in literature the most usual procedure is polynomial interpolation [10, 14, 16, 22, 28]. Hypothesis H4 is omitted in some practical algorithms that have good numerical results [8, 14], but in our algorithm it is a fundamental property to establish the global convergence.

For the purpose of our analysis, we shall consider the set of indices of the *successful* iterations defined by

$$\mathcal{S} = \{k \in \mathbb{N} \mid \gamma_k \geq \eta\}. \quad (5)$$

In the following lemma, the constants c_1 , L , β and c_2 are the ones defined in (3) and in Hypotheses H1, H3, H4, respectively. The lemma establishes that if the trust-region radius is sufficiently small, then the algorithm will perform a successful iteration.

Lemma 3.1. Suppose that Hypotheses H1, H3 and H4 hold. Consider the set

$$\mathcal{K} = \left\{k \in \mathbb{N} \mid \Delta_k \leq \min \left\{ \frac{\pi_k}{\beta}, \frac{\pi_k}{4c}, \alpha\pi_k, 1 \right\} \right\}, \quad (6)$$

where $c = \frac{L + c_2 + \frac{\beta}{2}}{c_1}$. If $k \in \mathcal{K}$, then $k \in \mathcal{S}$.

Proof. By the Mean Value Theorem, there exists $t_k \in (0, 1)$ such that

$$f(x^k + d^k) = f(x^k) + \nabla f(x^k + t_k d^k)^T d^k.$$

Therefore, by Hypotheses H1, H3 and H4,

$$\begin{aligned} |f(x^k) - f(x^k + d^k) - Q_k(0) + Q_k(d^k)| &= \left| -(\nabla f(x^k + t_k d^k) - g^k)^T d^k + \frac{1}{2}(d^k)^T G_k d^k \right| \\ &= \left| -(\nabla f(x^k + t_k d^k) - g^k - \nabla f(x^k) + \nabla f(x^k))^T d^k + \frac{1}{2}(d^k)^T G_k d^k \right| \\ &\leq (\|\nabla f(x^k + t_k d^k) - \nabla f(x^k)\| + \|\nabla f(x^k) - g^k\|) \|d^k\| + \frac{1}{2} \|d^k\|^2 \|G_k\| \\ &\leq (t_k L \|d^k\| + c_2 \Delta_k) \|d^k\| + \frac{1}{2} \beta \|d^k\|^2. \end{aligned}$$

Since $\|d^k\| \leq \Delta_k$ and $t_k \in (0, 1)$, we have that

$$|f(x^k) - f(x^k + d^k) - Q_k(0) + Q_k(d^k)| \leq c_0 \Delta_k^2, \quad (7)$$

where $c_0 = L + c_2 + \frac{\beta}{2}$.

From (6), for every $k \in \mathcal{K}$ we have that $\Delta_k \leq \alpha \pi_k$ and consequently $\pi_k > 0$. Then, it follows from (3) that $Q_k(0) - Q_k(d^k) \neq 0$. Thus, from expressions (4), (7) and (3), for all $k \in \mathcal{K}$

$$\begin{aligned} |\gamma_k - 1| &= \left| \frac{f(x^k) - f(x^k + d^k) - Q_k(0) + Q_k(d^k)}{Q_k(0) - Q_k(d^k)} \right| \\ &\leq \frac{c_0 \Delta_k^2}{c_1 \pi_k \min \left\{ \frac{\pi_k}{\beta}, \Delta_k, 1 \right\}} \\ &= \frac{c \Delta_k^2}{\pi_k \min \left\{ \frac{\pi_k}{\beta}, \Delta_k, 1 \right\}}. \end{aligned}$$

By (6),

$$\Delta_k = \min \left\{ \frac{\pi_k}{\beta}, \Delta_k, 1 \right\} \quad \text{and} \quad \frac{c \Delta_k}{\pi_k} \leq \frac{1}{4}.$$

Therefore $|\gamma_k - 1| \leq \frac{1}{4}$ and hence $\gamma_k \geq \frac{3}{4} > \eta$. Consequently the iteration k is successful. \square

The next lemma provides a weak convergence result for the problem of minimizing the model within the feasible set Ω . We prove that the sequence of projected gradients of the models onto Ω has a subsequence converging to zero.

Lemma 3.2. *Suppose that Hypotheses H1 to H4 hold. Then*

$$\liminf_{k \rightarrow \infty} \pi_k = 0.$$

Proof. The proof is by contradiction. Without loss of generality, suppose that there exists $\varepsilon > 0$ such that for all $k \in \mathbb{N}$, $\pi_k \geq \varepsilon$. Take $\tilde{\Delta} = \min \left\{ \frac{\varepsilon}{\beta}, \frac{\varepsilon}{4c}, \alpha\varepsilon, 1 \right\}$ where β is the constant of Hypothesis H3, c is defined in Lemma 3.1 and $\alpha > 0$ is given in the Algorithm 1.

If $\Delta_k \leq \tilde{\Delta}$, then $k \in \mathcal{K}$, with \mathcal{K} given in (6). By Lemma 3.1 the iteration k is successful and thus $\Delta_{k+1} = \Delta_k$. It follows that the radius of the trust region can only decrease if $\Delta_k > \tilde{\Delta}$, and in this case, $\Delta_{k+1} = \frac{\Delta_k}{2} > \frac{\tilde{\Delta}}{2}$.

Therefore, for all $k \in \mathbb{N}$,

$$\Delta_k \geq \min \left\{ \frac{\tilde{\Delta}}{2}, \Delta_0 \right\}. \quad (8)$$

If \mathcal{S} is finite, then Δ_k converges to zero, contradicting (8). On the other hand, if \mathcal{S} is infinite, we have from (3) that for all $k \in \mathcal{S}$,

$$\begin{aligned} f(x^k) - f(x^{k+1}) &= f(x^k) - f(x^k + d^k) \\ &\geq \eta [Q_k(0) - Q_k(d^k)] \\ &\geq \eta c_1 \pi_k \min \left\{ \frac{\pi_k}{\beta}, \Delta_k, 1 \right\}. \end{aligned}$$

Then, using (8) and the contradiction assumption that $\pi_k \geq \varepsilon$, we conclude that $f(x^k) - f(x^{k+1})$ is bounded away from zero. However, by Hypothesis H2, the sequence $(f(x^k))$ is bounded below, and as it is also nonincreasing, it follows that $f(x^k) - f(x^{k+1}) \rightarrow 0$, providing a contradiction what concludes the proof. \square

Hypothesis H4 says that the smaller Δ_k , the better the models represent the objective function. Therefore, it is reasonable to expect that the trust-region radius converges to zero. In the following lemma, we show that the proposed algorithm has this property.

Lemma 3.3. *Suppose that Hypotheses H1 to H4 hold. Then the sequence (Δ_k) converges to zero.*

Proof. Consider the set of indices

$$\mathbb{N}' = \{k \in \mathbb{N} \mid \Delta_k > \alpha\pi_k\}.$$

From the mechanism of Algorithm 1, for every $k \in \mathbb{N}'$, we have $\Delta_{k+1} = \Delta_k/2$. Suppose that \mathbb{N}' is infinite. Then, the trust-region radius is halved infinitely many times and, since (Δ_k) is nonincreasing, the whole sequence converges to zero. On the other hand, if \mathbb{N}' is finite, there exists $k_0 \in \mathbb{N}$ such that $\Delta_k \leq \alpha\pi_k$ for all $k \geq k_0$. But, by Lemma 3.2, (π_k) has a subsequence converging to zero. Therefore, the corresponding subsequence of (Δ_k) converges to zero. Again, the monotonicity ensures that $\Delta_k \rightarrow 0$. \square

Using the previous lemma, we can now prove that not only there exists a subsequence of π_k converging to zero, as stated in Lemma 3.2, but also that the convergence prevails for the whole sequence.

Lemma 3.4. *Suppose that Hypotheses H1 to H4 hold. Then*

$$\lim_{k \rightarrow \infty} \pi_k = 0.$$

Proof. Suppose by contradiction that for some $\varepsilon > 0$ the set

$$\mathbb{N}' = \{k \in \mathbb{N} \mid \pi_k \geq \varepsilon\} \quad (9)$$

is infinite. By Lemma 3.3, the sequence (Δ_k) converges to zero. Then, there exists $k_0 \in \mathbb{N}$ such that for all $k \geq k_0$,

$$\Delta_k \leq \min \left\{ \frac{\varepsilon}{\beta}, \frac{\varepsilon}{4c}, \alpha\varepsilon, \frac{\varepsilon}{8c_2}, 1 \right\},$$

where the constants β , c and c_2 are the ones given in Lemma 3.1 and $\alpha > 0$ is defined in Algorithm 1.

By (9), for all $k \in \mathbb{N}'$ with $k \geq k_0$,

$$\Delta_k \leq \min \left\{ \frac{\pi_k}{\beta}, \frac{\pi_k}{4c}, \alpha\pi_k, \frac{\pi_k}{8c_2}, 1 \right\}, \quad (10)$$

consequently by Lemma 3.1, $k \in \mathcal{S}$.

Given $k \in \mathbb{N}'$ with $k \geq k_0$, consider ℓ_k the first index such that $\ell_k > k$ and $\pi_{\ell_k} \leq \varepsilon/2$. The existence of ℓ_k is ensured by Lemma 3.2. So, $\pi_k - \pi_{\ell_k} \geq \varepsilon/2$. Using the definition of π_k , the triangle inequality and the contraction property of projections, we have that

$$\begin{aligned} \frac{\varepsilon}{2} &\leq \|P_\Omega(x^k - g^k) - x^k\| - \|P_\Omega(x^{\ell_k} - g^{\ell_k}) - x^{\ell_k}\| \\ &\leq \|P_\Omega(x^k - g^k) - x^k - P_\Omega(x^{\ell_k} - g^{\ell_k}) + x^{\ell_k}\| \\ &\leq 2\|x^k - x^{\ell_k}\| + \|g^k - g^{\ell_k}\| \\ &= 2\|x^k - x^{\ell_k}\| + \|g^k - \nabla f(x^k) + \nabla f(x^k) - \nabla f(x^{\ell_k}) + \nabla f(x^{\ell_k}) - g^{\ell_k}\| \\ &\leq 2\|x^k - x^{\ell_k}\| + \|g^k - \nabla f(x^k)\| + \|\nabla f(x^k) - \nabla f(x^{\ell_k})\| + \|\nabla f(x^{\ell_k}) - g^{\ell_k}\|. \end{aligned}$$

Using Hypotheses H1 and H4, one gets that

$$\frac{\varepsilon}{2} \leq 2\|x^k - x^{\ell_k}\| + c_2\Delta_k + L\|x^k - x^{\ell_k}\| + c_2\Delta_{\ell_k} = (2 + L)\|x^k - x^{\ell_k}\| + c_2(\Delta_k + \Delta_{\ell_k}).$$

Let us consider $C_k = \{i \in \mathcal{S} \mid k \leq i < \ell_k\}$. Note that, by (10), $k \in \mathcal{S}$, so $C_k \neq \emptyset$. Thus

$$\frac{\varepsilon}{2} \leq (2 + L) \sum_{i \in C_k} \|x^i - x^{i+1}\| + c_2(\Delta_k + \Delta_{\ell_k}).$$

As $\ell_k > k$, it follows that $\Delta_{\ell_k} \leq \Delta_k$ and

$$\frac{\varepsilon}{2} \leq (2 + L) \sum_{i \in C_k} \Delta_i + 2c_2 \Delta_k.$$

Since $\Delta_k \leq \frac{\varepsilon}{8c_2}$, we have that

$$\sum_{i \in C_k} \Delta_i \geq \frac{\varepsilon}{8 + 4L} > 0. \quad (11)$$

Using the fact that $i \in \mathcal{S}$, condition (3) and Hypothesis H3, we conclude that

$$\begin{aligned} f(x^k) - f(x^{\ell_k}) &= \sum_{i \in C_k} (f(x^i) - f(x^{i+1})) \\ &\geq \sum_{i \in C_k} \eta (Q_i(0) - Q_i(d^i)) \\ &\geq \sum_{i \in C_k} \eta c_1 \pi_i \min \left\{ \frac{\pi_i}{\beta}, \Delta_i, 1 \right\}. \end{aligned}$$

By the definition of ℓ_k , we have that $\pi_i > \varepsilon/2$, for all $i \in C_k$, and therefore

$$f(x^k) - f(x^{\ell_k}) \geq \frac{\eta c_1 \varepsilon}{4} \sum_{i \in C_k} \min \left\{ \frac{\varepsilon}{\beta}, \Delta_i, 1 \right\} \geq \frac{\eta c_1 \varepsilon}{4} \min \left\{ \frac{\varepsilon}{\beta}, \sum_{i \in C_k} \Delta_i, 1 \right\}.$$

Therefore, by the inequality (11), for all $k \in \mathbf{N}'$ with $k \geq k_0$,

$$f(x^k) - f(x^{\ell_k}) \geq \frac{\eta c_1 \varepsilon}{4} \min \left\{ \frac{\varepsilon}{\beta}, \frac{\varepsilon}{8 + 4L}, 1 \right\} > 0. \quad (12)$$

On the other hand, by Hypothesis H2, the sequence $(f(x^k))$ is bounded below, and since it is nonincreasing, $f(x^k) - f(x^{\ell_k}) \rightarrow 0$, contradicting (12) and completing the proof. \square

Now we have all the ingredients for proving global convergence to first-order stationary points. In the following theorem we establish a relation between the measure of stationarity for the original problem and the measure of stationarity given in Lemma 3.4, which provides a strong convergence result.

Theorem 3.5. *Suppose that Hypotheses H1 to H4 hold, then*

$$\lim_{k \rightarrow \infty} \|P_{\Omega}(x^k - \nabla f(x^k)) - x^k\| = 0.$$

Proof. By the triangle inequality, the contraction property of projections and Hypothesis H4, we have that

$$\begin{aligned} \|P_{\Omega}(x^k - \nabla f(x^k)) - x^k\| &= \|P_{\Omega}(x^k - \nabla f(x^k)) - P_{\Omega}(x^k - g^k) + P_{\Omega}(x^k - g^k) - x^k\| \\ &\leq \|P_{\Omega}(x^k - \nabla f(x^k)) - P_{\Omega}(x^k - g^k)\| + \|P_{\Omega}(x^k - g^k) - x^k\| \\ &\leq \|\nabla f(x^k) - g^k\| + \|P_{\Omega}(x^k - g^k) - x^k\| \\ &\leq c_2 \Delta_k + \pi_k. \end{aligned}$$

Using Lemmas 3.3 and 3.4, we complete the proof. \square

4 Conclusions

In this work we proposed a general derivative-free trust-region algorithm for minimizing a smooth objective function in a closed convex set. The algorithm has a very simple structure and it allows a certain degree of freedom on the choice of the models, as long as they approximate sufficiently well the objective function, in the sense of Hypothesis H4. Furthermore, any internal algorithm can be utilized for solving the subproblems, provided that it generates a sequence of points satisfying the efficiency condition (3). Under these hypotheses and other standard assumptions, we have established global convergence of the algorithm in a neat way. Further research is necessary in order to extend the analysis to derivative-free problems in general domains.

Acknowledgement. The authors are grateful to José Mario Martínez who encouraged them to pursue this investigation and to Luis Mauricio Graña Drummond for his valuable comments and suggestions.

References

- [1] B. M. Arouxét, N. Echebest, and A. Pilotta. Active-set strategy in Powell’s method for optimization without derivatives. *Computational and Applied Mathematics*, 30(1):171–196, 2011.
- [2] L. F. Bueno, A. Friedlander, J. M. Martínez, and F. N. C. Sobral. Inexact restoration method for derivative-free optimization with smooth constraints. Available in Optimization Online, 2011.
- [3] P. G. Ciarlet and P. A. Raviart. General Lagrange and Hermite interpolation in \mathbb{R}^n with applications to finite elements methods. *Arch. Rational Mech. Anal*, 46:177–199, 1972.
- [4] A. R. Conn, N. I. M. Gould, A. Sartenaer, and Ph. L. Toint. Convergence properties of minimization algorithms for convex constraints using a structured trust region. *SIAM Journal on Optimization*, 6(4):1059–1086, 1996.
- [5] A. R. Conn, N. I. M. Gould, and Ph. L. Toint. Global convergence of a class of trust region algorithms for optimization with simple bounds. *SIAM Journal on Numerical Analysis*, 25(2):433–460, 1988.
- [6] A. R. Conn, N. I. M. Gould, and Ph. L. Toint. *Trust-Region Methods*. MPS-SIAM Series on Optimization. SIAM, Philadelphia, 2000.
- [7] A. R. Conn, K. Scheinberg, and Ph. L. Toint. On the Convergence of Derivative-Free Methods for Unconstrained Optimization. In M. D. Buhmann and A. Iserles, editors, *Approximation Theory and Optimization: Tributes to M. J. D. Powell*, pages 83–108. Cambridge University Press, 1997.
- [8] A. R. Conn, K. Scheinberg, and Ph. L. Toint. A Derivative Free Optimization Algorithm in Practice. In *Proceedings of the AIAA Conference*, St Louis, 1998.
- [9] A. R. Conn, K. Scheinberg, and L. N. Vicente. Geometry of interpolation sets in derivative free optimization. *Mathematical Programming*, 111:141–172, 2008.
- [10] A. R. Conn, K. Scheinberg, and L. N. Vicente. *Introduction to Derivative-Free Optimization*. MPS-SIAM Series on Optimization. SIAM, Philadelphia, 2009.
- [11] A. R. Conn and Ph. L. Toint. An Algorithm using Quadratic Interpolation for Unconstrained Derivative Free Optimization. In G. Di Pillo and F. Gianessi, editors, *Nonlinear Optimization and Applications*, pages 27–47. Plenum, 1996.
- [12] P. J. Davis. *Interpolation and Approximation*. Blaisdell, New York, 1963.
- [13] M. A. Diniz-Ehrhardt, J. M. Martínez, and L. G. Pedroso. Derivative-free methods for nonlinear programming with general lower-level constraints. *Computational and Applied Mathematics*, 30:19–52, 2011.
- [14] G. Fasano, J. L. Morales, and J. Nocedal. On the geometry phase in model-based algorithms for derivative-free optimization. *Optimization Methods and Software*, 24:145–154, 2009.
- [15] C. C. Gonzaga, E. W. Karas, and M. Vanti. A globally convergent filter method for nonlinear programming. *SIAM J. Optimization*, 14(3):646–669, 2003.
- [16] S. Gratton, Ph. L. Toint, and A. Tröltzsch. An active set trust-region method for derivative-free nonlinear bound-constrained optimization. *Optimization Methods and Software*, 26:873–894, 2011.
- [17] T. G. Kolda, R. M. Lewis, and V. Torczon. A Generating Set Direct Search Augmented Lagrangian Algorithm for Optimization with a Combination of General and Linear Constraints. Technical Report SAND2006-5315, Sandia National Laboratories, August 2006.
- [18] T. G. Kolda, R. M. Lewis, and V. Torczon. Stationarity results for generating set search for linearly constrained optimization. *SIAM Journal on Optimization*, 17:943–968, 2006.

- [19] R. M. Lewis and V. Torczon. Pattern search algorithms for linearly constrained minimization. *SIAM Journal on Optimization*, 10:917–941, 2000.
- [20] J. Nocedal and S. J. Wright. *Numerical Optimization*. Springer Series in Operations Research. Springer-Verlag, 1999.
- [21] M. J. D. Powell. A Direct Search Optimization Method that Models the Objective and Constraint Functions by Linear Interpolation. In S. Gomez and J. P. Hennart, editors, *Advances in Optimization and Numerical Analysis*, pages 51–67. Kluwer Academic, Dordrecht, 1994.
- [22] M. J. D. Powell. The NEWUOA software for unconstrained optimization without derivatives. In G. Di Pillo and M. Roma, editors, *Large-Scale Nonlinear Optimization*, pages 255–297. Springer, New York, 2006.
- [23] M. J. D. Powell. Developments of NEWUOA for minimization without derivatives. *IMA Journal Numerical Analysis*, 28:649–664, 2008.
- [24] M. J. D. Powell. The BOBYQA Algorithm for Bound Constrained Optimization without Derivatives. Technical Report DAMTP 2009/NA06, Department of Applied Mathematics and Theoretical Physics, Cambridge, England, August 2009.
- [25] M. J. D. Powell. On Derivative-Free Optimization with Linear Constraints. In *21st ISMP*, Berlin, Germany, 2012.
- [26] M. J. D. Powell. On the convergence of trust region algorithms for unconstrained minimization without derivatives. *Computational Optimization and Applications*, pages 1–29, Online First, 2012.
- [27] A. A. Ribeiro and E. W. Karas. *Um Curso de Otimização*. Cengage Learning, São Paulo, Brazil. In Portuguese. To appear.
- [28] K. Scheinberg and Ph. L. Toint. Self-correcting geometry in model-based algorithms for derivative-free unconstrained optimization. *SIAM Journal on Optimization*, 20(6):3512–3532, 2010.
- [29] A. Tröltzsch. *An Active-set Trust-Region Method for Bound-Constrained Nonlinear Optimization without Derivatives Applied to Noisy Aerodynamic Design Problems*. PhD thesis, Université de Toulouse, 2011.
- [30] D. Winfield. Function minimization by interpolation in a data table. *IMA Journal of Applied Mathematics*, 12(3):339–347, 1973.