

The Generalized Trust Region Subproblem

Ting Kei Pong ^{*} Henry Wolkowicz [†]

November 17, 2012

Abstract

The *interval bounded generalized trust region subproblem* (GTRS) consists in minimizing a general quadratic objective, $q_0(x) \rightarrow \min$, subject to an upper and lower bounded general quadratic constraint, $\ell \leq q_1(x) \leq u$. This means that there are no definiteness assumptions on either quadratic function. We first study characterizations of optimality for this *implicitly* convex problem under a constraint qualification and show that it can be assumed without loss of generality. We next classify the GTRS into easy case and hard case instances, and demonstrate that the upper and lower bounded general problem can be reduced to an equivalent equality constrained problem after identifying suitable generalized eigenvalues and possibly solving a sparse system. We then discuss how the Rendl-Wolkowicz algorithm proposed in [11, 29] can be extended to solve the resulting equality constrained problem, highlighting the connection between the GTRS and the problem of finding minimum generalized eigenvalues of a parameterized matrix pencil. Finally, we present numerical results to illustrate this algorithm at the end of the paper.

Contents

1	Introduction	2
1.1	Outline	4
1.2	Notation	4
2	Duality and optimality conditions	4
2.1	Assumptions and properties	5
2.2	Weak duality	7
2.3	Strong duality and characterization of optimality	7
3	Easy and hard cases and regular pencils	11
3.1	Regular case	11
3.1.1	Three intervals for λ	16
4	A method for solving GTRS: the extended Rendl-Wolkowicz algorithm	17
4.1	Properties of $k_0(t)$	19
4.2	Recovering solution of GTRS from maximizing $k(t)$	25
4.3	Implementation details	28

^{*}Department of Combinatorics and Optimization, University of Waterloo, Ontario N2L 3G1, Canada. Research supported by AFOSR. Email: ptingkei@uwaterloo.ca

[†]Research supported by The Natural Sciences and Engineering Research Council of Canada and by AFOSR. Email: hwolkowicz@uwaterloo.ca

5 Numerical experiments with new RW algorithm	33
5.1 B is positive definite	34
5.2 B is indefinite	35
6 Conclusion	37
A Proof of Theorem 2.1(ii)	37
Index	42

List of Tables

1	The three different cases for GTRS.	15
2	Computational results for indefinite A and positive definite B	35
3	Computational results for positive definite A and positive definite B	36
4	Computational results for positive definite A and indefinite B	37

List of Figures

1	$\psi(\lambda)$ when B is indefinite. The dotted lines are $\lambda = \underline{\lambda}$ and $\lambda = \bar{\lambda}$	13
2	$\psi(\lambda)$ when B is positive definite. The dotted line is $\lambda = \bar{\lambda}$	14
3	$d(\lambda)$ when B is indefinite. The dotted lines are $\lambda = \underline{\lambda}$ and $\lambda = \bar{\lambda}$	22
4	$d(\lambda)$ when B is positive definite with $s_0 < -1$. The dotted line is $\lambda = \bar{\lambda}$	24
5	$d(\lambda)$ when B is positive definite with $s_0 > -1$. The dotted line is $\lambda = \bar{\lambda}$	24

1 Introduction

We consider the following quadratic minimization problem, which we call the *interval bounded generalized trust region subproblem* (GTRS):

$$\begin{aligned}
 \text{(GTRS)} \quad p^* &:= \inf \quad q_0(x) := x^T A x - 2a^T x \\
 &\text{s.t.} \quad \ell \leq \underbrace{x^T B x - 2b^T x}_{q_1(x)} \leq u.
 \end{aligned} \tag{1.1}$$

Here $A, B \in \mathcal{S}^n$, the space of real $n \times n$ symmetric matrices, $a, b \in \mathbb{R}^n$, and $-\infty < \ell \leq u < \infty$. To avoid trivialities, we assume that $B \neq 0$ and the GTRS is feasible.¹ We emphasize that the quadratic constraint is *two-sided*, and both the objective and constraint functions are possibly nonconvex. Thus we are essentially considering a general nonlinear program with a quadratic objective and *two* quadratic constraints.

Problem (1.1) is most studied in the *one-sided ball-constrained case*, i.e., the special case when $B = I$, $b = 0$, and $\ell = 0 < u$, known as the *trust region subproblem* (TRS):

$$\begin{aligned}
 \text{(TRS)} \quad &\inf \quad x^T A x - 2a^T x \\
 &\text{s.t.} \quad x^T x \leq u.
 \end{aligned} \tag{1.2}$$

These problems arise in regularization or trust region methods for unconstrained and constrained nonlinear programming; see, e.g., [7] for a comprehensive discussion. The TRS provides a quadratic model in a trust

¹Checking feasibility for GTRS is simple, see Theorem 2.1(i), below.

region around the current point. The use of scaled trust regions with a general $B \succ 0$ in the constraint (see, e.g. [15]) motivates our consideration of GTRS. In our general consideration of (1.1), our scaled trust region is defined with a possibly indefinite B ; hence, $x^T B x$ could be a quadratic form induced from the indefinite inner product $\langle x, y \rangle := x^T B y$; see, e.g. [13]. This together with the two-sided constraints model annular/hyperbolic type regions that allow minimum as well as maximum “steplengths”. In addition, the possibly nonzero b accounts for a shift of the “center” of the trust region, which allows trust regions to be built around previous iterates.

Problem (1.2) is explicitly nonconvex, since A is not necessarily positive semidefinite. Nevertheless, it is implicitly convex in that necessary and sufficient optimality conditions have been derived, see Gay [12] and Moré and Sorensen [25]. The conditions are rephrased in a modern primal-dual paradigm in, for example, [7, 10, 11, 15, 29, 34]. The optimality conditions of the TRS can also be derived using the S -lemma; see, e.g. [28]. The S -lemma, developed by Yakubovich [36] in 1971, states whether a quadratic (in)equality is a consequence of other quadratic (in)equalities. The earliest results of this kind date back to Finsler [9] and Hestenes and McShane [19] and we refer the readers to [28] for a detailed account of its history and its applications. Turning to the more general problem (1.1), necessary and sufficient optimality conditions have been studied in [24, 34], under certain constraint qualifications in the special cases when $b = 0$ or $u = \ell$. The general case (1.1) was considered in [37, Section 2.1] under a dual strict feasibility assumption; see also [3, 8, 20]. Further references are available in the online bibliography [17].

The necessary and sufficient optimality conditions for (1.2) have been the basis for developing efficient algorithms for solving TRS. The classical algorithm [25] by Moré and Sorensen (MS algorithm) applies Newton’s method with backtracking to the so-called secular function, and takes a primal step to the boundary for near hard case instances. This algorithm can be expensive since each iteration requires a full Cholesky factorization. The MS algorithm was later incorporated into the *generalized Lanczos trust-region* (GLTR) method [15, 16]. The GLTR algorithm first uses the Lanczos procedure to obtain a sequence of TRS on low dimensional subspaces with tridiagonal matrices as objectives. The resulting sequence of TRS can be solved efficiently by the MS algorithm. Another line of algorithms involves reformulating the TRS into a parameterized eigenvalue problem. This includes, e.g., the Rendl-Wolkowicz (RW) algorithm [11, 29], the *large-scale trust-region subproblem* (LSTRS) by Rojas et al. [31, 32] and its variant [21]. These algorithms are factorization free, and can take advantage of well developed eigensolvers for large, sparse matrices. Other algorithms for TRS can be found in, e.g. [18, 33, 35]. Despite the many algorithms developed for TRS, there are currently no algorithms specifically designed for the general problem (1.1); though the special case when $B \succ 0$, $b = 0$ and $\ell = 0$ can be solved by the GLTR method.

In this paper we present characterizations of optimality and propose an algorithm for solving large-scale instances of GTRS (1.1). Specifically, we obtain optimality conditions under a constraint qualification and show that it can be assumed without loss of generality. Our results show that even though the GTRS is a problem consisting of general, possibly *nonconvex*, quadratic functions, it has both necessary and sufficient optimality conditions as in *convex programming*, and it has strong duality results as in *linear programming*, in the sense that when the constraint qualification fails, the problem can be explicitly solved. Thus, the GTRS sits on the boundary between linear and nonlinear programming, and on the boundary between convex, and general nonconvex programming. We also discuss in detail the so-called *easy and hard cases* corresponding to (non)singularity of the Hessian of the Lagrangian, which was previously only studied for some special cases of (1.1) in [34]. Moreover, as in [11], we include a *shift and deflate* operation that finds an explicit solution in the hard case (case 2). We then demonstrate that the GTRS can be reduced to an equality constrained problem (i.e., an instance of GTRS with $u = \ell$) after finding some suitable generalized eigenvalues and possibly solving a sparse system. To solve this equality constrained problem, we generalize the ideas in [11, 29] and transform the problem into a parameterized *generalized eigenvalue problem*. The latter problem can then take advantage of specialized solvers for finding generalized eigenvalues/eigenvectors,

e.g., the `eigifp` developed in [14]. We compare this approach with the GLTR algorithm (when $B \succ 0$) and a simple implementation of Newton’s method with the Armijo line search rule as applied to solve the dual problem of (1.1). Our computational results on large-scale instances show that our approach is competitive with the Newton’s method and usually outperforms the GLTR algorithm for *random sparse* positive definite B in both runtime and solution accuracy. In the case when B is indefinite, our approach requires additional inputs for initialization; see Section 4.3 for details. Granting such inputs, our computational results show that our approach is competitive with the Newton’s method.

1.1 Outline

We complete this section with some preliminary notations in Section 1.2. We then present the characterizations of optimality in Section 2. This includes a constraint qualification that can be assumed without loss of generality in the sense that if it fails, then an explicit solution of the GTRS (1.1) can be obtained. In Section 3, we discuss the so-called easy and hard cases, the shift and deflation procedure to obtain an explicit solution in the hard case, and the reduction to an equality constrained problem. The algorithm for this equality constrained problem and its implementation details are discussed in Section 4. We present numerical tests in Section 5 and finally some concluding remarks in Section 6.

1.2 Notation

In this paper, the symbol \mathbb{R}^n denotes the n -dimensional vector space. For $v \in \mathbb{R}^n$, $\|v\|$ denotes the Euclidean norm of v . The space of $n \times n$ symmetric matrices equipped with the trace inner-product is denoted by \mathcal{S}^n . For $C \in \mathcal{S}^n$, $\lambda_{\max}(C)$ and $\lambda_{\min}(C)$ denote the largest and smallest eigenvalue of C , respectively. For $C, D \in \mathcal{S}^n$, $C \succeq D$, $C \succ D$ denote $C - D$ is positive semidefinite, and positive definite, respectively. For a (not necessarily symmetric) square matrix M , $\text{tr}(M)$ denotes the trace of M , M^\dagger denotes the Moore-Penrose generalized inverse of M , and $\text{Null}(M)$ and $\text{Range}(M)$ denote its null and range spaces, respectively. The identity matrix is denoted by I , whose dimension should be clear from the context.

For an interval J on the real line, the symbol $\text{cl}(J)$ denotes its closure, and $\text{ri}(J)$ denotes its relative interior. Finally, for a subspace V of \mathbb{R}^n , the orthogonal complement of V is denoted by V^\perp , and if a vector $v \in \mathbb{R}^n$ is orthogonal to V , we write $v \perp V$.

2 Duality and optimality conditions

In this section, we present the characterizations of optimality and duality properties for GTRS (1.1). In particular, we include assumptions that can be made without loss of generality. These assumptions include a constraint qualification, whose failure means that an explicit solution for GTRS can be obtained.

We start by writing down the Lagrangian dual of (1.1). The Lagrangian for (1.1) can be written with one *free* Lagrange multiplier λ , or with two nonnegative Lagrange multipliers²

$$\begin{aligned} \lambda_+ &:= \max\{\lambda, 0\}, \lambda_- := -\min\{\lambda, 0\}, \lambda = (\lambda_+ - \lambda_-) : & (2.1) \\ L(x, \lambda_+, \lambda_-) &= x^T A x - 2a^T x + \lambda_+(\ell - (x^T B x - 2b^T x)) + \lambda_-(x^T B x - 2b^T x - u). \\ &= x^T (A - (\lambda_+ - \lambda_-)B)x - 2(a - (\lambda_+ - \lambda_-)b)^T x + \lambda_+ \ell - \lambda_- u. \\ &= x^T (A - \lambda B)x - 2(a - \lambda b)^T x + \lambda_+ \ell - \lambda_- u. \end{aligned}$$

²Note that complementary slackness can be written with a single Lagrange multiplier $\lambda \in \mathbb{R}$ as follows: $(q_1(x) - \ell)\lambda \leq 0 \leq (u - q_1(x))\lambda$.

The Lagrangian dual problem of (1.1) can then be reduced to the following problem:

$$\begin{aligned} \text{(D-GTRS)} \quad d^* &:= \sup & h(\lambda) + \ell\lambda_+ - u\lambda_- \\ &\text{s.t.} & A - \lambda B \succeq 0, \end{aligned} \tag{2.2}$$

where the dual functional

$$\begin{aligned} h(\lambda) &= \inf_x x^T(A - \lambda B)x - 2(a - \lambda b)^T x \\ &= \begin{cases} -(a - \lambda b)^T(A - \lambda B)^\dagger(a - \lambda b) & \text{if } a - \lambda b \in \text{Range}(A - \lambda B), A - \lambda B \succeq 0, \\ -\infty & \text{otherwise.} \end{cases} \end{aligned}$$

Note that the objective function in D-GTRS is concave. In this paper, we will also look at the following closely related problem formed by enforcing positive definiteness in (2.2) and thus reducing the size of the feasible set:

$$\begin{aligned} \text{(D}_{\succ}\text{-GTRS)} \quad d_{\succ}^* &:= \sup & h_{\succ}(\lambda) + \ell\lambda_+ - u\lambda_- \\ &\text{s.t.} & A - \lambda B \succ 0, \end{aligned} \tag{2.3}$$

where

$$h_{\succ}(\lambda) := -(a - \lambda b)^T(A - \lambda B)^{-1}(a - \lambda b).$$

It is useful to study this problem since positive definiteness as in (2.3) is maintained in the algorithm presented below. Finally, we write down the semidefinite programming (SDP) relaxation of GTRS

$$\begin{aligned} \text{(SDP-GTRS)} \quad p_{SDP}^* &:= \inf & \text{tr}(AX) - 2a^T x \\ &\text{s.t.} & \ell \leq \text{tr}(BX) - 2b^T x \leq u, \\ & & X \succeq xx^T. \end{aligned} \tag{2.4}$$

By a Schur complement argument, the final inequality is equivalent to the linear constraint $\begin{bmatrix} 1 & x^T \\ x & X \end{bmatrix} \succeq 0$.

Proposition 2.1. *The dual of SDP-GTRS is*

$$\begin{aligned} \text{(DSDP-GTRS)} \quad d_{DSDP}^* &:= \sup & \ell\lambda_+ - u\lambda_- - \gamma \\ &\text{s.t.} & \begin{bmatrix} \gamma & -(a - \lambda b)^T \\ -(a - \lambda b) & A - \lambda B \end{bmatrix} \succeq 0, \end{aligned} \tag{2.5}$$

where the multipliers are defined in (2.1). Moreover, (2.5) is equivalent to (2.2) with $d_{DSDP}^* = d^*$.

Proof. It is routine to show that the dual of (2.4) is given by (2.5). Furthermore, by considering the Schur complement, for any $\gamma, \lambda \in \mathbb{R}$, we have

$$\begin{bmatrix} \gamma & -(a - \lambda b)^T \\ -(a - \lambda b) & A - \lambda B \end{bmatrix} \succeq 0 \Leftrightarrow \begin{cases} \gamma \geq (a - \lambda b)^T(A - \lambda B)^\dagger(a - \lambda b) \\ a - \lambda b \in \text{Range}(A - \lambda B) \\ A - \lambda B \succeq 0. \end{cases}$$

Thus, (2.5) is equivalent to (2.2). □

2.1 Assumptions and properties

We consider the following assumptions on the GTRS (1.1).

Assumption 2.1.

1. $B \neq 0$
2. GTRS (1.1) is feasible.
3. the following relative interior constraint qualification holds

$$(RICQ) \quad \text{tr}(B\hat{X}) - 2b^T\hat{x} \in \text{ri}([\ell, u]), \quad \text{for some } \hat{X} \succ \hat{x}\hat{x}^T. \quad (2.6)$$

4. GTRS is bounded below.
5. D-GTRS (2.2) is feasible.

Note that Assumption 2.1 together with weak duality yields

$$-\infty < d^* \leq p^* < +\infty. \quad (2.7)$$

We show in the next theorem that Assumption 2.1 is reasonable in the sense that they can be made without loss of generality, i.e., if an assumption fails then an explicit solution or a simplification can be easily obtained.

Theorem 2.1. *The following holds for the Items in Assumption 2.1.*

- (i) *The Items 1, 2, 3 in Assumption 2.1 can be made in the order given, without loss of generality, i.e., if an assumption fails then an explicit solution can easily be obtained.*
- (ii) *If Items 1, 2, 3 in Assumption 2.1 hold and $b = 0$, then Item 4 implies Item 5.*
- (iii) *If Item 4 in Assumption 2.1 fails, then Item 5 fails.*

Proof. Let $x^* := B^\dagger b$, $q_1^* := q_1(x^*)$. We now provide the details about how the assumptions hold in the order given.

- (i)
 - Suppose that $B = 0$, i.e., the constraint is linear. If $A \succeq 0$, $a \in \text{Range}(A)$ and the unconstrained minimum $\bar{x} = A^\dagger a$ satisfies $\ell \leq -2b^T(A^\dagger a) \leq u$, then \bar{x} solves GTRS. Otherwise, the optimum, if it exists, is on one of the two boundaries. Therefore, we can change the linear inequality constraint to an equality, and we can again check for an unconstrained minimum after the appropriate substitution using the linear constraint. Therefore, the assumption that $B \neq 0$ can be made without loss of generality.
 - First, GTRS is infeasible if, and only if, the following three conditions hold:

$$\begin{aligned} & B \text{ is semidefinite;} \\ & b \in \text{Range}(B) \quad (\text{equivalently } b = Bx^*); \\ & q_1^* < \ell \text{ if } B \preceq 0; \quad \text{or} \quad q_1^* > u \text{ if } B \succeq 0. \end{aligned}$$

More precisely, the characterization for infeasibility follows from the fact that q_1 is bounded below with minimum at x^* (resp. bounded above with maximum at x^*) if, and only if, $b \in \text{Range}(B)$ and $B \succeq 0$ (resp. $B \preceq 0$). We can verify the semidefiniteness of B by finding the largest and smallest eigenvalues. The range condition follows from finding the best least squares solution of $Bx = b$. If the range condition holds, then the final inequalities can be checked by evaluating q_1 at x^* . Thus, we conclude that feasibility can be verified by finding $\lambda_{\max}(B)$ and $\lambda_{\min}(B)$ and solving a system of equations, and hence, Assumption 2.1, Item 2, can be made without loss of generality.

- Suppose that Assumption 2.1, Items 1-2 hold, but the RICQ Assumption 2.1, Item 3 fails. Then we can find an explicit solution for GTRS. More precisely, since the RICQ fails, we have $b \in \text{Range}(B)$, and either $B \preceq 0$ with $\ell = \sup x^T Bx - 2b^T x$ or $B \succeq 0$ with $u = \inf x^T Bx - 2b^T x$. In either case, the conditions imply that the feasible set is $x^* + \text{Null}(B)$. We can then use a nullspace representation and substitute into the objective function to obtain an explicit solution or realize that the problem is unbounded below. In conclusion, if RICQ fails, we can obtain an explicit solution or realize that the problem is unbounded.

(ii) This conclusion is proved as Proposition A.1 in the appendix.

(iii) If Assumption 2.1, Item 5 holds, then $d^* > -\infty$. We thus see from (2.7) that $p^* > -\infty$, i.e., Item 4 holds. □

Remark 2.1. *From the proof of Theorem 2.1(i) above, it is not hard to see that if we assume Items 1 and 2 of Assumption 2.1, then the RICQ (2.6) may fail only when the constraint is convex and no Slater point (strict feasibility) exists. Comparing this with [24, Lemma 3.1], we see that the RICQ is equivalent to condition (3.2) in that paper, proposed for GTRS (1.1) with $u = \ell$.*

For the rest of the paper, we assume that Assumption 2.1 holds.

2.2 Weak duality

We have the following weak duality result describing the relationship between the optimal values of the above optimization problems.

Proposition 2.2. *The optimal values satisfy*

$$-\infty \leq d_{\gamma}^* \leq d^* = d_{SDP}^* \leq p_{SDP}^* \leq p^* < +\infty. \quad (2.8)$$

Moreover, if $d_{\gamma}^* > -\infty$, then $d_{\gamma}^* = d^*$.

Proof. The proof of (2.8) follows from the feasibility Assumption 2.1, weak duality, Proposition 2.1 and the definitions.

Now suppose that d_{γ}^* is finite and suppose to the contrary that $d_{\gamma}^* < d^*$. Then there exists λ_1 and λ_2 feasible for (2.2) and (2.3), respectively, such that $f(\lambda_1) > d^* - \frac{d^* - d_{\gamma}^*}{2}$ and $f(\lambda_2) > d_{\gamma}^* - \frac{d^* - d_{\gamma}^*}{2}$, where $f(\lambda)$ is the common objective function for both problems. Since $f(\lambda)$ is clearly a concave function and $\frac{\lambda_1 + \lambda_2}{2}$ is feasible for (2.3), we obtain that

$$d_{\gamma}^* \geq f\left(\frac{\lambda_1 + \lambda_2}{2}\right) \geq \frac{f(\lambda_1) + f(\lambda_2)}{2} > \frac{d^* + d_{\gamma}^*}{2} - \frac{d^* - d_{\gamma}^*}{2} = d_{\gamma}^*,$$

a contradiction. This completes the proof. □

2.3 Strong duality and characterization of optimality

We show below that equality holds for four of the five finite optimal values in (2.8); and, moreover, two of them are attained. We start with the following technical lemma where the quadratic forms in (1.1) are linearized. Recall that a Schur complement argument implies that the quadratic constraint $X \succeq xx^T$ in (2.9) is equivalent to the linear constraint $\begin{bmatrix} 1 & x^T \\ x & X \end{bmatrix} \succeq 0$.

Lemma 2.1. *Let $C \succ A$ and let*

$$\begin{aligned} p_{C,SDP}^* &:= \inf \quad \text{tr}(CX) - 2a^T x \\ \text{s.t.} \quad &\ell \leq \text{tr}(BX) - 2b^T x \leq u, \\ &X \succeq xx^T. \end{aligned} \tag{2.9}$$

Then, the optimal value $p_{C,SDP}^$ is finite and attained.*

Proof. The programs (D-GTRS) and (SDP-GTRS) are dual to each other. Assumption 2.1 implies that (2.9) is feasible and that there exists $\hat{\lambda}$ satisfying the Slater condition $C - \hat{\lambda}B \succ A - \hat{\lambda}B \succeq 0$, i.e., a Slater point exists for (D-GTRS) when A is replaced by C . This means that the dual program (2.9) is feasible and attained. \square

We next prove strong duality between (1.1) and (2.2). This technical result is used repeatedly throughout this paper. Similarly as in [6], we make use of bounds on the rank of the extreme points of SDP representable sets [1, 2, 27] to prove the exactness of the SDP relaxations.

Theorem 2.2. *Recall that Assumption 2.1 holds. Then the following holds for GTRS:*

- (i) *The optimal values of GTRS and its SDP relaxation are equal,*

$$p_{SDP}^* = p^*.$$

- (ii) *Strong duality holds for GTRS, i.e., $p^* = d^*$ and the dual optimal value d^* is attained. Moreover, equality holds for four of the five optimal values in (2.8),*

$$d_{\succ}^* \leq d^* = d_{DSDP}^* = p_{SDP}^* = p^*. \tag{2.10}$$

Moreover, if $d_{\succ}^ > -\infty$, then all quantities in (2.10) are equal.*

Proof. (i) For each $\epsilon \geq 0$, let $A_\epsilon := A + \epsilon I$. Consider the following perturbation of (1.1)

$$p_\epsilon^* = \inf \{x^T A_\epsilon x - 2a^T x : \ell \leq x^T B x - 2b^T x \leq u\},$$

and its SDP relaxation

$$\begin{aligned} v_\epsilon^* &= \inf \{ \text{tr}(A_\epsilon X) - 2a^T x : \ell \leq \text{tr}(BX) - 2b^T x \leq u, X \succeq xx^T \} \\ &= \inf \left\{ \text{tr}(A_\epsilon X) - 2a^T x : \ell \leq \text{tr}(BX) - 2b^T x \leq u, U = \begin{bmatrix} 1 & x^T \\ x & X \end{bmatrix} \succeq 0 \right\}. \end{aligned}$$

Then $p_\epsilon^* \geq v_\epsilon^*$, for all $\epsilon \geq 0$.

We proceed by first showing that $p_\epsilon^* = v_\epsilon^*$ for each fixed $\epsilon > 0$. We start with the case when $\ell = u$. In this case, the SDP relaxation can be written as

$$\begin{aligned} \inf \quad &\text{tr} \left(\begin{bmatrix} 0 & -a^T \\ -a & A_\epsilon \end{bmatrix} U \right) \\ \text{s.t.} \quad &\text{tr} \left(\begin{bmatrix} 0 & -b^T \\ -b & B \end{bmatrix} U \right) = u, \\ &U_{1,1} = 1, U \succeq 0. \end{aligned}$$

The optimal set of the above SDP is nonempty by Lemma 2.1. Since the cone of positive semidefinite matrices does not contain lines, the optimal value must be attained at an extreme point U^* of the

feasible set. Since there are two equality constraints, by [27, Theorem 2.2], the rank r_U of the extreme point satisfies

$$r_U(r_U + 1) \leq 4.$$

Since $U^* \neq 0$, we must have $r_U = 1$ and hence $U^* = \begin{bmatrix} 1 \\ x^* \end{bmatrix} \begin{bmatrix} 1 & x^{*T} \end{bmatrix}$, for some x^* . Then x^* is feasible for (1.1) and we conclude that $p_\epsilon^* = v_\epsilon^*$.

Next, we consider the case when $\ell < u$. In this case, the SDP relaxation can be written as

$$\begin{aligned} \inf \quad & \text{tr} \left(\begin{bmatrix} 0 & -a^T \\ -a & A_\epsilon \end{bmatrix} U \right) \\ \text{s.t.} \quad & \text{tr} \left(\begin{bmatrix} 0 & -b^T \\ -b & B \end{bmatrix} U \right) - \alpha = l, \\ & \text{tr} \left(\begin{bmatrix} 0 & -b^T \\ -b & B \end{bmatrix} U \right) + \beta = u, \\ & U_{1,1} = 1, U \succeq 0, \alpha \geq 0, \beta \geq 0. \end{aligned}$$

The optimal set of the above SDP is nonempty, again by Lemma 2.1. Hence, the optimal value must be attained at an extreme point (U^*, α^*, β^*) of the feasible set. Since there are three equality constraints, by [27, Theorem 2.2], the ranks r_U, r_α and r_β of this extreme point have to satisfy

$$r_U(r_U + 1) + r_\alpha(r_\alpha + 1) + r_\beta(r_\beta + 1) \leq 6. \quad (2.11)$$

Notice that at optimality, α^* and β^* cannot both be zero. This fact together with (2.11) and the fact $U^* \neq 0$ shows that $r_U = 1$ and hence $U^* = \begin{bmatrix} 1 \\ x^* \end{bmatrix} \begin{bmatrix} 1 & x^{*T} \end{bmatrix}$, for some x^* . Then x^* is feasible for (1.1) and we again conclude that $p_\epsilon^* = v_\epsilon^*$.

Hence, we have shown that $p_\epsilon^* = v_\epsilon^*$, for all $\epsilon > 0$. Now, let (x, X) be feasible for the SDP relaxation (2.4). Then we have

$$v_0^* \leq p_0^* \leq \lim_{\epsilon \downarrow 0} p_\epsilon^* = \lim_{\epsilon \downarrow 0} v_\epsilon^* \leq \lim_{\epsilon \downarrow 0} \text{tr}(A_\epsilon X) - 2a^T x = \text{tr}(AX) - 2a^T x.$$

Taking the infimum over the feasible set of the SDP relaxation gives the desired equality $v_0^* = p_0^*$. This completes the proof.

- (ii) Recall that the Lagrangian dual of the SDP relaxation (2.4) is given by (2.5). Moreover, from RICQ (2.6), the generalized Slater condition for (2.4) holds with the Slater point $\begin{bmatrix} 1 & \hat{x}^T \\ \hat{x} & \hat{X} \end{bmatrix}$. Hence $p_0^* = v_0^* = d_{DSDP}^* = d^*$ and the dual optimal values are attained. This proves (2.10). The rest of the claim follows from Proposition 2.2. □

We remark that, when $\ell < u$ in the above proof, the equality $p_\epsilon^* = v_\epsilon^*$, for all $\epsilon > 0$, can also be obtained as a consequence of [37, Theorem 2.3]. We are now ready to characterize optimality for GTRS. We note that our constraint qualification (2.6) is different from that of [34, Theorem 2.1]. In particular, we do not require $\ell < u$ when B is indefinite with the optimal solution in its kernel and $b = 0$. Moreover, as seen in Theorem 2.1, Assumption 2.1 can be made without loss of generality.

Theorem 2.3. *Recall that Assumption 2.1 holds. A point x^* is a solution to GTRS (1.1) if, and only if, for some (Lagrange multiplier) $\lambda^* \in \mathbb{R}$, we have*

$$\left. \begin{aligned} (A - \lambda^* B)x^* &= a - \lambda^* b, \\ A - \lambda^* B &\succeq 0, \end{aligned} \right\} \quad \text{dual feasibility}$$

$$\ell \leq x^{*T} Bx^* - 2b^T x^* \leq u, \quad \text{primal feasibility} \quad (2.12)$$

$$\left. \begin{aligned} (\lambda^*)_+(\ell - x^{*T} Bx^* + 2b^T x^*) &= 0, \\ (\lambda^*)_-(x^{*T} Bx^* - 2b^T x^* - u) &= 0. \end{aligned} \right\} \quad \text{complementary slackness}$$

Proof. Suppose first that x^* is a solution to (1.1). Then the third relation in (2.12) holds. Furthermore, by Theorem 2.2, strong duality holds. Hence, there exists λ^* such that

$$p^* = x^{*T} Ax^* - 2a^T x^* = \inf_{\ell \leq x^T Bx - 2b^T x \leq u} x^T Ax - 2a^T x \quad (2.13)$$

$$\begin{aligned} &= \sup_{\lambda} \inf_x x^T Ax - 2a^T x + \lambda_+(\ell - x^T Bx + 2b^T x) + \lambda_-(x^T Bx - 2b^T x - u) \\ &= \inf_x x^T Ax - 2a^T x + \lambda_+(\ell - x^T Bx + 2b^T x) + \lambda_-(x^T Bx - 2b^T x - u). \end{aligned} \quad (2.14)$$

The first two relations in (2.12) follow immediately from the optimality condition of the unconstrained optimization problem (2.14). Moreover, from (2.14) and primal feasibility of x^* , we have

$$\begin{aligned} p^* &= \inf_x x^T Ax - 2a^T x + \lambda_+(\ell - x^T Bx + 2b^T x) + \lambda_-(x^T Bx - 2b^T x - u) \\ &\leq x^{*T} Ax^* - 2a^T x^* + \lambda_+(\ell - x^{*T} Bx^* + 2b^T x^*) + \lambda_-(x^{*T} Bx^* - 2b^T x^* - u) \leq x^{*T} Ax^* - 2a^T x^*. \end{aligned}$$

Comparing this last relation with (2.13), we obtain the complementary slackness expressions in the last two relations of (2.12).

Next, assume that x^* is primal feasible and that there exists λ^* so that (2.12) holds. Then we have the following chain of inequalities.

$$\begin{aligned} p^* &\geq d^* = \sup_{\lambda} \inf_x x^T Ax - 2a^T x + \lambda_+(\ell - x^T Bx + 2b^T x) + \lambda_-(x^T Bx - 2b^T x - u) \\ &\geq \inf_x x^T Ax - 2a^T x + \lambda_+(\ell - x^T Bx + 2b^T x) + \lambda_-(x^T Bx - 2b^T x - u) \\ &= x^{*T} Ax^* - 2a^T x^* + \lambda_+(\ell - x^{*T} Bx^* + 2b^T x^*) + \lambda_-(x^{*T} Bx^* - 2b^T x^* - u) \\ &= x^{*T} Ax^* - 2a^T x^* \geq p^*, \end{aligned}$$

where the first equality follows from the definition of the Lagrangian dual problem, the second equality follows from the first two relations of (2.12), the third equality follows from the last two relations of (2.12), while the last inequality follows from the primal feasibility of x^* . Thus, in particular, we have $x^{*T} Ax^* - 2a^T x^* = p^*$, and so x^* solves (1.1). \square

Remark 2.2. *We note that in proving the optimality conditions (i.e., version of Theorem 2.3) in [24, 34] for the special cases of GTRS (1.1), the authors did not assume Items 4 and 5 of Assumption 2.1. Indeed, suppose we only assume Items 1 through 3 of Assumption 2.1. When $b = 0$, if x^* is known to be a solution of the GTRS (1.1), then the problem is bounded below and thus by Theorem 2.1(ii), we conclude that Assumption 2.1 holds. On the other hand, if there exists λ^* and x^* such that (2.12) holds, then we conclude from Theorem 2.1(iii) that Assumption 2.1 holds. Thus, our result is a strict improvement of [34, Theorem 2.1], which assumes in addition $b = 0$, since our RICQ (2.6) is weaker than condition (2.5) in [34, Theorem 2.1]. However, since it is not known to us whether Item 4 implies Item 5 when $b \neq 0$, we cannot assert that our Theorem 2.3 reduces to [24, Theorem 3.2], even though our RICQ is equivalent to their constraint qualifications.*

Example 2.1. We illustrate that (2.10) can fail if Assumption 2.1 is violated. Moreover, attainment for GTRS can fail. Consider

$$\begin{aligned} \inf \quad & x_1 x_2 \\ \text{s.t.} \quad & x_1^2 = 0. \end{aligned}$$

Then clearly, the optimal value of the above optimization problem is zero. Moreover, the optimal value of its SDP relaxation, which is given below, is also zero:

$$\begin{aligned} \inf \quad & X_{12} \\ \text{s.t.} \quad & X_{11} = 0, X \succeq 0. \end{aligned}$$

However, it is not hard to observe that the dual problem is infeasible and thus the dual optimal value $d^* = -\infty$. Hence, we have $p^* = p_{SDP}^* > d^*$. Indeed, Assumption 2.1 Item 3 and Item 5 are violated and thus Theorem 2.2 does not apply.

In the unconstrained case, boundedness below of $q_0(x)$ yields attainment. This is not the case for GTRS. Consider

$$\begin{aligned} \inf \quad & x_2^2 \\ \text{s.t.} \quad & x_1 x_2 = 1. \end{aligned}$$

The optimal value 0 is unattained. Note that $\lambda = 0$ is the unique dual feasible point. Further results on attainment can be found in, e.g., [23].

3 Easy and hard cases and regular pencils

From the first two relations in (2.12), it is obvious that we have to work with $A - \lambda I$ and the eigenvalues of A for TRS. Similarly, for GTRS, we have to deal with generalized eigenvalues and the matrix pencil $A - \lambda B$ corresponding to the matrix pair (A, B) . In the latter case, due to the arbitrariness of B , several ‘singular’ cases may arise, see e.g., [26]. Below are some prototypical examples.

1. Consider the case when $A = B = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$, where $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ is a *common nullspace vector*. Then the corresponding generalized eigenvalues include all of \mathbb{R} , i.e. the determinant $\det(A - tB) = 0$ is identically 0 in t . This case is problematic since a small perturbation in the matrices can cause a big change in the solution set.
2. Let $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$. The eigenpairs include complex eigenvalues. This case is problematic as the dual problem is infeasible.

3.1 Regular case

To avoid the aforementioned singular matrix cases, we assume from now on that the matrix pencil is definite. This corresponds to the so-called (positively) regular case in [34].

$$\text{Regular Pencil} \quad \widehat{C} := A - \widehat{\lambda}B \succ 0, \text{ for some } \widehat{\lambda} \in \mathbb{R}. \quad (3.1)$$

Condition (3.1) covers most of the interesting cases. Indeed, it is shown in [34, Lemma 2.3] that if (3.1) fails and $\det(A - tB)$ is not an identically zero polynomial in t , then the feasible set of (2.2) is a singleton set. Condition (3.1) also guarantees the existence of a solution to (1.1); the proof of this fact follows similarly to that of Lemma 2.1. For convenience in the discussions below, we state this result as a proposition.

Proposition 3.1. *If condition (3.1) holds, then the GTRS problem (1.1) has an optimal solution.*

Proof. See the proofs of Lemma 2.1 and Theorem 2.2(i). \square

Under the additional regularity assumption (3.1), we would like to classify the GTRS into the so-called easy case and hard cases analogously to the classical trust region subproblem. To this end, we follow a similar line of arguments as in [34, Section 3], which analyzed the case when $\ell = u = 1$ and $b = 0$. We have to make use of the fact that under (3.1), the matrix pencil $A - \lambda B$ is diagonalizable by a congruence, see e.g., [22, Theorem 10.1]. We include a proof in Theorem 3.1 below for the convenience of the readers.

Theorem 3.1. *Recall that the pencil is (positively) regular. Then, there exists an invertible matrix S and diagonal matrices D_1 and D_2 such that*

$$A = SD_1S^T \quad \text{and} \quad B = SD_2S^T.$$

Furthermore, $A - \lambda B \succeq 0$ if, and only if,

$$\underline{\lambda} := \max_{\{i:\beta_i < 0\}} \frac{\alpha_i}{\beta_i} \leq \lambda \leq \min_{\{i:\beta_i > 0\}} \frac{\alpha_i}{\beta_i} =: \bar{\lambda}, \quad (3.2)$$

where α_i and β_i are the i th diagonal entry of D_1 and D_2 , respectively. Moreover, $(\underline{\lambda}, \bar{\lambda})$ is a nonempty open interval; at least one of the end points of the interval has to be finite; and,

$$|\alpha_i| + |\beta_i| > 0, \quad \forall i.$$

Proof. Let \widehat{C} be as in (3.1). Without loss of generality, we can assume $\widehat{\lambda} \neq 0$. Let $\widehat{C} = PDP^T$ denote the spectral decomposition (orthogonal diagonalization) of \widehat{C} . Therefore the diagonal D satisfies $0 \prec D = P^T(A - \widehat{\lambda}B)P$ and, $I = D^{-1/2}P^T(A - \widehat{\lambda}B)PD^{-1/2}$. Let Q give the orthogonal diagonalization of $D^{-1/2}P^TAPD^{-1/2}$, i.e. $Q^TD^{-1/2}P^TAPD^{-1/2}Q = D_1$. Define $S = PD^{1/2}Q$. Then it follows that $A = SD_1S^T$ and $B = -\frac{1}{\widehat{\lambda}}S(I - D_1)S^T$.

Since positive semidefiniteness is preserved under congruence, we see immediately that $A - \lambda B \succeq 0$ if, and only if,

$$D_1 - \lambda D_2 \succeq 0.$$

Since D_1 and D_2 are diagonal, we obtain further that $A - \lambda B \succeq 0$ is equivalent to

$$\underline{\lambda} := \max_{\{i:\beta_i < 0\}} \frac{\alpha_i}{\beta_i} \leq \lambda \leq \min_{\{i:\beta_i > 0\}} \frac{\alpha_i}{\beta_i} =: \bar{\lambda},$$

where α_i and β_i are the i th diagonal entry of D_1 and D_2 , respectively. Next, note that $\lambda \in (\underline{\lambda}, \bar{\lambda})$ if, and only if, $A - \lambda B \succ 0$. Thus, by (3.1), $(\underline{\lambda}, \bar{\lambda})$ is a nonempty open interval. Furthermore, since $B \neq 0$, it follows that at least one of the end points of the interval has to be finite. Finally, we note that α_i and β_i cannot both be zero, because of (3.1). \square

We remark that the equivalence in (3.2) was established earlier in [24, Theorem 5.3]; see also [34, Lemma 3.1]. Next, for each real $\lambda \in \text{cl}(\underline{\lambda}, \bar{\lambda})$, the closure of the interval, we define the first order stationary point

$$x(\lambda) := (A - \lambda B)^\dagger(a - \lambda b), \quad (3.3)$$

and the constraint evaluated at $x(\lambda)$

$$\begin{aligned} \psi(\lambda) &:= q_1(x(\lambda)) \\ &= (a - \lambda b)^T(A - \lambda B)^\dagger B(A - \lambda B)^\dagger(a - \lambda b) - 2b^T(A - \lambda B)^\dagger(a - \lambda b). \end{aligned} \quad (3.4)$$

Recall that in the TRS case where $B = I$ and $b = 0$, we can orthogonally diagonalize A and get $\underline{\lambda} = -\infty < \lambda^* \leq \bar{\lambda} = \lambda_{\min}(A)$. In addition, if we also have $a = 0$ (homogeneous case), then both $x(\lambda)$ and $\psi(\lambda)$ are constant functions, identically 0. This property extends to the general case as follows, whose proof can be found in [24, Page 202].

Lemma 3.1. Let $\lambda \in (\underline{\lambda}, \bar{\lambda})$. Then the functions $x(\lambda)$ and $\psi(\lambda)$ in (3.3) and (3.4), respectively, are differentiable with derivatives given by

$$\begin{aligned} x'(\lambda) &= (A - \lambda B)^{-1}(Bx(\lambda) - b), \\ \psi'(\lambda) &= 2x'(\lambda)^T(A - \lambda B)x'(\lambda). \end{aligned}$$

Moreover, on $(\underline{\lambda}, \bar{\lambda})$, $\psi(\lambda)$ is either constant or strictly increasing with

$$\begin{aligned} \psi \text{ constant in } (\underline{\lambda}, \bar{\lambda}) &\Leftrightarrow x'(\lambda) = 0, \text{ for some } \lambda \in (\underline{\lambda}, \bar{\lambda}) \\ &\Leftrightarrow x'(\lambda) = 0, \forall \lambda \in (\underline{\lambda}, \bar{\lambda}) \\ &\Leftrightarrow \begin{pmatrix} a \\ b \end{pmatrix} \in \text{Range} \left(\begin{bmatrix} A \\ B \end{bmatrix} \right). \end{aligned} \tag{3.5}$$

Remark 3.1. Notice that if (3.5) holds, then $(a - \lambda b) = (A - \lambda B)v$, for some v . Therefore, the objective in (2.2) becomes $-v^T(A - \lambda B)v + \ell\lambda_+ - u\lambda_-$. Thus (2.2) reduces to a linear programming problem.


We now define the easy case and the hard cases. Unlike the case for TRS, we have to consider separately the cases when $\bar{\lambda}$ or $\underline{\lambda}$ is infinite.

Definition 3.1. The easy case occurs if one of the following three cases holds:

1. both $\bar{\lambda}$ and $\underline{\lambda}$ are finite, and $-\lim_{\lambda \downarrow \underline{\lambda}} \psi(\lambda) = \lim_{\lambda \uparrow \bar{\lambda}} \psi(\lambda) = +\infty$;
2. only $\bar{\lambda}$ is finite, and $\lim_{\lambda \uparrow \bar{\lambda}} \psi(\lambda) = +\infty$;
3. only $\underline{\lambda}$ is finite, and $\lim_{\lambda \downarrow \underline{\lambda}} \psi(\lambda) = -\infty$.


Otherwise, we have the hard case.

Figure 1: $\psi(\lambda)$ when B is indefinite. The dotted lines are $\lambda = \underline{\lambda}$ and $\lambda = \bar{\lambda}$.



From the above definition and Lemma 3.1, we see immediately that in the easy case, the function ψ has to be strictly increasing on $(\underline{\lambda}, \bar{\lambda})$. Sample shapes of the function ψ in the easy and hard cases are shown in Figures 1 and 2. The next lemma and the discussion following it justify the terminology “easy case”.

Figure 2: $\psi(\lambda)$ when B is positive definite. The dotted line is $\lambda = \bar{\lambda}$.



Lemma 3.2. *If $\bar{\lambda} = +\infty$, then $\lim_{\lambda \uparrow \bar{\lambda}} \psi(\lambda) = \sup x^T Bx - 2b^T x > \ell$. Similarly, if $\underline{\lambda} = -\infty$, then $\lim_{\lambda \downarrow \underline{\lambda}} \psi(\lambda) = \inf x^T Bx - 2b^T x < u$.*

Proof. Suppose first that $\bar{\lambda} = +\infty$. Then, from the definition of $\bar{\lambda}$, we see that $B \preceq 0$, is negative semidefinite. We consider two cases: $b \in \text{Range}(B)$ and $b \notin \text{Range}(B)$.

Suppose that $b \in \text{Range}(B)$. Notice first from the definition of $\underline{\lambda}$ and $\bar{\lambda}$ that $A - \lambda B \succ 0$, and so also $B - \frac{1}{\lambda}A \prec 0$, for all sufficiently large λ . From this and the definition of $x(\lambda)$, for all sufficiently large λ and for any $x \in \mathbb{R}^n$, we see that

$$\psi(\lambda) \geq x^T \left(B - \frac{1}{\lambda}A \right) x - 2 \left(b - \frac{1}{\lambda}a \right)^T x + \frac{1}{\lambda}q_0(x(\lambda)). \quad (3.6)$$

Furthermore,

$$\frac{1}{\lambda}q_0(x(\lambda)) = \sum_{i=1}^n \left(\frac{\alpha_i(\xi_i - \lambda\gamma_i)^2}{\lambda(\alpha_i - \lambda\beta_i)^2} - \frac{2\xi_i^2 - 2\lambda\xi_i\gamma_i}{\lambda(\alpha_i - \lambda\beta_i)} \right),$$

where α_i and β_i are defined as above, while $\xi = S^{-1}a$ and $\gamma = S^{-1}b$. Since $b \in \text{Range}(B)$ means that $\gamma_i = 0$ whenever $\beta_i = 0$, we see immediately that

$$\lim_{\lambda \uparrow \bar{\lambda}} \frac{1}{\lambda}q_0(x(\lambda)) = 0.$$

Taking limits in λ on both sides of (3.6) and then taking the supremum over all $x \in \mathbb{R}^n$, we obtain that $\lim_{\lambda \uparrow \bar{\lambda}} \psi(\lambda) \geq \sup x^T Bx - 2b^T x > \ell$, where the last strict inequality is a consequence of RICQ (2.6). Hence, the first conclusion follows when $b \in \text{Range}(B)$.

Next, suppose $b \notin \text{Range}(B)$. Then by the definition of $\psi(\lambda) = q_1(x(\lambda))$, we have

$$q_1(x(\lambda)) = \sum_{i=1}^n \left(\frac{\beta_i(\xi_i - \lambda\gamma_i)^2}{(\alpha_i - \lambda\beta_i)^2} - \frac{2\gamma_i\xi_i - 2\lambda\gamma_i^2}{(\alpha_i - \lambda\beta_i)} \right).$$

The assumption on b implies that there exists i_0 such that $\beta_{i_0} = 0$ but $\gamma_{i_0} \neq 0$ (notice that by (3.1), necessarily the corresponding $\alpha_{i_0} > 0$). Thus, $\lim_{\lambda \uparrow \bar{\lambda}} \psi(\lambda) = +\infty$ and the first conclusion holds trivially in this case.

The case when $\underline{\lambda} = -\infty$ can be proved similarly. \square

From Lemmas 3.1 and 3.2, we see that, in the easy case, $\psi(\lambda) = s$ has a unique solution in the open interval $(\underline{\lambda}, \bar{\lambda})$ for any $s \in (\inf \psi, \sup \psi)$. We next show that we can alternatively characterize the easy case and hard cases via the null space of $A - \bar{\lambda}B$ or $A - \underline{\lambda}B$; see also [34, Section 3.1] for a similar analysis on the special case of GTRS (1.1) with $u = \ell = 1$ and $b = 0$.

Lemma 3.3. *If $\underline{\lambda}$ is finite, then $\lim_{\lambda \downarrow \underline{\lambda}} \psi(\lambda) = -\infty$ if, and only if, $a - \underline{\lambda}b \notin \text{Range}(A - \underline{\lambda}B)$. Similarly, if $\bar{\lambda}$ is finite, then $\lim_{\lambda \uparrow \bar{\lambda}} \psi(\lambda) = +\infty$ if, and only if, $a - \bar{\lambda}b \notin \text{Range}(A - \bar{\lambda}B)$.*

Proof. Suppose first that $\underline{\lambda}$ is finite. Then the set of indices \underline{I} such that $\alpha_i - \underline{\lambda}\beta_i = 0$ is nonempty. Moreover, from the definition of $\underline{\lambda}$ and the fact that $\underline{\lambda} < \bar{\lambda}$, we obtain that $\beta_i < 0$ for all $i \in \underline{I}$. Next, notice that for any $\lambda \in (\underline{\lambda}, \bar{\lambda})$, we have from (3.4) that

$$q_1(x(\lambda)) = \sum_{i \in \underline{I}} \left(\frac{\beta_i(\xi_i - \lambda\gamma_i)^2}{(\alpha_i - \lambda\beta_i)^2} - \frac{2\gamma_i\xi_i - 2\lambda\gamma_i^2}{(\alpha_i - \lambda\beta_i)} \right) + \sum_{i \notin \underline{I}} \left(\frac{\beta_i(\xi_i - \lambda\gamma_i)^2}{(\alpha_i - \lambda\beta_i)^2} - \frac{2\gamma_i\xi_i - 2\lambda\gamma_i^2}{(\alpha_i - \lambda\beta_i)} \right),$$

with α_i, β_i defined as above and $\xi = S^{-1}a, \gamma = S^{-1}b$. Since $a - \underline{\lambda}b \notin \text{Range}(A - \underline{\lambda}B)$ if, and only if, there exists $i_0 \in \underline{I}$ with $\xi_{i_0} - \underline{\lambda}\gamma_{i_0} \neq 0$, the first conclusion now follows immediately. The second statement can be proved similarly. \square

Using Lemma 3.3, we see that the hard and easy cases can be alternatively characterized as in Table 1.

	1. Easy case	2 (a). Hard case 1	2 (b). Hard case 2
$-\infty < \underline{\lambda} < \bar{\lambda} < \infty$	$a - \bar{\lambda}b \notin \text{Range}(A - \bar{\lambda}B)$ and $a - \underline{\lambda}b \notin \text{Range}(A - \underline{\lambda}B)$ (implies $\underline{\lambda} < \lambda^* < \bar{\lambda}$)	$a - \bar{\lambda}b \in \text{Range}(A - \bar{\lambda}B)$ or $a - \underline{\lambda}b \in \text{Range}(A - \underline{\lambda}B)$ $\underline{\lambda} < \lambda^* < \bar{\lambda}$	$a - \bar{\lambda}b \in \text{Range}(A - \bar{\lambda}B)$ $\lambda^* = \bar{\lambda}$ or $a - \underline{\lambda}b \in \text{Range}(A - \underline{\lambda}B)$ $\lambda^* = \underline{\lambda}$
$-\infty = \underline{\lambda} < \bar{\lambda} < \infty$	$a - \bar{\lambda}b \notin \text{Range}(A - \bar{\lambda}B)$ (implies $-\infty < \lambda^* < \bar{\lambda}$)	$a - \bar{\lambda}b \in \text{Range}(A - \bar{\lambda}B)$ $-\infty < \lambda^* < \bar{\lambda}$	$a - \bar{\lambda}b \in \text{Range}(A - \bar{\lambda}B)$ $\lambda^* = \bar{\lambda}$
$-\infty < \underline{\lambda} < \bar{\lambda} = \infty$	$a - \underline{\lambda}b \notin \text{Range}(A - \underline{\lambda}B)$ (implies $\underline{\lambda} < \lambda^* < \infty$)	$a - \underline{\lambda}b \in \text{Range}(A - \underline{\lambda}B)$ $\underline{\lambda} < \lambda^* < \infty$	$a - \underline{\lambda}b \in \text{Range}(A - \underline{\lambda}B)$ $\lambda^* = \underline{\lambda}$

Table 1: The three different cases for GTRS.

Before describing how the easy case and hard cases can be tackled, we need the following technical lemma concerning the smallest and largest generalized eigenvalues $\underline{\lambda}$ and $\bar{\lambda}$. This result was briefly discussed in [24, Section 5]. We include a proof here for the convenience of the readers.

Lemma 3.4. *If $\bar{\lambda}$ is finite, then $v^T Bv > 0$ for all $v \in \text{Null}(A - \bar{\lambda}B) \setminus \{0\}$. Similarly, if $\underline{\lambda}$ is finite, then $v^T Bv < 0$ for all $v \in \text{Null}(A - \underline{\lambda}B) \setminus \{0\}$.*

Proof. Suppose first that $\bar{\lambda}$ is finite and let $v \in \text{Null}(A - \bar{\lambda}B) \setminus \{0\}$. Take any $\tilde{\lambda} \in (\underline{\lambda}, \bar{\lambda})$. Then $A - \tilde{\lambda}B \succ 0$ and hence we have

$$0 < v^T(A - \tilde{\lambda}B)v = v^T(A - \bar{\lambda}B)v + (\bar{\lambda} - \tilde{\lambda})v^TBv = (\bar{\lambda} - \tilde{\lambda})v^TBv.$$

Since $\bar{\lambda} > \tilde{\lambda}$, we conclude that $v^TBv > 0$. This proves the first part. The second conclusion can be proved similarly. \square

3.1.1 Three intervals for λ

We are now ready to describe how GTRS can be tackled. Our discussion is different from that in the concluding remarks of [24] in several aspects. First, our approach does not require solving two GTRS with equality constraints. Second, we include detailed discussion about the easy case and the hard cases, with hard case, case 2, solved explicitly. In what follows, we consider three cases dependent on where 0 is located relative to the interval $(\underline{\lambda}, \bar{\lambda})$.

Case 1: $\underline{\lambda} < 0 < \bar{\lambda}$

- **Easy case:** If $\ell \leq \psi(0) \leq u$, then it is easy to check that the optimality conditions of GTRS are satisfied with $\lambda^* = 0$ and $x^* = A^{-1}a$. We have an interior solution in this case.

Otherwise, suppose we have $\psi(0) < \ell$ instead. Since ψ is strictly increasing in $(\underline{\lambda}, \bar{\lambda})$, we observe that the optimal Lagrange multiplier $\lambda^* > 0$, i.e., it has to be positive. From the complementary slackness condition in (2.12), we observe further that such a multiplier is the unique solution of $\psi(\lambda) = \ell$. The optimal solution is then given by $x(\lambda^*)$.

The case when $\psi(0) > u$ can be considered similarly, where the optimal Lagrange multiplier $\lambda^* < 0$ solves $\psi(\lambda) = u$, and the optimal solution is $x(\lambda^*)$.

- **Hard case:** If $\ell \leq \psi(0) \leq u$, then $x(0) = x^* = A^{-1}a$ and we again obtain an interior solution.

Otherwise, we take a primal step to the boundary. First, suppose we have $\psi(0) < \ell$. Since ψ is increasing, we necessarily have $\lambda^* > 0$. If $\lim_{\lambda \uparrow \bar{\lambda}} \psi(\lambda) > \ell$, then the equation $\psi(\lambda) = \ell$ is solvable and gives the optimal Lagrange multiplier $\lambda^* < \bar{\lambda}$. The optimal solution is given by $x(\lambda^*)$. On the other hand, consider $\lim_{\lambda \uparrow \bar{\lambda}} \psi(\lambda) \leq \ell$. Then $\bar{\lambda}$ is finite by Lemma 3.2 and thus $q_1(x(\bar{\lambda})) = \lim_{\lambda \uparrow \bar{\lambda}} \psi(\lambda)$. Hence Lemma 3.4 implies that there exists $v \in \text{Null}(A - \bar{\lambda}B)$ satisfying $v^TBv > 0$. Scale such v so that $q_1(x(\bar{\lambda}) + v) = \ell$. Then an optimal solution is given explicitly by $x(\bar{\lambda}) + v$, with optimal Lagrange multiplier equal to $\bar{\lambda}$.

Finally, suppose we have $\psi(0) > u$ instead. Then we necessarily have $\lambda^* < 0$. If $\lim_{\lambda \downarrow \underline{\lambda}} \psi(\lambda) < u$, then the equation $\psi(\lambda) = u$ is solvable and gives the optimal Lagrange multiplier $\lambda^* > \underline{\lambda}$. The optimal solution is given by $x(\lambda^*)$. On the other hand, if $\lim_{\lambda \downarrow \underline{\lambda}} \psi(\lambda) \geq u$, then $\underline{\lambda}$ is finite by Lemma 3.2 and thus $q_1(x(\underline{\lambda})) = \lim_{\lambda \downarrow \underline{\lambda}} \psi(\lambda)$. From Lemma 3.4, there exists $v \in \text{Null}(A - \underline{\lambda}B)$ satisfying $v^TBv < 0$. Scale such v so that $q_1(x(\underline{\lambda}) + v) = u$. Then an optimal solution is given explicitly by $x(\underline{\lambda}) + v$, with the corresponding Lagrange multiplier being $\underline{\lambda}$.

Case 2: $\bar{\lambda} \leq 0$. Notice that the optimal Lagrange multiplier has to be nonpositive and $\bar{\lambda}$ has to be finite.

- **Easy case:** We see that the optimal Lagrange multiplier λ^* has to be negative, and thus, a solution of $\psi(\lambda) = u$. The optimal solution is then given by $x(\lambda^*)$.

- **Hard case:** If $\lim_{\lambda \downarrow \underline{\lambda}} \psi(\lambda) < u < \lim_{\lambda \uparrow \bar{\lambda}} \psi(\lambda)$, then $\psi(\lambda) = u$ is solvable with the optimal Lagrange multiplier λ^* as the unique solution. The optimal solution is again given by $x(\lambda^*)$.

On the other hand, if $\lim_{\lambda \downarrow \underline{\lambda}} \psi(\lambda) \geq u$, then by Lemma 3.2, $\underline{\lambda}$ is finite and thus $q_1(x(\underline{\lambda})) = \lim_{\lambda \downarrow \underline{\lambda}} \psi(\lambda)$. Hence, by Lemma 3.4 there exists $v \in \text{Null}(A - \underline{\lambda}B)$ satisfying $v^T Bv < 0$. Scale such v so that $q_1(x(\underline{\lambda}) + v) = u$. Then an optimal solution is given explicitly by $x(\underline{\lambda}) + v$, and $\lambda^* = \underline{\lambda} < 0$. Finally, if $\lim_{\lambda \uparrow \bar{\lambda}} \psi(\lambda) \leq u$, then $q_1(x(\bar{\lambda})) = \lim_{\lambda \uparrow \bar{\lambda}} \psi(\lambda)$. Thus, by Lemma 3.4 there exists $v \in \text{Null}(A - \bar{\lambda}B)$ satisfying $v^T Bv > 0$. Scale such v so that $q_1(x(\bar{\lambda}) + v) = u$. Then an optimal solution is given explicitly by $x(\bar{\lambda}) + v$, and $\lambda^* = \bar{\lambda} \leq 0$.

Case 3: $\underline{\lambda} \geq 0$. Notice that the optimal Lagrange multiplier has to be nonnegative and $\underline{\lambda}$ has to be finite.

- **Easy case:** Arguing similarly as in Case 2, we conclude that in the easy case, the optimal Lagrange multiplier λ^* is a solution of $\psi(\lambda) = \ell$ and the optimal solution is given by $x(\lambda^*)$.
- **Hard case:** If $\lim_{\lambda \downarrow \underline{\lambda}} \psi(\lambda) < \ell < \lim_{\lambda \uparrow \bar{\lambda}} \psi(\lambda)$, then the optimal Lagrange multiplier $\lambda^* > 0$ solves $\psi(\lambda) = \ell$ and the optimal solution is given by $x(\lambda^*)$. Furthermore, if $\lim_{\lambda \downarrow \underline{\lambda}} \psi(\lambda) \geq \ell$, then an optimal solution is $x(\underline{\lambda}) + v$, where $v \in \text{Null}(A - \underline{\lambda}B)$ satisfies $v^T Bv < 0$ and $q_1(x(\underline{\lambda}) + v) = \ell$, and $\lambda^* = \underline{\lambda}$. Finally, if $\lim_{\lambda \uparrow \bar{\lambda}} \psi(\lambda) \leq \ell$, then an optimal solution is $x(\bar{\lambda}) + v$, where $v \in \text{Null}(A - \bar{\lambda}B)$ satisfies $v^T Bv > 0$ and $q_1(x(\bar{\lambda}) + v) = \ell$, and $\lambda^* = \bar{\lambda}$.

From the above discussion, we see immediately that, unless $A \succ 0$ and $A^{-1}a$ is an interior solution, the GTRS always has a solution on the boundary of the feasible set. Following the above procedures, we either end up solving an equality constrained problem, or obtain a closed form solution by moving a point to the suitable boundary along a suitable generalized eigenvector. In the next section, we will adapt the Rendl-Wolkowicz (RW) algorithm for TRS in [29] to tackle the equality constrained problem.

Before closing this section, we discuss how to compute $\lim_{\lambda \uparrow \bar{\lambda}} \psi(\lambda)$ and $\lim_{\lambda \downarrow \underline{\lambda}} \psi(\lambda)$ efficiently, which are important for the above case analysis. We only outline the procedure for computing $\lim_{\lambda \uparrow \bar{\lambda}} \psi(\lambda)$. The one for computing $\lim_{\lambda \downarrow \underline{\lambda}} \psi(\lambda)$ is similar.

Recall that $A - \bar{\lambda}B$ is singular and positive semidefinite. Thus, a nullspace vector of $A - \bar{\lambda}B$ can be found by finding an eigenvector corresponding to the smallest eigenvalue, i.e., 0.

Procedure for computing $\lim_{\lambda \uparrow \bar{\lambda}} \psi(\lambda)$.

Step 1. Take $v \in \text{Null}(A - \bar{\lambda}B) \setminus \{0\}$, $\|v\| = 1$.

If $v^T(a - \bar{\lambda}b) \neq 0$, this means $(a - \bar{\lambda}b) \notin \text{Range}(A - \bar{\lambda}B)$ and thus $\lim_{\lambda \uparrow \bar{\lambda}} \psi(\lambda) = +\infty$; quit. Else, update/deflate $A \leftarrow A + \alpha v v^T$ for some $\alpha > 0$. Repeat Step 1 if $\text{Null}(A - \bar{\lambda}B) \neq \{0\}$.

Step 2. Solve for $\bar{x} = (A - \bar{\lambda}B)^{-1}(a - \bar{\lambda}b)$. Then $\lim_{\lambda \uparrow \bar{\lambda}} \psi(\lambda) = \psi(\bar{x})$.

4 A method for solving GTRS: the extended Rendl-Wolkowicz algorithm

In this section, we discuss how the Rendl-Wolkowicz (RW) algorithm proposed in [29] can be adapted to solve the following equality constrained problem:

$$(GTRS_{=}) \quad \begin{aligned} q^* &= \inf && q_0(x) \\ &\text{s.t.} && x^T Bx - 2b^T x = s, \end{aligned} \quad (4.1)$$

where $s \in \mathbb{R}$. Recall that we assume Assumption 2.1 holds and that the regularity in (3.1) holds. Thus, by Proposition 3.1, (4.1) has an optimal solution. Following [29], we consider the following chain of inequalities.

$$\begin{aligned}
q^* &= \inf_{\substack{x^T Bx - 2b^T xy_0 = s \\ y_0^2 = 1}} x^T Ax - 2a^T xy_0 \\
&= \sup_t \inf_{\substack{x^T Bx - 2b^T xy_0 = s \\ y_0^2 = 1}} x^T Ax - 2a^T xy_0 + t(y_0^2 - 1) \\
&\geq \sup_t \inf_{x^T Bx - 2b^T xy_0 + y_0^2 = s+1} x^T Ax - 2a^T xy_0 + t(y_0^2 - 1) \\
&\geq \sup_{r,t} \inf_{x,y_0} x^T Ax - 2a^T xy_0 + t(y_0^2 - 1) + r(x^T Bx - 2b^T xy_0 + y_0^2 - s - 1) \\
&= \sup_{r,\tau} \inf_{x,y_0} x^T Ax - 2a^T xy_0 + \tau(y_0^2 - 1) + r(x^T Bx - 2b^T xy_0 - s) \\
&= \sup_r \left(\sup_{\tau} \inf_{x,y_0} x^T Ax - 2a^T xy_0 + \tau(y_0^2 - 1) + r(x^T Bx - 2b^T xy_0 - s) \right)
\end{aligned} \tag{4.2}$$

Let Ω denote the set of all real numbers r such that $A + rB \succeq 0$ and $a + rb \in \text{Range}(A + rB)$. This set is nonempty by Assumption 2.1, Item 5. Furthermore, if $r \notin \Omega$, then it is easy to see that

$$\begin{aligned}
&\inf_{x,y_0} x^T Ax - 2a^T xy_0 + \tau(y_0^2 - 1) + r(x^T Bx - 2b^T xy_0 - s) \\
&= \inf_{y_0} \inf_x x^T Ax - 2a^T xy_0 + \tau(y_0^2 - 1) + r(x^T Bx - 2b^T xy_0 - s) = -\infty,
\end{aligned} \tag{4.3}$$

regardless of τ . Thus, continuing from (4.2) we have

$$\begin{aligned}
&\sup_r \left(\sup_{\tau} \inf_{x,y_0} x^T Ax - 2a^T xy_0 + \tau(y_0^2 - 1) + r(x^T Bx - 2b^T xy_0 - s) \right) \\
&= \sup_{r \in \Omega} \left(\sup_{\tau} \inf_{x,y_0} x^T Ax - 2a^T xy_0 + \tau(y_0^2 - 1) + r(x^T Bx - 2b^T xy_0 - s) \right) \\
&= \sup_{r \in \Omega} \inf_{x, y_0^2 = 1} x^T Ax - 2a^T xy_0 + r(x^T Bx - 2b^T xy_0 - s) \\
&= \sup_{r \in \Omega} \inf_x x^T Ax - 2a^T x + r(x^T Bx - 2b^T x - s) \\
&= \sup_r \inf_x x^T Ax - 2a^T x + r(x^T Bx - 2b^T x - s) \\
&= \inf_{x^T Bx - 2b^T x = s} x^T Ax - 2a^T x = q^*,
\end{aligned}$$

where: the first equality follows from the observation in (4.3); while the second equality follows from Theorem 2.2, since for any $r \in \Omega$, a Schur complement argument implies that there exists sufficiently large τ such that

$$\begin{bmatrix} \tau & -(a + rb)^T \\ -(a + rb) & A + rB \end{bmatrix} \succeq 0;$$

the third equality follows from the homogenization; the fourth equality follows from an observation similar to (4.3) for the inner optimization problem; while the last equality follows from Theorem 2.2. Thus, equality holds throughout in (4.2). In particular, from the third line in (4.2), we have

$$q^* = \sup_t \underbrace{k_0(t) - t}_{k(t)},$$

where

$$\begin{aligned}
k_0(t) &= \inf_{x^T Bx - 2b^T xy_0 + y_0^2 = s+1} x^T Ax - 2a^T xy_0 + ty_0^2 \\
&= \inf \left\{ y^T \begin{bmatrix} t & -a^T \\ -a & A \end{bmatrix} y : y^T \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} y = s + 1 \right\}.
\end{aligned}$$

The RW algorithm adapted to our context amounts to solving (4.1) via maximizing $k(t)$.

4.1 Properties of $k_0(t)$

In this subsection, we discuss differentiability of the function $k_0(t)$ and show that a maximizer of $k(t) := k_0(t) - t$ necessarily exists. In addition to Assumption 2.1 and the regularity in (3.1), we also assume that $s + 1 \neq 0$, which can always be satisfied by scaling B and b . Furthermore, to avoid triviality, we assume that (3.5) fails so that ψ is strictly increasing. Indeed, in view of Remark 3.1, if (3.5) holds, (2.2) is a simple linear programming problem and can be readily solved.

Our analysis is based on the following function:

$$d(\lambda) := \lambda + (a - \lambda b)^T (A - \lambda B)^\dagger (a - \lambda b).$$

This function reduces to the $d(\lambda)$ considered in [29] for problem (1.2). Since $d'(\lambda) = 1 + \psi(\lambda)$ on $(\underline{\lambda}, \bar{\lambda})$ and (3.5) fails, we see that this function is strictly convex with strictly increasing derivative on $(\underline{\lambda}, \bar{\lambda})$. With a Schur complement argument, we get the following result.

Lemma 4.1. *The following four properties hold.*

(i) *Suppose $\lambda \in (\underline{\lambda}, \bar{\lambda})$. Then*

$$\begin{aligned} \begin{bmatrix} t & -a^T \\ -a & A \end{bmatrix} - \lambda \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} \succ 0 &\Leftrightarrow t > d(\lambda), \\ \begin{bmatrix} t & -a^T \\ -a & A \end{bmatrix} - \lambda \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} \succeq 0 &\Leftrightarrow t \geq d(\lambda). \end{aligned}$$

(ii) *Suppose $\lim_{\lambda \downarrow \underline{\lambda}} d(\lambda)$ is finite and $\underline{\lambda}$ is finite. Then*

$$\begin{bmatrix} t & -a^T \\ -a & A \end{bmatrix} - \underline{\lambda} \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} \succeq 0 \Leftrightarrow t \geq d(\underline{\lambda}).$$

(iii) *Suppose $\lim_{\lambda \uparrow \bar{\lambda}} d(\lambda)$ is finite and $\bar{\lambda}$ is finite. Then*

$$\begin{bmatrix} t & -a^T \\ -a & A \end{bmatrix} - \bar{\lambda} \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} \succeq 0 \Leftrightarrow t \geq d(\bar{\lambda}).$$

(iv) *For any $s \in \mathbb{R}$, we have*

$$\sup \left\{ (s+1)\lambda : \begin{bmatrix} t & -a^T \\ -a & A \end{bmatrix} - \lambda \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} \succeq 0 \right\} = \sup \{ (s+1)\lambda : \lambda \in (\underline{\lambda}, \bar{\lambda}), t \geq d(\lambda) \}.$$

Proof. For part (i), notice that $\lambda \in (\underline{\lambda}, \bar{\lambda})$ if, and only if, $A - \lambda B \succ 0$. Hence the conclusion follows immediately from an application of the Schur complement. For part (ii), notice first that $\lim_{\lambda \downarrow \underline{\lambda}} d(\lambda)$ being finite implies $d(\underline{\lambda}) = \lim_{\lambda \downarrow \underline{\lambda}} d(\lambda)$ and hence d is continuous and convex on $[\underline{\lambda}, \bar{\lambda})$. Next, by taking $\lambda \in (\underline{\lambda}, \bar{\lambda})$ and $r > d(\lambda)$, we have from part (i) that

$$\begin{bmatrix} r & -a^T \\ -a & A \end{bmatrix} - \lambda \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} \succeq 0.$$

Hence, the convexity of $d(\cdot)$ and the continuity of $d(\cdot)$ at $\underline{\lambda}$ yields

$$\begin{aligned} \begin{bmatrix} t & -a^T \\ -a & A \end{bmatrix} - \underline{\lambda} \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} \succeq 0 &\Leftrightarrow \begin{bmatrix} (1-\epsilon)t + \epsilon r & -a^T \\ -a & A \end{bmatrix} - [(1-\epsilon)\underline{\lambda} + \epsilon\lambda] \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} \succeq 0, \quad \forall 0 < \epsilon < 1, \\ t \geq d(\underline{\lambda}) &\Leftrightarrow (1-\epsilon)t + \epsilon r \geq d((1-\epsilon)\underline{\lambda} + \epsilon\lambda), \quad \forall 0 < \epsilon < 1. \end{aligned}$$

Since $(1-\epsilon)\underline{\lambda} + \epsilon\lambda \in (\underline{\lambda}, \bar{\lambda})$ for all $0 < \epsilon < 1$, the conclusion of part (ii) now follows immediately from the above two relations and part (i). Part (iii) can be proved similarly. Finally, part (iv) follows immediately from parts (i), (ii), (iii) and the fact that

$$A - \lambda B \succeq 0 \Leftrightarrow \lambda \in \text{cl}(\underline{\lambda}, \bar{\lambda}).$$

□

We next consider three cases: B is indefinite, positive semidefinite and negative semidefinite. We only discuss the first two cases in detail since the third case is analogous to the second case. For this purpose, we will need to consider the following quantities:

$$\begin{aligned} t_0 &:= \inf_{\underline{\lambda} < \lambda < \bar{\lambda}} d(\lambda), & \bar{t} &:= \begin{cases} \lim_{\lambda \uparrow \bar{\lambda}} d(\lambda) & \text{if } \bar{\lambda} < \infty, \\ \infty & \text{else,} \end{cases} \\ s_0 &:= \inf x^T B x - 2b^T x. & \underline{t} &:= \begin{cases} \lim_{\lambda \downarrow \underline{\lambda}} d(\lambda) & \text{if } \underline{\lambda} > -\infty, \\ \infty & \text{else.} \end{cases} \end{aligned} \tag{4.4}$$

We remark that these four quantities are not necessarily finite. Furthermore, it follows from weak duality that

$$t_0 \geq - \inf_{x^T B x - 2b^T x = -1} x^T A x - 2a^T x.$$

Hence, t_0 is finite when $s_0 \leq -1$.

Case 1: B is indefinite Notice that in this case, the quantities t_0 , $\underline{\lambda}$ and $\bar{\lambda}$ are finite. In the next theorem, we show that $k_0(t)$ is well-defined on a closed interval, and is differentiable in the interior of that interval, except possibly at one point.

Theorem 4.1. *The function $k_0(t)$ is well-defined and continuous on $t \geq t_0$. Moreover:*

- (i) *when $s+1 > 0$, $k_0(t)$ is differentiable on $(t_0, \bar{t}) \cup (\bar{t}, \infty)$, and $k_0(t) = (s+1)\bar{\lambda}$ on $t > \bar{t}$;*
- (ii) *when $s+1 < 0$, $k_0(t)$ is differentiable on $(t_0, \underline{t}) \cup (\underline{t}, \infty)$, and $k_0(t) = (s+1)\underline{\lambda}$ on $t > \underline{t}$.*

Proof. Suppose first that $t > t_0$. Then from the definition of t_0 , there exists λ so that

$$\lambda \in (\underline{\lambda}, \bar{\lambda}), \quad t - d(\lambda) > 0.$$

It follows from this, Lemma 4.1(i), (iv) and Theorem 2.2 that

$$\begin{aligned} k_0(t) &= \inf \left\{ y^T \begin{bmatrix} t & -a^T \\ -a & A \end{bmatrix} y : y^T \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} y = s+1 \right\} \\ &= \sup \left\{ (s+1)\lambda : \begin{bmatrix} t & -a^T \\ -a & A \end{bmatrix} - \lambda \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} \succeq 0 \right\} > -\infty. \end{aligned} \tag{4.5}$$

Thus, $k_0(t)$ is finite. Next, suppose $t = t_0$. Notice that the infimum value t_0 in (4.4) has to be attained at some $\lambda_0 \in [\underline{\lambda}, \bar{\lambda}]$. Thus, we have from Lemma 4.1 that

$$\begin{bmatrix} t_0 & -a^T \\ -a & A \end{bmatrix} - \lambda_0 \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} \succeq 0.$$

Hence, from Theorem 2.2, we conclude that (4.5) holds with $t = t_0$. Finally, since k_0 is a one variable concave function, it is continuous in the interior of its domain [30, Theorem 10.1] and is lower semicontinuous up to the boundary [30, Theorem 10.2]. Furthermore, since k_0 is the infimum of a family of continuous functions

$$t \mapsto y^T \begin{bmatrix} t & -a^T \\ -a & A \end{bmatrix} y,$$

it also follows that k_0 has to be upper semicontinuous. Thus, k_0 is continuous on $t \geq t_0$.

We now turn to part (i). Since $d(\lambda)$ is a strictly convex function, we deduce that

$$d(\lambda) = t \tag{4.6}$$

has at most 2 solutions in $(\underline{\lambda}, \bar{\lambda})$ for any $t > t_0$, at which d has opposite slopes.

We first consider the case when \bar{t} is infinite. In this case, there must be a unique solution $\mu \in (\underline{\lambda}, \bar{\lambda})$ to (4.6) with $d'(\mu) > 0$. Since $s + 1 > 0$, we see from Lemma 4.1(iv) and (4.5) that $k_0(t) = (s + 1)\mu$. Moreover, since $d'(\mu) > 0$, by the inverse function theorem, μ is differentiable at t and $\mu'(t) = \frac{1}{d'(\mu)}$. This shows that k_0 is differentiable at t .

We next assume that \bar{t} is finite. When $t_0 < t < \bar{t}$, it still holds true that (4.6) has a unique solution $\mu \in (\underline{\lambda}, \bar{\lambda})$ satisfying $d'(\mu) > 0$ and $k_0(t) = (s + 1)\mu$ as above. Similarly, it can be shown that $k_0(t)$ is differentiable on $t_0 < t < \bar{t}$.

On the other hand, when $t \geq \bar{t} = d(\bar{\lambda})$, we see from Lemma 4.1(iii) that

$$\begin{bmatrix} t & -a^T \\ -a & A \end{bmatrix} - \bar{\lambda} \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} \succeq 0.$$

Hence, from Lemma 4.1(iv) and (4.5), we conclude that $k_0(t) = (s + 1)\bar{\lambda}$. This completes the proof of part (i).

The cases when $s + 1 < 0$ in part (ii) can be proved similarly, by noting that $k_0(t) = (s + 1)\nu$ for $t_0 < t < \underline{t}$, where $\nu = \nu(t)$ is the unique root of (4.6) in $(\underline{\lambda}, \bar{\lambda})$ with $d'(\nu) < 0$. \square

Before proceeding, we collect several facts that are readily obtained from the proof of Theorem 4.1.

Corollary 4.1. (i) *If $s + 1 > 0$ and $t_0 < t < \bar{t}$, then $k_0(t) = (s + 1)\mu$, where $\mu = \mu(t)$ is the unique root of (4.6) on $(\underline{\lambda}, \bar{\lambda})$ with $d'(\mu) > 0$; moreover, $\mu'(t) = \frac{1}{d'(\mu)} > 0$.*


(ii) *If $s + 1 < 0$ and $t_0 < t < \underline{t}$, then $k_0(t) = (s + 1)\nu$, where $\nu = \nu(t)$ is the unique root of (4.6) on $(\underline{\lambda}, \bar{\lambda})$ with $d'(\nu) < 0$; moreover, $\nu'(t) = \frac{1}{d'(\nu)} < 0$.*

In the next proposition, we confirm that the domain of $k_0(t)$ is actually $t \geq t_0$, i.e., $k_0(t) = -\infty$ whenever $t < t_0$. This is a consequence of the more general result from Proposition A.1.

Corollary 4.2. *When $t < t_0$, $k_0(t) = -\infty$.*

Notice that $k'(t) = k'_0(t) - 1$ whenever $t > t_0$ and k_0 is differentiable at t . Moreover, $\lim_{t \uparrow \infty} k'_0(t) = 0$, in view of Corollary 4.1 and Theorem 4.1. Thus, it holds true that a maximizer t^* of $k(t)$ must exist.

Figure 3: $d(\lambda)$ when B is indefinite. The dotted lines are $\lambda = \underline{\lambda}$ and $\lambda = \bar{\lambda}$.



Case 2: B is positive semidefinite In this case, we have $\bar{\lambda}$ being finite and $\underline{\lambda} = -\infty$. Moreover, we have the following result concerning the behavior of $d(\lambda)$ as λ goes to $-\infty$.

Proposition 4.1. *It holds that $\lim_{\lambda \rightarrow -\infty} d'(\lambda) = 1 + s_0$. Hence*

- (i) *If $s_0 < -1$, then $\lim_{\lambda \rightarrow -\infty} d(\lambda) = \infty$;*
- (ii) *If $s_0 = -1$, then $\lim_{\lambda \rightarrow -\infty} d(\lambda) = t_0 > -\infty$;*
- (iii) *If $s_0 > -1$, then $\lim_{\lambda \rightarrow -\infty} d(\lambda) = -\infty$.*

Proof. The fact that $\lim_{\lambda \rightarrow -\infty} d'(\lambda) = 1 + s_0$ follows from $d'(\lambda) = 1 + \psi(\lambda)$ and Lemma 3.2. The rest of the conclusion then follows immediately. \square

In view of the definition of s_0 and Assumption 2.1, Item 3, we only need to consider $s > s_0$ in subsequent analysis. We discuss the differentiability property of $k_0(t)$ in the next theorem.

Theorem 4.2. (i) *Suppose that $s_0 < -1$. Then $k_0(t)$ is continuous and well-defined on $t \geq t_0$. Furthermore, when $s + 1 > 0$, $k_0(t)$ is differentiable on $(t_0, \bar{t}) \cup (\bar{t}, \infty)$ and $k_0(t) = (s + 1)\bar{\lambda}$ on $t > \bar{t}$; when $s + 1 < 0$, $k_0(t)$ is differentiable on (t_0, ∞) .*

(ii) *Suppose that $s_0 = -1$. Then $k_0(t)$ is continuous and well-defined on $t > t_0$. Furthermore, when $s + 1 > 0$, $k_0(t)$ is differentiable on $(t_0, \bar{t}) \cup (\bar{t}, \infty)$ and $k_0(t) = (s + 1)\bar{\lambda}$ on $t > \bar{t}$.*

(iii) *Suppose that $s_0 > -1$. Then $k_0(t)$ is continuous and well-defined everywhere. Furthermore, when $s + 1 > 0$, $k_0(t)$ is differentiable on $(t_0, \bar{t}) \cup (\bar{t}, \infty)$ and $k_0(t) = (s + 1)\bar{\lambda}$ on $t > \bar{t}$.*

Proof. The theorem can be proved similarly as Theorem 4.1 once we show that (4.5) holds for each case. To prove this, it suffices to show that Assumption 2.1, Item 3 is satisfied for the corresponding GTRS in (4.5)

in each case. Suppose first that $s_0 + 1 < 0$. Then it is easy to see that $\begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix}$ has to be indefinite and so Assumption 2.1 Item 3 is satisfied. On the other hand, if $s_0 + 1 \geq 0$, then

$$\begin{aligned} \inf_y y^T \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} y &= \min \left\{ \inf_{y_1 \neq 0} y^T \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} y, \inf_y y^T B y \right\} \\ &\geq \min \left\{ \inf_{y_1 \neq 0} y_1^2 (s_0 + 1), \inf_y y^T B y \right\} \geq 0, \end{aligned}$$

showing that $\begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix}$ is positive semidefinite. Since $s + 1 > s_0 + 1$, we see that Assumption 2.1 Item 3 is also satisfied in this case. \square

As in Corollary 4.1, we have the following relationship between $k_0(t)$ and the roots of (4.6), which can be obtained from a detailed proof of Theorem 4.2.

- Corollary 4.3.** (i) *If $s + 1 > 0$ and $t_0 < t < \bar{t}$, then $k_0(t) = (s + 1)\mu$, where $\mu = \mu(t)$ is the unique root of (4.6) on $(\underline{\lambda}, \bar{\lambda})$ with $d'(\mu) > 0$; moreover, $\mu'(t) = \frac{1}{d'(\mu)} > 0$;*
- (ii) *If $s + 1 < 0$ and $t_0 < t$, then $k_0(t) = (s + 1)\nu$, where $\nu = \nu(t)$ is the unique root of (4.6) on $(\underline{\lambda}, \bar{\lambda})$ with $d'(\nu) < 0$; moreover, $\nu'(t) = \frac{1}{d'(\nu)} < 0$.*

The existence of a maximizer t^* of $k(t)$ in this case is less trivial and is obtained as a consequence of the next theorem.

- Theorem 4.3.** (i) *Suppose that $s_0 < -1$. When $s + 1 > 0$, we have $\lim_{t \rightarrow \infty} k'(t) = -1$. When $s + 1 < 0$, we have $\lim_{t \rightarrow \infty} k'(t) = \frac{s+1}{s_0+1} - 1 < 0$. Furthermore, in either case, $k(t) = -\infty$ for $t < t_0$.*
- (ii) *Suppose that $s_0 = -1$. When $s + 1 > 0$, we have $\lim_{t \rightarrow \infty} k'(t) = -1$ and $\lim_{t \downarrow t_0} k'(t) = \infty$. Furthermore, $k(t) = -\infty$ for $t \leq t_0$.*
- (iii) *Suppose that $s_0 > -1$. When $s + 1 > 0$, we have $\lim_{t \rightarrow \infty} k'(t) = -1$ and $\lim_{t \rightarrow -\infty} k'(t) = \frac{s+1}{s_0+1} - 1 > 0$.*

Proof. First of all, in all three cases, when $s + 1 > 0$, the limit of $k'(t)$ as t goes to ∞ can be found similarly as in the case when B is indefinite.

We now consider part (i). When $s + 1 < 0$, by Corollary 4.3, we have $k'(t) = \frac{s+1}{d'(\nu)} - 1$. Moreover, from Proposition 4.1, we see that $\lim_{\lambda \rightarrow -\infty} d(\lambda) = \infty$. Thus, ν is defined for all sufficiently large t and $\nu \rightarrow -\infty$ as $t \rightarrow \infty$. We conclude from these results and Proposition 4.1 that $\lim_{t \rightarrow \infty} k'(t) = \frac{s+1}{s_0+1} - 1 < 0$. The conclusion that $k(t) = -\infty$ for $t < t_0$ follows from Proposition A.1.

We next turn to part (ii). Since d is strictly increasing, it follows from Proposition 4.1 that

$$\bar{t} = \lim_{\lambda \uparrow \bar{\lambda}} d(\lambda) > \lim_{\lambda \rightarrow -\infty} d(\lambda) = t_0.$$

By Corollary 4.3, we have $k'(t) = \frac{s+1}{d'(\mu)} - 1$ for $\bar{t} > t > t_0$. Furthermore, from Proposition 4.1, we see that $\lim_{\lambda \rightarrow -\infty} d(\lambda) = t_0 > -\infty$ and hence $\mu \rightarrow -\infty$ as $t \rightarrow t_0$. We conclude from this, Proposition 4.1 and the fact that $s_0 + 1 = 0$ that $\lim_{t \downarrow t_0} k'(t) = \infty$. Furthermore, since $k(t) = (s + 1)\mu(t) - t$ for $\bar{t} > t > t_0$, we have by Corollary 4.3 that

$$\lim_{t \downarrow t_0} k(t) = (s + 1) \lim_{t \downarrow t_0} \mu(t) - t_0 = -\infty.$$

It then follows from concavity of k that $k(t) = -\infty$ when $t \leq t_0$.

Finally, for part (iii), note that from Proposition 4.1, we have $\lim_{\lambda \rightarrow -\infty} d(\lambda) = -\infty$ and hence $\mu \rightarrow -\infty$ as $t \rightarrow -\infty$. Thus, we conclude further from Proposition 4.1 that $\lim_{t \rightarrow -\infty} k'(t) = \frac{s+1}{s_0+1} - 1 > 0$. \square

Figure 4: $d(\lambda)$ when B is positive definite with $s_0 < -1$. The dotted line is $\lambda = \bar{\lambda}$.





Figure 5: $d(\lambda)$ when B is positive definite with $s_0 > -1$. The dotted line is $\lambda = \bar{\lambda}$.



4.2 Recovering solution of GTRS from maximizing $k(t)$

In this subsection, we discuss how one can obtain a solution to (4.1) after obtaining a maximizer t^* of $k(t)$. In addition, we will argue that such a maximizer has to be unique.

We focus on the case when $s + 1 > 0$. We will briefly comment on the case when $s + 1 < 0$ at the end of this subsection. In this case, for any $t \in (t_0, \bar{t})$, we see from the definition of t_0 that the maximization problem in (4.5) is strictly feasible, and hence the infimum in (4.5) is attained. By Theorem 2.3, the infimum is attained at some generalized eigenvector of the matrix pair

$$\left(\begin{bmatrix} t & -a^T \\ -a & A \end{bmatrix}, \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} \right), \quad (4.7)$$

with corresponding eigenvalue μ . Recall from Lemma 4.1 and (4.5) that μ is the unique root of (4.6) with $d'(\mu) > 0$. Let $y(t) = \begin{bmatrix} y_0(t) \\ z(t) \end{bmatrix}$ be a generalized eigenvector attaining the infimum in (4.5). Since $\mu \in (\underline{\lambda}, \bar{\lambda})$ by Corollary 4.1, it follows that the first coordinate $y_0(t)$ of $y(t)$ must be nonzero. Thus, we may further scale the vector so that $y_0(t) > 0$. Moreover, since $s + 1 > 0$, we can scale the vector again so that $y^T \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} y = 1$.

We claim that such a vector is unique. To see this, notice first that for any $\lambda \in (\underline{\lambda}, \bar{\lambda})$, we have

$$\det \left(\begin{bmatrix} t & -a^T \\ -a & A \end{bmatrix} - \lambda \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} \right) = (t - d(\lambda)) \det(A - \lambda B).$$

It thus follows from this, $\det(A - \mu B) > 0$ and $d'(\mu) > 0$ that μ is a root of multiplicity one of the polynomial

$$\lambda \mapsto \det \left(\begin{bmatrix} t - \lambda & -(a - \lambda b)^T \\ -(a - \lambda b) & A - \lambda B \end{bmatrix} \right).$$

Hence, the dimension of the nullspace of $\begin{bmatrix} t - \mu & -(a - \mu b)^T \\ -(a - \mu b) & A - \mu B \end{bmatrix}$ is one; i.e., any generalized eigenvector corresponding to μ differs only by a scaling. Since the scaling is uniquely determined by the constraints $y^T \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} y = 1$ and $y_0 > 0$, the generalized eigenvector constructed under these two restrictions is unique. Using standard arguments and the implicit function theorem, one can show in addition that $y(t)$ is differentiable for $t_0 < t < \bar{t}$.

By [5, Theorem 4.13] and using the above notations, the derivative of $k(t)$ at any $t \in (t_0, \bar{t})$ is given by

$$k'(t) = (s + 1)y_0^2(t) - 1 = (s + 1)\mu'(t) - 1. \quad (4.8)$$

We next analyze the following cases. Recall that we currently assume $s + 1 > 0$ and that t^* is a maximizer of $k(t)$.

Case 1: $t^* \in (t_0, \bar{t})$. In this case, k is differentiable at t^* . Hence, necessarily

$$y_0^2(t^*) = \frac{1}{s + 1}.$$

It follows, after a simple calculation, that $x^* := \frac{z(t^*)}{y_0(t^*)}$ satisfies $x^{*T} B x^* - 2b^T x^* = s$, from which it follows easily by checking optimality conditions that x^* is an optimal solution to (4.1), with the Lagrange multiplier given by $\mu(t^*) \in (\underline{\lambda}, \bar{\lambda})$.

Case 2: $t^* = t_0 < \bar{t}$. In this case, $k'(t) \leq 0$ for all $t > t_0$ sufficiently close to t_0 . Hence

$$\limsup_{t \downarrow t_0} y_0^2(t) \leq \frac{1}{s+1}. \quad (4.9)$$

Furthermore, since $t^* \in \text{dom}(k)$ by definition of maximizer, the assumption implies that $t_0 \in \text{dom}(k)$ and thus k_0 is continuous at t_0 . Then, by continuity of $k_0(t)$ and hence of $\mu(t)$, we see that $\lim_{t \downarrow t_0} \mu(t)$ exists (and is finite). We claim that $\lim_{t \downarrow t_0} \mu(t) = \underline{\lambda}$. To see this, note that if $\lim_{t \downarrow t_0} \mu(t) \in (\underline{\lambda}, \bar{\lambda})$, then $d'(\lim_{t \downarrow t_0} \mu(t)) = 0$, which implies that $\mu'(t) = \frac{1}{d'(\mu(t))} > 0$ is arbitrarily large as $t \downarrow t_0$. In view of (4.8), this contradicts the fact that $k'(t) \leq 0$ for t close to t_0 . On the other hand, if $\lim_{t \downarrow t_0} \mu(t) = \bar{\lambda}$, then $\bar{\lambda}$ is finite. But by definition of $\mu(t)$, it is the unique root of $d(\lambda) = t$ on $(\underline{\lambda}, \bar{\lambda})$ with positive slope and thus $\mu(t) > \lim_{t \downarrow t_0} \mu(t) = \bar{\lambda}$ for all $\bar{t} > t > t_0$. But $\mu(t) < \bar{\lambda}$ by definition, and we arrive at a contradiction. Thus, we must have $\lim_{t \downarrow t_0} \mu(t) = \underline{\lambda}$. In particular, $\underline{\lambda}$ is finite.

We next claim that there exists a generalized eigenvector $y^* = \begin{bmatrix} y_0^* \\ z^* \end{bmatrix}$ of the matrix pair

$$\left(\begin{bmatrix} t_0 & -a^T \\ -a & A \end{bmatrix}, \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} \right)$$

corresponding to $\underline{\lambda}$ such that $y_0^* \neq 0$ and $y^{*T} \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} y^* = 1$.

To prove this, fix $t_1 \in (t_0, \bar{t})$ and consider $\{y(t) : t_0 < t < t_1\}$. Since $t_1 > t_0$, by the definition of t_0 , there exists $\hat{\lambda} > \underline{\lambda}$ such that

$$\begin{bmatrix} t_1 & -a^T \\ -a & A \end{bmatrix} - \hat{\lambda} \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} \succeq \epsilon I, \quad (4.10)$$

for some $\epsilon > 0$. Then, for any $t_0 < t < t_1$, we have that

$$\begin{aligned} 0 &= y(t)^T \left(\begin{bmatrix} t & -a^T \\ -a & A \end{bmatrix} - \mu(t) \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} \right) y(t) = y(t)^T \begin{bmatrix} t & -a^T \\ -a & A \end{bmatrix} y(t) - \mu(t) \\ &= y_0^2(t)(t - t_1) + y(t)^T \begin{bmatrix} t_1 & -a^T \\ -a & A \end{bmatrix} y(t) - \mu(t) \\ &= y_0^2(t)(t - t_1) + y(t)^T \left(\begin{bmatrix} t_1 & -a^T \\ -a & A \end{bmatrix} - \hat{\lambda} \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} \right) y(t) - (\mu(t) - \hat{\lambda}) \\ &\geq y_0^2(t)(t - t_1) - (\mu(t) - \hat{\lambda}) + \epsilon \|y(t)\|^2, \end{aligned}$$

where the last inequality follows from (4.10). Thus,

$$y_0^2(t)(t_1 - t) + (\mu(t) - \hat{\lambda}) \geq \epsilon \|y(t)\|^2. \quad (4.11)$$

This together with (4.9) shows that $\{y(t) : t_0 < t < t_1\}$ is bounded. Consider any cluster point y^* of $\{y(t)\}$ as $t \downarrow t_0$. Then $y^{*T} \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} y^* = 1$, and hence in particular $y^* \neq 0$. Moreover, since $\mu(t) - \hat{\lambda} \rightarrow \underline{\lambda} - \hat{\lambda} < 0$, we see further from (4.11) that $y_0^* > 0$. Finally, it is easy to check that y^* is a generalized eigenvector of the matrix pair

$$\left(\begin{bmatrix} t_0 & -a^T \\ -a & A \end{bmatrix}, \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} \right)$$

corresponding to the eigenvalue $\underline{\lambda}$.

Next, define $x^* := \frac{1}{y_0^*} z^*$. Then it follows from (4.9) that $x^{*T} B x^* - 2b^T x^* \geq s$ and from the definition of $y(t)$ that $(A - \underline{\lambda} B)x^* = a - \underline{\lambda} b$. By Lemma 3.4, there exists $v \in \text{Null}(A - \underline{\lambda} B)$ with $v^T B v < 0$, and thus $x^* + \alpha v$ would solve (4.1), for some suitable $\alpha > 0$.

Case 3: $t^* = \bar{t} > t_0$. In this case, we must have $k'(t) \geq 0$ for all $t_0 < t < \bar{t}$. Hence

$$\liminf_{t \uparrow \bar{t}} y_0^2(t) \geq \frac{1}{s+1}. \quad (4.12)$$

Furthermore, since \bar{t} is finite by assumption, we must have $\bar{\lambda}$ finite from the definition of \bar{t} and thus $d(\bar{\lambda}) = \lim_{\lambda \uparrow \bar{\lambda}} d(\lambda) = \bar{t}$. This, together with the continuity of $\mu(t)$ and the fact that $d(\lambda)$ is strictly increasing when $\lambda > \lim_{t \downarrow t_0} \mu(t)$, implies that $\lim_{t \uparrow \bar{t}} \mu(t) = \bar{\lambda}$. We claim that there exists a generalized eigenvector $y^* = \begin{bmatrix} y_0^* \\ z^* \end{bmatrix}$ of the matrix pair

$$\left(\begin{bmatrix} \bar{t} & -a^T \\ -a & A \end{bmatrix}, \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} \right)$$

corresponding to $\bar{\lambda}$ such that $y_0^* \neq 0$ and $y^{*T} \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} y^* = 1$.

We first show that the parametrized set of eigenvectors $\{y(t) : t_1 \leq t < \bar{t}\}$ is bounded for any fixed $t_1 \in (t_0, \bar{t})$. Since $t_1 \in (t_0, \bar{t})$, by definition of t_0 and Lemma 4.1(i), there exists $\hat{\lambda} < \bar{\lambda}$ such that

$$\begin{bmatrix} t_1 & -a^T \\ -a & A \end{bmatrix} - \hat{\lambda} \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} \succeq \epsilon I, \quad (4.13)$$

for some $\epsilon > 0$. Using this and the definition of generalized eigenvector and eigenvalue, we have, for any $t_1 \leq t < \bar{t}$, that

$$\begin{aligned} 0 &= y(t)^T \left(\begin{bmatrix} t & -a^T \\ -a & A \end{bmatrix} - \mu(t) \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} \right) y(t) = y(t)^T \begin{bmatrix} t & -a^T \\ -a & A \end{bmatrix} y(t) - \mu(t) \\ &= y_0^2(t)(t - t_1) + y(t)^T \begin{bmatrix} t_1 & -a^T \\ -a & A \end{bmatrix} y(t) - \mu(t) \\ &= y_0^2(t)(t - t_1) + y(t)^T \left(\begin{bmatrix} t_1 & -a^T \\ -a & A \end{bmatrix} - \hat{\lambda} \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} \right) y(t) - (\mu(t) - \hat{\lambda}) \\ &\geq -(\mu(t) - \hat{\lambda}) + \epsilon \|y(t)\|^2, \end{aligned}$$

where the last inequality follows from (4.13). This shows that $\{y(t) : t_1 \leq t < \bar{t}\}$ is bounded. Consider any cluster point y^* of $\{y(t)\}$ as $t \uparrow \bar{t}$. Due to (4.12), the first coordinate y_0^* of y^* is nonzero. It is easy to check that $y^{*T} \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} y^* = 1$ and that y^* is a generalized eigenvector of the matrix pair

$$\left(\begin{bmatrix} \bar{t} & -a^T \\ -a & A \end{bmatrix}, \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} \right)$$

corresponding to the eigenvalue $\bar{\lambda}$.

Next, define $x^* := \frac{1}{y_0^*} z^*$. Then it follows from (4.12) that $x^{*T} B x^* - 2b^T x^* \leq s$ and from the definition of $y(t)$ that $(A - \bar{\lambda} B)x^* = a - \bar{\lambda} b$. By Lemma 3.4, there exists $v \in \text{Null}(A - \bar{\lambda} B)$ with $v^T B v > 0$, and thus $x^* + \alpha v$ would solve (4.1), for some suitable $\alpha > 0$.

Case 4. $t^* = t_0 = \bar{t}$. In this case, from the definitions of t_0 and \bar{t} , we have $\bar{\lambda}$ is finite and the infimum of d is attained at the right end point of the interval $(\underline{\lambda}, \bar{\lambda})$, i.e., at $\bar{\lambda}$. From the definition of d , this implies in particular that $a - \bar{\lambda} b \in \text{Range}(A - \bar{\lambda} B)$. Furthermore,

$$0 \geq d'(\bar{\lambda}) = 1 + \psi(\bar{\lambda}).$$

Hence, using the fact that $s + 1 > 0$, we see that

$$x^{*T} B x^* - 2b^T x^* \leq -1 < s,$$

where $x^* := x(\bar{\lambda})$. Finally, recall from Lemma 3.4 that there exists $v \in \text{Null}(A - \bar{\lambda}B)$ with $v^T B v > 0$. Combining these few facts, we conclude that $x^* + \alpha v$ solves (4.1), for some suitable $\alpha > 0$.

Remark 4.1. Uniqueness of maximizer of k . *In passing, we remark that $k(t)$ must have a unique maximizer. First of all, it is easy to see that the four cases analyzed above are mutually exclusive for a maximizer t^* of $k(t)$. If case 4 happens, then clearly the maximizer is unique and equals $t_0 = \bar{t}$. Otherwise, in all other three cases, we see that for each maximizer t^* , there corresponds at least one dual solution λ^* of the GTRS (4.1). Such λ^* constructed in those cases are different for different t^* , thanks to the strict monotonicity of $\mu(t)$ on (t_0, \bar{t}) . Since the GTRS (4.1) must have a unique dual optimal solution in view of the analysis in Section 3.1.1 and our assumption that ψ is strictly increasing on $\text{cl}(\underline{\lambda}, \bar{\lambda})$, we conclude that $k(t)$ must have a unique maximizer.*

The above arguments for $s + 1 < 0$ are completely analogous. Note that we need to scale by $\sqrt{-(s + 1)}$ instead of $\sqrt{s + 1}$ in the constraint of the minimization problem in (4.5). Hence $y(t)^T \begin{bmatrix} 1 & -b^T \\ -b & B \end{bmatrix} y(t) = -1$ and moreover $k'(t) = -(s + 1)y_0^2(t) - 1$.

In this section, we just gave a complete analysis on how a solution of (4.1) can be recovered after maximizing $k(t)$. Comparing this with the procedures outlined in Section 3, we see that if the problem (4.1) is obtained from those procedures, then $\lambda^* \in (\underline{\lambda}, \bar{\lambda})$. Thus, we must be in Case 1 and hence $k(t)$ has to be differentiable at t^* . On the other hand, it is easy to see that closed form solutions are obtained for the other cases using the procedures in Section 3.

4.3 Implementation details

In this subsection, we discuss implementation details of our algorithm. For simplicity, we only consider the case when B is nonsingular. In this case, by translating the optimization variable by $B^{-1}b$ if necessary, we may further assume without loss of generality that $b = 0$. Moreover, we only need to consider the cases when B is positive definite or indefinite. Furthermore, we still assume Assumption 2.1, the regularity (3.1) and $s + 1 \neq 0$. To explicitly guarantee that (3.5) never holds, we also assume $a \neq 0$.

When B is positive definite, $\underline{\lambda}$ is negative infinity and $\bar{\lambda}$ is the minimum generalized eigenvalue of the matrix pair (A, B) , which can be computed efficiently. On the other hand, when B is indefinite, the interval $(\underline{\lambda}, \bar{\lambda})$ cannot be easily determined in general. Thus, in this case, we consider the further subcase that A is positive definite. Then we know that $0 \in (\underline{\lambda}, \bar{\lambda})$. Observe that for any $\lambda \in [\underline{\lambda}, 0)$, we have

$$A - \lambda B = -\lambda \left(B - \frac{1}{\lambda} A \right). \quad (4.14)$$

Since $\lambda \in [\underline{\lambda}, 0)$ if, and only if, $\frac{1}{\lambda} \in (-\infty, \frac{1}{\underline{\lambda}}]$, we conclude from (4.14) that $\underline{\lambda} = \frac{1}{\bar{\rho}_1}$, where $\bar{\rho}_1$ is the minimum generalized eigenvalue of the matrix pair (B, A) . Similarly, one can show that $\bar{\lambda} = -\frac{1}{\bar{\rho}_2}$, with $\bar{\rho}_2$ being the minimum generalized eigenvalue of the matrix pair $(-B, A)$. The quantities $\bar{\rho}_1$ and $\bar{\rho}_2$ can both be computed efficiently.

Case check. We first carry out the case check as described in Section 3.1.1. We obtain an explicit solution when the GTRS (1.1) has an interior solution $A^{-1}a$ or when the GTRS instance is a hard case, case 2 instance.

Shift and deflation for the hard case, case 1, when B is positive definite. After the case check, the bound side has been determined and so we focus on the equality constrained case $x^T Bx = s$. In the next proposition, we describe a shift and deflation technique that transforms a GTRS instance from hard case, case 1, to the easy case, when B is positive definite.

Proposition 4.2. *Suppose that we consider the GTRS with the equality constraint $x^T Bx = s$ with $B \succ 0$. Let $\bar{\lambda}$ be the smallest generalized eigenvalue of the matrix pair (A, B) and suppose that a corresponding nonzero eigenvector v satisfies $a^T v = 0$. Furthermore, let $\alpha > 0$. Then*

$$\begin{aligned} & (x^*, \lambda^*) \text{ solves GTRS} \\ & \text{if, and only if,} \\ & (x^*, \lambda^*) \text{ solves GTRS with } A \text{ replaced by } A - \bar{\lambda}B + \alpha(Bv)(Bv)^T. \end{aligned}$$

Proof. First, since $x^T Bx = s$, it is clear that (x^*, λ^*) solves GTRS if, and only if, (x^*, λ^*) solves GTRS with A replaced by $A - \bar{\lambda}B$. Next, consider the GTRS with A replaced by $A - \bar{\lambda}B$. Let $y = B^{\frac{1}{2}}x$ and use the substitution $x = B^{-\frac{1}{2}}y$. This results in the following TRS:

$$\begin{aligned} \inf \quad & y^T \bar{A}y - 2\bar{a}^T y \\ \text{s.t.} \quad & y^T y = s, \end{aligned}$$

where $\bar{A} = B^{-\frac{1}{2}}(A - \bar{\lambda}B)B^{-\frac{1}{2}}$ and $\bar{a} = B^{-\frac{1}{2}}a$.

We now apply the shift and deflate lemma in [11] stated for the above TRS and obtain that

$$\begin{aligned} & (y^*, \lambda^*) \text{ solves TRS} \\ & \text{if, and only if,} \\ & (y^*, \lambda^*) \text{ solves TRS when } \bar{A} \text{ is replaced by } \bar{A} + \alpha w w^T, \end{aligned}$$

where $w = B^{\frac{1}{2}}v$ is the corresponding eigenvector and hence satisfies $w^T \bar{a} = 0$ by assumption and the definition of \bar{a} . After substituting again using $y = B^{\frac{1}{2}}x$, we see that

$$y^T \left(B^{-\frac{1}{2}}(A - \bar{\lambda}B)B^{-\frac{1}{2}} + \alpha w w^T \right) y = x^T (A - \bar{\lambda}B + \alpha(Bv)(Bv)^T) x.$$

The linear term and constraint follow similarly. □

Initialization. After the case check and the possible shift and deflation, we proceed to solve the equality constrained case by the RW algorithm. We discuss initialization in this subsection.

We first consider the case when B is positive definite. In this case, $k(t)$ is well-defined for all t by Theorem 4.2. Let t^* be the optimal solution of $\max k(t)$. As in [29], we derive an interval that contains t^* in the next proposition for initializing the RW algorithm.

Proposition 4.3. *Suppose that $\lambda^* < \bar{\lambda}$. Then*

$$\bar{\lambda} - \sqrt{\frac{a^T B^{-1} a}{s}} \leq t^* \leq \bar{\lambda} + \sqrt{s a^T B^{-1} a}. \quad (4.15)$$

Proof. The assumption $\lambda^* < \bar{\lambda}$ implies that we can recover a solution of GTRS from the maximizer of $k(t)$.

Let x^* be the solution of GTRS thus recovered. Then $\begin{bmatrix} 1 \\ x^* \end{bmatrix}$ is a generalized eigenvector of the matrix pair

$$\left(\begin{bmatrix} t^* & -a^T \\ -a & A \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & B \end{bmatrix} \right)$$

with generalized eigenvalue λ^* . Hence, in particular,

$$t^* - a^T x^* = \lambda^* \quad \Rightarrow \quad t^* = \lambda^* + a^T x^* \leq \bar{\lambda} + \sqrt{x^{*T} B x^*} \sqrt{a^T B^{-1} a} = \bar{\lambda} + \sqrt{s a^T B^{-1} a}.$$

This proves the upper bound in (4.15). We next derive the lower bound. In this case, we define $\delta = \bar{\lambda} - \lambda^* > 0$. Then we have

$$A - \lambda^* B \succeq \delta B \succ 0 \tag{4.16}$$

and hence

$$t^* - \lambda^* = a^T x^* = x^{*T} (A - \lambda^* B) x^* \geq \delta x^{*T} B x^* = \delta s. \tag{4.17}$$

In addition, from the definition of $d(\lambda)$, λ^* and (4.16), we have

$$t^* - \lambda^* = a^T (A - \lambda^* B)^{-1} a \leq \frac{1}{\delta} a^T B^{-1} a. \tag{4.18}$$

Combining (4.17) and (4.18), we see that

$$\delta \leq \sqrt{\frac{a^T B^{-1} a}{s}}.$$

Finally, we observe from (4.18) that $t^* \geq \lambda^*$ and hence $\delta \geq \bar{\lambda} - t^*$. The lower bound in (4.15) now immediately follows. \square

On the other hand, when B is indefinite, according to Theorem 4.1 and Corollary 4.2, we conclude that t_0 is finite and the function $k(t)$ would have t_0 as a left end point of its domain. We currently do not know of any efficient way for computing/estimating t_0 . However, if t_0 and the corresponding λ_0 such that $d(\lambda_0) = t_0$ are *known*, then the function value and the derivative of k at any $t > t_0$ can be efficiently computed; see the paragraph below Proposition 4.4, with $\mu = \lambda_0$ in (4.23). Furthermore, for initialization, one can find an interval containing t^* as follows:

Procedure for finding an interval containing t^ when B is indefinite.*

Step 1. Take $\epsilon > 0$ and compute $k'(t_0 + \epsilon)$.

Step 2. If $k'(t_0 + \epsilon) > 0$, set $t_+ = t_0 + \epsilon$. Otherwise, set $t_- = t_0 + \epsilon$ and $i = 1$.

Step 3. While $k'(t_-) < 0$, update $t_+ \leftarrow t_-$ and $t_- \leftarrow t_0 + \zeta^i \epsilon$ for some fixed $\zeta \in (0, 1)$. Update $i \leftarrow i + 1$. Repeat this step.

Step 4. While $k'(t_+) > 0$, update $t_- \leftarrow t_+$ and $t_+ \leftarrow \kappa(t_+ - \lambda_0) + \lambda_0$ for some fixed $\kappa > 1$. Repeat this step.

Step 5. By construction, $k'(t_-) > 0$ and $k'(t_+) < 0$. Thus, $t^* \in (t_-, t_+)$.

Computing $k_0(t)$ and $k'_0(t)$. We next discuss how $k_0(t)$ and $k'_0(t)$ for $t > t_0$, when they exist, can be computed efficiently. Recall from the definition of $k_0(t)$ and (4.8) that

$$k_0(t) = \inf \left\{ u^T \begin{bmatrix} t & -a^T \\ -a & A \end{bmatrix} u : u^T \begin{bmatrix} 1 & 0 \\ 0 & B \end{bmatrix} u = s + 1 \right\}, \quad k'_0(t) = u_0^2(t),$$

where $u_0(t) \geq 0$ is the first coordinate of a vector $u(t)$ (unique when $u_0(t) > 0$) attaining the infimum defining $k_0(t)$. In the case when B is positive definite, $k_0(t)$ can be computed from the minimum generalized eigenvalue of the matrix pair

$$\left(\begin{bmatrix} t & -a^T \\ -a & A \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & B \end{bmatrix} \right),$$

and $u_0(t)$ is obtained from the corresponding eigenvector. However, when B is not positive definite, $k_0(t)$ and its derivative cannot be obtained directly from an extremal generalized eigenpair. We now discuss how to compute k_0 and its derivative in this case.

To this end, we first consider a closely related problem. Let C and D be symmetric matrices. Consider the following program.

$$\begin{aligned} \text{val} &:= \inf && x^T C x \\ &\text{s.t.} && x^T D x = 1. \end{aligned} \tag{4.19}$$

Then we have the following result.

Proposition 4.4. *Suppose that (4.19) is feasible and that there exists λ so that $C - \lambda D \succeq 0$. Suppose further that $\text{val} \neq 0$.*

(i) *If $\text{val} > 0$ and if x^* is a solution to (4.19), then $\frac{1}{\sqrt{\text{val}}}x^*$ is a solution to the following optimization problem, whose optimal value is $-\frac{1}{\text{val}}$:*

$$\begin{aligned} \inf &&& -x^T D x \\ \text{s.t.} &&& x^T C x = 1. \end{aligned} \tag{4.20}$$

(ii) *If $\text{val} < 0$ and if x^* is a solution to (4.19), then $\frac{1}{\sqrt{-\text{val}}}x^*$ is a solution to the following optimization problem, whose optimal value is $-\frac{1}{\text{val}}$:*

$$\begin{aligned} \inf &&& x^T D x \\ \text{s.t.} &&& -x^T C x = 1. \end{aligned} \tag{4.21}$$

Proof. By Theorem 2.3, we see that x^* is an optimal solution to (4.19) if, and only if, there exists λ^* such that

$$\begin{aligned} (C - \lambda^* D)x^* &= 0, \\ C - \lambda^* D &\succeq 0, \\ x^{*T} D x^* &= 1. \end{aligned} \tag{4.22}$$

Using the first and third relations of (4.22), we see that

$$\text{val} = x^{*T} C x^* = \lambda^* x^{*T} D x^* = \lambda^*.$$

Since $\text{val} \neq 0$, we obtain that $\lambda^* = \text{val} \neq 0$. For part (i), we have $\lambda^* > 0$. From (4.22) and the relation $\text{val} = x^{*T} C x^*$, we see that

$$\begin{aligned} \left(\frac{1}{\lambda^*} C - D \right) \frac{x^*}{\sqrt{\lambda^*}} &= 0, \\ \frac{1}{\lambda^*} C - D &\succeq 0, \\ \left(\frac{x^*}{\sqrt{\lambda^*}} \right)^T C \left(\frac{x^*}{\sqrt{\lambda^*}} \right) &= \frac{\text{val}}{\lambda^*} = 1. \end{aligned}$$

Hence, $\frac{x^*}{\sqrt{\lambda^*}} = \frac{x^*}{\sqrt{\text{val}}}$ solves (4.20). On the other hand, for part (ii), $\lambda^* < 0$. Hence, we have

$$\begin{aligned} \left(-\frac{1}{\lambda^*} C + D \right) \frac{x^*}{\sqrt{-\lambda^*}} &= 0, \\ -\frac{1}{\lambda^*} C + D &\succeq 0, \\ \left(\frac{x^*}{\sqrt{-\lambda^*}} \right)^T C \left(\frac{x^*}{\sqrt{-\lambda^*}} \right) &= \frac{\text{val}}{-\lambda^*} = -1. \end{aligned}$$

Thus, $\frac{x^*}{\sqrt{-\lambda^*}} = \frac{x^*}{\sqrt{-\text{val}}}$ solves (4.21). □

We now complete our discussion on efficient computation of $k_0(t)$ and its derivative when B is indefinite. Suppose that we know a μ such that

$$\begin{bmatrix} t & -a^T \\ -a & A \end{bmatrix} - \mu \begin{bmatrix} 1 & 0 \\ 0 & B \end{bmatrix} \succ 0. \quad (4.23)$$

Since $\begin{bmatrix} 1 & 0 \\ 0 & B \end{bmatrix}$ is indefinite, we see immediately that

$$\inf \left\{ -\text{sign}(s+1)y^T \begin{bmatrix} 1 & 0 \\ 0 & B \end{bmatrix} y : y^T \left(\begin{bmatrix} t & -a^T \\ -a & A \end{bmatrix} - \mu \begin{bmatrix} 1 & 0 \\ 0 & B \end{bmatrix} \right) y = 1 \right\} < 0. \quad (4.24)$$

Thus, in view of Proposition 4.4, we see that $k_0(t)$ and $k'_0(t)$, upon scaling, can be computed by considering the optimization problem (4.24). Since the quadratic objective in the constraint of (4.24) is positive definite, the optimal value and solution of (4.24) can be obtained from an extremal generalized eigenpair.

Techniques for maximizing $k(t)$. These techniques are adapted from the original Rendl-Wolkowicz algorithm [11] for solving TRS (1.2).

Vertical Cut

Suppose there exist t_g with $k'(t_g) < 0$ and t_b with $k'(t_b) > 0$. Suppose further that $k(t_g) < k(t_b)$. Then we can use the so-called vertical cut to reduce the interval that contains t^* . More precisely, we obtain our new approximation t_+ by intersecting the tangent line of k at t_g with the horizontal line $k = k(t_b)$. Theoretically, we always get $t_+ \in [t_b, t_g]$ and that $k'(t_+) < 0$. However, approximate evaluation of slopes and function values of k can cause $t_+ \notin [t_b, t_g]$. When this happens, the vertical cut fails. The case when $k(t_g) > k(t_b)$ can be treated similarly.

Triangle Interpolation

Suppose there exist t_g with $k'(t_g) < 0$ and t_b with $k'(t_b) > 0$. We then intersect the tangent lines of $k(t)$ at t_g and t_b to obtain a new estimate t_+ . Theoretically, we always get $t_+ \in [t_b, t_g]$. However, since we only compute the slope and the function value of k approximately, it can happen that $t_+ \notin [t_b, t_g]$. When this happens, the triangle interpolation fails.

Inverse Linear Interpolation

Notice that $k'(t) = |s+1|y_0^2(t) - 1$, with $y_0(t)$ defined as in Section 4.2. Thus, $k'(t) = 0$ is equivalent to

$$\phi(t) := \sqrt{|s+1|} - \frac{1}{y_0(t)} = 0.$$

We consider approximating the inverse function of $\phi(t)$ by a linear function, i.e.,

$$t(\phi) = a\phi + b,$$

for some real numbers a, b , and then set our next approximation as $t_+ = t(0)$ if $t(0) \in [t_b, t_g]$. This technique is different from the previous techniques in the sense that it can also operate on two points whose slopes have the same sign.

Primal step to feasibility. As in the RW algorithm for solving TRS, we keep track of the point $x(t) := \frac{1}{y_0(t)}z(t)$, where $y(t) = \begin{bmatrix} y_0(t) \\ z(t) \end{bmatrix}$ is defined as in Section 4.2. In the easy case/hard case, case 1, $y_0(t)$ is nonzero as the algorithm proceeds and one can show that $x(t)$ converges to the optimal solution of GTRS (1.1) as t converges to t^* . Notice that if there exist t_g with $k'(t_g) < 0$ and t_b with $k'(t_b) > 0$, then $(x(t_g)^T Bx(t_g) - s)(x(t_b)^T Bx(t_b) - s) < 0$. Thus, by choosing a suitable α , the point $\alpha x(t_g) + (1 - \alpha)x(t_b)$ will be primal feasible. Furthermore, the resulting sequence is still convergent to $x(t^*)$.

Before closing this section, we summarize our algorithm in the following flowchart.

Flowchart for the extended RW algorithm

- Check for the hard cases/interior solution; shift and deflate; find initial interval containing t^* .
- Main loop: iterate until a termination criterion is met:
 1. update t .
 - (a) Set t to be the midpoint of the interval of uncertainty for t .
 - (b) If points at which k has positive and negative slope, respectively, are known:
 - i. Do vertical cut; reduce the interval of uncertainty for t if possible. Update t if vertical cut is successful.
 - ii. Do triangle interpolation. Update t if triangle interpolation is successful.
 - (c) Do inverse linear interpolation. Update t if inverse linear interpolation is successful.
 2. At the new t , compute $k(t)$ and $k'(t)$ using techniques discussed above and `eigifp` [14].³ Take a primal step to satisfy feasibility if iterates at which k has positive and negative slopes are available.
- End loop.

5 Numerical experiments with new RW algorithm

In this section, we study the numerical performance of our new RW algorithm for GTRS. We compute $\lim_{\lambda \uparrow \bar{\lambda}} \psi(\lambda)$ and $\lim_{\lambda \downarrow \underline{\lambda}} \psi(\lambda)$ as described in Section 3.1.1 with $\alpha = \frac{\|A\|_2}{\sqrt{n}}$, where $\|A\|_2$ is the (approximate) spectral norm found using the MATLAB command `normest(A, 1)`. In particular, we check for $\frac{|a^T v|}{\|a\|} < 1e-8$ for every vector v from an orthonormal basis of the nullspaces $\text{Null}(A - \bar{\lambda}B)$ or $\text{Null}(A - \underline{\lambda}B)$. If the instance is not a hard case, case 2, instance, we perform shift and deflation as described in Proposition 4.2 using these vectors v and the same α as above, when $B \succ 0$. We then initialize the RW algorithm (for the equivalent equality constrained problem) as discussed in the previous section, and terminate the algorithm when

$$\max \left\{ \frac{|k'(t)|^2}{|s+1|^2}, \frac{|q_0(x) - k(t)|}{|q_0^{\text{best}} + 1|}, \frac{|q_1(x) - s|}{|s+1|}, \frac{\|Ax - \lambda Bx - a\|^2}{(\|A\|_2 + \|a\| + 1)^2} \right\} < 10^{-13}, \quad (5.1)$$

or

$$\frac{|\text{high} - \text{low}|}{|\text{high}| + |\text{low}|} < 10^{-15},$$

or when the number of iterations hits 30; here x is recovered from a generalized eigenvector and then shifted to become approximately feasible, $\|A\|_2$ is the (approximate) spectral norm found using the MATLAB

³ EIGIFP is implemented in MATLAB and finds the minimum generalized eigenvalue and eigenvector of the matrix pair (A, B) for positive definite B . It does not involve factorization of the matrices A or B .

command `normest(A, 1)`, q_0^{best} is the smallest primal objective value up to the current iteration. Our code is written in MATLAB. All numerical experiments are performed on an SGI XE340 system, with two 2.4 GHz quad-core Intel E5620 Xeon 64-bit CPUs and 48 GB RAM, equipped with SUSE Linux Enterprise server 11 SP1 and MATLAB 7.14 (R2012a). All routines are timed using the tic-toc function in MATLAB.

5.1 B is positive definite

In this subsection, we consider GTRS (1.1) with a positive definite B . For simplicity, we also assume without loss of generality that $b = 0$. We consider two cases: A is not positive semidefinite or A is positive definite.

A is not positive semidefinite. In this case, the GTRS is equivalent to (4.1) with $s = u$, which is also equivalent to (1.1) with $l = 0$. We compare three algorithms:

- The RW algorithm for solving (1.1);
- The GLTR algorithm [15], an algorithm in the Galahad package (Version 2.50000 pre-release), to solve (1.1) with $l = 0$. The GLTR algorithm is written in Fortran and implemented as a MATLAB mex file;
- The Newton’s method with (Armijo) line search for maximizing the dual objective function to solve (4.1) with $s = u$. We code this in MATLAB.

For the GLTR algorithm, we use its default settings in our experiments. For the Newton’s method, we initialize at $\lambda^0 = \bar{\lambda} - 1$ and terminate when

$$\max \left\{ \frac{|q_0(x) - d(\lambda)|}{|q_0(x)| + 1}, \frac{|q_1(x) - s|}{|s| + 1}, \frac{\|Ax - \lambda Bx - a\|^2}{(\|A\|_2 + \|a\| + 1)^2} \right\} < 10^{-12},$$

or when the stepsize falls below $1e - 10$ or the number of iterations hits 10; here x is generated from λ via $x = (A - \lambda B)^{-1}a$. The Newton direction in the Newton’s method is computed via the MATLAB built-in function `pcg`, with termination tolerance $1e - 14$ and iteration bound 1000.

In our tests, we generate both easy and hard case instances, for dimensions $n = 10000, 15000$ and 20000 .

To generate an easy case instance, we first generate randomly a sparse symmetric matrix A and a sparse positive definite matrix B via `A = sprandsym(n, 1e-2)` and `B = sprandsym(n, 1e-2, 0.1, 2)`. We then compute $\bar{\lambda}$, the minimum generalized eigenvalue of the matrix pair (A, B) . We next generate x_0 with Gaussian entries of mean 0 and standard deviation 0.1 and set $a = (A - \tilde{\lambda}B)x_0$, $s = x_0^T Bx_0$, where $\tilde{\lambda} = \bar{\lambda} - r$ for some r chosen uniformly from $[5, 10]$. Finally, we set $u = 1.2s$ and $l = 0.8s$. From the optimality conditions and the fact that $r > 0$, we see that the above construction likely gives an easy case instance.

To generate a hard case instance, we follow the same procedure as above to obtain $A, B, x_0, \bar{\lambda}$ and s but we set $a = (A - \bar{\lambda}B)x_0$ instead. If we take $u = 0.6s$ and $l = 0.6u$, then we likely end up with a hard case, case 1 instance. On the other hand, by setting $u = 1.2s$, $l = 1.1s$, we likely end up with a hard case, case 2 instance.

For each n , we generate 10 easy, 10 hard case, case 1 and 10 hard case, case 2 instances as described above. The computational results are reported in Table 2, where we report the number of iterations (iter), CPU time in seconds (cpu), primal objective value (fval) at termination and also the primal infeasibility for the equivalent equality constrained problem measured by $|x^T Bx - u|$ (feas_{eq}),⁴ averaged over the 10 random instances. We observe the RW algorithm is faster than Newton+Armijo on easy case instances, but is slower on hard case, case 1 instances because of the extra time taken for preprocessing. The preprocessing is justified by the observation that, the simple Newton+Armijo strategy fails to obtain even two digits of

⁴Comparing with $|x^T Bx - \max\{\min\{x^T Bx, u\}, l\}|$, feas_{eq} is less sensitive to the quantity $|u - l|$, the “thickness” of the feasible region.

accuracy for hard case, case 2 instances. We also note that the GLTR algorithm is the slowest algorithm mainly because of the cost for factorizing the matrix B , whose Cholesky factorization has a high percentage fill-in.⁵

	n	RW	GLTR	Newton+Armijo
		iter/cpu/fval/feas _{eq}	iter/cpu/fval/feas _{eq}	iter/cpu/fval/feas _{eq}
Easy Case	10000	6/4.83/-3.889344396e+04/5.5e-13	7/188.75/-3.889344396e+04/1.3e-09	9/6.82/-3.889344397e+04/5.6e-07
	15000	6/10.91/-8.101943616e+04/6.4e-13	7/668.40/-8.101943616e+04/4.0e-09	9/15.09/-8.101943616e+04/4.1e-11
	20000	6/20.33/-1.573159441e+05/1.1e-12	7/1654.31/-1.573159441e+05/3.2e-09	9/27.57/-1.573159441e+05/1.8e-09
Hard Case 1	10000	7/18.68/-3.336135050e+03/2.0e-13	23/196.09/-3.336135050e+03/1.8e-09	7/8.25/-3.336135050e+03/9.4e-08
	15000	7/45.54/-6.039038487e+03/5.9e-13	24/678.88/-6.039038487e+03/1.1e-09	7/20.93/-6.039038488e+03/6.2e-07
	20000	7/100.60/-9.140937609e+03/1.8e-13	24/1753.84/-9.140937609e+03/1.3e-09	6/36.05/-9.140937609e+03/4.5e-13
Hard Case 2	10000	0/10.26/-4.724115040e+03/4.5e-13	211/229.18/-4.723248192e+03/2.2e-04	10/38.48/-4.293815820e+03/4.6e+02
	15000	0/25.06/-8.484568849e+03/1.2e-12	249/812.57/-8.482991578e+03/1.5e-05	10/98.79/-7.712953798e+03/1.0e+03
	20000	0/55.02/-1.290858124e+04/9.1e-13	282/2108.92/-1.290723940e+04/1.0e-02	10/214.12/-1.173381072e+04/1.7e+03

Table 2: Computational results for indefinite A and positive definite B .

A is positive definite. In this case, it is not certain whether the GTRS (1.1) is equivalent to (4.1) with $s = u$ or $s = l$, or the GTRS may have an interior solution. Hence, for comparison, we first solve the GTRS using the RW algorithm, from which we can determine whether the GTRS has a boundary solution. If the GTRS is equivalent to an instance of (4.1), we solve this instance of (4.1) using the Newton’s method that maximizes its dual. Furthermore, if the GTRS is equivalent to (4.1) with $s = u$, then the GLTR algorithm can be applied directly to solve the instance. On the other hand, even if the GTRS turns out to be equivalent to (4.1) with $s = l$ instead, it is not hard to see that the instance can be solved by applying the GLTR algorithm with $A \leftarrow A - (\bar{\lambda} + 1)B$, $u \leftarrow l$ and $l \leftarrow 0$. We initialize and terminate all three algorithms as in the previous test.

In our tests, similarly as above, we generate both easy and hard case instances, for dimensions $n = 10000$, 15000 and 20000. We follow the same procedure as in the previous case, except that instead of a random sparse symmetric A , we generate another matrix C similarly as we generated B , and then set $A = C + 10B$ so that the resulting GTRS (1.1) likely has a solution on the lower boundary of the feasible region.

Test results are reported in Table 3, averaged over 10 random instances.⁶ We note that the CPU time for GLTR+shift includes both the GLTR algorithm run time and the time taken to find $\bar{\lambda}$ for the transformation $A \leftarrow A - (\bar{\lambda} + 1)B$, with the latter time also reported separately in parenthesis. We again observe that the RW algorithm is faster than Newton+Armijo on easy case instances, and is slower on hard case, case 1 instances due to the extra preprocessing time. The failure of the simple Newton+Armijo strategy on hard case, case 2 instances again justifies the use of preprocessing. We also note that the GLTR algorithm is the slowest due to the cost for factorizing B .

5.2 B is indefinite

In this subsection, we consider GTRS (1.1) with an indefinite (and nonsingular) B . For simplicity, we also assume without loss of generality that $b = 0$. Furthermore, in order that the interval $(\underline{\lambda}, \bar{\lambda})$ can be located efficiently, we restrict our attention to the case when A is positive definite; see Section 4.3. There are

⁵Indeed, when B is structured, say, block diagonal with small blocks (size 100 by 100), the GLTR algorithm tends to terminate within 1 second for the easy case and hard case 1 instances, with high accuracy; and within a couple seconds (2 to 10) for hard case 2 instances, with moderate accuracy. In contrast, the other two codes are at most 50% faster than the time reported in Table 2 for such B .

⁶For consistency, we regenerate the problem instance to make sure that the instances considered in this subsection have minimizer on the lower boundary.

	n	RW	GLTR+shift	Newton+Armijo
		iter/cpu/fval/feas _{eq}	iter/cpu/fval/feas _{eq}	iter/cpu/fval/feas _{eq}
Easy Case	10000	6/3.26/-1.049837986e+04/1.4e-13	6/188.62 (1.32)/-1.049837986e+04/8.8e-10	9/5.76/-1.049837986e+04/2.1e-09
	15000	6/6.98/-1.466449774e+04/5.5e-13	6/672.60 (3.08)/-1.466449774e+04/2.8e-09	10/12.71/-1.466449774e+04/4.3e-07
	20000	6/13.21/-4.406550240e+04/6.4e-13	6/1742.12 (5.97)/-4.406550240e+04/3.7e-09	10/24.45/-4.406550240e+04/1.4e-10
Hard Case 1	10000	7/12.00/7.464921449e+03/1.8e-13	20/189.43 (1.42)/7.464921449e+03/8.6e-10	8/6.59/7.464921449e+03/7.1e-09
	15000	7/26.06/1.661401688e+04/2.3e-13	19/677.80 (2.95)/1.661401688e+04/1.3e-09	7/15.32/1.661401688e+04/1.8e-09
	20000	7/55.61/2.919616156e+04/3.2e-13	18/1752.79 (6.25)/2.919616156e+04/1.9e-09	8/34.74/2.919616156e+04/1.5e-08
Hard Case 2	10000	0/6.24/2.496011798e+04/1.8e-13	120/209.60 (1.46)/2.496177644e+04/6.0e-08	10/22.33/2.256793156e+04/2.3e+02
	15000	0/11.72/5.437098121e+04/7.3e-13	95/768.72 (2.80)/5.437462396e+04/6.0e-08	10/46.41/4.918446583e+04/5.0e+02
	20000	0/21.34/9.623939924e+04/5.5e-13	93/1807.07 (5.73)/9.624434210e+04/6.9e-08	10/92.46/8.710211303e+04/8.7e+02

Table 3: Computational results for positive definite A and positive definite B .

currently no algorithms in the literature specialized at solving such GTRS instances. Thus, we only compare our algorithm with the Newton’s method. Specifically, we first use our algorithm to determine whether the GTRS is equivalent to an instance of (4.1) with $s = u$ or $s = l$. We then solve this instance of (4.1) using the Newton’s method that maximizes its dual.

For simplicity, we only generate easy case and hard case, case 2, instances in our numerical experiments. For dimensions $n = 10000, 15000$ and 20000 , we first generate sparse symmetric matrices B_1 and A_1 via the commands `A_1 = sprandsym(n, 1e-2, 0.1, 2)` and `B_1 = sprandsym(n, 1e-2)`. We then locate the interval $(\underline{\lambda}, \bar{\lambda})$ for the matrix pair (A_1, B_1) .

To generate an easy case instance, we first set $\tilde{\lambda} = \frac{\bar{\lambda} + \underline{\lambda}}{2}$. We next generate x_0 with Gaussian entries of mean 0 and standard deviation 0.1, and set $a = (A_1 - \tilde{\lambda}B_1)x_0$ and $s_1 = x_0^T B_1 x_0$. We now further modify A_1 and B_1 so that we obtain an instance with explicit knowledge of t_0 and the corresponding $\lambda_0 \in (\underline{\lambda}, \bar{\lambda})$ with $d(\lambda_0) = t_0$. To this end, define $B := B_1 / (-|s_1|)$ and $A := A_1$. Then by construction, we have

$$\inf_{x^T B x = -1} x^T A x - 2a^T x = x_0^T A x_0 - 2a^T x_0 = -d(\lambda_0),$$

where $\lambda_0 = -|s_1| \tilde{\lambda}$. Finally, with the above A , a and B , setting $s = \alpha \cdot \sqrt{n}$ and $l = u - 1 = s$, where α follows standard Gaussian distribution, we likely obtain an easy case instance with explicitly known $t_0 = d(\lambda_0)$. The t_0 can be used for initialization of the RW algorithm, and λ_0 is used as in (4.23) in place of μ . In particular, we find t_- and t_+ as described in Section 4.3 with $\epsilon = 1$, $\zeta = 0.2$ and $\kappa = 1.2$. We note that computation of $k'(t_-)$ outlined in Section 4.3 can become inefficient when t_- gets too close to t_0 .⁷

Generating a hard case, case 2 instance with known t_0 is a bit more complicated. We first describe how to generate such an instance so that the optimal solution x_* satisfies $x_*^T B x_* = l$. To this end, we set $\tilde{\lambda} = \underline{\lambda}$. We then generate x_0 with Gaussian entries of mean 0 and standard deviation 0.1, and set $a = (A_1 - \tilde{\lambda}B_1)x_0$ and $s_1 = x_0^T B_1 x_0$. We now further modify A_1 and B_1 as in the previous paragraph to obtain A , B , t_0 and λ_0 . Finally, with these A , a and B , setting $s = \alpha \cdot \sqrt{n}$ and $l = u - 1 = s$, where α is uniformly chosen from $[1, 2]$, we likely obtain a hard case, case 2 instance with optimal solution x_* satisfies $x_*^T B x_* = l$. Next, to obtain a hard case, case 2 instance with optimal solution x_* satisfies $x_*^T B x_* = u$, we follow the same procedure as above except that we set $\tilde{\lambda} = \bar{\lambda}$ and choose α uniformly from $[-2, -1]$.

Test results are reported in Table 4, averaged over 10 random instances.⁸ We observe that the RW algorithm is slightly faster than Newton+Armijo on easy case instances, and it produces a solution with better quality in terms of feasibility (of the equivalent equality constrained GTRS). Furthermore, we again observe that the simple Newton+Armijo strategy fails on hard case, case 2 instances.

⁷In all our test instances below, they all turned out to have a valid left end point t_- lying in $[t_0 + (0.2)^3, t_0 + 1]$.

⁸We regenerate another instance if $A^{-1}b$ is an interior solution.

	n	RW	Newton+Armijo
		iter/cpu/fval/feas _{eq}	iter/cpu/fval/feas _{eq}
Easy Case	10000	6/9.75/-2.034919097e+03/1.1e-14	6/9.79/-2.034919097e+03/1.7e-08
	15000	7/24.17/-3.385643481e+03/1.4e-14	6/24.40/-3.385643481e+03/8.9e-12
	20000	7/47.29/-7.195068138e+03/1.1e-14	6/50.18/-7.195068132e+03/1.7e-07
Hard Case 2	10000	0/13.34/-6.405913926e+02/1.7e-14	10/41.62/-2.270662258e+03/1.4e+02
	15000	0/30.71/-4.882640374e+02/2.3e-14	10/97.86/-4.974215533e+03/1.8e+02
	20000	0/63.55/-9.857960694e+02/2.6e-14	10/213.46/-8.637983865e+03/1.8e+02

Table 4: Computational results for positive definite A and indefinite B .

6 Conclusion

We have presented optimality conditions for GTRS (1.1) that hold under a constraint qualification, the RICQ (2.6). Furthermore, if the RICQ fails, we showed how the GTRS can be explicitly solved. In this sense, GTRS has strong duality results as in *linear programming*. We also demonstrated that the GTRS can be classified into easy and hard cases as the classical TRS, and the problem can be preprocessed to identify explicit solution in hard case/interior solution and reduce the instance into an equality constrained GTRS (4.1). We then discussed in detail how the Rendl-Wolkowicz algorithm in [29] can be extended to solve such instances, which involved transforming the instance into a parameterized *generalized eigenvalue problem*. We also illustrated our algorithm numerically in various cases, including cases when B is indefinite. When B is a random sparse positive definite matrix, our algorithm is competitive with a simple Newton's method implementation, and is faster than the GLTR algorithm. On the other hand, when B is indefinite, our algorithm requires additional inputs for initialization. Given such inputs, our algorithm is also competitive with the Newton's method.

A Proof of Theorem 2.1(ii)

Proposition A.1. *Suppose that $b = 0$ and Items 1, 2 and 3 of Assumption 2.1 are satisfied. If GTRS (1.1) is bounded below, then D-GTRS (2.2) is feasible.*

Proof. We first consider the case when B is positive semidefinite. Then there exists an invertible matrix P such that

$$B = P \begin{bmatrix} 0 & 0 \\ 0 & I \end{bmatrix} P^T.$$

Thus, after a change of variables $y = P^T x$, GTRS (1.1) can be equivalently written as

$$\begin{aligned} \inf \quad q_0(y) &:= \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}^T \begin{bmatrix} \bar{A}_1 & \bar{A}_2^T \\ \bar{A}_2 & \bar{A}_3 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} - 2 \begin{bmatrix} \bar{a}_1 \\ \bar{a}_2 \end{bmatrix}^T \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} \\ \text{s.t.} \quad &\ell \leq \|y_2\|^2 \leq u, \end{aligned} \tag{A.1}$$

where

$$y = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}, \quad \bar{A} = \begin{bmatrix} \bar{A}_1 & \bar{A}_2^T \\ \bar{A}_2 & \bar{A}_3 \end{bmatrix} = P^{-1} A P^{-T}, \quad \bar{a} = \begin{bmatrix} \bar{a}_1 \\ \bar{a}_2 \end{bmatrix} = P^{-1} a. \tag{A.2}$$

By our assumptions, the program (A.1) is bounded below. Since

$$q_0(y) = y_1^T \bar{A}_1 y_1 + 2y_2^T \bar{A}_2 y_1 + y_2^T \bar{A}_3 y_2 - 2\bar{a}_1^T y_1 - 2\bar{a}_2^T y_2,$$

and the constraint does not involve y_1 , we must have

$$\bar{A}_1 \succeq 0, \quad \bar{A}_2^T y_2 - \bar{a}_1 \in \text{Range}(\bar{A}_1), \quad \forall \ell \leq \|y_2\|^2 \leq u. \tag{A.3}$$

Notice that $u > 0$ since the RICQ (2.6) holds. Hence, the second relation in (A.3) implies

$$\text{Range}(\bar{A}_2^T) \subseteq \text{Range}(\bar{A}_1) \quad \text{and} \quad \bar{a}_1 \in \text{Range}(\bar{A}_1). \quad (\text{A.4})$$

By (A.3), (A.4) and a consideration of the Schur complement, we conclude that for all sufficiently large $\lambda > 0$, $\bar{A}_3 + \lambda I \succ 0$ and

$$\bar{A} + \lambda \begin{bmatrix} 0 & 0 \\ 0 & I \end{bmatrix} = \begin{bmatrix} \bar{A}_1 & \bar{A}_2^T \\ \bar{A}_2 & \bar{A}_3 + \lambda I \end{bmatrix} \succeq 0. \quad (\text{A.5})$$

Moreover, from (A.4), we see that, for all sufficiently large $\lambda > 0$,

$$\text{Range}(\bar{A}_1 - \bar{A}_2^T[\bar{A}_3 + \lambda I]^{-1}\bar{A}_2) = \text{Range}(\bar{A}_1),$$

and hence $\bar{a}_1 \in \text{Range}(\bar{A}_1 - \bar{A}_2^T[\bar{A}_3 + \lambda I]^{-1}\bar{A}_2)$. Thus, for sufficiently large $\lambda > 0$, the system of equations

$$\begin{bmatrix} \bar{A}_1 - \bar{A}_2^T[\bar{A}_3 + \lambda I]^{-1}\bar{A}_2 & 0 \\ 0 & \bar{A}_3 + \lambda I \end{bmatrix} \begin{bmatrix} w_1 \\ w_2 \end{bmatrix} = \begin{bmatrix} \bar{a}_1 - \bar{A}_2^T[\bar{A}_3 + \lambda I]^{-1}\bar{a}_2 \\ \bar{a}_2 \end{bmatrix} \quad (\text{A.6})$$

is consistent. Since

$$\begin{bmatrix} \bar{A}_1 & \bar{A}_2^T \\ \bar{A}_2 & \bar{A}_3 + \lambda I \end{bmatrix} = \begin{bmatrix} I & \bar{A}_2^T[\bar{A}_3 + \lambda I]^{-1} \\ 0 & I \end{bmatrix} \begin{bmatrix} \bar{A}_1 - \bar{A}_2^T[\bar{A}_3 + \lambda I]^{-1}\bar{A}_2 & 0 \\ 0 & \bar{A}_3 + \lambda I \end{bmatrix} \begin{bmatrix} I & 0 \\ [\bar{A}_3 + \lambda I]^{-1}\bar{A}_2 & I \end{bmatrix},$$

and

$$\begin{bmatrix} I & \bar{A}_2^T[\bar{A}_3 + \lambda I]^{-1} \\ 0 & I \end{bmatrix}^{-1} \begin{bmatrix} \bar{a}_1 \\ \bar{a}_2 \end{bmatrix} = \begin{bmatrix} I & -\bar{A}_2^T[\bar{A}_3 + \lambda I]^{-1} \\ 0 & I \end{bmatrix} \begin{bmatrix} \bar{a}_1 \\ \bar{a}_2 \end{bmatrix} = \begin{bmatrix} \bar{a}_1 - \bar{A}_2^T[\bar{A}_3 + \lambda I]^{-1}\bar{a}_2 \\ \bar{a}_2 \end{bmatrix},$$

we see from the consistency of (A.6) that $\bar{a} \in \text{Range}\left(\bar{A} + \lambda \begin{bmatrix} 0 & 0 \\ 0 & I \end{bmatrix}\right)$ for sufficiently large $\lambda > 0$. From the definition of \bar{a} and \bar{A} in (A.2), we obtain immediately that $a \in \text{Range}(A + \lambda B)$ for sufficiently large $\lambda > 0$. This together with (A.5) proves that D-GTRS is feasible.

The case when B is negative semidefinite can be tackled similarly.

Next, we consider the case when B is indefinite. Suppose to the contrary that D-GTRS is infeasible. Then either $A - \lambda B$ is not positive semidefinite for any $\lambda \in \mathbb{R}$, or there exists λ with $A - \lambda B \succeq 0$ but all such λ satisfy $a \notin \text{Range}(A - \lambda B)$.

Case 1: $A - \lambda B$ is not positive semidefinite for all $\lambda \in \mathbb{R}$. By [24, Theorem 2.3], there exists u^* satisfying $u^{*T}Au^* < 0$ and $u^{*T}Bu^* = 0$.

Fix any $s \in [\ell, u]$. For the above u^* and each $t > 0$, we would like to find a solution x_t of $(x + tu^*)^T B(x + tu^*) = s$ that is uniformly bounded in t . For $s = 0$, we just take $x_t = 0$. In the case when $s > 0$, the x we seek has to satisfy

$$x^T Bx + 2tx^T Bu^* = s.$$

Let $B = PDP^T$ for some orthogonal matrix P and diagonal matrix D , and let β_i denote the i th diagonal entry of D . Also, let \tilde{u}^* denote $P^T u^*$. Since B is indefinite, there exists an index i_0 such that $\beta_{i_0} > 0$. Consider the quadratic equation

$$\beta_{i_0} r^2 + 2t\beta_{i_0}\tilde{u}_{i_0}^* r - s = 0.$$

It is clear that the solutions of the above quadratic equation are

$$r_t^1 = \frac{-2t\beta_{i_0}\tilde{u}_{i_0}^* + \sqrt{(2t\beta_{i_0}\tilde{u}_{i_0}^*)^2 + 4s\beta_{i_0}}}{2\beta_{i_0}} = \frac{2s}{2t\beta_{i_0}\tilde{u}_{i_0}^* + \sqrt{(2t\beta_{i_0}\tilde{u}_{i_0}^*)^2 + 4s\beta_{i_0}}},$$

$$r_t^2 = \frac{-2t\beta_{i_0}\tilde{u}_{i_0}^* - \sqrt{(2t\beta_{i_0}\tilde{u}_{i_0}^*)^2 + 4s\beta_{i_0}}}{2\beta_{i_0}} = \frac{2s}{2t\beta_{i_0}\tilde{u}_{i_0}^* - \sqrt{(2t\beta_{i_0}\tilde{u}_{i_0}^*)^2 + 4s\beta_{i_0}}}.$$

Moreover, it is easy to see that r_t^1 is uniformly bounded for $t > 0$ when $\tilde{u}_{i_0}^* \geq 0$, while r_t^2 is uniformly bounded for $t > 0$ when $\tilde{u}_{i_0}^* \leq 0$. Define

$$x_t = \begin{cases} r_t^1 P e_{i_0} & \text{if } \tilde{u}_{i_0}^* \geq 0, \\ r_t^2 P e_{i_0} & \text{otherwise,} \end{cases}$$

where e_{i_0} is the vector which is one at the i_0 th entry and is zero otherwise. Then $\{x_t\}$ is uniformly bounded in $t > 0$ and satisfies $x_t^T B x_t + 2t x_t^T B u^* = s$. The case when $s < 0$ can be considered similarly, by picking the index such that $\beta_i < 0$.

Next, since $\{x_t\}$ is bounded for $t > 0$ and $u^{*T} A u^* < 0$, we see that

$$(x_t + t u^*)^T A (x_t + t u^*) - 2a^T (x_t + t u^*) = x_t^T A x_t + 2t x_t^T A u^* + t^2 u^{*T} A u^* - 2a^T (x_t + t u^*) \rightarrow -\infty$$

as $t \rightarrow \infty$. This together with the feasibility of $x_t + t u^*$ for all $t > 0$ contradicts the boundedness of GTRS.

Case 2: There exists λ with $A - \lambda B \succeq 0$ but all such λ satisfy $a \notin \text{Range}(A - \lambda B)$. In this case, we apply [4, Theorem A.2] to derive a contradiction. To this end, let $s \in [\ell, u]$ be such that the set $\{x : x^T B x = s\}$ is nonempty. Then, by assumption, we see that the equality constrained GTRS (4.1), with this s , is bounded below. Thus, $f_2(x) := x^T A x - 2a^T x - L$ is nonnegative on the set $\{x : f_1(x) = 0\}$, where L is a lower bound of the optimal value of the equality constrained GTRS, and $f_1(x) := x^T B x - s$. Furthermore, since B is indefinite, there exist x_1, x_2 such that

$$f_1(x_1) > 0 \quad \text{and} \quad f_1(x_2) < 0. \tag{A.7}$$

Using these and invoking [4, Theorem A.2], we conclude that there exists μ such that

$$\begin{pmatrix} -L + \mu s & -a^T \\ -a & A - \mu B \end{pmatrix} \succeq 0.$$

From Schur complement, this implies that $A - \mu B \succeq 0$ and $a \in \text{Range}(A - \mu B)$. Thus, D-GTRS is feasible, a contradiction to our assumption.

Combining Cases 1 and 2, we conclude that D-GTRS is also feasible when B is indefinite. \square

Acknowledgement. The first author would like to thank Nicholas I. M. Gould for help concerning the GLTR algorithm and for providing the pre-release version of the software.

References

- [1] A. Barvinok. Problems of distance geometry and convex properties of quadratic maps. *Discrete Comput. Geom.*, 13(2):189–202, 1995. 8
- [2] A. Barvinok. A remark on the rank of positive semidefinite matrices subject to affine constraints. *Discrete Comput. Geom.*, 25(1):23–31, 2001. 8
- [3] A. Beck. Quadratic matrix programming. *SIAM J. Optim.*, 17(4):1224–1238, 2006. 3
- [4] A. Beck and Y.C. Eldar. Strong duality in nonconvex quadratic optimization with two quadratic constraints. *SIAM J. Optim.*, 17(3):844–860, 2006. 39
- [5] J. F. Bonnans and A. Shapiro. *Perturbation analysis of optimization problems*. Springer Series in Operations Research. Springer-Verlag, New York, 2000. 25

- [6] S. Burer and K.M. Anstreicher. Second-order-cone constraints for extended trust-region subproblems. Technical report, Department of Management Sciences, University of Iowa, 2011. www.optimization-online.org/DB_HTML/2011/03/2957.html. 8
- [7] A.R. Conn, N.I.M. Gould, and Ph.L. Toint. *Trust-Region Methods*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2000. 2, 3
- [8] Y. Ding, D. Ge, and H. Wolkowicz. On equivalence of semidefinite relaxations for quadratic matrix programming. *Math. Oper. Res.*, 36(1):88–104, 2011. 3
- [9] P. Finsler. Über das Vorkommen definiter und semidefiniter Formen in Scharen quadratischer Formen. *Comment. Math. Helv.*, 9:188–192, 1937. 3
- [10] O.E. Flippo and B. Jansen. Duality and sensitivity in nonconvex quadratic optimization over an ellipsoid. *European J. Oper. Res.*, 94(1):167–178, 1996. 3
- [11] C. Fortin and H. Wolkowicz. The trust region subproblem and semidefinite programming. *Optim. Methods Softw.*, 19(1):41–67, 2004. 1, 3, 29, 32
- [12] D.M. Gay. Computing optimal locally constrained steps. *SIAM J. Sci. Statist. Comput.*, 2:186–197, 1981. 3
- [13] I. Gohberg, P. Lancaster, and L. Rodman. *Matrices and Indefinite Scalar Products*. Birkhauser, Verlag Basel, 1983. 3
- [14] G.H. Golub and Qiang Ye. An inverse free preconditioned Krylov subspace method for symmetric generalized eigenvalue problems. *SIAM J. Sci. Comput.*, 24(1):312–334 (electronic), 2002. 4, 33
- [15] N.I.M. Gould, S. Lucidi, M. Roma, and Ph.L. Toint. Solving the trust-region subproblem using the Lanczos method. *SIAM J. Optim.*, 9(2):504–525, 1999. 3, 34
- [16] N.I.M. Gould, Daniel P. Robinson, and H. Sue Thorne. On solving trust-region and other regularised subproblems in optimization. *Math. Program. Comput.*, 2(1):21–57, 2010. 3
- [17] N.I.M. Gould and Ph.L. Toint. A quadratic programming bibliography. Technical report, Rutherford Appleton Laboratory, England, 2001. www.optimization-online.org/DB_HTML/2001/02/285.html. 3
- [18] W.W. Hager. Minimizing a quadratic over a sphere. *SIAM J. Optim.*, 12(1):188–208, 2001. 3
- [19] M.R. Hestenes and E.J. McShane. A theorem on quadratic forms and its application in the calculus of variations. *Trans. Amer. Math. Soc.*, 47:501–512, 1940. 3
- [20] Q. Jin, S.C. Fang, and W. Xing. On the global optimality of generalized trust region subproblems. *Optimization*, 59(8):1139–1151, 2010. 3
- [21] J. Lampe, M. Rojas, D.C. Sorensen, and H. Voss. Accelerating the LSTRS Algorithm. *SIAM J. Sci. Comput.*, 33(1):175–194, 2011. 3
- [22] P. Lancaster and L. Rodman. Canonical forms for Hermitian matrix pairs under strict equivalence and congruence. *SIAM Rev.*, 47(3):407–443, 2005. 12
- [23] Z-Q. Luo and S. Zhang. On extensions of the Frank-Wolfe theorems. *Comput. Optim. Appl.*, 13(1-3):87–110, 1999. Computational optimization—a tribute to Olvi Mangasarian, Part II. 11

- [24] J.J. Moré. Generalizations of the trust region problem. *Optim. Methods Software*, 2:189–209, 1993. 3, 7, 10, 12, 15, 16, 38
- [25] J.J. Moré and D.C. Sorensen. Computing a trust region step. *SIAM J. Sci. Statist. Comput.*, 4:553–572, 1983. 3
- [26] B.N. Parlett. *The symmetric eigenvalue problem*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1998. Corrected reprint of the 1980 original. 11
- [27] G. Pataki. On the rank of extreme matrices in semidefinite programs and the multiplicity of optimal eigenvalues. *Math. Oper. Res.*, 23(2):339–358, 1998. 8, 9
- [28] I. Pólik and T. Terlaky. A survey of the S-lemma. *SIAM Rev.*, 49(3):371–418 (electronic), 2007. 3
- [29] F. Rendl and H. Wolkowicz. A semidefinite framework for trust region subproblems with applications to large scale minimization. *Math. Programming*, 77(2, Ser. B):273–299, 1997. 1, 3, 17, 18, 19, 29, 37
- [30] R.T. Rockafellar. *Convex analysis*. Princeton Landmarks in Mathematics. Princeton University Press, Princeton, NJ, 1997. Reprint of the 1970 original, Princeton Paperbacks. 21
- [31] M. Rojas, S.A. Santos, and D.C. Sorensen. A new matrix-free algorithm for the large-scale trust-region subproblem. *SIAM J. Optim.*, 11(3):611–646 (electronic), 2000/01. 3
- [32] M. Rojas, S.A. Santos, and D.C. Sorensen. Algorithm 873: LSTRS: MATLAB software for large-scale trust-region subproblems and regularization. *ACM Trans. Math. Software*, 34(2):Art. 11, 28, 2008. 3
- [33] D.C. Sorensen. Minimization of a large-scale quadratic function subject to a spherical constraint. *SIAM Journal on Optimization*, 7(1):141–161, 1997. 3
- [34] R. Stern and H. Wolkowicz. Indefinite trust region subproblems and nonsymmetric eigenvalue perturbations. *SIAM J. Optim.*, 5(2):286–313, 1995. 3, 9, 10, 11, 12, 15
- [35] P.D. Tao and L.T.H. An. Difference of convex functions optimization algorithms (DCA) for globally minimizing nonconvex quadratic forms on Euclidean balls and spheres. *Oper. Res. Lett.*, 19(5):207–216, 1996. 3
- [36] V.A. Yakubovich. The S-procedure in nonlinear control theory. *Vestnik Leningrad. Univ.*, 4:73–93, 1977. English Translation, original Russian publication in Vestnik Leningradskogo Universiteta, Seriya Matematika 62-77, 1971. 3
- [37] Y. Ye and S. Zhang. New results on quadratic minimization. *SIAM J. Optim.*, 14(1):245–267 (electronic), 2003. 3, 9

Index

- M^\dagger , Moore-Penrose generalized inverse, 4
- λ_+ , positive part, 4
- λ_- , negative part, 4
- $\phi(t)$, 32
- $\psi(\lambda)$, constraint at $x(\lambda)$, 12
- ri, relative interior, 6
- \underline{t}, \bar{t} , 20
- $\underline{\lambda}, \bar{\lambda}$, 12
- $d(\lambda)$, 19
- $k(t)$, 18
- $k_0(t)$, 18
- $q_0(x)$, objective function, 2
- $q_1(x)$, constraint function, 2
- s_0 , 20
- t_0 , 20
- $x(\lambda)$, first order stationary point, 12
- cl, closure, 12

- closure, cl, 12
- constraint at $x(\lambda)$, $\psi(\lambda)$, 12
- constraint function, $q_1(x)$, 2
- constraint qualification, CQ, 6
- CQ, constraint qualification, 6

- easy/hard case instance, 13

- first order stationary point, $x(\lambda)$, 12

- generalized eigenvalues, $\bar{\lambda}, \underline{\lambda}$, 15
- generalized trust region subproblem, GTRS, 2
- GTRS, generalized trust region subproblem, 2

- hard/easy case instance, 13

- inverse linear interpolation, 32

- Lagrangian, 4

- matrix pencil, 11
- Moore-Penrose generalized inverse, M^\dagger , 4

- objective function, $q_0(x)$, 2
- optimality condition for GTRS, 9

- regular/singular pencil, 11
- relative interior constraint qualification, RICQ, 6
- relative interior, ri, 6

- RICQ, relative interior constraint qualification, 6
- RW algorithm, 17

- SDP relaxation, SDP-GTRS, 5
- SDP, semidefinite programming, 5
- SDP-GTRS, SDP relaxation, 5
- semidefinite programming, SDP, 5
- shift and deflation, 29
- singular/regular pencil, 11

- triangle interpolation, 32
- TRS, trust region subproblem, 2
- trust region subproblem, TRS, 2

- vertical cut, 32