

A FILTER METHOD WITH UNIFIED STEP COMPUTATION FOR NONLINEAR OPTIMIZATION ^{*}

NICHOLAS I. M. GOULD [†], YUELING LOH [‡], AND DANIEL P. ROBINSON [§]

Abstract. We present a filter linesearch method for solving general nonlinear and nonconvex optimization problems. The method is of the filter variety, but uses a robust (always feasible) subproblem based on an exact penalty function to compute a search direction. This contrasts traditional filter methods that use a (separate) restoration phase designed to reduce infeasibility until a feasible subproblem is obtained. Therefore, an advantage of our approach is that every trial step is computed from subproblems that value reducing both the constraint violation *and* the objective function. Moreover, our step computation involves subproblems that are computationally tractable and utilize second derivative information when it is available.

The formulation of each subproblem and the choice of weighting parameter is crucial for obtaining an efficient, robust, and practical method. Our strategy is based on steering methods designed for exact penalty functions, but fortified with a trial step convexification scheme that ensures that a single quadratic optimization problem is solved per iteration. Moreover, we use local feasibility estimates that emerge during the steering process to define a new and improved margin (envelope) of the filter. Under common assumptions, we show that the iterates converge to a local first-order solution of the optimization problem from an arbitrary starting point.

Key words. filter, restoration phase, large-scale, sequential quadratic programming, nonlinear programming

AMS subject classifications. 49M05, 49M15, 65K05, 65K10, 65K15

1. Introduction. This paper considers the general nonlinear optimization problem

$$\underset{x \in \mathbb{R}^n}{\text{minimize}} \quad f(x) \quad \text{subject to} \quad c(x) \geq 0, \quad (1.1)$$

where both the objective function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and the constraint function $c : \mathbb{R}^n \rightarrow \mathbb{R}^m$ are assumed to be twice continuously differentiable. We seek a first-order KKT point (x, y) that satisfies

$$F_{\text{KKT}}(x, y) := \begin{pmatrix} g(x) - J(x)^T y \\ \min [c(x), y] \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad (1.2)$$

where $g(x) := \nabla f(x) \in \mathbb{R}^n$ is the gradient of the objective function, $J(x) := \nabla c(x) \in \mathbb{R}^{m \times n}$ is the Jacobian of the constraint function, y is the Lagrange multiplier vector, and the minimum is taken component-wise. Our algorithm may easily handle constraints with general lower/upper bounds, and handle equality constraints directly, i.e., do not replace them with pairs of inequality constraints. Problems of this type arise naturally in many areas including optimal control [2, 3, 5, 22, 28], resource allocation [1, 27], solution of equilibrium models [17, 33], and structural engineering [4, 30], among others.

Popular methods for solving (1.1) can broadly be characterized as interior-point or active-set methods. Interior-point algorithms [38, 40, 41] offer polynomial-time complexity bounds in many cases and readily scale-up to problems involving millions of variables. Their main disadvantage is the inability to use effectively a good initial estimate of a solution. In fact, many interior-point methods immediately move the initial guess into the strict interior of the feasible region. It is from this interior location that future iterates are forced to remain and justifies the name “interior-point” methods; more modern “infeasible” interior-point methods avoid this defect to some degree.

^{*}Research supported by the National Science Foundation grant DMS-1217153 and EPSRC grant EP/I013067/1.

[†]Scientific Computing Dept., Rutherford Appleton Laboratory, Chilton, OX11-0QX, England.

[‡]Dept. of Applied Mathematics and Statistics, Johns Hopkins University, Baltimore, MD 21218, USA.

[§]Dept. of Applied Mathematics and Statistics, Johns Hopkins University, Baltimore, MD 21218, USA.

Active-set methods [8, 14, 23, 24, 25, 26, 32, 34] complement interior-point methods since they naturally utilize information derived from a good estimate of a solution. In fact, if the optimal active set (the set containing those constraints satisfied as equalities at a solution) was known in advance, then problem (1.1) could be solved as an equality constrained problem and its combinatorial nature would be eliminated. It is precisely this property that makes active-set methods widely used to solve the previously mentioned class of problems. The main weakness of active-set algorithms is that each subproblem typically requires the solution of a linear or quadratic program, which is often expensive when compared to interior-point methods that require a single linear system solve per iteration.

In this paper we describe an active-set method that generates a sequence of iterates from the solutions of subproblems defined by local models of the nonlinear problem functions. The subproblems are always feasible since they are based on an exact penalty function. To ensure that these models result in productive steps, we use steering techniques [9] to adaptively adjust the weighting (penalty) parameter. In contrast to original steering methods, we use a step convexification procedure similar to [] to avoid solving multiple quadratic programs during each iteration.

To provide convergence guarantees, we must include a mechanism for determining when one point is “better” than another. A merit (penalty) function or a filter is amongst the most common tools used for this purpose. A merit function combines the objective function and a measure of constraint violation into a single function, whereby their individual perceived importance is determined by a weighting parameter. The quality of competing points is then measured by comparing their respective merit function values. A potential weakness is that the quality of iterates depends on the value of the weighting parameter, which can make step acceptance sensitive to its value. In part, filter methods surfaced to mitigate this parameter dependence. In the context of nonlinear optimization, filter methods were introduced by Fletcher, Leyffer, and Toint [19, 20] and have since been rather popular [11, 12, 13, 18, 21, 38]. A filter views problem (1.1) as a multi-criterion optimization problem consisting of minimizing the objective function and minimizing some measure of the constraint violation, with certain preference given to the latter. Roughly, a trial iterate is then considered acceptable if it has a smaller value of *either* the objective function or the constraint violation compared to the previously encountered points. Consequently, it is often the case that filter methods accept more iterates and perform better. It should be mentioned, however, that every known provably convergent filter method has a weak dependence between these two criteria that is embedded in the step acceptance criteria. In fact, this observation partly motivated the work on *flexible* penalty methods by Curtis and Nocedal [16]. They describe how a *single* element filter is essentially equivalent to the union of points acceptable to the ℓ_1 -penalty function defined over an interval of weighting parameter values.

A great disadvantage of filter methods is that they (traditionally) require the use of a restoration phase. A restoration phase is (typically) entered when the subproblem used to compute trial steps is infeasible; some algorithms, e.g., [40], enter the restoration phase for additional reasons. When this situation occurs, the restoration phase is triggered and a sequence of iterates focused on reducing the constraint-violation is computed until the desired subproblem becomes feasible. During this phase, the objective function is essentially *ignored*, which is highly undesirable from both a practical and computational perspective.

Our active-set method is globalized by using a filter, but never needs to enter a (traditional) restoration phase. This is accomplished by using subproblems that are always feasible and, in certain instances, allowing for the acceptance of iterates that decrease both the exact penalty function and the constraint violation. In essence, we replace an undesirable restoration phase with an attractive penalty phase. Thus, we combine ideas from both filter and penalty methods to formulate a robust and effective method; we believe this further builds upon the basic observations in [16].

This paper contains three main contributions. First, we present a filter method that avoids a traditional and highly undesirable restoration phase. To this end, we utilize subproblems based on exact penalty functions that are *always* feasible and formed from models of *both* the objective function and

constraint violation. Second, our method incorporates second derivative information without requiring global minimizers of nonconvex constrained subproblems (c.f. [18]). Our step computation is most similar to [24, 26], which was described in the context of line-search and trust-region penalty methods. Third, we use local feasibility estimates that emerge during the steering step computation to define a new and improved margin (envelope) of the filter. This allows us to define an adaptive and practical margin.

Our work is not the only method designed to resolve weaknesses in traditional filter methods. Chen and Goldfarb [10] presented an interior point method that uses two penalty functions to determine step acceptance: a piecewise linear penalty function whose break points are essentially elements in the filter, and the ℓ_2 -penalty function. Under this scheme, a trial step is accepted if it provides sufficient reduction for either penalty function.

The remainder of this paper is organized as follows. In Section 2 we describe the algorithm in detail and in Section 3 prove that it is well-posed. In Section 4 we provide convergence results and conclude with final remarks in Section 5.

2. A filter sequential quadratic programming method. In this section we describe our new filter sequential quadratic programming method, FiSQP. The algorithm is iterative and relies on computing trial steps from carefully constructed subproblems. These subproblems and the resulting trial steps are explained in Sections 2.1–2.6. In Section 2.7 we introduce the filter construct and related terminology; we emphasize that acceptability to the filter is only a necessary condition for accepting a trial iterate. A full statement and description of the algorithm is given in Section 2.8.

Our step computation is based on the ℓ_1 -penalty function

$$\phi(x; \sigma) := f(x) + \sigma v(x), \quad (2.1)$$

where the constraint violation at x is defined by

$$v(x) := \|[c(x)]^-\|_1, \quad \text{with } [y]^- := \max(-y, 0). \quad (\text{minimum is component-wise}) \quad (2.2)$$

and σ is a positive weighting parameter. Appropriate linear and quadratic models of ϕ are given by

$$\ell^\phi(s; x, \sigma) := \ell^f(s; x) + \sigma \ell^v(s; x) \quad \text{and} \quad q^\phi(s; x, M, \sigma) := q^f(s; x, M) + \sigma \ell^v(s; x), \quad (2.3)$$

where

$$\ell^f(s; x) := f(x) + g(x)^T s \quad \text{and} \quad q^f(s; x, M) := \ell^f(s; x) + \frac{1}{2} s^T M s \quad (2.4)$$

are linear and quadratic model approximations, respectively, of the objective function f for a given symmetric matrix $M \in \mathbb{R}^{n \times n}$, and

$$\ell^v(s; x) := \|[c(x) + J(x)s]^-\|_1$$

is a piecewise-linear approximation to the constraint violation function v . Using these models we may predict the change in v with the function

$$\Delta \ell^v(s; x) := \ell^v(0; x) - \ell^v(s; x) = \|[c(x)]^-\|_1 - \|[c(x) + J(x)s]^-\|_1, \quad (2.5)$$

the change in f with the functions

$$\Delta \ell^f(s; x) := \ell^f(0; x) - \ell^f(s; x) = -g(x)^T s \quad \text{and} \quad (2.6a)$$

$$\Delta q^f(s; x, M) := q^f(0; x, M) - q^f(s; x, M) = \Delta \ell^f(s; x) - \frac{1}{2} s^T M s, \quad (2.6b)$$

and the change in the penalty function ϕ with the functions

$$\Delta \ell^\phi(s; x, \sigma) := \ell^\phi(0; x, \sigma) - \ell^\phi(s; x, \sigma) = \Delta \ell^f(s; x) + \sigma \Delta \ell^v(s; x) \quad \text{and} \quad (2.7)$$

$$\Delta q^\phi(s; x, M, \sigma) := q^\phi(0; x, M, \sigma) - q^\phi(s; x, M, \sigma) = \Delta \ell^\phi(s; x, \sigma) - \frac{1}{2} s^T M s. \quad (2.8)$$

For the remainder of this section, let (x_k, y_k) denote the current estimate of a solution to (1.1).

2.1. The steering step s_k^s . In order to strike a proper balance between reducing the objective function and the constraint violation, we compute a *steering* step s_k^s as a solution to the linear program

$$\underset{(s,r) \in \mathbb{R}^{n+m}}{\text{minimize}} \quad e^T r \quad \text{subject to} \quad c_k + J_k s + r \geq 0, \quad r \geq 0, \quad \|s\|_\infty \leq \delta_k, \quad (2.9)$$

where $c_k = c(x_k)$, $J_k = J(x_k)$, $\delta_k \in [\delta_{\min}, \delta_{\max}]$, and $0 < \delta_{\min} \leq \delta_{\max} < \infty$. Problem (2.9) is equivalent to the nonsmooth problem

$$\underset{s \in \mathbb{R}^n}{\text{minimize}} \quad \ell^v(s; x_k) \quad \text{subject to} \quad \|s\|_\infty \leq \delta_k, \quad (2.10)$$

since s solves (2.10) if and only if (s, r) solves (2.9), where $r = \max(-(c_k + J_k s), 0)$. Since $\ell^v(0; x_k) = v(x_k)$, ℓ^v is a convex function, and $s = 0$ is feasible for (2.10), it follows from (2.5) that $\Delta \ell^v(s_k^s; x_k) \geq 0$. The quantity $\Delta \ell^v(s_k^s; x_k)$ is the best local improvement in linearized constraint feasibility for steps of size δ_k .

All methods for nonconvex optimization may converge to an infeasible point that is a local minimizer of the constraint violation as measured by v . Points of this type are known as infeasible stationary points, which we now define by utilizing the steering subproblem.

DEFINITION 2.1 (Infeasible stationary point). *The vector x^i is an infeasible stationary point if $v(x^i) > 0$ and $\Delta \ell^v(s^i; x^i) = 0$, where $s^i = \operatorname{argmin}_{s \in \mathbb{R}^n} \ell^v(s; x^i)$ subject to $\|s\|_\infty \leq \delta$ for some $\delta > 0$.*

2.2. The predictor step s_k^p . The *predictor* step is computed as the *unique* solution to one of the following strictly convex minimization problems:

$$s_k^p = \begin{cases} \underset{s \in \mathbb{R}^n}{\operatorname{argmin}} \quad f_k + g_k^T s + \frac{1}{2} s^T B_k s \quad \text{subject to} \quad c_k + J_k s \geq 0, & \text{if } \Delta \ell^v(s_k^s; x_k) = v(x_k), \\ \underset{s \in \mathbb{R}^n}{\operatorname{argmin}} \quad q^\phi(s; x_k, B_k, \sigma_k), & \text{otherwise,} \end{cases} \quad (2.11a)$$

$$(2.11b)$$

where $\sigma_k > 0$ is the k th value of the penalty parameter, $f_k = f(x_k)$, $g_k = \nabla f(x_k)$, $c_k = c(x_k)$, $J_k = \nabla c(x_k)$, B_k is a positive-definite matrix that we are free to choose such that $B_k \approx \nabla_{xx}^2 L(x_k, y_k)$, and the Lagrangian L is defined by $L(x, y) = f(x) - c(x)^T y$. Analogous to the steering subproblem, the nonsmooth minimization problem (2.11b) is equivalent to the smooth problem

$$\underset{s \in \mathbb{R}^n, r \in \mathbb{R}^n}{\text{minimize}} \quad f_k + g_k^T s + \frac{1}{2} s^T B_k s + \sigma_k e^T r \quad \text{subject to} \quad c_k + J_k s + r \geq 0, \quad r \geq 0, \quad (2.12)$$

which is the problem solved in practice. We use y_k^p to denote the Lagrange multiplier vector for the constraint $c_k + J_k s \geq 0$ in (2.11a) and $c_k + J_k s + r \geq 0$ in (2.12) (equivalently (2.11b)). A trivial choice for the positive-definite matrix is $B_k = I$, but other choices based on quasi-Newton updates such as BFGS [35] or L-BFGS [29] are also possible.

The next result shows how convergence to KKT points may be deduced from the predictor problem.

LEMMA 2.2. *Suppose that x_* satisfies*

$$v(x_*) = 0 \quad \text{and} \quad 0 = \underset{s \in \mathbb{R}^n}{\operatorname{argmin}} \quad f(x_*) + g(x_*)^T s + \frac{1}{2} s^T B s \quad \text{subject to} \quad c(x_*) + J(x_*) s \geq 0 \quad (2.13)$$

for some positive definite matrix B , and let y_* denote the associated Lagrange multiplier vector. Then, it follows that (x_*, y_*) is a KKT point for problem (1.1) as defined by (1.2).

Proof. Since B is positive definite, $s = 0$ is the unique solution to the optimization problem in (2.13). It then follows from the first-order necessary optimality conditions at $s = 0$ that

$$g(x_*) = J(x_*)^T y_*, \quad \text{and} \quad \min(c(x_*), y_*) = 0,$$

where y_* is the Lagrange multiplier for the constraint $c(x_*) + J(x_*) s \geq 0$. It now follows from Definition 1.2 that (x_*, y_*) is a KKT point for problem (1.1). \square

2.3. The search direction s_k . The steering direction s_k^s provides a measure of local progress in infeasibility. Since we desire a search direction s_k that makes progress towards feasibility, we define

$$s_k := (1 - \tau_k)s_k^s + \tau_k s_k^p \quad (2.14)$$

where τ_k is the largest number on $[0, 1]$ such that

$$\Delta \ell^v(s_k; x_k) \geq \eta_v \Delta \ell^v(s_k^s; x_k) \geq 0 \text{ for some } \eta_v \in (0, 1). \quad (2.15)$$

The next lemma shows that $\tau_k > 0$ when x_k is not an infeasible stationary point. This is important since the step s_k then has a significant contribution from s_k^p , which was computed from a subproblem that modeled *both* the objective and constraint functions; this contrasts traditional filter methods when restoration is entered since the subproblem formulations then focus solely on the constraint violation.

LEMMA 2.3. *If x_k is not an infeasible stationary point as given by Definition 2.1, then $\tau_k > 0$.*

Proof. If $v(x_k) = 0$, then $\Delta \ell^v(s_k^s; x_k) = 0$. It then follows from (2.11a) that $c_k + J_k s_k^p \geq 0$, which in turn implies that $\Delta \ell^v(s_k^p; x_k) = 0$. Thus, the choice $\tau_k = 1$ satisfies (2.14) and (2.15).

Now suppose that $v(x_k) > 0$ and define

$$s(\tau) = (1 - \tau)s_k^s + \tau s_k^p$$

so that $\lim_{\tau \downarrow 0} s(\tau) = s_k^s$. It then follows from continuity of $\Delta \ell^v(\cdot; x_k)$ and the fact that $\Delta \ell^v(s_k^s; x_k) > 0$ since x_k is not an infeasible stationary point by assumption, that

$$\lim_{\tau \downarrow 0} \Delta \ell^v(s(\tau); x_k) = \Delta \ell^v(s_k^s; x_k) > 0.$$

Therefore, there exists $\tau' > 0$ such that

$$|\Delta \ell^v(s(\tau); x_k) - \Delta \ell^v(s_k^s; x_k)| < (1 - \eta_v) \Delta \ell^v(s_k^s; x_k) \text{ for all } \tau \in [0, \tau']$$

since $\eta_v \in (0, 1)$ in (2.15) and $\Delta \ell^v(s_k^s; x_k) > 0$. However, this implies that

$$\Delta \ell^v(s(\tau); x_k) \geq \eta_v \Delta \ell^v(s_k^s; x_k) \text{ for all } \tau \in [0, \tau'],$$

which guarantees that $t_k \geq \tau' > 0$. \square

We now proceed to show that if $\Delta \ell^v(s_k^s; x_k) > 0$, then s_k is a descent direction for $v(\cdot)$. We require the definition of the directional derivative of a function.

DEFINITION 2.4. *The directional derivative of a function $h(\cdot)$ in the direction d and at the point x is defined (when it exists) as*

$$[D_d h](x) := \lim_{t \downarrow 0} \frac{h(x + td) - h(x)}{t}.$$

We now show that the directional derivative is bounded by the negative of the change in its model.

LEMMA 2.5. *At any point x and for any direction d , it follows that*

$$[D_d v](x) \leq -\Delta \ell^v(d; x),$$

where the function $D_d v$ is the directional derivative of v in the direction d .

Proof. Since ℓ^v is a convex function and $\ell^v(0; x)$ is finite, it follows from [37, Theorem 23.1] that

$$\frac{\ell^v(td; x) - \ell^v(0; x)}{t}$$

is monotonically non-decreasing with t , $[D_d \ell^v](0; x)$ exists, and

$$[D_d \ell^v](0; x) = \inf_{t>0} \frac{\ell^v(td; x) - \ell^v(0; x)}{t}. \quad (2.16)$$

It then follows from [7, Lemma 3.1], (2.16) and the definition of $\Delta \ell^v$ that

$$[D_d v](x) = [D_d \ell^v](0; x) \leq \ell^v(d; x) - \ell^v(0; x) = -\Delta \ell^v(d; x),$$

which is the desired result. \square

Thus the search direction s_k is a descent direction for v when our infeasibility measure is positive.

LEMMA 2.6. *If $\Delta \ell^v(s_k^s; x_k) > 0$, the direction s_k is a descent direction for v at the point x_k , i.e.,*

$$[D_{s_k} v](x_k) \leq -\Delta \ell^v(s_k; x_k) \leq -\eta_v \Delta \ell^v(s_k^s; x_k) < 0, \text{ where } \eta_v \text{ is defined in (2.15).}$$

Proof. It follows directly from Lemma 2.5, (2.15), and $\Delta \ell^v(s_k^s; x_k) > 0$ that

$$[D_{s_k} v](x_k) \leq -\Delta \ell^v(s_k; x_k) \leq -\eta_v \Delta \ell^v(s_k^s; x_k) < 0,$$

which implies that s_k is a descent direction for v at the point x_k . \square

We now consider the case when our infeasibility measure is zero.

LEMMA 2.7. *Suppose $\Delta \ell^v(s_k^s; x_k) = 0$, then one of the following must occur:*

- (i) $v(x_k) > 0$ and x_k is an infeasible stationary point; or
- (ii) $v(x_k) = 0$ and $\Delta \ell^\phi(s_k; x_k, \sigma) \geq \frac{1}{2} s_k^{pT} B_k s_k^p$ for all $0 < \sigma < \infty$.

Proof. If $v(x_k) > 0$, then by Definition 2.1, x_k is an infeasible stationary point which is part (i). Now, suppose that $v(x_k) = 0$. As in the proof of Lemma 2.3, it follows that

$$\Delta \ell^v(s_k^p; x_k) = 0, \quad \tau_k = 1, \quad \text{and} \quad s_k = s_k^p. \quad (2.17)$$

We may then use the definition of s_k^p in (2.11a), (2.17), and (2.6b) to conclude that

$$0 \leq \Delta q^\phi(s_k^p; x_k, B_k, \sigma_k) = \Delta q^f(s_k^p; x_k, B_k) = \Delta \ell^f(s_k^p; x_k) - \frac{1}{2} s_k^{pT} B_k s_k^p,$$

which yields $\Delta \ell^f(s_k^p; x_k) \geq \frac{1}{2} s_k^{pT} B_k s_k^p$. Combining this with (2.7) and (2.17), we have that

$$\begin{aligned} \Delta \ell^\phi(s_k; x_k, \sigma) &= \Delta \ell^f(s_k; x_k) + \sigma \Delta \ell^v(s_k; x_k) \\ &= \Delta \ell^f(s_k; x_k) = \Delta \ell^f(s_k^p; x_k) \geq \frac{1}{2} s_k^{pT} B_k s_k^p \text{ for all finite } \sigma, \end{aligned}$$

which completes the proof. \square

2.4. Updating the weighting parameter. By design, the trial step s_k is a descent direction for v when local improvement in feasibility is possible. Since the weighting parameter provides a balance between reducing the objective function and the constraint violation, it makes sense to adjust the weighting parameter so that s_k is also a descent direction for ϕ . This is accomplished by defining

$$\sigma_{k+1} = \begin{cases} \sigma_k & \text{if } \Delta \ell^\phi(s_k; x_k, \sigma_k) \geq \sigma_k \eta_\sigma \Delta \ell^v(s_k^s; x_k) \\ \max \left\{ \sigma_k + \sigma_{\text{inc}}, \frac{-\Delta \ell^f(s_k; x_k)}{\Delta \ell^v(s_k; x_k) - \eta_\sigma \Delta \ell^v(s_k^s; x_k)} \right\} & \text{otherwise} \end{cases} \quad (2.18)$$

for some $\sigma_{\text{inc}} > 0$ and η_σ satisfying $0 < \eta_\sigma < \eta_v < 1$, where η_v is defined in (2.15).

LEMMA 2.8. *If x_k is not an infeasible stationary point, then the parameter update (2.18) is well defined and ensures that*

$$\Delta \ell^\phi(s_k; x_k, \sigma_{k+1}) \geq \sigma_{k+1} \eta_\sigma \Delta \ell^v(s_k^s; x_k) \geq 0 \quad \text{for all } k \geq 0. \quad (2.19)$$

Proof. If $\Delta\ell^\phi(s_k; x_k, \sigma_k) \geq \sigma_k \eta_\sigma \Delta\ell^v(s_k^s; x_k)$, then the desired result immediately follows from the update $\sigma_{k+1} = \sigma_k$. Thus, for the remainder of the proof we assume that

$$\Delta\ell^\phi(s_k; x_k, \sigma_k) < \sigma_k \eta_\sigma \Delta\ell^v(s_k^s; x_k). \quad (2.20)$$

Suppose, for a contradiction, that $\Delta\ell^v(s_k^s; x_k) = 0$. Since x_k is not an infeasible stationary point by assumption, it follows that $v(x_k) = 0$. Then, it follows from Lemma 2.7 and the fact that B_k is positive definite by assumption that $\Delta\ell^\phi(s_k; x_k, \sigma_k) \geq \frac{1}{2} s_k^p{}^T B_k s_k^p \geq 0$, which contradicts (2.20) since $\Delta\ell^v(s_k^s; x_k) = 0$. Thus, we conclude that $\Delta\ell^v(s_k^s; x_k) > 0$. Combining this with the choice $0 < \eta_\sigma < \eta_v < 1$ in (2.18) and (2.15) we conclude that $\Delta\ell^v(s_k; x_k) \geq \eta_v \Delta\ell^v(s_k^s; x_k) > \eta_\sigma \Delta\ell^v(s_k^s; x_k) > 0$, and thus

$$\eta_\sigma \Delta\ell^v(s_k^s; x_k) - \Delta\ell^v(s_k; x_k) < 0. \quad (2.21)$$

It then follows from (2.7), (2.20), (2.21), and the fact that $\sigma_k > 0$ that

$$\Delta\ell^f(s_k; x_k) = \Delta\ell^\phi(s_k; x_k, \sigma_k) - \sigma_k \Delta\ell^v(s_k; x_k) < \sigma_k [\eta_\sigma \Delta\ell^v(s_k^s; x_k) - \Delta\ell^v(s_k; x_k)] < 0. \quad (2.22)$$

Inequalities (2.21) and (2.22) imply that the penalty parameter update (2.18) is well-defined and positive.

It now follows from (2.18) that

$$\sigma_{k+1} \geq \frac{-\Delta\ell^f(s_k; x_k)}{\Delta\ell^v(s_k; x_k) - \eta_\sigma \Delta\ell^v(s_k^s; x_k)},$$

which may then be combined with (2.21) to yield

$$\sigma_{k+1} \eta_\sigma \Delta\ell^v(s_k^s; x_k) \leq \Delta\ell^f(s_k; x_k) + \sigma_{k+1} \Delta\ell^v(s_k; x_k) = \Delta\ell^\phi(s_k; x_k, \sigma_{k+1}),$$

which is the desired result (2.19). \square

The next result will allow us to show that s_k is a descent direction for ϕ under certain assumptions.

LEMMA 2.9. *For any given value of the penalty parameter σ , point x , direction d , and positive-definite matrix B , it follows that*

$$[D_d\phi](x; \sigma) \leq -\Delta\ell^\phi(d; x, \sigma) \leq -\Delta q^\phi(d; x, B, \sigma).$$

Proof. Linearity of the directional derivative, (2.6a), Lemma 2.5, (2.7), (2.8), and the fact that B_k is positive definite by choice, imply that

$$\begin{aligned} [D_d\phi](x; \sigma) &= [D_d f](x) + \sigma [D_d v](x) = -g(x)^T d + \sigma [D_d v](x) \leq -\Delta\ell^f(d; x) - \sigma \Delta\ell^v(d; x) \\ &= -\Delta\ell^\phi(d; x, \sigma) = -\Delta q^\phi(d; x, B, \sigma) - \frac{1}{2} d^T B d \leq -\Delta q^\phi(d; x, B, \sigma), \end{aligned}$$

which is the desired result. \square

In most situations, we may now show that s_k is a descent direction for the penalty function.

LEMMA 2.10. *If x_k is neither an infeasible stationary point nor a KKT point for problem (1.1), then the direction s_k is a descent direction for $\phi(x; \sigma_{k+1})$ at the point x_k , i.e.,*

$$[D_{s_k}\phi](x_k; \sigma_{k+1}) \leq -\Delta\ell^\phi(s_k; x_k, \sigma_{k+1}) < 0.$$

Proof. If $\Delta\ell^v(s_k^s; x_k) > 0$, then x_k cannot be an infeasible stationary point, and it follows from Lemma 2.9, Lemma 2.8, and (2.19) that $[D_{s_k}\phi](x_k; \sigma_{k+1}) \leq -\Delta\ell^\phi(s_k; x_k, \sigma_{k+1}) < 0$, which is the desired result. Conversely, if $\Delta\ell^v(s_k^s; x_k) = 0$, then $v(x_k) = 0$ since x_k is not an infeasible stationary point by assumption. It now follows from Lemma 2.9, $v(x_k) = 0$, Lemma 2.7, the fact that B_k is positive definite, and $s_k^p \neq 0$ since x_k is not a KKT point for problem (1.1) by assumption (see Lemma 2.2), that $[D_{s_k}\phi](x_k; \sigma_{k+1}) \leq -\Delta\ell^\phi(s_k; x_k, \sigma_{k+1}) \leq -\frac{1}{2} s_k^p{}^T B_k s_k^p < 0$, which completes the proof. \square

2.5. The accelerator step s_k^a . To improve performance, we compute an additional “acceleration” step; here we consider a single (simple) possibility, but other variants may be used [24].

Under common assumptions, the predictor step s_k^p will ultimately correctly identify those constraints that are active at a local solution of (1.1) [36]. A prediction based on s_k^p is formulated by

$$\mathcal{A}_k := \{i : [c_k + J_k s_k^p]_i = 0\}. \quad (2.23)$$

It is then natural to compute an *accelerator* step s_k^a as the solution to

$$\underset{s \in \mathbb{R}^n}{\text{minimize}} \quad q^f(s_k^p + s; x_k, H_k) \quad \text{subject to} \quad [J_k s]_{\mathcal{A}_k} = 0, \quad \|s\|_2 \leq \delta_k^a, \quad (2.24)$$

where $\delta_k^a > 0$ is the trust-region radius, H_k is a symmetric and uniformly bounded approximation of $\nabla_{xx}^2 L(x_k, y_k)$, and y_k is a suitable Lagrange multiplier vector such as those from the predictor subproblem. We note that subproblem (2.24) may be solved, for example, with the projected GLTR algorithm (see [15, Section 7.5.4] and the notes at the end that describe how to cope with the affine constraints $[J_k s]_{\mathcal{A}_k} = 0$). It can be shown that if $c_k + J_k s \geq 0$ is feasible, σ_k is sufficiently large, and x_k is “close enough” to a solution of (1.1) that satisfies certain second-order sufficient optimality conditions, then $s_k^p + s_k^a$ is the solution to

$$\underset{s \in \mathbb{R}^n}{\text{minimize}} \quad q^f(s; x_k, H_k) \quad \text{subject to} \quad c_k + J_k s \geq 0, \quad (2.25)$$

which is the traditional SQP subproblem. However, our method of step computation is robust whereas the generally nonconvex subproblem (2.25) introduces many points of contention such as multiple solutions, unboundedness, and inconsistent constraints.

2.6. The Cauchy steps s_k^{cf} and $s_k^{c\phi}$. Since the matrix B_k is positive definite by construction and the exact second-derivative matrix H_k is generally an indefinite matrix, they may differ dramatically. To account for this when assessing overall step acceptance, we define and use a *Cauchy-f* step s_k^{cf} and *Cauchy- ϕ* step $s_k^{c\phi}$ as follows.

Given the search direction s_k , we define the Cauchy-f step as

$$s_k^{cf} := \alpha_k^f s_k, \quad \text{where} \quad \alpha_k^f := \underset{0 \leq \alpha \leq 1}{\operatorname{argmin}} \quad q^f(\alpha s_k; x_k, H_k). \quad (2.26)$$

Similarly, we define the Cauchy- ϕ step as

$$s_k^{c\phi} := \alpha_k^\phi s_k, \quad \text{where} \quad \alpha_k^\phi := \underset{0 \leq \alpha \leq 1}{\operatorname{argmin}} \quad q^\phi(\alpha s_k; x_k, H_k, \sigma_{k+1}). \quad (2.27)$$

The step size α_k^ϕ may be found efficiently by examining the piecewise quadratic function $q^\phi(\alpha s_k; x_k, H_k, \sigma_{k+1})$ segment-by-segment between each derivative discontinuity.

2.7. The filter. We ensure global convergence of our method by maintaining/updating a filter \mathcal{F}_k during each iteration. A filter is defined as follows, where \mathbb{R}^+ denotes the positive real numbers.

DEFINITION 2.11 (filter). A filter is any finite set of points in $\mathbb{R}^+ \times \mathbb{R}$.

The initial filter is defined to be $\mathcal{F}_0 = \emptyset$ and then sequentially updated in a manner that guarantees that $\mathcal{F}_k \subseteq \{(v_j, f_j) : 0 \leq j < k\}$. The decision to add certain ordered pairs to the filter depends on the concept of trial points being acceptable to the filter, which we now define.

DEFINITION 2.12 (acceptable to \mathcal{F}_k). We say that the point x is acceptable to \mathcal{F}_k if its associated ordered pair $(v(x), f(x))$ satisfies

$$v(x) \leq \max \{v_i - \alpha_i \eta_v \Delta \ell^v(s_i^s; x_i), \beta v_i\} \quad \text{or} \quad f(x) \leq f_i - \gamma \min \{v_i - \alpha_i \eta_v \Delta \ell^v(s_i^s; x_i), \beta v_i\} \quad (2.28)$$

for all $0 \leq i < k$ such that $(v_i, f_i) \in \mathcal{F}_k$ and some constants $\{\eta_v, \beta, \gamma\} \subset (0, 1)$.

The first inequality in (2.28) ensures that the constraint violation has been sufficiently reduced. We note that previous filter methods have not used the first quantity in the max on the right-hand side. Our improved condition takes advantage of the information supplied by the steering steps s_k^s . Previous filter methods may easily have requested a decrease in the constraint violation that was unreasonable. In these circumstances, the trust-region radius would be decreased until the subproblem became infeasible and then a feasibility restoration phase would be entered. Our modified definition provides a practical target constraint violation based on local information derived from the steering step s_k^s . The second inequality in (2.28) guarantees that the objective function is sufficiently smaller at the point x than at points x_i whose ordered pair is in the current filter \mathcal{F}_k . These two conditions provide a so-called *margin* around the elements of the filter.

Note that Definition 2.12 does not require and does not imply that the current vector x_k is in \mathcal{F}_k when determining acceptability. During our search for an improved estimate of a solution to (1.1), it often does not make sense to accept a new point unless it is acceptable to the current filter *and* better than the current point x_k . This leads to the following definition.

DEFINITION 2.13 (acceptable to \mathcal{F}_k augmented by x_k). *We say that x is acceptable to \mathcal{F}_k augmented by x_k if x is acceptable to \mathcal{F}_k as given by Definition 2.12 and (2.28) holds with $i = k$.*

In the next section we present our main filter SQP method. Each iteration requires the search for a new point that must satisfy a subset of specified conditions. We stress that the updated point x_{k+1} is not necessarily acceptable to \mathcal{F}_k . Moreover, the vector x_{k+1} being acceptable to \mathcal{F}_k (possibly augmented by x_k) is a necessary, but not sufficient condition, for adding the ordered pair (v_{k+1}, f_{k+1}) to the filter \mathcal{F}_k . Details of how we update \mathcal{F}_k are described in the next section.

2.8. The algorithm. In this section we formally state and describe our filter trust-region algorithm. Specific termination tests are not stated, but would be included in practice.

Algorithm 1 on page 12 begins by defining step acceptance parameters $\{\eta_v, \eta_\sigma, \sigma_{\text{inc}}, \gamma, \gamma_v, \gamma_f, \gamma_\phi\} \subset (0, 1)$, steering trust region parameters $0 < \delta_{\min} \leq \delta_{\max} < \infty$, an initial weighting parameter σ_0 , an initial trust region radius for the steering subproblem $\delta_0 \in [\delta_{\min}, \delta_{\max}]$, and an initial positive definite matrix B_0 . It then sets the iteration index $k = 0$ and flag *P-mode* to false, which indicates that we begin in what we shall call *filter* (as opposed to *penalty*) mode.

The main loop is now entered and a sequence of trial steps is computed. First, a steering step s_k^s is computed as a solution of (2.10), which is then used to calculate $\Delta \ell^v(s_k^s; x_k)$. This quantity gives us a tangible quantity that predicts the decrease in feasibility that one might expect to acquire. In particular, the quantity allows us to determine whether x_k is an infeasible stationary point. If x_k is not an infeasible stationary point, then the predictor step s_k^p is computed as the unique solution to the strictly convex subproblem (2.11). The search direction s_k is then defined by (2.14) to satisfy (2.15). By construction this definition ensures that s_k is a descent direction for v whenever $v(x_k) > 0$ (see Lemma 2.6). Next, we adjust the penalty parameter using (2.18) so that s_k is also a decent direction for the penalty function ϕ (see Lemma 2.10). To accelerate convergence, we compute an accelerator step s_k^a as an (approximate) solution to subproblem (2.24), which requires an active set prediction \mathcal{A}_k given by (2.23). We complete the step calculations by computing the Cauchy steps s_k^{cf} and $s_k^{c\phi}$ defined by (2.26) and (2.27).

If *P-mode* has the value true, we perform a backtracking linesearch until we find a p-pair (α_k, \hat{s}_k) for some $\hat{s}_k \in \{s_k^a, s_k\}$. We define a p-pair as follows.

P-PAIR	
<i>The pair (α, s) constitutes a p-pair if</i>	
$\phi(x_k + \alpha s; \sigma_{k+1}) \leq \phi(x_k; \sigma_{k+1}) - \gamma_\phi \alpha \rho_k^\phi \quad (2.29)$	
<i>for some $\gamma_\phi \in (0, 1)$, where</i>	
$\rho_k^\phi := \min \left[\Delta \ell^\phi(s_k; x_k, \sigma_{k+1}), \Delta q^\phi(s_k^{c\phi}; x_k, H_k, \sigma_{k+1}) \right]. \quad (2.30)$	

If (α_k, \hat{s}_k) is a p-pair, then $\phi(x_k + \alpha_k \hat{s}_k; \sigma_{k+1})$ is sufficiently smaller than $\phi(x_k; \sigma_{k+1})$, and we say that the k th iterate is a p-iterate. Moreover, if $x_k + \alpha_k \hat{s}_k$ is acceptable to the current filter, we signal a return to filter mode by setting \mathcal{P} -mode to false. Otherwise, \mathcal{P} -mode remains true and penalty mode continues.

In contrast, if \mathcal{P} -mode has the value false, we perform a backtracking linesearch until we find either a pair (α_k, \hat{s}_k) with $\hat{s}_k \in \{s_k, s_k^a\}$ that is a v-pair or o-pair, or a pair (α_k, s_k) that is a b-pair. A v-pair is defined as follows.

V-PAIR	
<i>The pair (α, s) constitutes a v-pair if $x_k + \alpha s$ is acceptable to \mathcal{F}_k augmented by x_k and</i>	
$\Delta \ell^f(s_k; x_k) < \gamma_v \Delta \ell^v(s_k; x_k) \text{ for some } \gamma_v \in (0, 1). \quad (2.31)$	

A v-pair (α_k, \hat{s}_k) earns its name since the step $x_k + \alpha_k \hat{s}_k$ is acceptable to the current filter augmented by x_k , but the step s_k did not predict sufficient decrease in f ; we say that k is a v-iterate. Note that the focus of the iteration is on reducing the violation v and that we add the pair (v_k, f_k) to the filter \mathcal{F}_k . An o-pair is characterized as follows.

O-PAIR	
<i>The pair (α, s) constitutes an o-pair if $x_k + \alpha s$ is acceptable to \mathcal{F}_k,</i>	
$\Delta \ell^f(s_k; x_k) \geq \gamma_v \Delta \ell^v(s_k; x_k), \text{ and} \quad (2.32a)$	
$f(x_k + \alpha s) \leq f(x_k) - \gamma_f \alpha \rho_k^f, \quad (2.32b)$	
<i>where $\gamma_v \in (0, 1)$ is the same constant used to define a v-pair and</i>	
$\rho_k^f := \min \left[\Delta \ell^f(s_k; x_k), \Delta q^f(s_k^{cf}; x_k, H_k) \right]. \quad (2.33)$	

An o-pair (α_k, \hat{s}_k) is so designated since $x_k + \alpha_k \hat{s}_k$ is acceptable to the filter, s_k predicts decrease in the objective function, and a sufficient decrease in the objective is realized; we say that k is an o-iterate. Finally, a b-pair is formalized as follows.

B-PAIR	
<i>The pair (α, s) constitutes a b-pair if (2.29) holds and</i>	
$v(x_k + \alpha s) < v(x_k). \quad (2.34)$	

An iterate $x_k + \alpha_k s_k$ associated with a b-pair (α_k, s_k) decreases both the constraint violation and penalty function, and thus suggests that one or more filter entries is blocking a productive step; we say that k is

a b-iterate. Our strategy is to accept the point, add (v_k, f_k) to the filter, and enter penalty mode. We view this as a satisfying alternative to a traditional restoration phase. Note that we only add (v_k, f_k) to the filter for v- and b-iterates. Moreover, if $\mathcal{P}\text{-mode}$ has the value false at the beginning of the k th iterate, steps are always tested for acceptability based on the filter criteria (o- and v-pairs) before checking for decrease in the constraint violation and penalty function (b-pairs). In this manner, we give clear preference to staying in filter mode.

Finally, we increase the penalty parameter if

$$\Delta q^\phi(s_k; x_k, B_k, \sigma_{k+1}) < \eta_\phi \Delta q^\phi(s_k^p; x_k, B_k, \sigma_{k+1}), \quad (2.35)$$

since this indicates that τ_k is very small and the search direction s_k does not adequately reflect the decrease predicted by s_k^p in the penalty function.

We find it useful to define the following index sets:

$$\begin{aligned} \mathcal{S}_v &= \{k : k \text{ is a v-iterate}\}, & \mathcal{S}_o &= \{k : k \text{ is an o-iterate}\}, \\ \mathcal{S}_p &= \{k : k \text{ is a p-iterate}\}, & \mathcal{S}_b &= \{k : k \text{ is an b-iterate}\}. \end{aligned}$$

These definitions and the construction of Algorithm 1 allow us to prove the following important result.

LEMMA 2.14. *If $\mathcal{P}\text{-mode} = \text{false}$ at the beginning of iteration k , then x_k is acceptable to \mathcal{F}_k .*

Proof. The result immediately follows from the construction of Algorithm 1 and consideration of the possible outcomes associated with iteration $k - 1$. \square

3. Well-posedness. In this section we verify that every step of the method is well-posed under the following assumption, which we do not explicitly state for each result.

ASSUMPTION 3.1. *The functions f and c are both differentiable with Lipschitz continuous derivatives in the neighborhood of the point x_k .*

We begin by observing that the steering problem (2.10) is convex, always feasible, and the objective function is bounded below by zero, i.e., it is well-defined. Next, we argue that the predictor problem (2.11) is well-defined. This is obvious when $\Delta \ell^v(s_k^s) \neq v(x_k)$ since then the strictly convex problem (2.11b) is always feasible. On the other hand, if $\Delta \ell^v(s_k^s) = v(x_k)$, then it follows that $\|[c(x_k) + J(x_k)s_k^s]^-\|_1 = 0$, which implies that $c_k + J_k s_k^s \geq 0$. Thus, $s = s_k^s$ is feasible for (2.11a), and the predictor problem is well-defined. Lemma 2.3 shows that $\tau_k > 0$ and Lemma 2.8 shows that the update to the weighting parameter is well-defined. The accelerator problem (2.24) does not cause difficulties since by construction it is feasible, has bounded solutions, and may be solved (approximately) as noted in Section 2.5. It is also easy to see that both Cauchy step problems (2.26) and (2.27) are well-defined.

We now proceed to show that the linesearch terminates finitely. To this end, we first show that feasible iterates are never added to the filter.

LEMMA 3.1. *Algorithm 1 ensures that if (v_k, f_k) is added to the filter, then $v_k > 0$.*

Proof. For a proof by contradiction, suppose that $v(x_k) = 0$. It follows from $v(x_k) = 0$ and the fact that ℓ^v is a convex function that $\Delta \ell^v(s_k^s, x_k) = 0$, and we may then use (2.11a), (2.14), (2.15), and the fact that x_k is not a KKT point for (1.1) (otherwise we would already have exited on Line 10 of Algorithm 1) to show that

$$\tau_k = 1, \quad s_k = s_k^p \neq 0, \quad \text{and} \quad \Delta \ell^v(s_k; x_k) = \Delta \ell^v(s_k^p; x_k) = 0. \quad (3.1)$$

It then follows from (3.1), (2.7), Lemma 2.7, $v(x_k) = 0$, and the fact that $B_k \succ 0$ that

$$\Delta \ell^f(s_k; x_k) = \Delta \ell^f(s_k^p; x_k) = \Delta \ell^\phi(s_k^p; x_k, \sigma_{k+1}) = \Delta \ell^\phi(s_k; x_k, \sigma_{k+1}) \geq \frac{1}{2} s_k^{pT} B_k s_k^p > 0. \quad (3.2)$$

Since (v_k, f_k) was added to the filter, it follows from the construction of Algorithm 1 that either (α_k, \hat{s}_k) is a v-iterate or (α_k, s_k) is a b-pair, which implies that at least one of $v(x_k + \alpha_k s_k) < v(x_k)$ or $\Delta \ell^f(s_k; x_k) <$

Algorithm 1 Filter sequential quadratic programming algorithm.

```

1: Input an initial primal-dual pair  $(x_0, y_0)$ .
2: Choose parameters  $\{\eta_v, \eta_\sigma, \eta_\phi, \sigma_{\text{inc}}, \gamma, \gamma_v, \gamma_f, \gamma_\phi\} \subset (0, 1)$  and  $0 < \delta_{\min} \leq \delta_{\max} < \infty$ .
3: Set  $k \leftarrow 0$ ,  $\mathcal{F}_0 \leftarrow \emptyset$ ,  $\mathcal{P}\text{-mode} \leftarrow \text{false}$ , and then choose  $\sigma_0 > 0$  and  $\delta_0 \in [\delta_{\min}, \delta_{\max}]$ .
4: loop
5:   Compute  $s_k^s$  as a solution of (2.9), and then calculate  $\Delta\ell^v(s_k^s; x_k)$  from (2.5).
6:   if  $\Delta\ell^v(s_k^s; x_k) = 0$  and  $v(x_k) > 0$ , then
7:     return with the infeasible stationary point  $x_k$  for problem (1.1).
8:   Choose  $B_k \succ 0$  and then compute  $s_k^p$  as the unique solution of (2.11) with multiplier  $y_k^p$ .
9:   if  $\Delta q^\phi(s_k^p; x_k, \sigma_k) = v(x_k) = 0$ , then
10:    return with the KKT point  $(x_k, y_k^p)$  for problem (1.1).
11:   Compute  $s_k$  from (2.14) such that (2.15) is satisfied.
12:   Compute the new weight  $\sigma_{k+1}$  from (2.18).
13:   Choose  $\delta_k^a > 0$  and then compute  $s_k^a$  as an (approximate) solution of (2.24).
14:   Compute  $s_k^{c\phi}$  from (2.27) and then calculate  $\Delta q^\phi(s_k^{c\phi}; x_k, H_k, \sigma_{k+1})$  from (2.8).
15:   if  $\mathcal{P}\text{-mode}$  then
16:     for  $j = 0, 1, 2, \dots$  do
17:       Set  $\alpha_k = \gamma^j$ .
18:       for  $\hat{s}_k \in \{s_k^a, s_k\}$  do
19:         if  $(\alpha_k, \hat{s}_k)$  is a p-pair then
20:           Set  $\mathcal{F}_{k+1} \leftarrow \mathcal{F}_k$  and go to Line 21. ▷ p-iterate
21:         if  $x_k + \alpha_k \hat{s}_k$  is acceptable to  $\mathcal{F}_k$  then
22:           Set  $\mathcal{P}\text{-mode} \leftarrow \text{false}$ .
23:     else
24:       Compute  $s_k^{cf}$  from (2.26) and then calculate  $\Delta q^f(s_k^{cf}; x_k, H_k)$  from (2.6b).
25:       for  $j = 0, 1, 2, \dots$  do
26:         Set  $\alpha_k \leftarrow \gamma^j$ .
27:         for  $\hat{s}_k \in \{s_k^a, s_k\}$  do
28:           if  $(\alpha_k, \hat{s}_k)$  is a v-pair then
29:             Set  $\mathcal{F}_{k+1} \leftarrow \mathcal{F}_k \cup \{(v_k, f_k)\}$  and go to Line 34. ▷ v-iterate
30:           if  $(\alpha_k, \hat{s}_k)$  is an o-pair then
31:             Set  $\mathcal{F}_{k+1} \leftarrow \mathcal{F}_k$  and go to Line 34. ▷ o-iterate
32:           if  $(\alpha_k, s_k)$  is a b-pair then
33:             Set  $\mathcal{F}_{k+1} \leftarrow \mathcal{F}_k \cup \{(v_k, f_k)\}$ ,  $\mathcal{P}\text{-mode} \leftarrow \text{true}$ , and go to Line 34. ▷ b-iterate
34:       if  $\Delta q^\phi(s_k; x_k, B_k, \sigma_{k+1}) < \eta_\phi \Delta q^\phi(s_k^p; x_k, B_k, \sigma_{k+1})$  then
35:         Set  $\sigma_{k+1} \leftarrow \sigma_{k+1} + \sigma_{\text{inc}}$ .
36:       Set  $x_{k+1} \leftarrow x_k + \alpha_k \hat{s}_k$ ,  $y_{k+1} \leftarrow y_k^p$ ,  $\delta_{k+1} \in [\delta_{\min}, \delta_{\max}]$ , and  $k \leftarrow k + 1$ .

```

$\gamma_v \Delta\ell^v(s_k; x_k)$ holds, amongst other requirements. However, since $v(x_k + \alpha_k s_k) < v(x_k) = 0$ is not possible, we conclude that $\Delta\ell^f(s_k; x_k) < \gamma_v \Delta\ell^v(s_k; x_k) = 0$, where we have also used (3.1); this contradicts (3.2) and proves the result. \square

The next two results show that our linesearch procedure terminates anytime $\mathcal{P}\text{-mode}$ has the value false at the beginning of the k th iteration. We first consider the case when x_k is feasible.

LEMMA 3.2. *If $\mathcal{P}\text{-mode} = \text{false}$ at the beginning of the k th iteration, $v(x_k) = 0$, and x_k is not a first-order solution to problem (1.1), then the pair (α, s_k) is an o-pair for all $\alpha > 0$ sufficiently small. Moreover, $k \in \mathcal{S}_o$.*

Proof. As in the proof of Lemma 3.1, it follows that $v(x_k) = \Delta\ell^v(s_k^s; x_k) = 0$. This may be

combined with the fact that x_k is assumed to not be a first-order solution to (1.1), (2.11a), (2.14), (2.15), Lemma 2.7, $B_k \succ 0$, and the definition of $\Delta\ell^\phi$ to conclude that

$$s_k = s_k^p \neq 0, \quad c_k + J_k s_k \geq 0, \quad \text{and} \quad \Delta\ell^f(s_k; x_k) = \Delta\ell^\phi(s_k; x_k, \sigma_{k+1}) > 0 = \gamma_v \Delta\ell^v(s_k; x_k). \quad (3.3)$$

Next, $v(x_k) = 0$ and (3.3) imply that $c_k + \alpha J_k s_k \geq 0$ for all $\alpha \in [0, 1]$. Combining this fact with Taylor's Theorem, Assumption 3.1, and (3.3) yields

$$v(x_k + \alpha s_k) = \|[c(x_k + \alpha s_k)]^-\|_1 = \|[c_k + \alpha J_k s_k + O(\alpha^2)]^-\|_1 \leq O(\alpha^2) \quad \text{for } \alpha \in [0, 1]. \quad (3.4)$$

Since Lemma 3.1 implies that $v_i > 0$ for all $(v_i, f_i) \in \mathcal{F}_k$, we may conclude from (3.4) that

$$v(x_k + \alpha s_k) \leq \min_{(v_i, f_i) \in \mathcal{F}_k} \beta v_i \quad \text{for all } \alpha > 0 \text{ sufficiently small,}$$

where $\beta \in (0, 1)$ is defined in (2.28), so that

$$x_k + \alpha s_k \text{ is acceptable to the filter for all } \alpha > 0 \text{ sufficiently small.} \quad (3.5)$$

Next, Taylor's Theorem, Assumption 3.1, the definition of $\Delta\ell^f$, and (3.3) imply that

$$\begin{aligned} f(x_k + \alpha s_k) &= f_k + \alpha g_k^T s_k + O(\alpha^2) = f_k - \alpha \Delta\ell^f(s_k; x_k) + O(\alpha^2) \\ &\leq f_k - \gamma_f \alpha \Delta\ell^f(s_k; x_k) \quad \text{for all } \alpha > 0 \text{ sufficiently small,} \end{aligned} \quad (3.6)$$

where $\gamma_f \in (0, 1)$ is defined in (2.32b). It follows from (3.3), (3.5), and (3.6) that (α, s_k) is an o-pair for all $\alpha > 0$ sufficiently small, which proves the first result of this lemma.

We just proved that the **for** loop on line 25 in Algorithm 1 always terminates. Moreover, it can never terminate as a result of the **if** on line 32 since $v(x_k + \alpha s_k) < v(x_k) = 0$ is impossible for all α . Moreover, it can not terminate on line 28 since (3.3) holds. Therefore, the linesearch must terminate with an o-pair (α_k, \hat{s}_k) , which implies that $k \in \mathcal{S}_o$. \square

We now consider the case when x_k is infeasible.

LEMMA 3.3. *If \mathcal{P} -mode = **false** at the beginning of iteration k , $v(x_k) > 0$, and x_k is not an infeasible stationary point, then (α, s_k) is a b-pair for all $\alpha > 0$ sufficiently small.*

Proof. It follows from the assumptions of this lemma and Lemma 2.10 that

$$[D_{s_k} \phi](x_k; \sigma_{k+1}) \leq -\Delta\ell^\phi(s_k; x_k, \sigma_{k+1}) < 0 \quad (3.7)$$

so that the direction s_k is a strict descent direction for ϕ at x_k with penalty parameter σ_{k+1} . Using the definition of the directional derivative, (3.7), $\gamma_\phi \in (0, 1)$ defined in (2.29), and (2.30) we conclude that

$$\begin{aligned} \phi(x_k + \alpha s_k; \sigma_{k+1}) &\leq \phi(x_k; \sigma_{k+1}) + \alpha \gamma_\phi [D_{s_k} \phi](x_k; \sigma_{k+1}) \\ &\leq \phi(x_k; \sigma_{k+1}) - \alpha \gamma_\phi \Delta\ell^\phi(s_k; x_k, \sigma_{k+1}) \\ &\leq \phi(x_k; \sigma_{k+1}) - \alpha \gamma_\phi \rho_k^\phi \quad \text{for all } \alpha > 0 \text{ sufficiently small.} \end{aligned} \quad (3.8)$$

Since $v(x_k) \neq 0$ and x_k is not an infeasible stationary point, we know that $\Delta\ell^v(s_k^s; x_k) > 0$. Lemma 2.6 then implies that

$$[D_{s_k} v](x_k) \leq -\Delta\ell^v(s_k; x_k) \leq -\eta_v \Delta\ell^v(s_k^s; x_k) < 0$$

so that s_k is a descent direction for v at x_k . A similar argument as the one that lead to (3.8), yields

$$v(x_k + \alpha s_k) < v(x_k) \quad \text{for all } \alpha > 0 \text{ sufficiently small.} \quad (3.9)$$

It follows from (3.8) and (3.9) that (α, s_k) is a b-pair for all $\alpha > 0$ sufficiently small. \square

The next lemma considers the case when \mathcal{P} -mode is true at the beginning of the k th iteration, and shows that successful trial iterates may be obtained through backtracking as performed in Algorithm 1.

LEMMA 3.4. *If \mathcal{P} -mode = **true** at the beginning of the k th iteration and x_k is neither an infeasible stationary point nor a first-order solution to problem (1.1), then (α, s_k) is a p-pair for all $\alpha > 0$ sufficiently small.*

Proof. The proof follows exactly as in the first part of Lemma 3.3. \square

We now combine these results to prove that Algorithm 1 is well-posed.

THEOREM 3.5. *Algorithm 1 is well-posed.*

Proof. As described in the first paragraph of Section 3, every subproblem and step computation is well defined, and Lemma 2.8 ensures that the update to the weighting parameter is well defined.

All that remains is to prove that the linesearch terminates. First, if \mathcal{P} -mode has the value false at the beginning of iteration x_k and $v(x_k) = 0$, then Lemma 3.2 guarantees finite termination and that $k \in \mathcal{S}_o$. Second, if \mathcal{P} -mode has the value false and $v(x_k) > 0$, then Lemma 3.3 ensures that the backtracking linesearch will terminate finitely. Finally, suppose that \mathcal{P} -mode has the value true at the beginning of iteration k . It then follows from Lemma 3.4 that the backtracking terminates finitely. \square

4. Global Convergence. In this section we prove that limit points of the iterates generated by Algorithm 1 have desirable properties. To this end, we use the following common assumptions.

ASSUMPTION 4.1. *The iterates $\{x_k\}$ lie in an open, bounded, convex set \mathcal{X} .*

ASSUMPTION 4.2. *The problem functions $f(x)$ and $c(x)$ are twice continuously differentiable on \mathcal{X} .*

ASSUMPTION 4.3. *The matrices B_k are uniformly positive definite and bounded, i.e., there exists values $0 < \lambda_{\min} < \lambda_{\max} < \infty$ such that $\lambda_{\min} \|s\|_2^2 \leq s^T B_k s \leq \lambda_{\max} \|s\|_2^2$ for all $s \in \mathbb{R}^n$ and all B_k .*

ASSUMPTION 4.4. *The matrices H_k are uniformly bounded, i.e., $\|H_k\|_2 \leq \mu_{\max}$ for some $\mu_{\max} \geq 1$.*

For clarity and motivational purposes, we immediately state our main convergence theorem that makes use of the Mangasarian-Fromovitz constraint qualification (MFCQ) [31].

THEOREM 4.1. *If Assumptions 4.1–4.4 hold, then one of the following must occur.*

- (i) *Algorithm 1 terminates finitely with either a first-order KKT point or an infeasible stationary point in lines 10 or 7, respectively, for problem (1.1).*
- (ii) *Algorithm 1 generates infinitely many iterations $\{x_k\}$, $\sigma_k = \bar{\sigma} < \infty$ for all k sufficiently large, and there exists a limit point x_* of $\{x_k\}$ that is either a first-order KKT point or an infeasible stationary point for problem (1.1).*
- (iii) *Algorithm 1 generates infinitely many iterations $\{x_k\}$, $\lim_{k \rightarrow \infty} \sigma_k = \infty$, and there exists a limit point x_* of $\{x_k\}$ that is either an infeasible stationary point or a feasible point at which the MFCQ fails.*

Proof. The result follows from the following analysis that considers the various cases that can occur. In particular, it follows from Theorems 4.11, 4.14, 4.17, 4.20, and the construction of Algorithm 1. \square

We now present a sequence of lemmas that will be useful in the convergence analysis. The first result is adapted from [7, Theorem 3.6] and provides a bound on the trial step s_k .

LEMMA 4.2. *If Assumptions 4.1–4.3 hold and x_k and s_k are generated by Algorithm 1, then*

$$\|s_k\|_2 \leq \max \left\{ 1, \frac{2}{\lambda_{\min}} [\|g_k\|_2 + \sigma_k v(x_k)], \sqrt{n} \delta_{\max} \right\}. \quad (4.1)$$

Furthermore, if $\{\sigma_k\}$ is bounded, then there exists a constant $M_s > 0$ such that $\|s_k\|_2 \leq M_s$ for all k .

Proof. First, we claim that the predictor step s_k^p must satisfy

$$\|s_k^p\|_2 \leq \max \left\{ 1, \frac{2}{\lambda_{\min}} [\|g_k\|_2 + \sigma_k v(x_k)] \right\}, \quad (4.2)$$

which can be seen as follows. Suppose that (4.2) is not satisfied so that

$$\|s_k^p\|_2 > 1 \quad \text{and} \quad \frac{1}{2} \lambda_{\min} \|s_k^p\|_2 > \|g_k\|_2 + \sigma_k v(x_k). \quad (4.3)$$

It then follows from the definitions of Δq^ϕ and ℓ^v , the Cauchy-Schwarz inequality, Assumption 4.3, and (4.3) that

$$\begin{aligned}\Delta q^\phi(s_k^p; x_k, B_k, \sigma_k) &= -g_k^T s_k^p - \frac{1}{2} s_k^{pT} B_k s_k^p + \sigma_k (\ell^v(0; x_k) - \ell^v(s_k^p; x_k)) \\ &\leq \|g_k\|_2 \|s_k^p\|_2 - \frac{1}{2} \lambda_{\min} \|s_k^p\|_2^2 + \sigma_k v(x_k) \\ &\leq \|g_k\|_2 \|s_k^p\|_2 - \frac{1}{2} \lambda_{\min} \|s_k^p\|_2^2 + \|s_k^p\|_2 \sigma_k v(x_k) \\ &= \|s_k^p\|_2 \left(\|g_k\|_2 - \frac{1}{2} \lambda_{\min} \|s_k^p\|_2 + \sigma_k v(x_k) \right) < 0,\end{aligned}$$

which contradicts the fact that s_k^p is the unique global minimizer to the strictly convex predictor problem. Thus, (4.2) must hold and when combined with (2.14), the use of the triangle-inequality, the use of the trust-region radius $\delta_k \in [\delta_{\min}, \delta_{\max}]$ in the steering problem, implies that

$$\|s_k\|_2 \leq \max \left\{ 1, \frac{2}{\lambda_{\min}} [\|g_k\|_2 + \sigma_k v(x_k)], \sqrt{n} \delta_{\max} \right\} \quad (4.4)$$

which proves (4.1). Since g_k and $v(x_k)$ are uniformly bounded as a result of Assumptions 4.1 and 4.2, it is clear that if $\{\sigma_k\}$ is bounded, then there exists $M_s < \infty$ such that $\|s_k\|_2 \leq M_s$ for all k . \square

The following result provides a relationship between the predicted change in the linear model and the change achieved in the line search process for both the objective function and the constraint violation.

LEMMA 4.3. (Equivalent to [39, Lemma 3]) Suppose that Assumptions 4.1 and 4.2 hold. Then, there exist constants $\{C_f, C_v\} > 0$ such that for all k and $\alpha \in (0, 1]$, we have

$$f(x_k + \alpha s) \leq f(x_k) - \alpha \Delta \ell^f(s; x_k) + \alpha^2 C_f \|s\|_2^2 \quad (4.5)$$

and

$$v(x_k + \alpha s) \leq v(x_k) - \alpha \Delta \ell^v(s; x_k) + \alpha^2 C_v \|s\|_2^2. \quad (4.6)$$

Proof. Inequality (4.5) is a direct result of Taylor's theorem and Assumption 4.2.

For (4.6), it follows from the integral mean-value theorem, Assumptions 4.1 and 4.2 and their the implied Lipschitz continuity of $J(x)$, the triangle inequality, and the convexity of ℓ^v , that for some constant Lipschitz constant C ,

$$\begin{aligned}v(x_k + \alpha s) &= \|[c(x_k + \alpha s)]^-\|_1 = \left\| \left[c(x_k) + \alpha J_k s + \alpha \int_0^1 [J(x_k + \theta \alpha s) - J(x_k)] s d\theta \right]^-\right\|_1 \\ &\leq \|[c(x_k) + \alpha J_k s]^-\|_1 + \alpha^2 \sqrt{n} C \|s\|_2^2 \\ &\leq (1 - \alpha) \|[c(x_k)]^-\|_1 + \alpha \|[c(x_k) + J_k s]^-\|_1 + \alpha^2 \sqrt{n} C \|s\|_2^2 \\ &= v(x_k) - \alpha \Delta \ell^v(s; x_k) + \alpha^2 \sqrt{n} C \|s\|_2^2 \quad \text{for all } \alpha \in (0, 1].\end{aligned}$$

This proves (4.6) by defining $C_v := \sqrt{n} C$. \square

The next two lemmas provide a relationship between the predicted linear decrease in the objective function and the quantity ρ_k^f defined by (2.33).

LEMMA 4.4. If Assumption 4.4 holds and $\Delta \ell^f(s_k; x_k) \geq 0$, then

$$\Delta q^f(s_k^{cf}; x_k, H_k) \geq \frac{1}{2} \Delta \ell^f(s_k; x_k) \min \left\{ \frac{\Delta \ell^f(s_k; x_k)}{\mu_{\max} \|s_k\|_2^2}, 1 \right\}. \quad (4.7)$$

Proof. If $\Delta \ell^f(s_k; x_k) = 0$, then the result follows immediately from the definition of s_k^{cf} in (2.26). Now, suppose that $\Delta \ell^f(s_k; x_k) > 0$. It follows from (2.26) and the definition of Δq^f that

$$\Delta q^f(s_k^{cf}; x_k, H_k) \geq \Delta q^f(\alpha s_k; x_k, H_k) = -\alpha g_k^T s_k - \frac{1}{2} \alpha^2 s_k^T H_k s_k \text{ for all } 0 \leq \alpha \leq 1.$$

The right hand side of the previous equation may be written as

$$q(\alpha) = a\alpha^2 + b\alpha, \text{ where } a = -\frac{1}{2} s_k^T H_k s_k \text{ and } b = \Delta \ell^f(s_k; x_k) = -g_k^T s_k > 0.$$

We wish to maximize q on the interval $[0, 1]$ so we differentiate $q(\alpha)$ with respect to α and set the result to zero to obtain a stationary point at $-\frac{b}{2a}$. Now, consider three cases.

Case 1: ($a < 0$ and $-\frac{b}{2a} \leq 1$) The maximum of $q(\alpha)$ on the interval $[0, 1]$ is achieved at $\alpha = -\frac{b}{2a}$. Note that $\alpha > 0$, since $b = \Delta \ell^f(s_k; x_k) > 0$ by assumption. Then, we have

$$q\left(-\frac{b}{2a}\right) = a \frac{b^2}{4a^2} - b \frac{b}{2a} = -\frac{b^2}{4a}.$$

It follows from the definition of a and b , the Cauchy-Schwarz inequality, and Assumption 4.4 that

$$q\left(-\frac{b}{2a}\right) = \frac{\Delta \ell^f(s_k; x_k)^2}{2s_k^T H_k s_k} \geq \frac{\Delta \ell^f(s_k; x_k)^2}{2\|H_k\|_2 \|s_k\|_2^2} \geq \frac{\Delta \ell^f(s_k; x_k)^2}{2\mu_{\max} \|s_k\|_2^2}.$$

Case 2: ($a < 0$ and $-\frac{b}{2a} > 1$) The maximum of $q(\alpha)$ on the interval $[0, 1]$ is achieved at $\alpha = 1$, where

$$q(1) = a + b > -\frac{1}{2}b + b = \frac{1}{2}b = \frac{1}{2}\Delta \ell^f(s_k; x_k).$$

Case 3: ($a \geq 0$) The maximum of $q(\alpha)$ on the interval $[0, 1]$ is achieved at $\alpha = 1$ so that

$$q(1) = a + b > b > \frac{1}{2}b = \frac{1}{2}\Delta \ell^f(s_k; x_k).$$

Finally, combining all three cases and defining $\alpha' = \arg \max_{\alpha \in [0, 1]} q(\alpha)$, it follows that

$$\Delta q^f(s_k^{cf}; x_k, H_k) = q(\alpha') \geq \min \left\{ \frac{\Delta \ell^f(s_k; x_k)^2}{2\mu_{\max} \|s_k\|_2^2}, \frac{1}{2}\Delta \ell^f(s_k; x_k) \right\} = \frac{1}{2}\Delta \ell^f(s_k; x_k) \min \left\{ \frac{\Delta \ell^f(s_k; x_k)}{\mu_{\max} \|s_k\|_2^2}, 1 \right\}$$

as desired. \square

LEMMA 4.5. *Suppose that the Assumptions 4.1–4.4 are satisfied and that $\{\sigma_k\}$ is bounded. Then, there exists a constant $C_\rho > 0$ such that whenever $\Delta \ell^f(s_k; x_k) \geq 0$, it follows that*

$$\rho_k^f \geq \min \left[C_\rho \Delta \ell^f(s_k; x_k)^2, \frac{1}{2} \Delta \ell^f(s_k; x_k) \right]. \quad (4.8)$$

Proof. It follows from (2.33), Lemma 4.4, Lemma 4.2 and the assumption $\Delta \ell^f(s_k; x_k) \geq 0$ that

$$\begin{aligned} \rho_k^f &= \min \left[\Delta \ell^f(s_k; x_k), \Delta q^f(s_k^{cf}; x_k, H_k) \right] \\ &\geq \min \left[\Delta \ell^f(s_k; x_k), \min \left\{ \frac{\Delta \ell^f(s_k; x_k)^2}{2\mu_{\max} \|s_k\|_2^2}, \frac{1}{2}\Delta \ell^f(s_k; x_k) \right\} \right] \\ &\geq \min \left[\frac{\Delta \ell^f(s_k; x_k)^2}{2\mu_{\max} \|s_k\|_2^2}, \frac{1}{2}\Delta \ell^f(s_k; x_k) \right] \geq \min \left[\frac{\Delta \ell^f(s_k; x_k)^2}{2\mu_{\max} M_s^2}, \frac{1}{2}\Delta \ell^f(s_k; x_k) \right], \end{aligned}$$

where $\{M_s, \mu_{\max}\} \subset (0, \infty)$ are defined in (4.1) and Assumption 4.4, respectively. The result now follows by defining $C_\rho := 1/(2\mu_{\max}M_s^2)$. \square

The next two results provide a relationship between the predicted linear change in the penalty function and the quantity ρ_k^ϕ defined by (2.30).

LEMMA 4.6. *If Assumption 4.4 holds and x_k is not an infeasible stationary point, then*

$$\Delta q^\phi(s_k^{c\phi}; x_k, H_k, \sigma_{k+1}) \geq \frac{1}{2} \Delta \ell^\phi(s_k; x_k, \sigma_{k+1}) \min \left\{ \frac{\Delta \ell^\phi(s_k; x_k, \sigma_{k+1})}{\mu_{\max} \|s_k\|_2^2}, 1 \right\}. \quad (4.9)$$

Proof. Since x_k is not an infeasible stationary point by assumption, it follows from Lemma 2.8 that $\Delta \ell^\phi(s_k; x_k, \sigma_{k+1}) \geq 0$. If $\Delta \ell^\phi(s_k; x_k, \sigma_{k+1}) = 0$, then the result follows immediately. Therefore, for the remainder of the proof we assume that $\Delta \ell^\phi(s_k; x_k, \sigma_{k+1}) > 0$.

It follows from (2.27), the convexity of $\ell^v(\cdot)$, and simple algebra that

$$\begin{aligned} \Delta q^\phi(s_k^{c\phi}; x_k, H_k, \sigma_{k+1}) &\geq \Delta q^\phi(\alpha s_k; x_k, H_k, \sigma_{k+1}) \\ &= -\alpha g_k^T s_k - \frac{1}{2} \alpha^2 s_k^T H_k s_k + \sigma_{k+1} (\| [c_k]^- \|_1 - \| [c_k + \alpha J_k s_k]^- \|_1) \\ &\geq -\alpha g_k^T s_k - \frac{1}{2} \alpha^2 s_k^T H_k s_k + \sigma_{k+1} (\| [c_k]^- \|_1 - \alpha \| [c_k + J_k s_k]^- \|_1 - (1 - \alpha) \| [c_k]^- \|_1) \\ &= -\alpha g_k^T s_k - \frac{1}{2} \alpha^2 s_k^T H_k s_k + \alpha \sigma_{k+1} (\| [c_k]^- \|_1 - \| [c_k + J_k s_k]^- \|_1) \\ &= \alpha \Delta \ell^\phi(s_k; x_k, \sigma_{k+1}) - \frac{1}{2} \alpha^2 s_k^T H_k s_k \text{ for all } \alpha \in [0, 1]. \end{aligned}$$

The right hand side of the equation is a quadratic function of α :

$$q(\alpha) = a\alpha^2 + b\alpha, \text{ where } a = -\frac{1}{2} s_k^T H_k s_k \text{ and } b = \Delta \ell^\phi(s_k; x_k, \sigma_{k+1}) > 0.$$

Analysis similar to that used in the proof of Lemma 4.4 yields

$$\Delta q^\phi(s_k^{c\phi}; x_k, H_k, \sigma_{k+1}) \geq \min \left\{ \frac{\Delta \ell^\phi(s_k; x_k, \sigma_{k+1})^2}{2\mu_{\max} \|s_k\|_2^2}, \frac{1}{2} \Delta \ell^\phi(s_k; x_k, \sigma_{k+1}) \right\}, \quad (4.10)$$

where μ_{\max} is from Assumption 4.4, as desired. \square

LEMMA 4.7. *Suppose that Assumptions 4.1–4.4 are satisfied, that Algorithm 1 never encounters an infeasible stationary point, and $\{\sigma_k\}$ is bounded. Then, there exists a constant $C_\rho \in (0, \infty)$ such that*

$$\rho_k^\phi \geq \min [C_\rho \Delta \ell^\phi(s_k; x_k, \sigma_{k+1})^2, \frac{1}{2} \Delta \ell^\phi(s_k; x_k, \sigma_{k+1})] \text{ for all } k \geq 0.$$

Proof. The proof follows exactly as in Lemma 4.5. \square

4.1. Convergence analysis under bounded weighting parameter. In this section we study Algorithm 1 under the assumption that the weighting parameter stays bounded. It follows from this assumption and Lemma 4.2 that there exists some k' and $\bar{\sigma} < \infty$ such that

$$\|s_k\|_2 \leq M_s < \infty \quad \text{and} \quad \sigma_k = \bar{\sigma} < \infty \quad \text{for all } k \geq k'. \quad (4.11)$$

In certain situations, we can ensure that the line search step length is bounded away from zero.

LEMMA 4.8. *If Assumptions 4.1–4.3 and (4.11) hold and $\epsilon > 0$, then the following hold:*

(i) *There exists a constant $\alpha_P > 0$ such that $\alpha_k \geq \alpha_P > 0$ for all $k \in \mathcal{K}_P$, where*

$$\mathcal{K}_P = \{k \in \mathcal{S}_P : k \geq k' \text{ and } \Delta \ell^\phi(s_k; x_k, \bar{\sigma}) \geq \epsilon\}.$$

(ii) There exists a constant $\alpha_F > 0$ such that $\alpha_k \geq \alpha_F > 0$ for all $k \in \mathcal{K}_F$, where

$$\mathcal{K}_F = \{k \in \mathcal{S}_v \cup \mathcal{S}_o \cup \mathcal{S}_b : k \geq k' \text{ and } \Delta \ell^v(s_k^s; x_k) \geq \epsilon\}.$$

(iii) There exists a constant $\alpha_f > 0$ such that $(\alpha, s) = (\alpha, s_k)$ satisfies (2.32b) for all $0 < \alpha \leq \alpha_f$ and all $k \in \mathcal{K}_f$, where

$$\mathcal{K}_f = \{k \geq k' : \Delta \ell^f(s_k; x_k) \geq \epsilon\}.$$

Proof. From [6, Lemma 3.4], there exists some positive constant C_ϕ such that

$$|\phi(x_k + \alpha s_k; \bar{\sigma}) - \ell^\phi(\alpha s_k; x_k, \bar{\sigma})| \leq C_\phi \|\alpha s_k\|_2^2 \text{ for all } k \geq k' \text{ and } \alpha \in [0, 1]. \quad (4.12)$$

We first prove part (i). Suppose that α satisfies

$$0 \leq \alpha \leq \frac{(1 - \gamma_\phi)\epsilon}{C_\phi M_s^2}, \quad (4.13)$$

where $\gamma_\phi \in (0, 1)$ is set in Algorithm 1 and M_s is defined in (4.11). To simplify notation, we define $\ell_k^\phi(s) := \ell^\phi(s; x_k, \bar{\sigma})$ and $\Delta \ell_k^\phi(s) := \Delta \ell^\phi(s; x_k, \bar{\sigma})$. We then use $\phi(x_k; \bar{\sigma}) = \ell_k^\phi(0)$, the convexity of $\ell_k^\phi(\cdot)$, (4.12), $\Delta \ell_k^\phi(s_k) \geq \epsilon$ for $k \in \mathcal{K}_P$, (4.11), (4.13), and (2.30) to conclude that

$$\begin{aligned} \phi(x_k; \bar{\sigma}) - \phi(x_k + \alpha s_k; \bar{\sigma}) &= [\ell_k^\phi(0) - \ell_k^\phi(\alpha s_k)] - [\phi(x_k + \alpha s_k; \bar{\sigma}) - \ell_k^\phi(\alpha s_k)] \\ &\geq [\ell_k^\phi(0) - \alpha \ell_k^\phi(s_k) - (1 - \alpha)\ell_k^\phi(0)] - C_\phi \alpha^2 \|s_k\|_2^2 \\ &= \alpha [\ell_k^\phi(0) - \ell_k^\phi(s_k)] - C_\phi \alpha^2 \|s_k\|_2^2 \\ &= \gamma_\phi \alpha \Delta \ell_k^\phi(s_k) + (1 - \gamma_\phi) \alpha \Delta \ell_k^\phi(s_k) - C_\phi \alpha^2 \|s_k\|_2^2 \\ &\geq \gamma_\phi \alpha \Delta \ell_k^\phi(s_k) + (1 - \gamma_\phi) \alpha \epsilon - C_\phi \alpha^2 \|s_k\|_2^2 \\ &\geq \gamma_\phi \alpha \Delta \ell_k^\phi(s_k) \geq \gamma_\phi \alpha \rho_k^\phi \text{ for all } k \in \mathcal{K}_P, \end{aligned}$$

which with (2.29) implies that (α, s_k) is a p-pair. Thus, Algorithm 1 must select an α_k that satisfies

$$\alpha_k \geq \min \left\{ \frac{\gamma(1 - \gamma_\phi)\epsilon}{C_\phi M_s^2}, 1 \right\} =: \alpha_P, \quad (4.14)$$

where $\gamma \in (0, 1)$ is the backtracking parameter in Algorithm 1, which completes the proof of part (i).

We now prove part (ii). It follows from (2.19) that

$$\Delta \ell^\phi(s_k; x_k, \bar{\sigma}) \geq \bar{\sigma} \eta_\sigma \Delta \ell^v(s_k^s; x_k) \geq \bar{\sigma} \eta_\sigma \epsilon \text{ for } k \in \mathcal{K}_F. \quad (4.15)$$

If α satisfies

$$\alpha < \min \left[\frac{(1 - \gamma_\phi)\bar{\sigma} \eta_\sigma \epsilon}{C_\phi M_s^2}, \frac{\eta_v \epsilon}{C_v M_s^2} \right], \quad (4.16)$$

where C_v is defined in (4.6) and η_v is defined in (2.15), then we may use (4.15) and proceed as in the proof of part (i) to conclude that (α, s_k) is a p-pair, i.e., (2.29) holds. Moreover, we have from Lemma 4.3, (2.15), (4.16), $\Delta \ell^v(s_k^s; x_k) \geq \epsilon$ for $k \in \mathcal{K}_F$, and (4.11) that

$$\begin{aligned} v(x_k + \alpha s_k) - v(x_k) &\leq -\alpha \Delta \ell^v(s_k; x_k) + \alpha^2 C_v \|s_k\|_2^2 < -\alpha \eta_v \Delta \ell^v(s_k^s; x_k) + \alpha \frac{\eta_v \epsilon}{C_v M_s^2} C_v \|s_k\|_2^2 \\ &\leq -\alpha \eta_v \epsilon + \alpha \eta_v \epsilon = 0 \text{ for all } k \in \mathcal{K}_F, \end{aligned} \quad (4.17)$$

where the strict inequality holds since $s_k \neq 0$ as a result of (4.15). Combining (4.17) with (2.29) implies that (α, s_k) is a b-pair. Thus, we conclude from the structure of Algorithm 1 that

$$\alpha_k \geq \min \left\{ \frac{\gamma(1-\gamma_\phi)\bar{\sigma}\eta_\sigma\epsilon}{C_\phi M_s^2}, \frac{\gamma\eta_v\epsilon}{C_v M_s^2}, 1 \right\} =: \alpha_F > 0 \text{ for all } k \in \mathcal{K}_F, \quad (4.18)$$

where $\gamma \in (0, 1)$ is the backtracking parameter used in Algorithm 1.

Part (iii) is a standard result used in continuous unconstrained optimization that follows since $\Delta \ell^f(s_k; x_k) \geq \epsilon$ is equivalent to $g(x_k)^T s_k \leq -\epsilon < 0$ and s_k is uniformly bounded by (4.11). \square

The next lemma justifies the three cases that we consider when analyzing Algorithm 1.

LEMMA 4.9. *If Algorithm 1 does not terminate finitely, then one of the following scenarios occurs:*

Case 1: $k \in \mathcal{S}_p$ for all k sufficiently large;

Case 2: $k \in \mathcal{S}_o$ for all k sufficiently large; or

Case 3: $|\mathcal{S}_v \cup \mathcal{S}_b| = \infty$.

Proof. We proceed by contradiction and assume that none of the cases occur. In particular, since Case 3 does not hold it follows that $k \in \mathcal{S}_p \cup \mathcal{S}_o$ for all k sufficiently large. Combining this with the fact that Cases 1 and 2 do not hold implies that the iterates must oscillate between p- and o-iterates. However, this is not possible since there is no mechanism in Algorithm 1 that allows for iterate $k+1$ to be a p-iterate if iterate k is an o-iterate. \square

We now analyze Algorithm 1 for each of the three possible scenarios stated in the previous result.

Case 1: $k \in \mathcal{S}_p$ for all k sufficiently large. In this case, there exists k'' such that

$$k \in \mathcal{S}_p \text{ for all } k \geq k'' \geq k', \quad (4.19)$$

where k' is defined in (4.11). We first show that our measure of feasibility converges to zero.

LEMMA 4.10. *If Assumptions 4.1–4.4, (4.11), and (4.19) hold, then $\lim_{k \rightarrow \infty} \Delta \ell^v(s_k^s; x_k) = 0$.*

Proof. For a proof by contradiction, suppose that there exists an infinite subsequence

$$\mathcal{S}'' := \{k \geq k'' : \Delta \ell^v(s_k^s; x_k) \geq \epsilon''\}$$

for some constant $\epsilon'' > 0$. It follows from (4.19), (2.19), (4.11), and the definition of \mathcal{S}'' that

$$\Delta \ell^\phi(s_k; x_k, \bar{\sigma}) \geq \bar{\sigma}\eta_\sigma \Delta \ell^v(s_k^s; x_k) \geq \bar{\sigma}\eta_\sigma \epsilon'' =: \epsilon > 0 \text{ for all } k \in \mathcal{S}'', \quad (4.20)$$

which implies with (4.19) that

$$\mathcal{S}'' \subseteq \mathcal{K}_P := \{k \in \mathcal{S}_p : k \geq k' \text{ and } \Delta \ell^\phi(s_k; x_k, \bar{\sigma}) \geq \epsilon > 0\}.$$

Combining \mathcal{K}_P with Lemma 4.8 implies the existence of a positive α_P such that $\alpha_k \geq \alpha_P > 0$ for all $k \in \mathcal{S}''$, which used with the definitions of \mathcal{S}'' and \mathcal{S}_p , (4.19), Lemma 4.7, and (4.20) yield

$$\begin{aligned} \phi(x_k; \bar{\sigma}) - \phi(x_k + \alpha_k \hat{s}_k; \bar{\sigma}) &\geq \gamma_\phi \alpha_k \rho_k^\phi \geq \gamma_\phi \alpha_k \min [C_\rho \Delta \ell^\phi(s_k; x_k, \bar{\sigma})^2, \tfrac{1}{2} \Delta \ell^\phi(s_k; x_k, \bar{\sigma})] \\ &\geq \gamma_\phi \alpha_P \min [C_\rho \epsilon^2, \tfrac{1}{2} \epsilon] > 0 \text{ for all } k \in \mathcal{S}''. \end{aligned} \quad (4.21)$$

Now, for $k'' \leq k \in \mathcal{S}_p \setminus \mathcal{S}''$, it follows from (2.29), (2.19), and (2.27) that

$$\phi(x_k; \bar{\sigma}) - \phi(x_k + \alpha_k \hat{s}_k; \bar{\sigma}) \geq \gamma_\phi \alpha_k \min [\Delta \ell^\phi(s_k; x_k, \bar{\sigma}), \Delta q^\phi(s_k^{c\phi}; x_k, H_k, \bar{\sigma})] \geq 0. \quad (4.22)$$

It is now easy to see from (4.21), (4.22), and (4.19) that $\phi(x_k; \bar{\sigma}) \rightarrow -\infty$, which contradicts Assumptions 4.1 and 4.2. Thus, we conclude that $\lim_{k \rightarrow \infty} \Delta \ell^v(s_k^s; x_k) = 0$. \square

We now show that all limit points are infeasible stationary points for problem (1.1).

THEOREM 4.11. *Suppose that Assumptions 4.1–4.4, (4.11), and (4.19) hold. If x_* is any limit point of the sequence $\{x_k\}$ generated by Algorithm 1, then x_* is an infeasible stationary point for problem (1.1).*

Proof. Let $v_{\min} := \min\{v_j : (v_j, f_j) \in \mathcal{F}_{k''}\} \equiv \min\{v_j : (v_j, f_j) \in \mathcal{F}_k \text{ and } k \geq k''\}$, where the second equality holds since by assumption $k \in \mathcal{S}_p$ for all $k \geq k''$ and the filter is never expanded when $k \in \mathcal{S}_p$. It follows from Lemma 3.1 that $v_{\min} > 0$. But then if there was a feasible limit point x_* , there must be iterates x_k , $k > k''$ that are arbitrarily close to feasibility, and thus ultimately one such that x_k is acceptable to \mathcal{F}_k . Thus line 21 of Algorithm 1 implies that there will be an iterate $k > k''$ for which $k \notin \mathcal{S}_p$ which contradicts (4.19). Thus, all limit points are infeasible. It follows from this fact, Lemma 4.10, and Lemma 2.1 that all limit points are infeasible stationary points. \square

Case 2: $k \in \mathcal{S}_o$ for all k sufficiently large. In this case, there exists k'' such that

$$k \in \mathcal{S}_o \quad \text{for all } k \geq k'' \geq k', \quad (4.23)$$

where k' is defined in (4.11). We begin by showing that our feasibility measure converges to zero.

LEMMA 4.12. *If Assumptions 4.1–4.4, (4.11), and (4.23) hold, then $\lim_{k \rightarrow \infty} \Delta \ell^v(s_k^s; x_k) = 0$.*

Proof. For a contradiction, suppose that there exists $\epsilon'' > 0$ and an infinite subsequence

$$\mathcal{S}'' := \{k \geq k'' : \Delta \ell^v(s_k^s; x_k) \geq \epsilon''\} \subseteq \mathcal{S}_o,$$

where we have used k'' defined in (4.23). It then follows from the definition of \mathcal{S}_o , the o-pair (α_k, \hat{s}_k) selected in Algorithm 1, (2.32b), (4.11), Lemma 4.5, (2.32a), (2.15), and part (ii) of Lemma 4.8 that

$$\begin{aligned} f(x_k) - f(x_k + \alpha_k \hat{s}_k) &\geq \gamma_f \alpha_k \rho_k^f \geq \gamma_f \alpha_k \min \left\{ C_\rho \Delta \ell^f(s_k; x_k)^2, \frac{1}{2} \Delta \ell^f(s_k; x_k) \right\} \\ &\geq \gamma_f \alpha_k \min \left\{ C_\rho [\gamma_v \Delta \ell^v(s_k; x_k)]^2, \frac{1}{2} \gamma_v \Delta \ell^v(s_k; x_k) \right\} \\ &\geq \gamma_f \alpha_k \min \left\{ C_\rho [\gamma_v \eta_v \Delta \ell^v(s_k^s; x_k)]^2, \frac{1}{2} \gamma_v \eta_v \Delta \ell^v(s_k^s; x_k) \right\} \\ &\geq \gamma_f \alpha_F \min \left\{ C_\rho [\gamma_v \eta_v \epsilon'']^2, \frac{1}{2} \gamma_v \eta_v \epsilon'' \right\} \quad \text{for all } k \in \mathcal{S}'', \end{aligned}$$

for some $\alpha_F > 0$. Similarly, for $k'' \leq k \in \mathcal{S}_o \setminus \mathcal{S}''$, it follows from (2.32), (2.15), and (2.26) that

$$f(x_k) - f(x_k + \alpha_k \hat{s}_k) \geq \gamma_f \alpha_k \rho_k^f \geq \gamma_f \alpha_k \min \left\{ \gamma_v \Delta \ell^v(s_k; x_k), \Delta q^f(s_k^{cf}; x_k, H_k) \right\} \geq 0.$$

Combining the two previous inequalities with the definition of k'' yields $f(x_k) \rightarrow -\infty$, which contradicts the fact that f is bounded as a consequence of Assumptions 4.1 and 4.2. This proves the result. \square

We now show that feasible limit points are also first-order solutions of the penalty function.

LEMMA 4.13. *Suppose that Assumptions 4.1–4.4, (4.11), and (4.23) hold. If $x_* = \lim_{k \in \mathcal{S}} x_k$ for some subsequence \mathcal{S} and $v(x_*) = 0$, then $\lim_{k \in \mathcal{S}} \Delta q^\phi(s_k^p; x_k, B_k, \bar{\sigma}) = 0$.*

Proof. Suppose that there exists a constant $\epsilon'' > 0$ and an infinite subsequence

$$\mathcal{S}'' := \{k \in \mathcal{S} : k \geq k'' : \Delta q^\phi(s_k^p; x_k, B_k, \bar{\sigma}) \geq \epsilon''\},$$

where k'' is defined in (4.23). It follows from line 34 of Algorithm 1, (4.11), and (4.23) that

$$\Delta q^\phi(s_k; x_k, B_k, \bar{\sigma}) \geq \eta_\phi \Delta q^\phi(s_k^p; x_k, B_k, \bar{\sigma}) \geq \eta_\phi \epsilon'' \quad \text{for } k \in \mathcal{S}''. \quad (4.24)$$

From (2.5) and (2.15), we know that $v(x_k) \geq \Delta \ell^v(s_k; x_k) \geq \eta_v \Delta \ell^v(s_k^s; x_k) \geq 0$ for all k , which may be combined with $\lim_{k \in \mathcal{S}} v(x_k) = v(x_*) = 0$ (holds by assumption) to conclude that

$$\Delta \ell^v(s_k; x_k) \leq \frac{\eta_\phi \epsilon''}{\bar{\sigma} + \gamma_v} \quad \text{for } k \in \mathcal{S} \text{ sufficiently large,} \quad (4.25)$$

where $\gamma_v \in (0, 1)$ is defined in (2.32a). It follows from (2.7), (2.8), (4.24), $B_k \succ 0$, and (4.25) that

$$\begin{aligned} \Delta \ell^f(s_k; x_k) &\geq \frac{1}{2} s_k^T B_k s_k - \bar{\sigma} \Delta \ell^v(s_k; x_k) + \eta_\phi \epsilon'' > \eta_\phi \epsilon'' - \bar{\sigma} \Delta \ell^v(s_k; x_k) \\ &\geq \eta_\phi \epsilon'' - \bar{\sigma} \frac{\eta_\phi \epsilon''}{\bar{\sigma} + \gamma_v} = \frac{\gamma_v \eta_\phi \epsilon''}{\bar{\sigma} + \gamma_v} =: \epsilon^f > 0 \quad \text{for } k \in \mathcal{S}'' \text{ sufficiently large.} \end{aligned} \quad (4.26)$$

Combining this with part (iii) of Lemma 4.8, we know that there exists some $\alpha_f > 0$ such that (α, s_k) satisfies (2.32b) for all $k \in \mathcal{S}''$ sufficiently large and $\alpha \in (0, \alpha_f]$, since by assumption $\mathcal{S}_o = \mathcal{S}_v \cup \mathcal{S}_o \cup \mathcal{S}_b$ for $k \geq k''$.

Next, we define

$$\Phi_k := \min_{(v_i, f_i) \in \mathcal{F}_k} \left\{ \max \left[v_i - \alpha_i \eta_v \Delta \ell^v(s_i^s; x_i), \beta v_i \right] \right\} > 0, \quad (4.27)$$

where \mathcal{F}_k is the k th filter. The fact that $\Phi_k > 0$ follows since $v_i > 0$ for all $(v_i, f_i) \in \mathcal{F}_k$ as a consequence of Lemma 3.1. Moreover, it follows from (4.23) that $\mathcal{F}_k \equiv \mathcal{F}_{k''}$ for all $k \in \mathcal{S}''$ so that $\Phi_k \equiv \Phi_{k''} > 0$ for all $k \in \mathcal{S}''$. Now, pick $\epsilon^v > 0$ such that $\Phi_{k''} - C_v M_s^2 \leq \epsilon^v < \Phi_{k''}$ and consider α such that

$$0 < \alpha \leq \frac{\Phi_{k''} - \epsilon^v}{C_v M_s^2} \leq 1. \quad (4.28)$$

It then follows from Lemma 4.3, the fact that $\lim_{x \in \mathcal{S}} v(x_k) = 0$, (2.15), (4.23), and (4.28), that

$$\begin{aligned} v(x_k + \alpha s_k) &\leq v(x_k) - \alpha \Delta \ell^v(s_k; x_k) + \alpha^2 C_v \|s_k\|_2^2 \leq \epsilon^v + \alpha^2 C_v M_s^2 \\ &\leq \epsilon^v + \alpha C_v M_s^2 \leq \epsilon^v + \frac{\Phi_{k''} - \epsilon^v}{C_v M_s^2} C_v M_s^2 = \Phi_{k''} \quad \text{for all } k \in \mathcal{S}'' \text{ sufficiently large.} \end{aligned}$$

Thus, $x_k + \alpha s_k$ is acceptable to $\mathcal{F}_k \equiv \mathcal{F}_{k''}$ for all α satisfying (4.28) and $k \in \mathcal{S}''$ sufficiently large.

Combining the above, (4.23), and the structure of Algorithm 1, we conclude that

$$\alpha_k \geq \min \left\{ \gamma \frac{\Phi_{k''} - \epsilon^v}{C_v M_s^2}, \gamma \alpha_f, 1 \right\} =: \alpha_{\min} > 0 \quad \text{for all } k \in \mathcal{S}'' \text{ sufficiently large,} \quad (4.29)$$

where $\gamma \in (0, 1)$ is the backtracking parameter used in Algorithm 1. It then follows from (4.23), (2.32b), (2.15), Lemma 4.5, (2.32a), (4.29), and (4.26) that

$$\begin{aligned} f(x_k) - f(x_k + \alpha_k \hat{s}_k) &\geq \gamma_f \alpha_k \rho_k^f \geq \gamma_f \alpha_k \min \left[C_\rho \Delta \ell^f(s_k; x_k)^2, \frac{1}{2} \Delta \ell^f(s_k; x_k) \right] \\ &\geq \gamma_f \alpha_{\min} \min \left[C_\rho (\epsilon^f)^2, \frac{1}{2} \epsilon^f \right] > 0 \quad \text{for all } k \in \mathcal{S}'' \text{ sufficiently large.} \end{aligned} \quad (4.30)$$

However, for all $k \in \mathcal{S}_o$, it follows from (2.32b), (2.32a), (2.15), and (2.26) that $f(x_k) - f(x_k + \alpha_k \hat{s}_k) \geq 0$. This observation combined with (4.30) implies that $\lim_{k \rightarrow \infty} f(x_k) = -\infty$, which contradicts the fact that f is bounded as a consequence of Assumptions 4.1 and 4.2. This completes the proof. \square

We now show that limit points are either infeasible stationary points or KKT points for problem (1.1).

THEOREM 4.14. *Suppose that Assumptions 4.1–4.4, (4.11), and (4.23) hold. If x_* is a limit point of $\{x_k\}$, then either*

- (i) x_* is an infeasible stationary point; or
- (ii) x_* is a KKT point for problem (1.1).

Proof. Suppose that $\lim_{k \in \mathcal{S}} x_k = x_*$ for some subsequence \mathcal{S} . It follows from Lemma 4.12 that $\lim_{k \rightarrow \infty} \Delta \ell^v(s_k^s; x_k) = 0$ so that if $v(x_*) > 0$, then x_* is an infeasible stationary point (see Definition 2.1). Otherwise, we have that $v(x_*) = 0$. In this case, it follows from Lemma 4.13 and (4.11) that $\lim_{k \in \mathcal{S}} \Delta q^\phi(s_k^p; x_k, B_k, \bar{\sigma}) = 0$. It follows from this fact, $v(x_*) = 0$, and Lemma 2.2 that x_* is a KKT point for problem (1.1). \square

Case 3: $|\mathcal{S}_v \cup \mathcal{S}_b| = \infty$. We first show that the feasibility measure converges to zero along $\mathcal{S}_v \cup \mathcal{S}_b$.

LEMMA 4.15. *If Assumptions 4.1–4.3 hold and $|\mathcal{S}_v \cup \mathcal{S}_b| = \infty$, then $\lim_{k \in \mathcal{S}_v \cup \mathcal{S}_b} \Delta \ell^v(s_k^s; x_k) = 0$.*

Proof. To reach a contradiction, suppose that we have the infinite subsequence

$$\mathcal{S} := \{k \in \mathcal{S}_v \cup \mathcal{S}_b : \Delta \ell^v(s_k^s; x_k) \geq \epsilon\}$$

for some constant $\epsilon > 0$. It follows from the definition of \mathcal{S} , Lemma 2.14 and (2.28) that

$$v_k \leq \max \{v_j - \alpha_j \eta_v \Delta \ell^v(s_j^s; x_j), \beta v_j\} \quad \text{or} \quad f_k \leq f_j - \gamma \min \{v_j - \alpha_j \eta_v \Delta \ell^v(s_j^s; x_j), \beta v_j\} \quad (4.31)$$

for $k \in \mathcal{S}$ and $(v_j, f_j) \in \mathcal{F}_k$; note that by construction $(v_k, f_k) \in \mathcal{F}_{k+1}$ for all $k \in \mathcal{S}$. Moreover, it follows from the definitions of $\Delta \ell^v$ and \mathcal{S} that $v_k \geq \Delta \ell^v(s_k^s; x_k) \geq \epsilon$ for $k \in \mathcal{S}$. Using Assumptions 4.1 and 4.2 we have a subsequence $\mathcal{S}' \subseteq \mathcal{S}$ so that

$$\lim_{k \in \mathcal{S}'} \Delta \ell^v(s_k^s; x_k) = \theta_\ell \quad \text{and} \quad \lim_{k \in \mathcal{S}'} v_k = \theta_v \quad \text{for constants } \theta_v \geq \theta_\ell \geq \epsilon > 0.$$

For any $\epsilon_\ell \in (0, \theta_\ell)$ and $\epsilon_v \in (0, \theta_v)$, it follows that

$$|\Delta \ell^v(s_k^s; x_k) - \theta_\ell| < \epsilon_\ell \quad \text{and} \quad |v_k - \theta_v| < \epsilon_v \quad \text{for all } k \in \mathcal{S}' \subseteq \mathcal{S} \text{ sufficiently large.} \quad (4.32)$$

Using (4.32), the definitions of ϵ_ℓ , η_v , $\Delta \ell^v$ and \mathcal{S} , $\alpha_k \in (0, 1]$, $\mathcal{S}' \subseteq \mathcal{S}$, and part (ii) of Lemma 4.8 gives

$$0 \leq v_k - \alpha_k \eta_v \Delta \ell^v(s_k^s; x_k) < v_k - \alpha_F \eta_v (\theta_\ell - \epsilon_\ell) \leq \beta_2 v_k \quad \text{for all } k \in \mathcal{S}' \text{ sufficiently large} \quad (4.33)$$

and some $\alpha_F > 0$, where

$$\beta_2 := \frac{(\theta_v + \epsilon_v) - \alpha_F \eta_v (\theta_\ell - \epsilon_\ell)}{(\theta_v + \epsilon_v)} \in (0, 1)$$

and β_2 may be forced to lie in $(0, 1)$ by choosing ϵ_v sufficiently close to zero and ϵ_ℓ sufficiently close to θ_ℓ . Now define $\beta^* := \max\{\beta_2, \beta\} \in (0, 1)$,

$$\epsilon^* = \min \left\{ \frac{1 - \beta^*}{1 + \beta^*} \theta_v, \epsilon_v \right\} > 0,$$

and the subsequence $\mathcal{S}'' = \{k \in \mathcal{S}' : |v_k - \theta_v| < \epsilon^*\}$ so that

$$\frac{2\beta^*}{1 + \beta^*} \theta_v < v_k < \frac{2}{1 + \beta^*} \theta_v \quad \text{for all } k \in \mathcal{S}'' \subseteq \mathcal{S}' \text{ sufficiently large.} \quad (4.34)$$

Given $k \in \mathcal{S}''$, define $k^+ \in \mathcal{S}''$ to be the successor to k in \mathcal{S}'' . It then follows from (4.34), the definition of β^* , and (4.33) that

$$v_{k^+} > \frac{2\beta^*}{1 + \beta^*} \theta_v > \beta^* v_k = \max\{\beta_2, \beta\} v_k \geq \max \{v_k - \alpha_k \eta_v \Delta \ell^v(s_k^s; x_k), \beta v_k\} \quad \text{for all } k \in \mathcal{S}''.$$

Since $\mathcal{S}'' \subseteq \mathcal{S}' \subseteq \mathcal{S}$, it follows from the previous inequality, the definition of k^+ , the fact that $(v_k, f_k) \in \mathcal{F}_{k^+}$, (4.31), the definition of $\Delta \ell^v(s_k^s; x_k)$, $\alpha_k \in (0, 1]$, $\eta_v \in (0, 1)$, $\beta \in (0, 1)$, $\gamma \in (0, 1)$, $\theta_v > \epsilon_v \geq \epsilon^*$ and the definition of \mathcal{S}'' that

$$\begin{aligned} f_k - f_{k^+} &\geq \gamma \min \{v_k - \alpha_k \eta_v \Delta \ell^v(s_k^s; x_k), \beta v_k\} \\ &= \gamma \min \{(1 - \alpha_k \eta_v) v_k + \alpha_k \eta_v \| [c(x_k) + J(x_k) s_k^s]^- \|_1, \beta v_k\} \\ &\geq \gamma \min \{1 - \alpha_k \eta_v, \beta\} v_k \geq \gamma \min \{1 - \eta_v, \beta\} (\theta_v - \epsilon^*) > 0 \quad \text{for all } k \text{ in } \mathcal{S}''. \end{aligned}$$

Summing over $k \in \mathcal{S}''$, we deduce that $\{f_k\}_{k \in \mathcal{S}''} \rightarrow -\infty$, which contradicts Assumptions 4.1 and 4.2. \square

We now prove that our optimality measure for ϕ converges to zero along a certain subsequence.

LEMMA 4.16. *Suppose that Assumptions 4.1–4.4 and (4.11) hold, and that $|\mathcal{S}_v \cup \mathcal{S}_b| = \infty$.*

(i) If $|\mathcal{S}_v| = \infty$ and $\lim_{k \in \mathcal{S}_v} x_k = x_*$ for some x_* satisfying $v(x_*) = 0$, then

$$\lim_{k \in \mathcal{S}_v} \Delta q^\phi(s_k^p; x_k, B_k, \bar{\sigma}) = 0.$$

(ii) If $|\mathcal{S}_v| < \infty$ and $\lim_{k \in \mathcal{S}_b} x_k = x_*$ for some x_* satisfying $v(x_*) = 0$, then

$$\liminf_{k \in \mathcal{S}_b} \Delta q^\phi(s_k^p; x_k, B_k, \bar{\sigma}) = 0.$$

Proof. We first prove part (i). To obtain a contradiction, suppose that there exists the subsequence

$$\mathcal{S}' := \{k \in \mathcal{S}_v : k \geq k' \text{ and } \Delta q^\phi(s_k^p; x_k, B_k, \bar{\sigma}) \geq \epsilon'\}$$

for some constant $\epsilon' > 0$ and k' defined in (4.11). It then follows from line 34 of Algorithm 1 that

$$\Delta q^\phi(s_k; x_k, B_k, \bar{\sigma}) \geq \eta_\phi \Delta q^\phi(s_k^p; x_k, B_k, \bar{\sigma}) \geq \eta_\phi \epsilon' \text{ for } k \in \mathcal{S}'. \quad (4.35)$$

Then, since $v(x_*) = 0$ by assumption, we may use (4.35) (analogous to (4.24)) and follow the same steps that led to (4.26) to show that

$$\Delta \ell^f(s_k; x_k) \geq \epsilon^f \geq \gamma_v \Delta \ell^v(s_k; x_k) \text{ for } k \in \mathcal{S}' \text{ sufficiently large and some } \epsilon^f > 0,$$

where the second inequality follows from $\lim_{k \in \mathcal{S}_v} x_k = x_*$, $v(x_*) = 0$, and the definition of $\Delta \ell^v$. Thus, (2.31) does not hold and implies that $k \notin \mathcal{S}_v$. This is a contradiction and proves part (i).

We now prove part (ii), where $|\mathcal{S}_v| < \infty = |\mathcal{S}_b|$. To obtain a contradiction, suppose that

$$\Delta q^\phi(s_k^p; x_k, B_k, \bar{\sigma}) \geq \epsilon' \text{ for } k \in \mathcal{S}_b \text{ sufficiently large}$$

and some constant $\epsilon' > 0$. It then follows from line 34 of Algorithm 1 that

$$\Delta q^\phi(s_k; x_k, B_k, \bar{\sigma}) \geq \eta_\phi \Delta q^\phi(s_k^p; x_k, B_k, \bar{\sigma}) \geq \eta_\phi \epsilon' \text{ for } k \in \mathcal{S}_b \text{ sufficiently large.} \quad (4.36)$$

Since (4.36) is analogous to (4.35), we may again conclude as above that

$$\Delta \ell^f(s_k; x_k) \geq \epsilon^f \geq \gamma_v \Delta \ell^v(s_k; x_k) \text{ for } k \in \mathcal{S}_b \text{ sufficiently large and some } \epsilon^f > 0. \quad (4.37)$$

Using (4.37), $\lim_{k \in \mathcal{S}_b} v(x_k) = v(x_*) = 0$, and part (iii) of Lemma 4.8, we may conclude that there exists $\alpha_f > 0$ such that (α, s_k) satisfies (2.32b) for all $\alpha \in (0, \alpha_f]$ and $k \in \mathcal{S}_b$ sufficiently large. Now, if $\alpha_k \rightarrow 0$ along some subsequence $\mathcal{S}'_b \subseteq \mathcal{S}_b$, then it follows from the previous sentence and (4.37) that (α_k, s_k) satisfies (2.32a) and (2.32b) for all $k \in \mathcal{S}_b$ sufficiently large. We now show that $x_k + \alpha_k s_k$ is also acceptable to the filter \mathcal{F}_k for all $k \in \mathcal{S}'_b$ sufficiently large.

To this end, let $(v_i, f_i) \in \mathcal{F}_k$ for some $k \in \mathcal{S}'_b$. It then follows from Lemma 2.14 that either $v_k \leq \max\{v_i - \alpha_i \eta_v \Delta \ell^v(s_i^s; x_i), \beta v_i\}$ or $f_k \leq f_i - \gamma \min\{v_i - \alpha_i \eta_v \Delta \ell^v(s_i^s; x_i), \beta v_i\}$. In this first case, it follows from the definition of a b-pair that $v(x_k + \alpha_k s_k) \leq v_k \leq \max\{v_i - \alpha_i \eta_v \Delta \ell^v(s_i^s; x_i), \beta v_i\}$ for all $k \in \mathcal{S}_b$. In the second case, we have from the fact that (2.32b) holds for $k \in \mathcal{S}'_b$ sufficiently large (recall that $\alpha_k \rightarrow 0$ on \mathcal{S}'_b), (4.37), and Lemma 4.5 that $f(x_k + \alpha_k s_k) \leq f_k \leq f_i - \gamma \min\{v_i - \alpha_i \eta_v \Delta \ell^v(s_i^s; x_i), \beta v_i\}$. Thus, in either case we have that $(v(x_k + \alpha_k s_k), f(x_k + \alpha_k s_k))$ is acceptable to the single element filter $\{(v_i, f_i)\}$ for all $k \in \mathcal{S}'_b$ sufficiently large. Since this element (v_i, f_i) of the filter \mathcal{F}_k was arbitrary, we may conclude that $(v(x_k + \alpha_k s_k), f(x_k + \alpha_k s_k))$ is, in fact, acceptable to the filter \mathcal{F}_k for all $k \in \mathcal{S}'_b$ sufficiently large.

To summarize, we have shown that (α_k, s_k) is an o-pair for $k \in \mathcal{S}'_b$ sufficiently large. This is a contradiction since Algorithm 1 would have labeled such an iterate as an o-iterate, not a b-iterate.

Thus, there exists α_b such that $\alpha_k \geq \alpha_b > 0$ for all $k \in \mathcal{S}_b$ sufficiently large. Combining this with (2.1), (2.29), $v(\cdot) \geq 0$, Lemma 4.7, (2.7), (2.15), and (4.37) gives

$$\begin{aligned}
f(x_k) - f(x_k + \alpha_k \hat{s}_k) &= \phi(x_k; \bar{\sigma}) - \phi(x_k + \alpha_k \hat{s}_k; \bar{\sigma}) - \bar{\sigma}(v(x_k) - v(x_k + \alpha_k \hat{s})) \geq \gamma_\phi \alpha_k \rho_k^\phi - \bar{\sigma}v(x_k) \\
&\geq \gamma_\phi \alpha_b \min \left[C_\rho \Delta \ell^\phi(s_k; x_k, \bar{\sigma})^2, \frac{1}{2} \Delta \ell^\phi(s_k; x_k, \bar{\sigma}) \right] - \bar{\sigma}v(x_k) \\
&= \gamma_\phi \alpha_b \min \left[C_\rho (\Delta \ell^f(s_k; x_k)^2 + 2\bar{\sigma} \Delta \ell^f(s_k; x_k) \Delta \ell^v(s_k; x_k) + \bar{\sigma}^2 \Delta \ell^v(s_k; x_k)^2), \right. \\
&\quad \left. \frac{1}{2} (\Delta \ell^f(s_k; x_k) + \bar{\sigma} \Delta \ell^v(s_k; x_k)) \right] - \bar{\sigma}v(x_k) \\
&\geq \gamma_\phi \alpha_b \min \left[C_\rho \Delta \ell^f(s_k; x_k)^2, \frac{1}{2} \Delta \ell^f(s_k; x_k) \right] - \bar{\sigma}v(x_k) \\
&\geq \gamma_\phi \alpha_b \min \left[C_\rho (\epsilon^f)^2, \frac{1}{2} \epsilon^f \right] - \bar{\sigma}v(x_k) \text{ for } k \in \mathcal{S}_b \text{ sufficiently large.} \tag{4.38}
\end{aligned}$$

Since $|\mathcal{S}_b| = \infty > |\mathcal{S}_v|$, we may define k^+ as the first iteration greater than k such that $k^+ \in \mathcal{S}_b \cup \mathcal{S}_o$. It then follows from the construction of Algorithm 1 and $|\mathcal{S}_v| < \infty$ that

if $k \in \mathcal{S}_b$ is sufficiently large, then $k^+ \in \mathcal{S}_b \cup \mathcal{S}_o$ and $l \in \mathcal{S}_p$ for all $k < l < k^+$.

Using (2.29), $\alpha_i \geq 0$, (2.19), and (2.27) we conclude that

$$\begin{aligned}
\phi(x_{k+1}; \bar{\sigma}) - \phi(x_{k^+}; \bar{\sigma}) &= \sum_{i=k+1}^{k^+-1} \phi(x_i; \bar{\sigma}) - \phi(x_i + \alpha_i \hat{s}_i; \bar{\sigma}) \geq \sum_{i=k+1}^{k^+-1} \gamma_\phi \alpha_i \rho_i^\phi \\
&= \sum_{i=k+1}^{k^+-1} \gamma_\phi \alpha_i \min \left[\Delta \ell^\phi(s_i; x_i, \bar{\sigma}), \Delta q^\phi(s_i^{c\phi}; x_i, H_i, \bar{\sigma}) \right] \\
&\geq 0 \text{ for } k \in \mathcal{S}_b \text{ sufficiently large,}
\end{aligned}$$

which may be combined with (2.1), $v(\cdot) \geq 0$, and (2.34) to conclude that

$$f(x_{k+1}) - f(x_{k^+}) \geq \bar{\sigma}(v(x_{k^+}) - v(x_{k+1})) \geq -\bar{\sigma}v(x_{k+1}) > -\bar{\sigma}v(x_k) \text{ for } k \in \mathcal{S}_b \text{ sufficiently large.} \tag{4.39}$$

It then follows from (4.38) and (4.39) that

$$\begin{aligned}
f(x_k) - f(x_{k^+}) &= (f(x_k) - f(x_k + \alpha_k \hat{s}_k)) + (f(x_{k+1}) - f(x_{k^+})) \\
&> \gamma_\phi \alpha_b \min \left[C_\rho (\epsilon^f)^2, \frac{1}{2} \epsilon^f \right] - 2\bar{\sigma}v(x_k) \text{ for } k \in \mathcal{S}_b \text{ sufficiently large.} \tag{4.40}
\end{aligned}$$

Next, since $\lim_{k \in \mathcal{S}_b} v(x_k) = 0$ we know that

$$v(x_k) \leq \frac{1}{4\bar{\sigma}} \gamma_\phi \alpha_b \min \left[C_\rho (\epsilon^f)^2, \frac{1}{2} \epsilon^f \right] \text{ for } k \in \mathcal{S}_b \text{ sufficiently large,}$$

which may be combined with (4.40) to deduce that

$$f(x_k) - f(x_{k^+}) > \frac{1}{2} \gamma_\phi \alpha_b \min \left[C_\rho (\epsilon^f)^2, \frac{1}{2} \epsilon^f \right] =: \epsilon^v > 0 \text{ for } k \in \mathcal{S}_b \text{ sufficiently large.} \tag{4.41}$$

If we define \hat{k}^+ to be the first b-iteration greater than k (thus, $\hat{k}^+ \geq k^+$), it follows from (4.41), the fact that Algorithm 1 does not allow further p-iterations until it has its next b-iteration, and the fact that the objective f is decreased during o-iterations that $f(x_k) - f(x_{\hat{k}^+}) > \epsilon^v$ for $k \in \mathcal{S}_b$ sufficiently large. Since $|\mathcal{S}_b| = \infty$, this implies that $f(x_k) \rightarrow -\infty$, which contradicts the fact that f is bounded as a consequence of Assumptions 4.1 and 4.2. This proves the result. \square

We now show that limit points of $\{x_k\}_{\mathcal{S}_v \cup \mathcal{S}_b}$ are infeasible stationary or KKT point for problem (1.1).

THEOREM 4.17. *Suppose that the Assumptions 4.1–4.4, (4.11), and $|\mathcal{S}_v \cup \mathcal{S}_b| = \infty$ hold. Then, there exists a limit point x_* of $\{x_k\}_{\mathcal{S}_v \cup \mathcal{S}_b}$ such that either*

- (i) x_* is a KKT point of problem (1.1) or
- (ii) x_* is an infeasible stationary point.

Proof. From Assumptions 4.1 and 4.2 we that there exists a limit point x_* of $\{x_k\}_{\mathcal{S}_v \cup \mathcal{S}_b}$. First, if $v(x_*) > 0$, then it follows from Lemma 4.15 and Lemma 2.1 that x_* is an infeasible stationary point, which is part (ii) of this theorem. Second, if $v(x_*) = 0$ and $|\mathcal{S}_v| = \infty$, then it follows from part (i) of Lemma 4.16 and Lemma 2.2 that x_* is a KKT point of problem (1.1). This is case (i) of this theorem. Finally, if $v(x_*) = 0$ and $|\mathcal{S}_v| < \infty$ (so that $|\mathcal{S}_b| = \infty$), then it follows from part (ii) of Lemma 4.16 and Lemma 2.2 that x_* is a KKT point of problem (1.1), which once again is case (i) of the theorem. \square

4.2. Convergence analysis under unbounded weighting parameter. We now consider the situation when the weighting parameter increases without bound, i.e, that

$$\lim_{k \rightarrow \infty} \sigma_k = \infty. \quad (4.42)$$

Our analysis begins with the following lemma, which is similar to [7, Lemma 3.8].

LEMMA 4.18. *Suppose that Assumptions 4.1–4.4 are satisfied, (4.42) holds, x_* is a limit point of $\{x_k\}$ satisfying $v(x_*) > 0$, and $\Delta \ell^v(s_*^s; x_*) > 0$, where s_*^s is the solution to*

$$\underset{(s,r) \in \mathbb{R}^{n+m}}{\text{minimize}} \quad e^T r \quad \text{subject to} \quad c(x_*) + J(x_*)s + r \geq 0, \quad r \geq 0, \quad \|s\|_\infty \leq \delta,$$

for some $\delta \in [\delta_{\min}, \delta_{\max}]$. Then, along any subsequence $\{x_k\}_{k \in \mathcal{K}}$ that converges to x_* , the weighting parameter is updated only a finite number of times.

Proof. We begin by defining

$$s_k^\phi(\sigma) := \underset{s \in \mathbb{R}^n}{\text{argmin}} \quad q^\phi(s; x_k, B_k, \sigma) \quad (4.43)$$

and

$$\mu := \mu(\sigma) := \left(1 - \frac{\eta_\sigma}{\eta_v}\right) \sigma < \sigma, \quad (4.44)$$

where we used the fact that $0 < 1 - \eta_\sigma/\eta_v < 1$ holds since $0 < \eta_\sigma < \eta_v < 1$ is defined in (2.18).

Using the fact that $\Delta q^\phi(s_k^\phi(\mu); x_k, B_k, \mu) \geq 0$, (2.8), and the definition of $\mu = \mu(\sigma)$, we can see that

$$\begin{aligned} \Delta q^\phi(s_k^\phi(\mu); x_k, B_k, \mu) &= \Delta q^f(s_k^\phi(\mu); x_k, B_k) + \left(1 - \frac{\eta_\sigma}{\eta_v}\right) \sigma \Delta \ell^v(s_k^\phi(\mu); x_k) \\ &= \Delta q^\phi(s_k^\phi(\mu); x_k, B_k, \sigma) - \frac{\eta_\sigma}{\eta_v} \sigma \Delta \ell^v(s_k^\phi(\mu); x_k) \geq 0 \quad \text{for } \mu = \mu(\sigma) \text{ and all } \sigma > 0, \end{aligned}$$

which implies that

$$\Delta q^\phi(s_k^\phi(\mu); x_k, B_k, \sigma) \geq \frac{\eta_\sigma}{\eta_v} \sigma \Delta \ell^v(s_k^\phi(\mu); x_k) \quad \text{for } \mu = \mu(\sigma) \text{ and all } \sigma > 0. \quad (4.45)$$

Since $\Delta \ell^v(s_*^s; x_*) > 0$ and $\lim_{k \in \mathcal{K}} x_k = x_*$ by assumption, it follows from [15, Theorem 3.2.8] that there exists $\epsilon \in (0, 1)$ and k' such that

$$\Delta \ell^v(s_k^s; x_k) > \epsilon \quad \text{for all } k' \leq k \in \mathcal{K}. \quad (4.46)$$

Moreover, since the Newton step $-B_k^{-1}g_k$ minimizes $q^f(s; x_k, B_k)$, it follows from Assumption 4.3 that

$$q^f(s_k^\phi(\sigma); x_k, B_k) \geq q^f(-B_k^{-1}g_k; x_k, B_k) = f_k - \frac{1}{2}g_k^T B_k^{-1}g_k \geq f_k - \frac{\|g_k\|_2^2}{2\lambda_{\min}} \quad \text{for all } \sigma > 0. \quad (4.47)$$

Next, it follows from (2.9), the choice $\delta_k \in [\delta_{\min}, \delta_{\max}]$, norm inequalities, and Assumption 4.3 that

$$q^f(s_k^s; x_k, B_k) \leq f_k + \|g_k\|_2 \delta_{\max} + \frac{1}{2} \lambda_{\max} \delta_{\max}^2. \quad (4.48)$$

Then, (4.47), (4.48), and Assumptions 4.1 and 4.2 imply the existence of a constant $C_{\text{qf}} > 0$ such that

$$q^f(s_k^s; x_k, B_k) - q^f(s_k^\phi(\sigma); x_k, B_k) \leq C_{\text{qf}} \text{ for all } \sigma > 0. \quad (4.49)$$

We now define

$$\sigma_{\text{crit}} := \frac{C_{\text{qf}}}{\epsilon(1 - \eta_v) \left(1 - \frac{\eta_\sigma}{\eta_v}\right)} > \mu(\sigma_{\text{crit}}) = \frac{C_{\text{qf}}}{\epsilon(1 - \eta_v)} > 0, \quad (4.50)$$

and the associated infinite subsequence

$$\mathcal{S}' = \{k \in \mathcal{K} : k \geq k' \text{ and } \sigma_k \geq \sigma_{\text{crit}}\}. \quad (4.51)$$

It follows from the fact that $\Delta q^\phi(s_k^\phi(\sigma); x_k, B_k, \sigma) \geq \Delta q^\phi(s_k^s; x_k, B_k, \sigma)$ (by the definition of $s_k^\phi(\sigma)$), (2.8), (4.49), (4.46), and (4.50) that

$$\begin{aligned} \Delta \ell^v(s_k^\phi(\sigma); x_k) &\geq \Delta \ell^v(s_k^s; x_k) - \frac{1}{\sigma} \left(q^f(s_k^s; x_k, B_k) - q^f(s_k^\phi(\sigma); x_k, B_k) \right) \\ &\geq \Delta \ell^v(s_k^s; x_k) - \frac{1}{\sigma} C_{\text{qf}} = \Delta \ell^v(s_k^s; x_k) \left(1 - \frac{C_{\text{qf}}}{\sigma \Delta \ell^v(s_k^s; x_k)} \right) \\ &\geq \eta_v \Delta \ell^v(s_k^s; x_k) \text{ for } \sigma \geq \mu(\sigma_{\text{crit}}) \text{ and } k' \leq k \in \mathcal{K}. \end{aligned} \quad (4.52)$$

We may now use the definition of \mathcal{S}' , (4.52), the fact that $s_k^\phi(\sigma_k) \equiv s_k^p$, (2.14), and (2.15) to show that

$$\tau_k = 1, \quad s_k = s_k^p, \quad \text{and} \quad \Delta q^\phi(s_k; x_k, B_k, \sigma_{k+1}) \geq \eta_\phi \Delta q^\phi(s_k^p; x_k, B_k, \sigma_{k+1}) \text{ for } k \in \mathcal{S}' \quad (4.53)$$

since $\eta_\phi \in (0, 1)$ in Algorithm 1. Next, it follows from (4.53), (2.8), $B_k \succ 0$, the fact that $s_k^p \equiv s_k^\phi(\sigma_k)$ and s_k^p minimizes $q^\phi(s; x_k, B_k, \sigma_k)$, (4.45), the fact that $\mu(\sigma_k) \geq \mu(\sigma_{\text{crit}})$ for $k \in \mathcal{S}'$, and (4.52) that

$$\begin{aligned} \Delta \ell^\phi(s_k; x_k, \sigma_k) &\geq \Delta q^\phi(s_k^p; x_k, B_k, \sigma_k) \geq \Delta q^\phi(s_k^\phi(\mu(\sigma_k)); x_k, B_k, \sigma_k) \geq \frac{\eta_\sigma}{\eta_v} \sigma_k \Delta \ell^v(s_k^\phi(\mu(\sigma_k)); x_k) \\ &\geq \frac{\eta_\sigma}{\eta_v} \sigma_k (\eta_v \Delta \ell^v(s_k^s; x_k)) = \sigma_k \eta_\sigma \Delta \ell^v(s_k^s; x_k) \text{ for } k \in \mathcal{S}'. \end{aligned} \quad (4.54)$$

We now conclude from (4.53), (4.54), (2.18), and the fact that the weighting parameter is only increased in lines 12 and 34 of Algorithm 1, that σ_k is increased a finite number of times on \mathcal{K} . \square

We now consider feasible limit points at which the MFCQ [31] holds.

LEMMA 4.19. *Suppose that Assumptions 4.1–4.4 are satisfied, (4.42) holds, x_* is a limit point of $\{x_k\}$ at which $v(x_*) = 0$ and the MFCQ holds. Then, the following hold for all x_k sufficiently close to x_* and σ_k sufficiently large: (i) $\Delta \ell^v(s_k^p; x_k) = v(x_k)$; (ii) $s_k = s_k^p$; and (iii) σ_k is not increased during iteration k .*

Proof. We may use [7, Lemmas 3.12 and 3.13] since the proofs only used the properties of the MFCQ, the continuity of the problem functions f and g , and the convexity of their penalty and steering subproblems. Their subproblem [7, Equations 2.7(a–d)] is equivalent to our predictor subproblem (2.11) and both methods minimize the same quadratic model of the penalty function. A small difference is that our predictor subproblem is designed so that if $\ell^v(s_k^s; x_k) = 0$, then $\ell^v(s_k^p; x_k) = 0$ as well; they satisfy this requirement by increasing their penalty parameter in Step 4a [7, Eqn 2.11] and re-solving for a new step. Their steering subproblem [7, Equations 2.9(a–e)] is equivalent to (2.9).

The assumptions of this lemma and [7, Lemma 3.12] imply the existence of $r > 0$ and $k' \geq 0$ so that

$$\ell^v(s_k^p, x_k) = v(x_k) \text{ for all } k \in \mathcal{S}', \quad (4.55)$$

where $\mathcal{S}' := \{k : \|x_k - x_*\| \leq r \text{ and } k \geq k'\}$, which proves part (i). The inequality $\Delta\ell^v(s_k^s; x_k) \geq 0$, (4.55), and the definition of $\Delta\ell^v$ imply

$$\Delta\ell^v(s_k^p; x_k) \geq v(x_k) - \ell^v(s_k^s; x_k) = \Delta\ell^v(s_k^s; x_k) \geq \eta_v \Delta\ell^v(s_k^s; x_k) \text{ for } k \in \mathcal{S}',$$

where $\eta_v \in (0, 1)$ is defined in (2.15). Thus, we conclude from (2.15) that $\tau_k = 1$ and $s_k = s_k^p$ for $k \in \mathcal{S}'$, which proves part (ii). Finally, it follows from [7, Lemma 3.13] and the assumptions of this lemma, that

$$\Delta q^\phi(s_k^p; x_k, B_k, \sigma_k) \geq \sigma_k \eta_\sigma v(x_k) \geq \sigma_k \eta_\sigma \Delta\ell^v(s_k^s; x_k) \text{ for } k \in \mathcal{S}', \quad (4.56)$$

where the last inequality follows from the definition of $\Delta\ell^v$. It then follows from part (ii) of this lemma, (2.8), $B_k \succ 0$, and (4.56) that

$$\Delta\ell^\phi(s_k; x_k, \sigma_k) = \Delta\ell^\phi(s_k^p; x_k, \sigma_k) \geq \Delta q^\phi(s_k^p; x_k, B_k, \sigma_k) \geq \sigma_k \eta_\sigma \Delta\ell^v(s_k^s; x_k) \text{ for } k \in \mathcal{S}'.$$

We may conclude from this inequality, (2.18), and the fact that σ_k will not be increased on Line 34 as a result of part (ii) of this lemma, that $\sigma_{k+1} = \sigma_k$ for $k \in \mathcal{S}'$, which proves part (iii). \square

THEOREM 4.20. *If Assumptions 4.1–4.4 and (4.42) hold, there is a limit point x_* such that either*

- (i) x_* is an infeasible stationary point; or
- (ii) x_* is feasible, but the MFCQ does not hold.

Proof. Let \mathcal{D} to be the infinite index set consisting of the iterations for which the weighting parameter is increased. Then, let x_* be a limit point of $\{x_k\}_{k \in \mathcal{D}}$, which must exist as a consequence of Assumptions 4.1 and 4.2. First, suppose that $v(x_*) > 0$. It then follows from Lemma 4.18 that if $\Delta\ell^v(s_*^s; x_*) > 0$ (s_*^s is defined in Lemma 4.18), then the weighting parameter is updated only a finite number of times along \mathcal{D} , which is a contradiction. Therefore, we deduce that $\Delta\ell^v(s_*^s; x_*) = 0$ and consequently that x_* is an infeasible stationary point. Second, suppose that $v(x_*) = 0$. It then follows from Lemma 4.19 that if the MFCQ holds at x_* , then σ_k only be increased a finite number of times along \mathcal{D} . This is a contradiction and, therefore, the MFCQ does not hold at x_* . \square

5. Conclusions. In this paper, we presented a new filter linesearch method that replaced the traditional restoration phase with a penalty mode that systematically decreased an exact penalty function. Importantly, we solved a *single* strictly convex quadratic program subproblem during each iteration that was *always* feasible. Each search direction was defined as a convex combination of a steering step (a solution of a linear program) that represented the best local improvement in constraint violation and a predictor step that reduced our strictly convex quadratic model of the exact penalty function. We also allowed for the computation of an accelerator step defined as a solution to a simple equality constrained quadratic program (plus trust-region constraint) to promote fast local convergence. In this manner, the trial step always incorporated information from both the objective function and constraint violation. To further promote step acceptance, we utilized second-order information in the computation of Cauchy steps that provided realistic measurements of the decrease one might expect from the nonlinear problem functions. An additional contribution was the use of local feasibility estimates that emerged during the steering process to define a new and improved margin (envelope) of the filter. This new definition encouraged the acceptance of steps that make reasonable progress, but might very well be considered inadmissible by a traditional filter. Under standard assumptions, we proved global convergence of our algorithm.

REFERENCES

- [1] G. ALARCÓN, C. TORRES, AND L. GÓMEZ, *Global optimization of gas allocation to a group of wells in artificial lift using nonlinear constrained programming*, Journal of energy resources technology, 124 (2002), p. 262.
- [2] A. BARCLAY, P. E. GILL, AND J. B. ROSEN, *SQP methods in optimal control*, in 12th Conference on Variational Calculus, Optimal Control and Applications, R. Bulirsch, L. Bittner, W. Schmidt, and K. Heier, eds., International Series of Numerical Mathematics, Basel, 1997, Birkhäuser.
- [3] A. BARCLAY, P. E. GILL, AND J. B. ROSEN, *SQP methods in optimal control*, in Variational Calculus, Optimal Control and Applications, R. Bulirsch, L. Bittner, W. H. Schmidt, and K. Heier, eds., vol. 124 of International Series of Numerical Mathematics, Basel, Boston and Berlin, 1998, Birkhäuser, pp. 207–222.
- [4] M. BAZARAA, H. SHERALI, AND C. SHETTY, *Nonlinear programming: theory and algorithms*, John Wiley and Sons, 2006.
- [5] A. BRYSON AND Y. HO, *Applied Optimal Control*, Taylor and Francis, New York, 1975.
- [6] R. H. BYRD, N. I. M. GOULD, J. NOCEDAL, AND R. A. WALTZ, *On the convergence of successive linear-quadratic programming algorithms*, SIAM Journal on Optimization, 16 (2006), pp. 471–489.
- [7] R. H. BYRD, G. LOPEZ-CALVA, AND J. NOCEDAL, *A line search exact penalty method using steering rules*, Mathematical Programming, 133 (2012), pp. 39–73.
- [8] R. H. BYRD, J. NOCEDAL, AND R. A. WALTZ, *KNITRO optimization software package*. <http://www.ziena.com/knitro/kindex.htm>.
- [9] ———, *Steering exact penalty methods for nonlinear programming*, Optim. Methods Softw., 23 (2008), pp. 197–213.
- [10] L. CHEN AND D. GOLDFARB, *An interior-point piecewise linear penalty method for nonlinear programming*, Math. Program., 128 (2011), pp. 73–122.
- [11] C. M. CHIN AND R. FLETCHER, *On the global convergence of an SLP-filter algorithm that takes EQP steps*, Math. Program., 96 (2003), pp. 161–177.
- [12] C. M. CHIN, A. H. A. RASHID, AND K. M. NOR, *A combined filter line search and trust region method for nonlinear programming*, WSEAS Trans. Math., 5 (2006), pp. 656–662.
- [13] ———, *Global and local convergence of a filter line search method for nonlinear programming*, Optim. Methods Softw., 22 (2007), pp. 365–390.
- [14] A. R. CONN, N. I. M. GOULD, AND PH. L. TOINT, *LANCELOT: a Fortran package for large-scale nonlinear optimization (Release A)*, Lecture Notes in Computation Mathematics 17, Springer Verlag, Berlin, Heidelberg, New York, London, Paris and Tokyo, 1992.
- [15] ———, *Trust-Region Methods*, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2000.
- [16] F. E. CURTIS AND J. NOCEDAL, *Flexible penalty functions for nonlinear constrained optimization*, IMA J. Numer. Anal., 28 (2008), pp. 749–769.
- [17] S. FLÅM AND A. ANTIPIN, *Equilibrium programming using proximal-like algorithms*, Mathematical Programming, 78 (1996), pp. 29–41.
- [18] R. FLETCHER, N. I. M. GOULD, S. LEYFFER, PH. L. TOINT, AND A. WÄCHTER, *Global convergence of a trust-region SQP-filter algorithm for general nonlinear programming*, SIAM J. Optim., 13 (2002), pp. 635–659 (electronic) (2003).
- [19] R. FLETCHER AND S. LEYFFER, *Nonlinear programming without a penalty function*, Math. Program., 91 (2002), pp. 239–269.
- [20] R. FLETCHER, S. LEYFFER, AND PH. L. TOINT, *On the global convergence of an SLP-filter algorithm*, Report 98/13, Department of Mathematics, University of Namur, Belgium, 1998.
- [21] R. FLETCHER, S. LEYFFER, AND PH. L. TOINT, *On the global convergence of a filter-SQP algorithm*, SIAM J. Optim., 13 (2002), pp. 44–59 (electronic).
- [22] P. E. GILL, L. O. JAY, M. W. LEONARD, L. R. PETZOLD, AND V. SHARMA, *An SQP method for the optimal control of large-scale dynamical systems*, J. Comput. Appl. Math., 120 (2000), pp. 197–213. SQP-based direct discretization methods for practical optimal control problems.
- [23] P. E. GILL, W. MURRAY, AND M. A. SAUNDERS, *SNOPT: An SQP algorithm for large-scale constrained optimization*, SIAM Rev., 47 (2005), pp. 99–131.
- [24] N. I. M. GOULD AND D. P. ROBINSON, *A second derivative SQP method: Global convergence*, SIAM J. Optim., 20 (2010), pp. 2023–2048.
- [25] ———, *A second derivative SQP method: Local convergence and practical issues*, SIAM J. Optim., 20 (2010), pp. 2049–2079.
- [26] ———, *A second derivative SQP method with a "trust-region-free" predictor step*, IMA J. Numer. Anal., 32 (2012), pp. 580–601.
- [27] T. JOHANSEN, T. FOSSEN, AND S. BERGE, *Constrained nonlinear control allocation with singularity avoidance using sequential quadratic programming*, Control Systems Technology, IEEE Transactions on, 12 (2004), pp. 211–216.
- [28] D. KRAFT, *On converting optimal control problems into nonlinear programming problems*, in Computational Mathematical Programming, K. Schittkowski, ed., NATO ASI Series F: Computer and Systems Sciences 15, Springer Verlag, Berlin and Heidelberg, 1985, pp. 261–280.
- [29] D. C. LIU AND J. NOCEDAL, *On the limited memory BFGS method for large scale optimization*, Math. Program., 45

- (1989), pp. 503–528.
- [30] P. LIU AND A. DER KIUREGHIAN, *Optimization algorithms for structural reliability*, Structural safety, 9 (1991), pp. 161–177.
 - [31] O. L. MANGASARIAN AND S. FROMOVITZ, *The Fritz John necessary optimality conditions in the presence of equality and inequality constraints*, J. Math. Anal. Appl., 17 (1967), pp. 37–47.
 - [32] J. MORALES, J. NOCEDAL, AND Y. WU, *A sequential quadratic programming algorithm with an additional equality constrained phase*, IMA Journal of Numerical Analysis, DOI: 10.1093/imanum/drq037 (2011).
 - [33] F. MURPHY, H. SHERALI, AND A. SOYSTER, *A mathematical programming approach for determining oligopolistic market equilibrium*, Mathematical Programming, 24 (1982), pp. 92–106.
 - [34] B. A. MURTAGH AND M. A. SAUNDERS, *MINOS 5.5 User's Guide*, Report SOL 83-20R, Department of Operations Research, Stanford University, Stanford, CA, Revised 1998.
 - [35] J. NOCEDAL AND S. J. WRIGHT, *Numerical Optimization*, Springer-Verlag, New York, 1999.
 - [36] S. M. ROBINSON, *Perturbed Kuhn-Tucker points and rates of convergence for a class of nonlinear programming algorithms*, Math. Program., 7 (1974), pp. 1–16.
 - [37] R. ROCKAFELLAR, *Convex Analysis*, Princeton University Press, 1970.
 - [38] M. ULBRICH, S. ULBRICH, AND L. VICENTE, *A globally convergent primal-dual interior-point filter method for nonlinear programming*, Mathematical Programming, 100 (2004), pp. 379–410.
 - [39] A. WACHTER AND L. T. BIEGLER, *Line search filter methods for nonlinear programming: motivation and global convergence*, SIAM Journal on Optimization, 16 (2006), pp. 1–31.
 - [40] A. WÄCHTER, L. T. BIEGLER, Y.-D. LANG, AND A. RAGHUNATHAN, *IPOPT: An interior point algorithm for large-scale nonlinear optimization*. <http://www.coin-or.org>, 2002.
 - [41] R. WALTZ, J. MORALES, J. NOCEDAL, AND D. ORBAN, *An interior algorithm for nonlinear optimization that combines line search and trust region steps*, Mathematical Programming, 107 (2006), pp. 391–408.