

Angewandte Mathematik und Optimierung Schriftenreihe
Applied Mathematics and Optimization Series
AMOS # 20(2015)

György Dósa, Armin Fügenschuh, Zhiyi Tan, Zsolt Tuza,
and Krzysztof Węsek

**New Lower Bounds for Semi-online Scheduling
on Two Uniform Machines with Known Optimum**

Herausgegeben von der
Professur für Angewandte Mathematik
Professor Dr. rer. nat. Armin Fügenschuh

Helmut-Schmidt-Universität / Universität der Bundeswehr Hamburg
Fachbereich Maschinenbau
Holstenhofweg 85
D-22043 Hamburg

Telefon: +49 (0)40 6541 3540
Fax: +49 (0)40 6541 3672

e-mail: appliedmath@hsu-hh.de
URL: <http://www.hsu-hh.de/am>

Angewandte Mathematik und Optimierung Schriftenreihe (AMOS), ISSN-Print 2199-1928
Angewandte Mathematik und Optimierung Schriftenreihe (AMOS), ISSN-Internet 2199-1936

New Lower Bounds for Semi-online Scheduling on Two Uniform Machines with Known Optimum

György Dósa ^{*} Armin Fügenschuh [†]

Zhiyi Tan (corresponding author) [‡] Zsolt Tuza ^{§¶}

Krzysztof Węsek ^{**}

January 10, 2015

Abstract

This problem is about to schedule a number of jobs of different lengths on two uniform machines with given speeds 1 and $s \geq 1$, so that the overall finishing time, i.e. the makespan, is earliest possible. We consider a semi-online variant introduced (for equal speeds) by Azar and Regev, where the jobs are arriving one after the other, while the scheduling algorithm knows the optimum value of the corresponding offline problem. It is desired to construct an algorithm that achieves a schedule close to this optimum value for any given sequence of incoming jobs. Furthermore, one can ask how close any potential algorithm could get to the optimum value, that is, to give a lower bound on the competitive ratio: the supremum over ratios between the value of the solution given by the algorithm and the optimal offline solution. For certain values of s , there are already algorithms known to be tight in the sense that they are scheduling not worse than this bound. For other values of s , this question remained open. We contribute to this question by constructing better lower bounds for some values of s . As a consequence, this proves that the two algorithms given by Ng et al. were in fact optimal, at least, for certain intervals of s .

Keywords: Semi-online scheduling, makespan minimization, machine scheduling, lower bounds.

^{*}Department of Mathematics, University of Pannonia, Veszprém, Hungary, dosagy@almos.vein.hu

[†]Helmut Schmidt University / University of the Federal Armed Forces Hamburg, Holstenhofweg 85, 22043 Hamburg, Germany, fuegenschuh@hsu-hh.de

[‡]Department of Mathematics, Zhejiang University, Hangzhou, Peoples Republic of China, tanzhy@zju.edu.cn

[§]Department of Computer Science, University of Pannonia, Veszprém, Hungary, tuza@isztaki.hu

[¶]Computer and Automation Institute, Hungarian Academy of Sciences, Budapest, Hungary

^{||}Helmut Schmidt University / University of the Federal Armed Forces Hamburg, Holstenhofweg 85, 22043 Hamburg, Germany, wesekk@hsu-hh.de

^{**}Faculty of Mathematics and Information Science, Warsaw University of Technology, ul. Koszykowa 75, 00-662 Warszawa, Poland, wesekk@mini.pw.edu.pl

1 Introduction

We deal with the problem of scheduling two uniform machines: Given are two machines, denoted by M1 and M2, that are both capable to process incoming jobs. They only differ in the processing speed. We assume that machine M1 is working at some unit speed 1, and machine M2 is s times faster, with $s \geq 1$. Hence when machine M1 processes a job of length L , then machine M2 can handle this job in L/s time. There is a number of incoming jobs (finitely or infinitely many) of various lengths. The task is to assign the jobs to the two machines. It is desired to finish all incoming jobs as early as possible, that is, to minimize the makespan. In the offline variant of this problem, all jobs to be assigned are fully known in advance. If nothing is known about the jobs beforehand, we are faced with an online problem. We deal here with a semi-online problem, that means, the jobs are still not known individually, but we assume to have some further overall knowledge. In particular, we assume that the value of the solution to the corresponding offline problem, which we denote by OPT, is known in advance.

Assume there is an algorithm \mathcal{A} that solves the semi-online variant of the problem. This algorithm receives, besides the value of OPT, one job after the other in an unknown order and must now immediately decide to which of the two machines this job should go to (later changes not possible). This algorithm \mathcal{A} finally arrives at a makespan value M greater or equal than OPT. Still, M can be compared to OPT by considering the ratio $\frac{M}{\text{OPT}}$. Of course, it is preferred to have an algorithm where this ratio is close to 1 for any given input data. Thus the question arises, how close to 1 can we get? Is it possible to construct an algorithm reaches this value, or is there a theoretical lower bound well above 1, that no algorithm will ever undercut, no matter how hard it tries?

The paradigm of revealing the instance of a problem in parts, and the decision has to be made as the part is revealed, is naturally motivated by many real-world applications. As it was mentioned before, if no information is given in advance, then we call such problem online, and if some partial information about the instance is known beforehand, then the scheme is called semi-online. The most common way of measuring quality of an online or semi-online algorithm uses the notion of the mentioned competitive ratio. Assume that we are dealing with a minimization problem and the (offline) optimal value for instance I is equal to $\text{OPT}(I)$. Formally, an online algorithm is said to be r -competitive if for any instance I the value of the result of the algorithm $A(I)$ satisfies $\frac{A(I)}{\text{OPT}(I)} \leq r$, and the competitive ratio of an algorithm is defined as infimum of such ratios. The question is: what is the best possible ratio for our (online or semi-online) problem? Formally, we would ask for the optimal competitive ratio, that is the infimum over all numbers r for which there exists an r -competitive algorithm. An algorithm is optimal if its competitive ratio matches the lower bound (value for which it is proven that no algorithm can have better competitive ratio).

1.1 Survey of the Literature

The on-line and various semi-online variations of the problem with a set of jobs to be scheduled on m (not necessarily uniform) machines with an objective to minimize the maximum completion time have been studied for decades. Here we will describe some results concerning deterministic algorithms, *uniformly related* machines, that is, every machine has its speed s and processing a job of length p takes $\frac{p}{s}$ time. We assume (except where noted) that any job has to be done on one machine, and jobs arrive in a list one after another (list-online scheme). For more information we refer to the survey of Tan and Zhang [25].

We will first deal with the basic case of identical machines (i.e. with the same speed) in the pure online scheme. A classic work by Graham [18] is probably the first step in this direction. It gives a $(2 - \frac{1}{m})$ -competitive algorithm, by using a heuristic of assigning every task to the least loaded machine. It was proved by Faigle et al. [16] that for m equal to 2 or 3 this algorithm is in fact optimal. In the case of arbitrary m , many papers appeared in order to decrease the gaps between lower and upper bounds. For the lower bound, Gormley et al. [17] showed that no online algorithm can have competitive ratio better than 1.852 generally (when m can be arbitrarily large). For the upper bound, Albers [1] proposed a 1.923-competitive algorithm for any m , and if m tends to infinity, then work of Fleischer and Wahl gives an algorithm with competitive ratio tending to 1.9201.

Recently, a new approach was presented for this problem. What if we try to compare with the best possible competitive ratio of any online algorithm, even if it is not known? Assume that the optimal competitive ratio is ρ^* . Megow and Wiese [23] and Chen et al. [9] independently provided competitive-ratio approximation schemes that compute algorithms with a competitive ratio not greater than $(1 + \varepsilon)\rho^*$ for any positive ε . However the best possible competitive ratio is still unknown.

In more general case of machines with arbitrary speeds, the best general bounds are the following. Berman et al. [7] provided an 5.828-competitive algorithm, and Ebenlendr and Sgall [12] proved a lower bound of 2.564. If $m = 2$ (and the speeds are 1 and s), then the greedy strategy of Graham is again useful - in every step we choose the machine which will finish the actual job as soon as possible. Epstein showed that such algorithm is optimal and has competitive ratio $\min\{1 + \frac{s}{1+s}, 1 + \frac{1}{s}\}$. We will later see that in semi-online variants the situation is much more complex.

How much can we gain if some information is known in advance? In this paper we are interested in the semi-online scheme when only the optimal offline value is known (OPT version, for short), although there are many other studied semi-online models. For example: known sum of jobs (SUM version) [20, 3, 4, 5, 24, 11, 21], known largest job [8, 2, 21], scheduling with a buffer [20], information that the last job will be the largest [15], combinations of them. Interestingly, the relation between SUM and OPT versions is very strong: for m (nonidentical) machines the optimal competitive ratio of OPT version is at most the optimal competitive ratio for SUM version (see Dósa et al. [11], first stated implicitly

for equal speeds by Chang et al. [10]). In fact, every algorithm for SUM version, can be modified and used for OPT version with the same competitive ratio.

Azar and Regev [6] were first to investigate the OPT version for identical machines (under the name of bin stretching), although the observation about the relation with the SUM version implies that the first upper bound of $\frac{4}{3}$ for the case of two identical machines follows from the work of Kellerer et al. [20] (now it is known to be optimal). Azar and Regev [6], Cheng et al. [10] and Lee and Lim [21] continued the progress for the case of more than 2 identical machines.

Since we are interested in the OPT version with non-identical speeds on two uniform machines, we will state the previous results in terms of s (recall that the speeds are 1 and s). Epstein [14] was first to investigate this problem. She proved the following bounds for the optimal competitive ratio $r^*(s)$:

$$r^*(s) : \begin{cases} r^*(s) \in \left[\frac{3s+1}{3s}, \frac{2s+2}{2s+1} \right] & \text{for } s \in [1, q_E \approx 1.1243] \\ r^*(s) \in \left[s \left(\frac{3}{4} + \frac{\sqrt{65}}{20} \right), \frac{2s+2}{2s+1} \right] & \text{for } s \in [q_E, \frac{1+\sqrt{65}}{8} \approx 1.1328] \\ r^*(s) = \frac{2s+2}{2s+1} & \text{for } s \in \left[\frac{1+\sqrt{65}}{8}, \frac{1+\sqrt{17}}{4} \approx 1.2808 \right] \\ r^*(s) = s & \text{for } s \in \left[\frac{1+\sqrt{17}}{4}, \frac{1+\sqrt{3}}{2} \approx 1.3660 \right] \\ r^*(s) \in \left[\frac{2s+1}{2s}, s \right] & \text{for } s \in \left[\frac{1+\sqrt{3}}{2}, \sqrt{2} \approx 1.4142 \right] \\ r^*(s) \in \left[\frac{2s+1}{2s}, \frac{s+2}{s+1} \right] & \text{for } s \in \left[\sqrt{2}, \frac{1+\sqrt{5}}{2} \approx 1.6180 \right] \\ r^*(s) \in \left[\frac{s+2}{s+1}, \frac{s+2}{s+1} \right] & \text{for } s \in \left[\frac{1+\sqrt{5}}{2}, \sqrt{3} \approx 1.7321 \right] \\ r^*(s) = \frac{s+2}{s+1} & \text{for } s \geq \sqrt{3} \end{cases}$$

Where q_E is the solution of $36x^2 - 135x^3 + 45x^2 + 60x + 10 = 0$. Ng et al. [24] presented algorithms giving the following upper bounds:

$$r^*(s) \leq \begin{cases} \frac{2s+1}{2s} & \text{for } s \in \left[\frac{1+\sqrt{3}}{2}, \frac{1+\sqrt{21}}{4} \approx 1.3956 \right] \\ \frac{6s+6}{4s+5} & \text{for } s \in \left[\frac{1+\sqrt{21}}{4}, \frac{1+\sqrt{13}}{3} \approx 1.5352 \right] \\ \frac{12s+10}{9s+7} & \text{for } s \in \left[\frac{1+\sqrt{13}}{3}, \frac{5+\sqrt{241}}{12} \approx 1.7104 \right] \\ \frac{2s+3}{4s+3} & \text{for } s \in \left[\frac{5+\sqrt{241}}{12}, \sqrt{3} \right] \end{cases}$$

and provided the following lower bounds:

$$r^*(s) \geq \begin{cases} \frac{3s+5}{2s+4} & \text{for } s \in [\sqrt{2}, \frac{\sqrt{21}}{3} \approx 1.5275] \\ \frac{3s+3}{3s+1} & \text{for } s \in [\frac{\sqrt{21}}{3}, \frac{5+\sqrt{193}}{12} \approx 1.5744] \\ \frac{4s+2}{2s+3} & \text{for } s \in [\frac{5+\sqrt{193}}{12}, \frac{7+\sqrt{145}}{12} \approx 1.5868] \\ \frac{5s+2}{4s+1} & \text{for } s \in [\frac{7+\sqrt{145}}{19}, \frac{9+\sqrt{193}}{14} \approx 1.6352] \\ \frac{7s+4}{7s} & \text{for } s \in [\frac{9+\sqrt{193}}{14}, \frac{5}{3}] \\ \frac{7s+4}{4s+5} & \text{for } s \in [\frac{5}{3}, \frac{5+\sqrt{73}}{8} \approx 1.6930] \end{cases}$$

Finally, Dósa et al. [11] provided following bounds:

$$r^*(s) \geq \begin{cases} \frac{8s+5}{5s+5} & \text{for } s \in [\frac{5+\sqrt{205}}{18}, \frac{1+\sqrt{31}}{6} \approx 1.0946] \\ \frac{2s+2}{2s+1} & \text{for } s \in [\frac{1+\sqrt{31}}{6}, \frac{1+\sqrt{17}}{4} \approx 1.2808] \end{cases}$$

$$r^*(s) \leq \begin{cases} \frac{3s+1}{3s} & \text{for } s \in [1, q_D \approx 1.071] \\ \frac{7s+6}{4s+6} & \text{for } s \in [q_D, \frac{1+\sqrt{145}}{12} \approx 1.0868] \end{cases}$$

Where q_D is the unique root of equation $3s^2(9s^2 - s - 5) = (3s + 1)(5s + 5 - 6s^2)$. For a visual summary (with our contribution included), see Figures 1–4. Whenever the dotted line (that represents an upper bound) is on an unbroken line (that represents a lower bound), the optimal competitive ratio is known. As a consequence of all these results, before this publication three intervals were open, namely $(q_d, \frac{1+\sqrt{31}}{6}) \approx (1.071, 1.0946)$ and $(\frac{1+\sqrt{21}}{4}, \sqrt{3}) \approx (1.3660, 1.7321)$. We will call the latter one the right interval.

There are also two variations of the aforementioned problems we find worth mentioning, as those variations are studied also in the case of OPT version. In the scheduling with *preemption* model it is allowed to split a job into multiple parts and assign to different machines, as long as those parts will be processed in disjoint time intervals. Ebenlendr and Sgall [13] presented one optimal preemptive algorithm working for various semi-online conditions and their combinations, including OPT version, SUM version, known longest job, jobs sorted in decreasing order - this seems to be much different from the situation in models without preemption. In the second variation, called *under a grade of service provision* (GoS), not every job can be processed by every machine: there is a level function for jobs, and job with level i can be processed only by machines with index at most i . GoS-OPT version for two (nonidentical) machines is also solved due to Lu and Liu [22], with the optimal competitive ratio being $\min\{\frac{1+2s}{1+s}, \frac{1+s}{s}\}$, again equal to the optimal competitive ratio of GoS-SUM version.

1.2 Our Contribution

We deal with the semi-online two uniform machines scheduling problem with a “known opt” condition, that is, $Q_2|OPT|C_{max}$ according to three field notation

introduced by Graham et al. [19]. This problem is studied on two parts of the right interval, $[\frac{1+\sqrt{21}}{4}, \frac{3+\sqrt{73}}{8}] \approx [1.3956, 1.443]$ and $[\frac{5}{3}, \frac{4+\sqrt{133}}{9}] \approx [\frac{5}{3}, 1.7258]$, for which we give new lower bound constructions. (Note that these numbers are solutions of certain equations, and will be formally introduced in the following section.) We apply an adversary strategy, that is, depending on the current assignment of a given job the adversary defines the next job that makes life complicated for the algorithm. We show that the input can always be continued in such malicious way, that any kind of algorithm will at some step exceed the lower bound on the makespan.

“Marry, and you will regret it; don’t marry, you will also regret it; marry or don’t marry, you will regret it either way” - says the Danish philosopher Søren Kierkegaard. He describes a decision situation from which two different possible choices will lead into the future. But no matter how the person decides, the continuation will not lead to a happy end. Both can be seen as “unhappy situations” (from the person’s point of view who has to make the decision). This in our scheduling setting will correspond to the three “Final Cases” that are described in Section 2.3. The scheduler (the algorithm) can make a decision, similar to the person deciding for or against a marriage. No matter how the algorithm decides, it will lead to an “unhappy situation”, since it is possible to generate further jobs, based on its decision, in such way that the competitive ratio of the algorithm is relatively high. It will not be obvious at first sight that the algorithm is always trapped in a situation that leads to an unhappy ending; it evolves over several rounds of further jobs that are determined in what we call “intermediate cases” or “final cases”. At most, eight jobs need to be generated to close the trap.

The moves of the algorithm and the generation of jobs can be seen as a two player game, such as chess. One player is the algorithm, and the other player is the adversary that constructs malicious jobs. What our result then shows is that the player of the algorithm is “checkmate” after at most eight moves. Our purpose was not to give a difficult construction, however, we were not able to get the tight bounds with less than eight jobs.

Together with the upper bound result of Ng et al. [24] we obtain that our new lower bound is tight (that is, the algorithms presented by Ng et al. are optimal) for any value of s in one of these intervals: $[\frac{1+\sqrt{21}}{4}, \frac{3+\sqrt{73}}{8}] \approx [1.3956, 1.443]$, $[\frac{5}{3}, \frac{13+\sqrt{1429}}{30}] \approx [\frac{5}{3}, 1.6934]$, and $[\frac{31+\sqrt{8305}}{72}, \frac{5+\sqrt{241}}{12}] \approx [1.6963, 1.7103]$, see also Figure 1.




Figure 1: Our new lower bound in comparison with existing lower and upper bounds from Epstein [14], Ng et al. [24], and Dósa et al. [11].




Figure 2: Zooming into the left part of Figure 1.




Figure 3: Zooming into the right part of Figure 1.




Figure 4: Zooming into the middle part of Figure 3.

2 Preliminaries and Notations

Let OPT and SUM mean, respectively, the known optimum value (given by some oracle), and the total size of the jobs. By \mathcal{J}_t we denote the family of all sets of jobs with optimum value of t . In other words, if the oracle returns a value of OPT , then there is a guarantee that the set of jobs belongs to \mathcal{J}_{OPT} . We denote the prescribed competitive ratio (that we do not want to violate) by r .

Lemma 1 *Given a set of jobs. Assume we are able to assign these jobs to the two machines in such way that M1 receives a load of t and M2 a load of $s \cdot t$, then this set of jobs belongs to \mathcal{J}_t .*

Proof. The assignment of jobs given in the formulation of the lemma is a feasible solution with a makespan of t . It remains to show that there is no better solution. The sum of jobs (the total load) is $(s + 1) \cdot t$. Assume there is a better assignment with makespan $t' < t$. Then the load on machine M1 can be at most t' . Hence the load on machine M2 is the remaining load, which is at least $st + (t - t') > st > st'$. This contradicts the assumption that t' is the makespan. ■

As a consequence of Lemma 1, we remark that $\text{SUM} \leq (s + 1) \cdot \text{OPT}$, and that the size of any job is at most $s \cdot \text{OPT}$. We denote $\overline{\text{SUM}} := (s + 1) \cdot \text{OPT}$.

2.1 Definitions

Let $q_1 := \frac{\sqrt{21}+1}{4} \approx 1.3956$, which is the positive solution of $\frac{2s+1}{2s} = \frac{6s+6}{4s+5}$.

Let $q_2 := \frac{\sqrt{73}+3}{8} \approx 1.443$, which is the positive solution of $\frac{6s+6}{4s+5} = \frac{5s+2}{4s+1}$.

Let $q_3 := \frac{13+\sqrt{1429}}{30} \approx 1.6934$, which is the positive solution of $\frac{12s+10}{9s+7} = \frac{18s+16}{16s+7}$.

Let $q_4 := \frac{30+7\sqrt{186}}{74} \approx 1.6955$, which is the positive solution of $\frac{18s+16}{16s+7} = \frac{8s+7}{3s+10}$.

Let $q_5 := \frac{31+\sqrt{8305}}{72} \approx 1.6963$, which is the positive solution of $\frac{8s+7}{3s+10} = \frac{12s+10}{9s+7}$.

Let $q_6 := \frac{4+\sqrt{133}}{9} \approx 1.7258$, which is the positive solution of $\frac{12s+10}{9s+7} = \frac{s+1}{2}$.

We note that in this paper we do not consider speeds between q_2 and $\frac{5}{3}$.

Let

$$r(s) = \begin{cases} r_1(s) := \frac{6s+6}{4s+5} & \text{if } q_1 \leq s \leq q_2 \approx 1.443 & \text{i.e. } s \text{ is small} \\ r_2(s) := \frac{12s+10}{9s+7} & \text{if } \frac{5}{3} \leq s \leq q_3 \approx 1.6934 & \text{i.e. } s \text{ is smaller regular} \\ r_3(s) := \frac{18s+16}{16s+7}, & \text{if } q_3 \leq s \leq q_4 \approx 1.6955 & \text{i.e. } s \text{ is smaller medium} \\ r_4(s) := \frac{8s+7}{3s+10}, & \text{if } q_4 \leq s \leq q_5 \approx 1.6963 & \text{i.e. } s \text{ is bigger medium} \\ r_2(s) := \frac{12s+10}{9s+7} & \text{if } q_5 \leq s \leq q_6 \approx 1.7258 & \text{i.e. } s \text{ is bigger regular} \end{cases}$$

As we will show in the very end (cf. Theorem 16), this function will be our lower bound on the optimal competitive ratio.




Figure 5: Safe sets.

As abbreviations, we call s being regular, if s is smaller regular or bigger regular, and we call s being medium, if s is smaller medium or bigger medium.

Now we define the so called "safe sets". A safe set is a time interval on some of the machines, and it is safe in sense, that if the load of the machine is in this interval, this enables a "smart" algorithm to finish the schedule by not violating the desired competitive ratio. In other words, from the view of point of a lower bound construction (the adversary), we must avoid that the algorithm can assign the actual job in a way that the increased load of some machine will be inside a safe set.

Safe sets S_1 and S_3 are defined on the second machine, while safe sets S_2 and S_4 are defined on the first machine. We introduce notations for the top, bottom, and length of the safe sets. Thus let $S_i = [B_i, T_i]$, and $D_i = T_i - B_i$, for any $1 \leq i \leq 4$, see Figure 5. We will show in Section 2.2 that these intervals are well-defined.

The optimum value is known. For the sake of simplicity let us assume without loss of generality that $\text{OPT} = 1$. (This can be done by normalization: If OPT differs from being unit, all values are multiplied by the value of OPT^{-1} .)

Then the boundaries of safe sets S_i are defined as follows. Below we show that the safe sets are properly defined.

1. $B_1 = s + 1 - r$, and $T_1 = rs$, thus $D_1 = (s + 1)(r - 1)$,
2. $B_2 = s + 1 - sr$, and $T_2 = r$, thus $D_2 = (s + 1)(r - 1)$,
3. $B_3 = 2s - 2r - rs + 2$, and $T_3 = s(r - 1)$, thus $D_3 = 2r - 3s + 2rs - 2$,
4. $B_4 = 4s - 2r - 3rs + 3$, and $T_4 = r - 1$, thus $D_4 = (3r - 4)(s + 1)$,

We introduce as abbreviations some expressions that are used in the sequel.

$$\begin{aligned} a &:= T_4 - B_3, \\ b &:= T_3 - B_2, \\ c &:= \text{OPT} - D_1, \end{aligned}$$

and, if $s \geq \frac{5}{3}$, let

$$d := b - c - B_4.$$

For bigger regular speeds, we will also need the next notations:

$$\begin{aligned} e &:= \frac{1}{2}(b - 2c - a - B_4), \\ f &:= \frac{1}{2}(a + b - B_4), \\ g &:= \frac{1}{2}(b - a - B_4). \end{aligned}$$

By easy calculations we get the following expressions:

$$\begin{aligned} a &= 3r - 2s + rs - 3, \\ b &= 2rs - 2s - 1, \\ c &= s - r - rs + 2, \\ d &= 3r - 7s + 6rs - 6. \end{aligned}$$

Note that

$$\begin{aligned} a + b &= D_4, \\ e + f &= d, \\ f + g &= b - B_4, \\ f - g &= a, \\ c + e &= g. \end{aligned}$$

And for the values of e , f and g , we get the following equalities:

$$\begin{aligned} e &= \frac{1}{2}r - 3s + 3rs - \frac{5}{2}, \\ f &= \frac{5}{2}r - 4s + 3rs - \frac{7}{2}, \\ g &= 2rs - 2s - \frac{1}{2}r - \frac{1}{2}. \end{aligned}$$

2.2 General Properties

In Figure 6 we show plots of the functions $r_2(s)$, $r_3(s)$, and $r_4(s)$. If s is medium, then both $r_3(s)$ and $r_4(s)$ are below $r_2(s)$. Moreover $r_3(s) \geq r_4(s)$, if s is small medium, and the opposite inequality holds if s is bigger medium. Note, that s is medium sized only on a very narrow interval.




Figure 6: Comparing $r_2(s)$ (red), $r_3(s)$ (green), and $r_4(s)$ (blue).

- Lemma 2**
1. $r_1(s) \leq r_2(s)$ if $s \leq q_2$,
 2. $r_2(s) \leq r_1(s)$ if $s \geq \frac{5}{3}$,
 3. $r_3(s) \leq r_2(s)$ if $s \geq q_3$,
 4. $r_4(s) \leq r_2(s)$ if $s \leq q_5$.

Proof.

1. This estimation was already proven in [24]. We repeat it here for the sake of completeness. The inequality $r_1(s) \leq r_2(s)$ is equivalent to $(12s + 10)(4s + 5) - (6s + 6)(9s + 7) = -6s^2 + 4s + 8 \geq 0$, which holds iff $\frac{1-\sqrt{13}}{3} \leq s \leq \frac{1+\sqrt{13}}{3} \approx 1.5352$. Hence it holds for all $s \leq q_2$.
2. Follows from the previous computations.
3. The inequality $r_3(s) \leq r_2(s)$ is equivalent to $(12s + 10)(16s + 7) - (18s + 16)(9s + 7) = 30s^2 - 26s - 42 \geq 0$, which holds iff $s \leq \frac{13-\sqrt{1429}}{30}$ or $s \geq \frac{13+\sqrt{1429}}{30} = q_3$.
4. The inequality $r_4(s) \leq r_2(s)$ is equivalent to $(12s + 10)(3s + 10) - (8s + 7)(9s + 7) = -36s^2 + 31s + 51 \geq 0$, which holds iff $\frac{31-\sqrt{8305}}{72} \leq s \leq \frac{31+\sqrt{8305}}{72} = q_5$.

■

In the next lemma we prove a lower and upper bound result on $r(s)$. These bounds are needed to show that the safe sets are well-defined.

Lemma 3 1. $\frac{3s+2}{2s+2} < \frac{4}{3} < 1.35 < r(s) < \min\left\{\frac{4s+3}{3s+2}, \frac{s+2}{s+1}\right\} < \frac{2s+2}{s+2} < \frac{2s+1}{s+1}$
hold in all considered domain of the function r , i.e., for all $s \in [q_1, q_2] \cup [\frac{5}{3}, q_6] =: \text{Dom}(r)$.

2. *If $s \geq \frac{5}{3}$, we have $r(s) \geq \frac{8s+7}{6s+5}$.*

Proof.

1. The leftmost lower bound holds as $\frac{3s+2}{2s+2} < \frac{4}{3}$ is equivalent to $4(2s+2) - 3(3s+2) > 0$, hence $s < 2$.

Now we show that $r(s) > 1.35$.

- For $q_1 \leq s \leq q_2$ (where $r(s) = r_1(s)$), we get $0 < \frac{6s+6}{4s+5} - \frac{135}{100} = \frac{12s-15}{80s+100}$, which is true since $s > \frac{5}{4}$.
- For $\frac{5}{3} \leq s \leq q_3$ (where $r(s) = r_2(s)$), we similarly obtain $0 < \frac{12s+10}{9s+7} - \frac{135}{100} = \frac{11-3s}{20(9s+7)}$, which is true since $s < \frac{11}{3}$. This also includes the case $q_5 \leq s \leq q_6$, where also $r(s) = r_2(s)$.
- For $q_3 \leq s \leq q_4$ (where $r(s) = r_3(s)$), we get $0 < \frac{18s+16}{16s+7} - \frac{135}{100} = \frac{131-72s}{20(16s+7)}$, which is true since $s < \frac{131}{72} \approx 1.8194$.
- For $q_4 \leq s \leq q_5$ (where $r(s) = r_4(s)$), we get $0 < \frac{8s+7}{3s+10} - \frac{135}{100} = \frac{79s-130}{20(3s+10)}$, which is true since $s > \frac{130}{79} \approx 1.6456$.

Hence $r(s) > 1.35$.

Regarding the rightmost upper bound, $\frac{2s+2}{s+2} < \frac{2s+1}{s+1}$ holds since $\frac{2s+1}{s+1} - \frac{2s+2}{s+2} = \frac{s}{(s+2)(s+1)} > 0$.

Moreover, $\frac{2s+2}{s+2} - \frac{4s+3}{3s+2} = \frac{2s^2-s-2}{(s+2)(3s+2)} > 0$, which holds since $2s^2 - s - 2 > 0$ for all $s < \frac{1-\sqrt{17}}{4}$ or $s > \frac{1+\sqrt{17}}{4} \approx 1.2808$. Thus, it holds that $\min\left\{\frac{4s+3}{3s+2}, \frac{s+2}{s+1}\right\} < \frac{2s+2}{s+2}$.

Thus it remained to show that $r < \min\left\{\frac{s+2}{s+1}, \frac{4s+3}{3s+2}\right\}$. Note that $\frac{4s+3}{3s+2} \leq \frac{s+2}{s+1}$ holds for all $s \in \text{Dom}(r)$, iff $(4s+3)(s+1) - (s+2)(3s+2) = s^2 - s - 1 \leq 0$, i.e., $\frac{1-\sqrt{5}}{2} \leq s \leq \frac{1+\sqrt{5}}{2} \approx 1.618$. Thus, if $s \leq q_2$, then we need to verify that $r < \frac{4s+3}{3s+2}$, otherwise, if $s \geq \frac{5}{3}$, we need to verify that $r < \frac{s+2}{s+1}$.

For $q_1 \leq s \leq q_2$ (where $r(s) = r_1(s)$), we get $\frac{4s+3}{3s+2} - \frac{6s+6}{4s+5} = \frac{2s+3-2s^2}{(4s+5)(3s+2)} > 0$, which holds since $2s+3-2s^2 > 0$ for all $\frac{1-\sqrt{7}}{2} < s < \frac{1+\sqrt{7}}{2} \approx 1.8229$.

Now let us consider the case when $\frac{5}{3} \leq s \leq q_6$. For the two cases where $r(s) = r_2(s)$ we have that $\frac{12s+10}{9s+7} < \frac{s+2}{s+1}$ holds since $\frac{s+2}{s+1} - \frac{12s+10}{9s+7} = \frac{-3s^2+3s+4}{(s+1)(9s+7)} > 0$, which holds since $-3s^2+3s+4 > 0$ for all $\frac{1}{2} - \frac{\sqrt{57}}{6} < s < \frac{1}{2} + \frac{\sqrt{57}}{6} \approx 1.7583$.

In Lemma 2 we showed that $r_3(s) \leq r_2(s)$ for all $s \geq q_3$ and that $r_4(s) \leq r_2(s)$ for all $s \leq q_5$. From this the claimed upper bound on $r(s)$ follows.

2. For $r(s) = r_2(s)$ we get $\frac{12s+10}{9s+7} - \frac{8s+7}{6s+5} = \frac{s+1}{(6s+5)(9s+7)} > 0$.

For $r(s) = r_3(s)$ we get $\frac{18s+16}{16s+7} - \frac{8s+7}{6s+5} = \frac{31+18s-20s^2}{(6s+5)(16s+7)} > 0$ for all $s < \frac{9+\sqrt{701}}{20} \approx 1.7738$.

For $r(s) = r_4(s)$ we get $\frac{8s+7}{3s+10} - \frac{8s+7}{6s+5} = \frac{(8s+7)(3s-5)}{(3s+10)(6s+5)} \geq 0$, because $s \geq \frac{5}{3}$.

■

Lemma 4 1. $D_1 = D_2$,

2. $T_1 - T_3 = s \cdot OPT$ and $T_2 - T_4 = OPT$,

3. $B_3 = B_1 - D_1$,

4. $B_4 = B_2 - D_3$,

5. $B_2 + T_3 = OPT$,

6. $T_1 + B_2 = T_2 + B_1 = \overline{SUM}$.

Proof. All properties are checked using the definition of the safe sets, where we make use of the assumption $OPT = 1$.

1. $D_1 = D_2$ holds directly by definition.

2. $T_1 - T_3 = rs - s(r-1) = s = s \cdot OPT$ and $T_2 - T_4 = r - (r-1) = 1 = OPT$.

3. $B_3 + D_1 = (2s - 2r - rs + 2) + (s+1)(r-1) = s - r + 1 = B_1$.

4. $B_4 + D_3 = (4s - 2r - 3rs + 3) + (2r - 3s + 2rs - 2) = s + 1 - rs = B_2$.

5. $T_3 + T_4 = s(r-1) + (r-1) = (s+1)(r-1) = D_1$.

6. $B_2 + T_3 = (s+1 - sr) + s(r-1) = 1 = OPT$.

7. $T_1 + B_2 = rs + (s+1 - sr) = s+1 = \overline{SUM}$. Moreover $T_2 + B_1 = r + (s+1 - r) = s+1 = \overline{SUM}$.

■

Now we show that the definition of the safe sets is of sense, these sets do not intersect each other, and they follow each other on the machines.

Lemma 5 1. $0 < B_3 < T_3 < B_1 < T_1$.

2. $0 < B_4 < T_4 < B_2 < T_2$.

Proof. In the calculations we generally use Lemma 3, if not otherwise stated.

1. From $r < \frac{2s+2}{s+2}$ follows $0 < 2s - 2r - rs + 2 = B_3$. From $r > \frac{3s+2}{2s+2}$ follows $0 < 2r + 2rs - 3s - 2 = D_3 = T_3 - B_3$. From $r < \frac{2s+1}{s+1}$ follows $0 < (s+1-r) - s(r-1) = B_1 - T_3$. From the definition we have that $0 < (s+1)(r-1) = D_1 = T_1 - B_1$.

2. From $r < \frac{4s+3}{3s+2}$ follows $0 < 4s + 3 - 3rs - 2r = B_4$. From $r > \frac{4}{3}$ and the definition we have that $0 < (3r-4)(s+1) = D_4 = T_4 - B_4$. From $r < \frac{s+2}{s+1}$ follows $0 < (s+1-sr) - (r-1) = B_2 - T_4$. From the definition we have that $0 < (s+1)(r-1) = T_2 - B_2$.

■

Now we have seen that the safe sets are properly defined. We will need some further bounds on r . With their help, we can prove several properties of the expressions we introduced.

Lemma 6 *The following bounds on r are valid:*

1. $\frac{2s+1}{2s} \leq r < s$.

2. If $s \geq \frac{5}{3}$, then $\frac{7s+6}{6s+3} \leq r \leq \frac{3s+5}{2s+4}$.

3. If s is regular, then we have $\max\left\{\frac{6s+5}{6s+1}, \frac{8s+7}{6s+5}\right\} \leq r$.

Proof.

1. This bound was already proven in [24] (cf. Figure 1 in [24]). We give it here for the sake of completeness. Regarding the lower bound, if s is small (i.e., $q_1 \leq s \leq q_2$), then $\frac{2s+1}{2s} \leq r = r_1(s) = \frac{6s+6}{4s+5}$ holds, since $2s(6s+6) - (2s+1)(4s+5) = 4s^2 - 2s - 5 \geq 0$, if $s \geq \frac{1+\sqrt{21}}{4} = q_1$. For $s \geq \frac{5}{3}$, we know from Lemma 3 that $\frac{4}{3} < r$, and thus get $\frac{4}{3} - \frac{2s+1}{2s} = \frac{2s-3}{6s} > 0$. Now let us consider the upper bound. For a small speed ratio s , we get $s - \frac{6s+6}{4s+5} = \frac{4s^2-s-6}{4s+5} > 0$, if $s \geq \frac{1+\sqrt{97}}{8} \approx 1.3561$, in particular, for $s \geq q_1$. For $s \geq \frac{5}{3}$, the statement follows from Lemma 2.

2. Let us consider the lower bound. In the case $r = r_2(s)$, we get $\frac{12s+10}{9s+7} - \frac{7s+6}{6s+3} = \frac{9s^2-7s-12}{3(9s+7)(2s+1)} \geq 0$, if $s \geq \frac{7+\sqrt{481}}{18} \approx 1.6073$. In the case $r = r_3(s)$, we get $\frac{18s+16}{16s+7} - \frac{7s+6}{6s+3} = \frac{(4s+3)(2-s)}{3(2s+1)(16s+7)} > 0$. Finally, in the case $r = r_4(s)$, we get $\frac{8s+7}{3s+10} - \frac{7s+6}{6s+3} = \frac{27s^2-22s-39}{3(3s+10)(2s+1)} \geq 0$, if $s \geq \frac{11+\sqrt{1174}}{27} \approx 1.6764$, which in particular holds for $s \geq q_4$. To prove the other upper bound, it suffices to show that $r_2(s) \leq \frac{3s+5}{2s+4}$, since $r \leq r_2(s)$ by Lemma 2. We get $\frac{3s+5}{2s+4} - \frac{12s+10}{9s+7} = \frac{(s+1)(3s-5)}{2(s+2)(9s+7)} \geq 0$.

3. If s is regular, we get $\frac{12s+10}{9s+7} - \frac{6s+5}{6s+1} = \frac{(6s+5)(3s-5)}{(9s+7)(6s+1)} \geq 0$, and we get $\frac{12s+10}{9s+7} - \frac{8s+7}{6s+5} = \frac{s+1}{(6s+5)(9s+7)} > 0$ (cf. proof of Lemma 3.2).

■

In the next lemma we prove properties regarding the just introduced expressions. Note that d is defined only if $s \geq \frac{5}{3}$, and e, f, g are defined only if s is bigger regular.

- Lemma 7**
1. $b, c, d, e, f, g \geq 0$,
 2. If $s \geq \frac{5}{3}$, then $a \leq c$,
 3. $s > T_2$,
 4. $B_1 > OPT$.

Proof.

1. In the proof we use Lemma 3 and Lemma 6.
 - $b \geq 0$, since $r \geq \frac{2s+1}{2s}$.
 - $c > 0$, since $r < \frac{s+2}{s+1}$.
 - $d \geq 0$, since $r \geq \frac{7s+6}{6s+3}$.
 - $e > 0$, since $r > \frac{6s+5}{6s+1}$. The strict inequality follows from the proof of Lemma 6, because $e = 0$ can only occur for $s = \frac{5}{3}$, which means, s would be smaller regular.
 - $f > 0$, since $r > \frac{8s+7}{6s+5}$. The strict inequality can also be seen in the proof of Lemma 6.
 - $g > 0$, since $g = c + e > 0$.
2. If $s \geq \frac{5}{3}$, then $a \leq c$, since $c - a = (s - r - rs + 2) - (3r - 2s + rs - 3) = 3s - 4r - 2rs + 5 \geq 0$, where the last estimation uses $r \leq \frac{3s+5}{2s+4}$.
3. Remind you that $OPT = 1$. We obtain $s - T_2 = s - r > 0$, since $r < s$ from Lemma 6.1.
4. $B_1 - OPT = B_1 - 1 = (s+1-r) - 1 = s - r > 0$, since $r < s$ by Lemma 6.1.

■

In fact, we did not forget to prove $a \geq 0$, as the following remark shows.

Remark 8 If $s > \frac{5+\sqrt{241}}{12} = 1.7103$, then a is negative.

Proof. If $s > \frac{5+\sqrt{241}}{12}$, then s is bigger regular. Hence $r = r_2(s) = \frac{12s+10}{9s+7}$. Substituting this expression, we thus get $a = 3r - 2s + rs - 3 = \frac{-6s^2+5s+9}{9s+7} > 0$, since $s > \frac{5+\sqrt{241}}{12}$. ■

2.3 General Subcases

The general idea to show that no semi-online algorithm knowing s and OPT can ever be better than the ratio $r(s)$ is to construct a malicious sequence of jobs that are in \mathcal{J}_{OPT} , but force any algorithm to schedule them in such way that they have a makespan of at least r . We construct this sequence iteratively, depending on the previous assignment choices of the algorithm. This leads to a number of cases that need to be considered separately. Some cases are “final” (or terminal): If such case is entered, the algorithm is trapped by adding just one or a few more jobs, which ends in a situation described in the assumption of the respective cases. When entering the case, the algorithm is trapped, because we can then construct one or a few more jobs, which make the algorithm to overshoot the desired makespan.

In this sense, the following three “G”-cases, G1, G2, and G3, are general, because most of the other cases, independent of the particular interval of s , will lead to them.

We will denote by L_1 and L_2 the current load of machine M1 and M2, respectively. Moreover let L'_1 and L'_2 denote the increased load of that machine, if the actual job is assigned there.

Final Case G1. Suppose $T_3 \leq L_2$ and $L_1 + L_2 \leq \text{OPT}$.

Note that $\text{OPT} = B_2 + T_3 > T_3$ holds by Lemma 4.5. In this situation, let the next job be $A = s \cdot \text{OPT}$ and $B = \text{OPT} - (L_1 + L_2) \geq 0$. If A is assigned to M2, we get that the new M2-load $L'_2 \geq T_3 + s \cdot \text{OPT} = T_1$ by Lemma 4.2, thus we are done. Otherwise A is assigned to M1. Then the new M1-load $L'_1 \geq s \cdot \text{OPT} = s > T_2$, by Lemma 7.3, we are done again.

The set of all jobs that were initially on M1 and M2, plus jobs A and B belong to \mathcal{J}_{OPT} by Lemma 1, if we assign A to M2 and all other jobs to M1.

Final Case G2. Suppose $L_1 = B_2$ and $0 \leq L_2 \leq B_3$, and there exists an already job with size c (where c as defined above.)

The next and last jobs are $B = D_1$ and

$$\begin{aligned} C &= \overline{\text{SUM}} - (L_1 + L_2 + B) \geq (T_2 + B_1) - (B_2 + D_2) - B_3 \\ &= B_1 - B_3 = D_1 = B \end{aligned}$$

using $\overline{\text{SUM}} = T_2 + B_1$, $D_1 = D_2$, and $B_1 - B_3 = D_1$ from Lemma 4.1, 4.3, and 4.6. If any of B and C is assigned to M1, the lower bound holds since $L'_1 \geq B_2 + D_1 = B_2 + D_2 = T_2$. Otherwise both jobs go to M2 and we are done again as $L'_2 = \overline{\text{SUM}} - B_2 = T_1$ by Lemma 4.6.

The set of jobs belong to \mathcal{J}_{OPT} by Lemma 1: assign jobs c and $B = D_1$ to machine M1, then $L'_1 = c + D_1 = \text{OPT}$ by the definition of c . All remaining jobs go to machine M2, then $L'_2 = B_2 + L_2 - c + \overline{\text{SUM}} - (L_1 + L_2 + B) = B_2 - c + \overline{\text{SUM}} - L_1 - B = -c + \overline{\text{SUM}} - B = \overline{\text{SUM}} - \text{OPT} = s \cdot \text{OPT}$.

Final Case G3. Suppose $L_1 = T_4$ and $L_2 = 0$.

The next and last two jobs are $B = \text{OPT}$ and $C = \overline{\text{SUM}} - (T_4 + \text{OPT}) = (T_2 + B_1) - T_4 - \text{OPT} = B_1 > \text{OPT}$, by Lemma 4.6, 4.2, and 7.4. If any of B or C is assigned to M1, then $L'_1 \geq T_4 + \text{OPT} = T_2$ by Lemma 4.2. Thus the lower bound holds. Otherwise both go to M2 and $L'_2 = \overline{\text{SUM}} - T_4 > \overline{\text{SUM}} - B_2 = T_1$ by Lemma 5.2 and 4.6, and we are done again.

The jobs belong to \mathcal{J}_{OPT} by Lemma 1, because we can assign $B = \text{OPT}$ to machine M1 and all other jobs to machine M2, which have a total sum of $T_4 + C = T_4 + \overline{\text{SUM}} - (T_4 + \text{OPT}) = \overline{\text{SUM}} - \text{OPT} = (s+1)\text{OPT} - \text{OPT} = s\text{OPT}$.

3 Lower Bound for Small s

At the end of this section we will prove that $r_1(s)$ is a lower bound on the competitive ratio for small s . This is done by constructing a sequence of jobs in \mathcal{J}_{OPT} that will force the algorithm to make bad decisions about their assignments, so that it soon will meet or overshoot the ratio r . Before giving this construction, we consider a number of cases, from which the lower bound can be soon achieved. We start with some further estimations.

3.1 Properties

Lemma 9 *If s is small, then $\frac{2s-1}{s} \leq r \leq \frac{5s+2}{4s+1} \leq \frac{2}{s}$.*

Proof.

- For the first estimation, we see that $\frac{6s+6}{4s+5} - \frac{2s-1}{s} = \frac{5-2s^2}{(4s+5)s} \geq 0$ holds for all $s \leq \frac{\sqrt{10}}{2} \approx 1.5811$, in particular, for small s .
- For the second estimation, we obtain that $\frac{5s+2}{4s+1} - \frac{6s+6}{4s+5} = \frac{-4s^2+3s+4}{(4s+1)(4s+5)} \geq 0$ for $s \leq \frac{3+\sqrt{73}}{8} = q_2$, in particular, for small s .
- For the third estimation, we compute that $\frac{2}{s} - \frac{5s+2}{4s+1} = \frac{-5s^2+6s+2}{s(4s+1)} \geq 0$ for all $s \leq \frac{3+\sqrt{19}}{5} \approx 1.4717$, in particular, for small s .

■

Lemma 10 1. $(T_3 - B_2) + c = b + c \leq B_4$,

2. $c \leq B_4$,

3. $2c \geq B_4$,

4. $2c \geq B_3$,

5. $B_3 = D_4$,

6. $c < B_3$,

7. $B_2 + B_4 < OPT$.

Proof.

1. $B_4 - b - c = (4s - 2r - 3rs + 3) - (2rs - 2s - 1) - (s - r - rs + 2) = 5s - r - 4rs + 2 \geq 0$, since $r \leq \frac{5s+2}{4s+1}$ by Lemma 9.
2. Follows from Lemma 10.1, together with $b \geq 0$ from Lemma 7.1.
3. $2c - B_4 = 2(s - r - rs + 2) - (4s - 2r - 3rs + 3) = rs - 2s + 1 \geq 0$, as $r \geq \frac{2s-1}{s}$ from Lemma 9.
4. $2c - B_3 = 2(s - r - rs + 2) - (2s - 2r - rs + 2) = 2 - rs \geq 0$, as $r \leq \frac{2}{s}$ from Lemma 9.
5. $B_3 - D_4 = (2s - 2r - rs + 2) - (3r - 4)(s + 1) = 6s + 6 - r(4s + 5) = 0$, since $r = r_1(s)$ for small s .
6. $B_3 - c = (2s - 2r - rs + 2) - (s - r - rs + 2) = s - r > 0$, as $r < s$ by Lemma 6.1.
7. Since $OPT = B_2 + T_3$ by Lemma 4.5, we have to show that $B_4 < T_3$. From Lemma 5.2 and the definitions of T_3, T_4 , it follows that $B_4 < T_4 = r - 1 < s(r - 1) = T_3$.

■

3.2 Subcases

The adversary constructs a sequence of jobs in such way that any assignment strategy of an arbitrary algorithm will lead to one of the ‘‘S’’-cases described below, which make use of the fact that s is small. When the assumptions of these cases are fulfilled, the adversary knows how to define the next jobs, so that the algorithm is trapped, and must return a solution having a ratio worse-or-equal than $r = r_1(s)$.

Case S1. Suppose $L_1 = 0$ and $L_2 = B_4 - c \geq 0$ (by Lemma 10.2).

Let the next job be $A = B_2$. Suppose A is assigned to M2. Then $L'_1 + L'_2 = B_2 + B_4 - c \leq B_2 + T_4 \leq B_2 + T_3 = OPT$ holds (applying $c \geq 0$, $T_4 = r - 1 \leq s(r - 1) = T_3$ and Lemma 4.3). Moreover by Lemma 10.1 we get $L'_2 = B_2 + B_4 - c \geq T_3$, thus case G1 holds for the new loads L'_1, L'_2 , and hence we are done.

Otherwise A is assigned to M1. At this moment $L'_1 = B_2$ and $L'_2 = B_4 - c$. Then we are in case S2.

Final Case S2. Assume that $L'_1 = B_2$ and $L'_2 = B_4 - c$.

Then the next jobs are $B = \text{OPT} - (B_4 - c) \geq c - (B_4 - c) = 2c - B_4 \geq 0$ (by Lemma 10.3) and $C = \text{SUM} - (L'_1 + L'_2 + B) = (B_2 + T_1) - (B_2 + \text{OPT}) = T_1 - \text{OPT} = (T_1 - D_1) - c = B_1 - c \geq D_2$, since $B_1 \geq D_2 + c = \text{OPT}$ holds by Lemma 7.4. Note that $B = \text{OPT} - (B_4 - c) = (D_2 + c) - (B_4 - c) = D_2 + 2c - B_4 \geq D_2$ also holds by Lemma 10.3. If any of B or C is assigned to M1, we are done: If B is assigned to M1, then $L''_1 = L'_1 + B \geq B_2 + D_2 = T_2$, and similar for C . Otherwise both B and C are assigned to M2, and we are done again: $L''_2 = B_4 - c + B + C = B_4 - c + (\text{OPT} - (B_4 - c)) + (T_1 - \text{OPT}) = T_1$.

The set of jobs A, B, C and the previous load of M2 belong to \mathcal{J}_{OPT} (by Lemma 1): assign the previous load of machine M2 and B to machine M1, then its load is $L_2 + B = (B_4 - c) + (\text{OPT} - (B_4 - c)) = \text{OPT}$. The remaining jobs A and C go to machine M2, which then has a load of $A + C = B_2 + (B_1 - c) = (s+1-r) + (s+1+sr) - (s-r-rs+2) = s = s \cdot \text{OPT}$.

Case S3. Suppose $L_1 = B_4 - c$ and $L_2 = c$.

Let the next job be $A = B_2 - (B_4 - c) = D_3 + c$ by Lemma 4.4. Suppose A is assigned to M2. Then $L'_1 + L'_2 = (B_4 - c) + c + B_2 - (B_4 - c) = B_2 + c \leq B_2 + B_4 \leq \text{OPT}$ by Lemma 10.7. Moreover $L'_2 = D_3 + 2c \geq D_3 + B_3 = T_3 - B_3 + B_3 = T_3$, since $2c \geq B_3$ holds by Lemma 10.4. Thus case G1 holds for L'_1, L'_2 , and we are done. Otherwise A is assigned to M1. Then the loads are $L'_1 = B_4 - c + A = B_4 - c + (B_2 - (B_4 - c)) = B_2$ and $L'_2 = c \leq B_3$ by Lemma 10.6. Hence, case G2 holds for loads L'_1, L'_2 , and we are done again.

3.3 The Construction

Assume that we have an algorithm to solve the semi-online scheduling problem, where the values s and OPT are known. This algorithm now has to schedule all incoming jobs in the best possible way. In the following construction, we take the point of view of an adversary, and try to make the algorithm's life as hard as possible. More formally, we will show that the algorithm will provide a schedule having the optimality ratio of at least $r_1(s)$. Although the whole family of jobs belongs to \mathcal{J}_{OPT} , the adversary still has enough freedom to force any algorithm to an assignment where it ends up with a load L_1 on machine M1 with $L_1 \geq T_2 = r$ or a load L_2 on machine M2 with $L_2 \geq T_1 = rs$. Recall that $\text{OPT} = 1$, so having a load of at least r on machine M_1 means that the makespan is (at least) r , so the optimality ratio is (at least) r . Similarly, having a load of at least rs on the s -times faster machine M_2 means that the makespan is also at least r , and again the optimality ratio is at least r . Note that the family of jobs has a total size of $(s+1)\text{OPT}$, hence by Lemma 1 we know that it belongs to \mathcal{J}_{OPT} .

The adversary decides that the first job shall be $J_1 = B_4 - c$. This job is non-negative by Lemma 10.2. Suppose J_1 goes to M2, then case S1 is satisfied, and

we are done (i.e., we trapped the algorithm as explained above). We conclude J_1 goes to M1.

The second job is $J_2 = c$. Suppose J_2 goes to M2. Then case S3 is satisfied, and we are done. We conclude J_2 goes to M1. At this moment the loads are $L_1 = B_4$ and $L_2 = 0$.

The third job is $J_3 = B_3$. Suppose J_3 goes to M1. Since $L'_1 = B_4 + B_3 = T_4$ holds by Lemma 10.5, we are in case G3, and thus we are done. We conclude J_3 goes to M2. At this moment the loads are $L_1 = B_4$ and $L_2 = B_3$.

Then the next (and final) job is $J_4 = D_3$. Suppose J_4 goes to M2. Then $L'_1 = B_4$ and $L'_2 = T_3$. We estimate that $L'_1 + L'_2 = T_3 + B_4 < T_3 + B_2 = \text{OPT}$, where we applied first Lemma 5.2 and then Lemma 4.5. Thus we showed that we are in case G1, and we are done. We conclude J_4 goes to M1. At this moment $L_1 = B_4 + D_3 = B_2$ by Lemma 4.4, and $L_2 = B_3$. Now we are in case G2, and we are done.

We remark that the sequence of jobs can be drawn as a decision tree, with the first job at its root node, and all other jobs at the subsequent nodes. A left branch means that the job at a node is assigned to machine M1, and a right branch means that it is assigned to machine M2. Note that this tree has a depth of 6 jobs.

4 Lower bounds for regular and medium s

Here we consider the four cases of s being small regular, small medium, bigger medium or bigger regular, respectively. We need several further properties regarding the lower bounds.

4.1 Properties

Lemma 11 *If $s \geq \frac{5}{3}$, then*

1. $\max \left\{ \frac{2}{s}, \frac{5s+6}{4s+4}, \frac{6s+5}{6s+1} \right\} \leq r \leq \frac{7s+5}{6s+2}$,
2. *if s is small medium, then also holds that $\frac{11s+8}{8s+6} \leq r$.*

Proof.

1. Regarding the lower bounds, applying from Lemma 3.1 that $\frac{4}{3} < 1.35 < r$, we get $\frac{4}{3} - \frac{2}{s} = \frac{2(2s-3)}{3s} > 0$, and $\frac{135}{100} - \frac{5s+6}{4s+4} = \frac{2s-3}{20(s+1)} > 0$, both inequalities are true since $s \geq \frac{5}{3}$. Let us see $r \geq \frac{6s+5}{6s+1}$. We already have seen this for regular speeds in Lemma 6.1. For smaller medium s we get $\frac{18s+16}{16s+7} - \frac{6s+5}{6s+1} = \frac{12s^2-8s-19}{(16s+7)(6s+1)} \geq 0$, which is true for $s \geq \frac{3+\sqrt{73}}{8} \approx 1.443$, in particular for $s \geq \frac{5}{3}$. Considering upper medium s , we get $\frac{8s+7}{3s+10} - \frac{6s+5}{6s+1} = \frac{30s^2-25s-43}{(3s+10)(6s+1)} \geq 0$, which is true for $s \geq \frac{25+\sqrt{5785}}{60} \approx 1.684$, in particular for $s \geq q_4$.

Regarding the upper bound, by Lemma 2 it is enough to show that $r_2(s)$ is at most the upper bound. We get $\frac{7s+4}{5s+3} - \frac{12s+10}{9s+7} = \frac{(3s+2)(s-1)}{(9s+7)(5s+3)} \geq 0$.

2. We get $\frac{18s+16}{16s+7} - \frac{11s+8}{8s+6} = \frac{-32s^2+31s+40}{(16s+7)(8s+6)} \geq 0$, which is true, since $-32s^2 + 31s + 40 \geq 0$ for $\frac{31-\sqrt{6081}}{64} \leq s \leq \frac{31+\sqrt{6081}}{64} \approx 1.703$, in particular for $q_3 \leq s \leq q_4$.

■

Lemma 12 1. $s \geq T_2 + c$.

2. $B_2 \leq (s-1) \cdot OPT$.
3. $T_4 + B_2 - 2c > T_3$, i.e., $T_4 - 2c > T_3 - B_2 = b$, i.e., $T_4 > b + 2c$.
4. $T_4 \geq d$.
5. $T_4 + c \leq B_3 + d$, i.e., $c + (T_4 - B_3) \leq d$, i.e., $c + a \leq d$.
6. $c \leq B_3$.
7. $T_4 + B_2 + c - d < OPT = B_2 + T_3$ (c.f. Lemma 4.5), i.e., $T_4 + c < T_3 + d$.
8. $2T_3 + 2B_4 \leq s \cdot OPT$.
9. If s is regular, then $3B_4 = T_4 < B_2$.
10. $T_4 \leq 3B_4 \leq B_2$, if s is small medium.
11. $B_3 + B_4 \leq (s-1)OPT$.

Proof. We apply Lemma 11, if not stated otherwise.

1. From the definitions of T_2 and c we get $s - T_2 - c = s - r - (s - r - rs + 2) = rs - 2 \geq 0$, which is true, because $r \geq \frac{2}{s}$.
2. As before, we obtain $s - 1 - B_2 = s - 1 - (s + 1 - sr) = rs - 2 \geq 0$.
3. We have from the definitions: $T_4 - b - 2c = (r - 1) - (2rs - 2s - 1) - 2(s - r - rs + 2) = 3r - 4 > 0$, where the last inequality was shown in Lemma 3.1.
4. We get $T_4 - d = (r - 1) - (3r - 7s + 6rs - 6) = 7s + 5 - r(6s + 2) \geq 0$, since $r \leq \frac{7s+5}{6s+2}$.
5. We compute $d - c - a = (3r - 7s + 6rs - 6) - (s - r - rs + 2) - (3r - 2s + rs - 3) = r - 6s + 6rs - 5 \geq 0$, which follows from $r \geq \frac{6s+5}{6s+1}$.
6. Applying Lemma 12.4 and 12.5, we get $c \leq B_3 + d - T_4 \leq B_3$.
7. Follows from Lemma 12.5 and Lemma 5.1.

8. From the definitions we obtain $s - 2T_3 - 2B_4 = s - 2s(r - 1) - 2(4s - 2r - 3rs + 3) = 4r - 5s + 4rs - 6 \geq 0$, since $r \geq \frac{5s+6}{4s+4}$.
9. From the definitions we have $3B_4 - T_4 = 3(4s - 2r - 3rs + 3) - (r - 1) = 12s + 10 - r(9s + 7) = 0$, since $r = r_2(s)$. Moreover, $T_4 < B_2$ by Lemma 5.2.
10. For small medium s , it holds that $3B_4 - T_4 = 12s + 10 - r(9s + 7) \geq 0$, since $r = r_3(s) \leq r_2(s)$ by Lemma 2.3. Moreover, $B_2 - 3B_4 = (s + 1 - sr) - 3(4s - 2r - 3rs + 3) = 6r - 11s + 8rs - 8 \geq 0$, as $r \geq \frac{11s+8}{8s+6}$.
11. We estimate $s - 1 - B_3 - B_4 = s - 1 - (2s - 2r - rs + 2) - (4s - 2r - 3rs + 3) = 4r - 5s + 4rs - 6 \geq 0$, since $r \geq \frac{5s+6}{4s+4}$.

■

In the next lemma we consider only the cases that s is smaller regular or smaller medium.

- Lemma 13**
1. $T_4 + c \geq b + 2d$ holds if s is smaller regular and $s \leq \frac{\sqrt{4633} + 23}{54} \approx 1.6864$,
 2. $2B_2 + c \geq T_3 + 2d$ holds if s is smaller regular and $1.6864 \leq s \leq \frac{13 + \sqrt{1429}}{30} \approx 1.6934$,
 3. $2B_2 + c = T_3 + 2d$ holds if s is smaller medium,
 4. $c + B_2 \leq B_3 + d$ holds if s is smaller regular and $s \geq 1.6864$,
 5. $c + B_2 \leq B_3 + d$ holds if s is smaller medium,
 6. $2B_2 + c - d \leq OPT = B_2 + T_3$, i.e. $B_2 + c \leq T_3 + d$, if s is smaller regular and $s \geq 1.6864$, or if s is smaller medium.

Proof.

1. $T_4 + c - (b + 2d) = (r - 1) + (s - r - rs + 2) - (2rs - 2s - 1) - 2(3r - 7s + 6rs - 6) = 17s + 14 - r(6 + 15s) = 17s + 14 - \frac{12s + 10}{9s + 7}(6 + 15s) = \frac{38 + 23s - 27s^2}{9s + 7} \geq 0$, which holds because $38 + 23s - 27s^2 \geq 0$ if and only if $\frac{23 - \sqrt{4633}}{54} \leq s \leq \frac{23 + \sqrt{4633}}{54}$.
2. $2B_2 + c - 2d - T_3 = 2(s + 1 - sr) + (s - r - rs + 2) - 2(3r - 7s + 6rs - 6) - s(r - 1) = 18s + 16 - r(7 + 16s) = 18s + 16 - (7 + 16s)\frac{12s + 10}{9s + 7} = \frac{2(21 + 13s - 15s^2)}{9s + 7} \geq 0$, which holds because $21 + 13s - 15s^2 \geq 0$ if and only if $\frac{13 - \sqrt{1429}}{30} \leq s \leq \frac{13 + \sqrt{1429}}{30}$.
3. $2B_2 + c - 2d - T_3 = 18s + 16 - (7 + 16s)r = 18s + 16 - (7 + 16s)\frac{18s + 16}{16s + 7} = 0$.
4. $B_3 + d - c - B_2 = (2s - 2r - rs + 2) + (3r - 7s + 6rs - 6) - (s - r - rs + 2) - (s + 1 - sr) = (7s + 2)r - 7s - 7 = (7s + 2)\frac{12s + 10}{9s + 7} - 7s - 7 = \frac{21s^2 - 18s - 29}{9s + 7} \geq 0$, since $21s^2 - 18s - 29 \geq 0$ if and only if $s \leq \frac{9 - \sqrt{690}}{21}$ or $s \geq \frac{9 + \sqrt{690}}{21} \approx 1.6794$.

5. $B_3 + d - c - B_2 = (7s+2)r - 7s - 7 = \frac{(7s+2)(18s+16)}{16s+7} - 7s - 7 = \frac{14s^2 - 13s - 17}{16s+7} \geq 0$, since $14s^2 - 13s - 17 \geq 0$ if and only if $s \leq \frac{13 - \sqrt{1121}}{28}$ or $s \geq \frac{13 + \sqrt{1121}}{28} \approx 1.660$.

6. Follows from Lemma 13.4 and Lemma 13.5 using Lemma 4.5.

■

In the next lemma we consider only bigger regular s .

Lemma 14 *Let s be bigger regular. Then*

1. $c + B_2 + T_4 - 2e \geq T_3$, i.e., $c + T_4 - 2e \geq T_3 - B_2 = b$.
2. $c + T_4 \leq B_3 + e$, i.e., $c + (T_4 - B_3) = c + a \leq e$.
3. $c + B_2 + T_4 - e < OPT = B_2 + T_3$, i.e., $c + T_4 < T_3 + e$.
4. $B_4 \geq d$.

Proof.

1. We derive that $c + T_4 - 2e - b = (s - r - rs + 2) + (r - 1) - 2(\frac{1}{2}r - 3s + 3rs - \frac{5}{2}) - (2rs - 2s - 1) = 11s + 6 - r(9s + 1) \geq 0$. The last estimation is true, if $\frac{11s+6}{9s+1} - \frac{12s+10}{9s+7} = \frac{-9s^2+29s+32}{(9s+1)(9s+7)} \geq 0$, which holds since $-9s^2 + 29s + 32 \geq 0$ if and only if $\frac{29 - \sqrt{1993}}{18} \leq s \leq \frac{29 + \sqrt{1993}}{18} \approx 4.0912$, in particular, for bigger regular s .
2. $e - c - a = (\frac{1}{2}r - 3s + 3rs - \frac{5}{2}) - (s - r - rs + 2) - (3r - 2s + rs - 3) = 3rs - 2s - \frac{3}{2}r - \frac{3}{2} \geq 0$, which is true for $r \geq \frac{4s+3}{6s-3}$. Hence we need to verify that $\frac{12s+10}{9s+7} - \frac{4s+3}{6s-3} = \frac{36s^2-31s-51}{3(9s+7)(2s-1)} \geq 0$, which holds for all $s \geq \frac{31 + \sqrt{8305}}{72} = q_5 \approx 1.6963$ (and some negative values for s , which we can ignore).
3. Follows from Lemma 14.2 and Lemma 5.1: $c + T_4 \leq B_3 + e < T_3 + e$.
4. Using the definitions of B_4 and d it is to show that $4s - 2r - 3rs + 3 \geq 3r - 7s + 6rs - 6$, which is equivalent to $-9rs - 5r + 11s + 9 \geq 0$. Taking into account that $r = r_2(s) = \frac{12s+10}{9s+7}$, we arrive at $-\frac{9s^2-8s-13}{9s+7} \geq 0$, which is in particular true for bigger regular values for s .

■

Now we consider the case of the bigger medium speeds.

Lemma 15 *If s is bigger medium, then*

1. $4c + 4a = B_4$,

$$2. b \leq 8c + 7a \leq B_3.$$

Proof.

$$1. \frac{B_4}{4} - c - a = \frac{4s-2r-3rs+3}{4} - (s-r-rs+2) - (3r-2s+rs-3) = \frac{-r(3s+10)-8s-7}{4} = 0.$$

$$2. \text{Left inequality: } 8c+7a-b = 8(s-r-rs+2) + 7(3r-2s+rs-3) - (2rs-2s-1) = r(13-3s) - 4s - 4 = \frac{8s+7}{3s+10}(13-3s) - 4s - 4 = \frac{31s+51-36s^2}{3s+10} \geq 0.$$

The inequality is satisfied since $31s + 51 - 36s^2 \geq 0$ holds if and only if $\frac{31-\sqrt{8305}}{72} \leq s \leq \frac{31+\sqrt{8305}}{72} = q_5$.

$$\text{Right inequality: } B_3 - 8c - 7a = (2s - 2r - rs + 2) - 8(s - r - rs + 2) - 7(3r - 2s + rs - 3) = 8s - 15r + 7 = 15\left(\frac{8s+7}{15} - r\right) = 15\left(\frac{8s+7}{15} - \frac{8s+7}{3s+10}\right) = \frac{(8s+7)(3s-5)}{15(3s+10)} \geq 0, \text{ which is true for } s \geq \frac{5}{3}.$$

■

4.2 Subcases

For s being regular or medium (“RM”), we consider several further situations, from which the lower bound can be quickly achieved directly, or that lead to other general cases we dealt with before.

Final Case RM1. Suppose $T_4 \leq L_1 \leq B_2$ and $0 \leq L_1 + L_2 \leq (s-1) \cdot \text{OPT}$.

Note that by Lemma 12.2, we have $(s-1) \cdot \text{OPT} > B_2$. The next and last two jobs are $B = \text{OPT}$ and $C = \overline{\text{SUM}} - (L_1 + L_2 + B) \geq (s+1)\text{OPT} - s\text{OPT} = \text{OPT} = B$. If any of B and C is assigned to M1, then $L'_1 \geq T_4 + \text{OPT} = T_2$ thus the lower bound holds, otherwise both go to M2 and $L'_2 \geq \overline{\text{SUM}} - B_2 = T_1$, we are done again.

The set of jobs belong to \mathcal{J}_{OPT} by Lemma 1: assign B to M1 and the remaining jobs to M2, which is a load of $L_1 + L_2 + C = L_1 + L_2 + \overline{\text{SUM}} - (L_1 + L_2 + B) = \overline{\text{SUM}} - B = (s+1)\text{OPT} - \text{OPT} = s \cdot \text{OPT}$.

Case RM2. Suppose $L_1 = c$ and $L_2 = 0$.

Let the next job be $A = T_4 - c$. This is nonnegative by Lemma 12.3, using that $b > 0$ from Lemma 7.1. Suppose A is assigned to M1. At this time $L'_1 = T_4$ and $L_2 = 0$. Case G3 holds, we are done. Now suppose A goes to M2, then let the next job be $B = B_2 - c$. This is nonnegative, as $c \leq T_4$ (which we already observed) and $T_4 \leq B_2$. If B goes to M2, then the load of M2 will be $L'_2 = A + B = T_4 + B_2 - 2c$. This is at least T_3 by Lemma 12.3. Moreover, using $T_3 = sT_4 > T_4$ (by the definitions of T_3 and T_4) and $c \geq 0$ (by Lemma 7.1), we have $L'_1 + L'_2 = T_4 + B_2 - c < T_3 + B_2 = \text{OPT}$. Thus we are in case G1, we are done. Otherwise B goes to M1. At this moment the loads are $L'_1 = B_2$ and $L'_2 = A = T_4 - c \leq B_3$, since $T_4 - B_3 = a \leq c$ by the definition of a and Lemma 7.2. Hence we are in case G2, and we are done.

Case RM3. Suppose $L_1 = T_4$ and $L_2 = c$. Applying $T_2 - T_4 = \text{OPT}$ by Lemma 4.2, we get $L_1 + L_2 + \text{OPT} = (T_4 + \text{OPT}) + c = T_2 + c \leq s = s \cdot \text{OPT}$, by Lemma 12.1 and the assumption $\text{OPT} = 1$. Thus $L_1 + L_2 \leq (s-1)\text{OPT}$. So we are in case RM1, and thus we are done.

Case RM4. Suppose $L_1 = d$ and $L_2 = c$ and s is smaller regular or smaller medium.

1. Assume $\frac{5}{3} \leq s \leq \frac{23 + \sqrt{4633}}{54} \approx 1.6864$.

Note that $L_1 = d > 0$. Let the next job be $A = T_4 - d$. This is positive by Lemma 12.4. If A is assigned to M1, we meet the prerequisites of case RM3, and we are done. Thus suppose A goes to M2. Let the next job be $B = B_2 - d$. (This is positive as $B_2 > T_4$.) If B goes to M2, then $L'_2 = c + A + B$. Moreover, $L'_1 + L'_2 = d + c + (T_4 - d) + (B_2 - d) = T_4 + B_2 + c - d < \text{OPT}$, applying Lemma 12.7. We state that $L'_2 = T_4 + B_2 + c - 2d \geq T_3$ holds in the considered interval. Since $b = T_3 - B_2$, it suffices to see $T_4 + c \geq b + 2d$, which holds by Lemma 13.1. Thus we are in case G1, and we are done. Otherwise B goes to M1. At this moment the loads are $L'_1 = B_2$ and $L'_2 = L_2 + A = c + T_4 - d \leq B_3$ by Lemma 12.5. Thus this is case G2, and we are done.

2. Assume $\frac{23 + \sqrt{4633}}{54} \approx 1.6864 \leq s \leq q_3$ or s is smaller medium.

Note that $0 < L_1 = d \leq T_4$ by Lemma 12.4. Let the next job be $A = B_2 - d$, which is positive, since $B_2 > T_4$ by Lemma 5.2. If A is assigned to M1, then the new load on this machine is $L'_1 = d + B_2 - d = B_2$, and $L'_2 = L_2 = c$. By Lemma 12.6 we get $c < B_3$, thus we are in case G2, and we are done. Thus suppose A goes to M2. Then, let the next job be $B = B_2 - d$. If B goes to M2, then the load of M2 will be $L'_2 = c + A + B = c + 2(B_2 - d)$. Then $L'_1 + L'_2 = 2B_2 + c - d$. This is at most OPT by Lemma 13.6. Moreover, $L'_2 = 2B_2 + c - 2d \geq T_3$, by Lemma 13.2 and 13.3. Thus we are at case G1, we are done. Otherwise, B goes to M1. In this moment, the loads are $L'_1 = d + (B_2 - d) = B_2$ and $L'_2 = L_2 + A = c + B_2 - d \leq B_3$, by Lemma 13.4 and 13.5. We are in case G2, and we are done.

Case RM5. Suppose $L_1 = B_4$, and $L_2 = b - B_4$, and s is small medium.

Let the next job be $A = 2B_4$. Suppose A is assigned to M1. At this time $L'_1 = 3B_4$ and $L_2 = b - B_4$. Note that $T_4 \leq 3B_4 \leq B_2$ by Lemma 12.10. Using Lemma 4.5, the definition of b and Lemma 12.8 we get $L'_1 + L'_2 + \text{OPT} = 3B_4 + (b - B_4) + (T_3 + B_2) = 2B_4 + (T_3 - B_2) + (T_3 + B_2) = 2T_3 + 2B_4 < s = s \cdot \text{OPT}$. Hence we are in case RM1, and we are done. We conclude A is assigned to M2. Let the next job be $B = B_2 - B_4$ (this is positive, since $B_2 > B_4$). Suppose B goes to M2, then

$$L'_2 = L_2 + A + B = (b - B_4) + 2B_4 + (B_2 - B_4) = (T_3 - B_2) + B_2 = T_3.$$

by the definition of b . Therefore $L'_1 + L'_2 = B_4 + T_3 < B_2 + T_3 = \text{OPT}$, by Lemma 5.2 and Lemma 4.5. Since L'_2 equals T_3 , we are in case G1, and we are done. Otherwise B goes to M1. At this moment the loads are $L'_1 = B_2$ and

$$\begin{aligned} L'_2 &= L_2 + A = (b - B_4) + 2B_4 = (T_3 - B_2) + B_4 \\ &= (B_3 + D_3) - B_2 + B_4 = B_3 + (B_2 - B_4) - B_2 + B_4 = B_3, \end{aligned}$$

using $T_3 = B_3 + D_3$ and $D_3 = B_2 - B_4$ (by Lemma 4.4). Hence we are in case G2, and we are done.

Case RM6. Suppose $L_1 = e$ and $L_2 = c$, and s is bigger regular.

Let the next job be $A = T_4 - e$. This job is non-negative by Lemma 12.4, $A \geq d - e$, and $d = e + f \geq e$ by Lemma 7.1. Suppose A is assigned to M1. Then $L'_1 = T_4$ and $L'_2 = c$, thus case RM3 holds, and so we are done. Otherwise, A goes to M2. Let the next job be $B = B_2 - e$. (This job is positive, since $B_2 > T_4$ from Lemma 5.2, and the observation that $T_4 \geq e$ from above.) If B goes to M2, then the load of M2 will be $L'_2 = c + (T_4 - e) + (B_2 - e) = c + B_2 + T_4 - 2e \geq T_3$ by Lemma 14.1. Then $L'_1 + L'_2 = c + B_2 + T_4 - e$. This is smaller than OPT by Lemma 14.3. Thus we are at case G1, we are done. Otherwise B goes to M1. At this moment, the loads are $L'_1 = B_2$ and $L'_2 = c + T_4 - e \leq B_3$ by Lemma 14.2. Hence we are in case G2, and we are done again.

Case RM7. Suppose $L_1 = L_2 + a \leq B_4$.

Let the next job be $A = T_4 - L_1$. This is positive as $L_1 \leq B_4$ and $B_4 < T_4$ (by Lemma 5.2). Suppose A is assigned to M1. Since $L'_1 + L'_2 \leq T_4 + B_4 - a = T_4 + B_4 - (T_4 - B_3) = B_3 + B_4 \leq (s - 1)\text{OPT}$ holds by the definition of a and Lemma 12.11, we are in case RM1, and we are done. Now suppose A goes to M2. Then let the next job be $B = B_2 - L_1$, which is positive by Lemma 5.2. Suppose B goes to M2. Then the load of M2 will be $L'_2 = (L_1 - a) + (T_4 - L_1) + (B_2 - L_1) = B_2 + T_4 - a - L_1 \geq B_2 + T_4 - (T_4 - B_3) - B_4 = (B_2 - B_4) + B_3 = D_3 + B_3 = T_3$, by the definition of a and Lemma 4.4. On the other hand, $L'_1 + L'_2 = B_2 + T_4 - a = B_2 + T_4 - (T_4 - B_3) = B_2 + B_3 < B_2 + T_3 = \text{OPT}$, by the definition of a and Lemma 5.2. Thus we are in case G1, and we are done. Otherwise B goes to M1. At this moment the loads are $L'_1 = B_2$ and $L'_2 = (L_1 - a) + (T_4 - L_1) = T_4 - a = T_4 - (T_4 - B_3) = B_3$ by the definition of a . Thus we meet case G2, and we are done again.

Case RM8. Suppose $L_1 = 0$, and $b \leq L_2 \leq B_3$.

Let the next job be $A = B_2$. Suppose job A goes to M2, then the increased load of M2 will be $L'_2 = L_2 + B_2 \geq b + B_2 = (T_3 - B_2) + B_2 = T_3$ (by the definition of b), and $L'_1 + L'_2 \leq B_2 + B_3 < B_2 + T_3 = \text{OPT}$ (by Lemma 5.1 and Lemma 4.5), thus case G1 is satisfied, and we are done. Otherwise, A is assigned to M1. We meet case G2, we are also done.

4.3 The Construction

Similarly to the construction for small s in Section 3.3, we construct a sequence of jobs such that any semi-online algorithm knowing s and OPT will assign in such way that the optimality ratio is at least $r(s)$, where s is between $\frac{5}{3}$ and q_6 . Again, the sequence of jobs belongs to \mathcal{J}_{OPT} , and the total size of the jobs is $\overline{\text{SUM}} = (s + 1)\text{OPT}$.

First, the adversary choses the job $J_1 = c$. Suppose that J_1 goes to $\mathbf{M1}$, then case RM2 is satisfied, and we are done. We conclude that J_1 goes to $\mathbf{M2}$.

We divide the further construction into two main cases, depending on the value of s .

Case 1: s is smaller regular, smaller medium, or bigger regular.

Case 1.1: s is smaller regular or smaller medium.

The second job is $J_2 = d$. Suppose J_2 goes to $\mathbf{M1}$, then case RM4 is satisfied, and we are done. Thus we conclude J_2 goes to $\mathbf{M2}$. At this point $L_1 = 0$ and $L_2 = d + c = (d - e) + (c + e) = f + g = b - B_4$. We continue the construction after case 1.2.

Case 1.2: s is bigger regular.

The second job is $J_{21} = e$. If J_{21} goes to $\mathbf{M1}$, then the assumption of case RM6 is satisfied, and we are done. We conclude that J_{21} goes to $\mathbf{M2}$. The next job is $J_{22} = f$. Suppose J_{22} goes to $\mathbf{M1}$. Then $L_1 = f$ and $L_2 = J_1 + J_{21} = c + e = g$. From Lemma 14.4 and Lemma 7.1 it follows that $B_4 \geq d = e + f \geq f$. Since we also have that $f - g = a$, we are altogether in case RM7 . Thus we are done. We conclude J_{22} goes to $\mathbf{M2}$. At this point $L_1 = 0$ and $L_2 = c + e + f = f + g = b - B_4$.

Now we join the treatments of these subcases, case 1.1 and case 1.2, and finish the construction. In both subcases now $L_1 = 0$ and $L_2 = b - B_4$. Then comes $J_3 = B_4$. Suppose J_3 goes to $\mathbf{M1}$. Then $L_1 = B_4$ and $L_2 = b - B_4$. If s is small medium, then we are in case RM5 , and we are done. Otherwise, if s is small regular or bigger regular, then we claim that we are in case RM7 , for which we have to show $L_2 + a = L_1 \leq B_4$. The inequality on the right follows already from the construction, and it remains to show $L_1 = L_2 + a$. Thus by the definitions of a and b and Lemma 4.4, we obtain $L_1 - L_2 - a = B_4 - (b - B_4) - a = 2B_4 - (T_3 - B_2) - (T_4 - B_3) = 2B_4 + (B_2 - D_3) - T_4 = 2B_4 + B_4 - T_4 = 3B_4 - T_4 = 0$, where the last equality was shown in Lemma 12.9. Thus we can enter case RM7 , and we are done.

We conclude J_3 goes to $\mathbf{M2}$, and at this moment $L_1 = 0, L_2 = b$. We claim that we are in case RM8 then, for which we need to show that $L_2 \leq B_3$. From Lemma 5.2 we know that $B_4 > 0$. Hence we obtain from Lemma 4.4 and the definition of D_3 that $B_2 = B_4 + D_3 > D_3 = T_3 - B_3$. We conclude that $B_3 > T_3 - B_2 = b$, by the definition of b .

Case 2: s is bigger medium.

The second job is $J_{21} = c + a$. If J_{21} goes to M1, then the assumptions of case RM7, $L_1 = L_2 + a \leq B_4$, are satisfied, because $B_4 = 4c + 4a > c + a > 0$ by Lemma 15.1 and Lemma 5.2, and we are done. We conclude J_{21} goes to M2. Then comes $J_{22} = 2c + 2a$. Suppose J_{22} goes to M1. Then $L_1 = 2c + 2a$ and $L_2 = J_1 + J_{21} = 2c + a$. Thus we are again in case RM7 (by repeating the previous arguments), and we are done again. We conclude J_{22} goes to M2. At this point $L_1 = 0$ and $L_2 = 4c + 3a$. Then comes $J_{23} = 4c + 4a$. Suppose J_{23} goes to M1. Then $L_1 = 4c + 4a$ and $L_2 = 4c + 3a$. Thus case RM7 is applicable again (by applying Lemma 15.1), and we are done. We conclude J_{23} goes to M2. At this moment $L_1 = 0, L_2 = 8c + 7a$. By Lemma 15.2 we know that $b \leq 8c + 7a \leq B_3$. Hence we are in case RM8, and we are done.

Again, it is possible to sketch the above assignment steps in a decision tree, as explained at the end of Section 3.3. The depth of this tree depends on the value of s . For smaller regular and smaller medium s , we have a depth of 7 jobs, and for bigger medium and bigger regular s , we have a depth of 8 jobs.

5 Main Theorem

The following theorem summarizes the work done before.

Theorem 16 *The function $r(s)$ (defined in Section 2.1) is a lower bound on the optimal competitive ratio for the two uniform machine semi-online scheduling problem with known optimal offline objective function value.*

Together with the algorithms from Ng et al. [24], we then obtain:

Corollary 17 *The lower bound given by $r(s)$ is tight for $[\frac{1+\sqrt{21}}{4}, \frac{3+\sqrt{73}}{8}] \approx [1.3956, 1.443]$, moreover for $[\frac{5}{3}, \frac{13+\sqrt{1429}}{30}] \approx [1.6666, 1.6934]$, and $[\frac{31+\sqrt{8305}}{72}, \frac{5+\sqrt{241}}{12}] \approx [1.6963, 1.7103]$.*

6 Conclusions and Outlook

Starting with the work of Epstein [14] on this semi-online two uniform machines scheduling problem with known-opt, researchers have continued to close the gap between lower and upper bounds. As one can deduce from Figure 1, this goal has been achieved for some large portions of the line $[1, \infty)$. We contributed to this ultimate goal by giving new lower bounds and thus showing that some already existing algorithms (of Ng et al. [24]) are in fact best-possible, so our bounds are tight. Our new results give insight into the difficulty of the problem: Why so hard to give the tight competitive ratio for this model? In part, an answer lies in the fact that not a single algebraic function can describe the tight lower bound. From what is known by now, at least six different

piecewise-defined algebraic functions are necessary. And still, the question of the optimal competitive ratio is open on certain parts of the "right" interval, namely $(q_2, \frac{5}{3})$, (q_3, q_5) , $(\frac{5+\sqrt{241}}{12}, \sqrt{3})$.

Acknowledgements. György Dósa and Zsolt Tuza were supported by research grant TÁMOP-4.2.2.A-11/1/KONV-2012-0072. Krzysztof Węsek's work was partially supported by the European Union in the framework of European Social Fund through the Warsaw University of Technology Development Programme, realized by Center for Advanced Studies. Furthermore, Węsek's work was conducted as a guest researcher at the Helmut Schmidt University.

References

- [1] Susanne Albers. Better bounds for online scheduling. *SIAM Journal of Computing*, 29:459–473, 1999.
- [2] Susanne Albers and Matthias Hellwig. Semi-online scheduling revisited. *Theoretical Computer Science*, 443:1 – 9, 2012.
- [3] Enrico Angelelli, A.B. Nagy, Maria Grazia Speranza, and Zsolt Tuza. The On-line Multiprocessor Scheduling Problem with Known Sum of the Tasks. *Journal of Scheduling*, 7:421 – 428, 2004.
- [4] Enrico Angelelli, Maria Grazia Speranza, and Zsolt Tuza. Semi on-line scheduling on three processors with known sum of the tasks. *Journal of Scheduling*, 10:263 – 269, 2007.
- [5] Enrico Angelelli, Maria Grazia Speranza, and Zsolt Tuza. Semi-online scheduling on two uniform processors. *Theoretical Computer Science*, 393:211 – 219, 2008.
- [6] Yossi Azar and Oded Regev. On-line bin-stretching. *Theoretical Computer Science*, 268(1):17–41, 2001.
- [7] Piotr Berman, Moses Charikar, and Marek Karpinski. On-line load balancing for related machines. *Journal of Algorithms*, 35:108 – 121, 2000.
- [8] Sheng-Yi Cai and Qi-Fan Yang. Semi-online scheduling on two uniform machines with the known largest size. *Journal of Combinatorial Optimization*, 21:393 – 408, 2011.
- [9] Lin Chen, Deshi Ye, and Guochuan Zhang. Approximating the optimal competitive ratio for an ancient online scheduling problem. Technical report, College of Computer Science, Zhejiang University, Hangzhou, 310027, China, 2013.
- [10] T. C. E. Cheng, H. Kellerer, and V. Kotov. Semi-on-line multi-processor scheduling with given total processing time. *Theoretical Computer Science*, 337:134 – 146, 2005.

- [11] György Dósa, Maria Grazia Speranza, and Zsolt Tuza. Two uniform machines with nearly equal speeds: unified approach to known sum and known optimum in semi on-line scheduling. *Journal of Combinatorial Optimization*, 21:458 – 480, 2011.
- [12] Tomas Ebenlendr and Jiří Sgall. A lower bound on deterministic online algorithms for scheduling unrelated machines without preemption. In *Proceeding of the 9th Workshop on Approximation and Online Algorithms*, Lecture Notes in Computer Science, pages 102 – 108, 2007.
- [13] Tomas Ebenlendr and Jiří Sgall. Semi-Online Preemptive Scheduling: One Algorithm for All Variants. *Theory of Computer Systems*, 48:577 – 613, 2011.
- [14] Leah Epstein. Bin stretching revisited. *Acta Informatica*, 39:97 – 117, 2003.
- [15] Leah Epstein and Deshi Ye. Semi-online scheduling with “end of sequence” information. *Journal of Combinatorial Optimization*, 14:45 – 61, 2007.
- [16] Ulrich Faigle, Walter Kern, and György Turán. On the performance of on-line algorithm for particular problem. *Acta cybernetica*, 9:107–119, 1989.
- [17] T. Gormley, N. Reingold, E. Torng, and J. Westbrook. Generating adversaries for request-answer games. In *Proceeding of the 11th ACM-SIAM Symposium on Discrete Algorithms*. ACM, New York/Society for Industrial and Applied Mathematics, Philadelphia, 2000.
- [18] R. L. Graham. Bounds for certain multiprocessing anomalies. *Bell System Technical Journal*, 45:1563 – 1581, 1966.
- [19] R. L. Graham, E. L. Lawler, J.K. Lenstra, and A.H.G. Rinnooy Kan. Optimization and approximation in deterministic sequencing and scheduling: A survey. *Annals of Discrete Mathematics*, 5, 1979.
- [20] Kellerer Hans, Kotov Vladimir, Maria Grazia, and Zsolt Tuza. Semi on-line algorithms for the partition problem. *Operations Research Letters*, 21:235 – 242, 1997.
- [21] Kangbok Lee and Kyungkuk Lim. Semi-online scheduling problems on a small number of machines. *Journal of Scheduling*, 16:461 – 477, 2013.
- [22] Xinrong Lu and Zhaohui Liu. Semi-online scheduling problems on two uniform machines under a grade of service provision. *Theoretical Computer Science*, 489 - 490:58 – 66, 2013.
- [23] Nicole Megow and Andreas Wiese. Competitive-Ratio Approximation Schemes for Minimizing the Makespan in the Online-List Model. Technical report, Department of Mathematics, Technische Universität Berlin, Germany, 2013.

- [24] C.T. Ng, Zhiyi Tan, Yong He, and T.C.E. Cheng. Two semi-online scheduling problems on two uniform machines. *Theoretical Computer Science*, 410(8 - 10):776 – 792, 2009.
- [25] Zhiyi Tan and An Zhang. Online and Semi-online Scheduling. In P. M. Pardalos et al., editor, *Handbook of Combinatorial Optimization*. Springer Science+Business Media New York, 2013.

