

Regularity of collections of sets and convergence of inexact alternating projections

Alexander Y. Kruger

*Centre for Informatics and Applied Optimization
Faculty of Science and Technology, Federation University Australia
POB 663, Ballarat, Vic, 3350, Australia
a.kruger@federation.edu.au*

Nguyen H. Thao

*Centre for Informatics and Applied Optimization
Faculty of Science and Technology, Federation University Australia
POB 663, Ballarat, Vic, 3350, Australia
hieuthaonguyen@students.federation.edu.au; nhthao@ctu.edu.vn*

Dedicated to the memory of Jean Jacques Moreau

Abstract

We study the usage of regularity properties of collections of sets in convergence analysis of alternating projection methods for solving feasibility problems. Several equivalent characterizations of these properties are provided. Two settings of inexact alternating projections are considered and the corresponding convergence estimates are established and discussed.

Keywords: Alternating projections, uniform regularity, normal cone, subdifferential

2000 Mathematics Subject Classification: 49J53, 49K27, 58E30

1 Introduction

In this article we study the usage of regularity properties of collections of sets in convergence analysis of alternating projection methods for solving *feasibility problems*, i.e., finding a point in the intersection of several sets.

Given a set A and a point x in a metric space, the (metric) *projection* of x on A is defined as follows:

$$P_A(x) := \{a \in A \mid d(x, a) = d(x, A)\},$$

where $d(x, A) := \inf_{a \in A} d(x, a)$ is the distance from x to A . If A is a closed subset of a finite dimensional space, then $P_A(x) \neq \emptyset$. If A is a closed convex subset of a Euclidean space, then $P_A(x)$ is a singleton.

Given a collection $\{A, B\}$ of two subsets of a metric space, we can talk about *alternating projections*.

Definition 1 (Alternating projections). $\{x_n\}$ is a sequence of alternating projections for $\{A, B\}$ if

$$x_{2n+1} \in P_B(x_{2n}) \quad \text{and} \quad x_{2n+2} \in P_A(x_{2n+1}) \quad (n = 0, 1, \dots).$$

Investigations of convergence of the alternating projections to a point in the intersection of closed sets in the setting of a Hilbert space, or more often a finite dimensional Euclidean space, have long history which can be traced back to von Neumann; see the historical comments in [10, 19, 22]. In the convex case, the key convergence estimates were established by Bregman [6] and Bauschke & Borwein [3]. In the nonconvex case, in the finite dimensional setting, linear convergence of the method was shown by Lewis et al. [19, Theorem 5.16] under the assumptions of the *uniform regularity* of the collection $\{A, B\}$ and *super-regularity* of one of the sets; see the definitions and characterizations of these properties in Section 2.

Throughout this paper, we assume that A and B are closed.

Theorem 2 (Linear convergence of alternating projections). *Let X be a finite dimensional Euclidean space. Suppose that*

- (i) $\{A, B\}$ is uniformly regular at $\bar{x} \in A \cap B$, i.e.,

$$\bar{c} := \sup \{ -\langle u, v \rangle \mid u \in \bar{N}_A(\bar{x}) \cap \bar{\mathbb{B}}, v \in \bar{N}_B(\bar{x}) \cap \bar{\mathbb{B}} \} < 1; \quad (1)$$

- (ii) A is super-regular at \bar{x} .

Then, for any $c \in (\bar{c}, 1)$, a sequence of alternating projections for $\{A, B\}$ with initial point sufficiently close to \bar{x} converges to a point in $A \cap B$ with R -linear rate \sqrt{c} .

$\bar{N}_A(\bar{x})$ and $\bar{N}_B(\bar{x})$ in (1) stand for the *limiting normal cones* to the corresponding sets at \bar{x} ; see definition (12) below.

Observe that $-\langle u, v \rangle$ in (1) can be interpreted as the cosine of the angle between the cones $\bar{N}_A(\bar{x})$ and $-\bar{N}_B(\bar{x})$.

The role of the regularity (transversality-like) property (i) of $\{A, B\}$ and convexity-like property (ii) of A in the convergence proof is analysed in Drusvyatskiy et al. [10] and Noll & Rondepierre [22]. It has well been recognized that the uniform regularity assumption is far from being necessary for the linear convergence of alternating projections. For example, as observed in [10], it fails when the affine span of $A \cup B$ is not equal to the whole space.

The drawback of the uniform regularity property as defined by (1) from the point of view of the alternating projections is the fact that it takes into account

all (limiting) normals to each of the sets while in many situations (like the one in the above example) some normals are irrelevant to the idea of projections.

Recently, there have been several successful attempts to relax the discussed above uniform regularity property by restricting the set of involved (normal) directions to only those relevant for characterizing alternating projections. All the newly introduced regularity properties still possess some uniformity in the sense that they take into account directions originated from points in a neighbourhood of the reference point and some estimate is required to hold uniformly over all such directions.

Bauschke et al. [4,5] suggested restricting the set of normals participating in (1) by replacing $\bar{N}_A(\bar{x})$ and $\bar{N}_B(\bar{x})$ with *restricted limiting normal cones* $\bar{N}_A^B(\bar{x})$ and $\bar{N}_B^A(\bar{x})$ depending on both sets and attuned to the method of alternating projections. For example, the cone $\bar{N}_A^B(\bar{x})$ consists of limits of sequences of the type $t_k(b_k - a_k)$ where $t_k > 0$, $b_k \in B$, a_k is a projection of b_k on A , and $a_k \rightarrow \bar{x}$; cf. definitions (17) and (18). Bauschke et al. also adjusted (weakened) the notion of super-regularity accordingly (by considering *joint super-restricted regularity* taking into account the other set) and, under these weaker assumptions, arrived at the same conclusion as in Theorem 2; cf. [4, Theorem 3.14] and Theorem 15 below.

The idea of Bauschke et al. has been further refined by Drusvyatskiy et al. [10, Definition 4.4] who observed that it is sufficient to consider only sequences $t_k(b_k - a_k)$ as above with $b_k \rightarrow \bar{x}$; cf. Definition 16 below. In this case, $a_k \rightarrow \bar{x}$ automatically.

In [10], the authors suggested also another way of weakening the uniform regularity condition (1). Instead of measuring the angles between (usual or restricted in some sense) normals (and negative normals) to the sets, they measure the angles between vectors of the type $a - b$ with $a \in A$ and $b \in B$ and each of the cones $N_B^{\text{prox}}(b)$ and $-N_A^{\text{prox}}(a)$. At least one of the angles must be sufficiently large when a and b are sufficiently close to \bar{x} ; cf. [10, Definition 2.2]. Assuming this property and using a different technique, Drusvyatskiy et al. produced a significant advancement in convergence analysis of projection algorithms by establishing (see [10, Corollary 4.2]) R -linear convergence of alternating projections without the assumption of super-regularity of one of the sets (and with a slightly different convergence estimate). The idea is closely related to the more general approach, where the feasibility problem is reformulated as a problem of minimizing a coupling function, and the property introduced in [10] is sufficient for the coupling function to satisfy the Kurdyka-Łojasiewicz inequality [1, Proposition 4.1].

The two relaxed regularity properties introduced in [10] are in general independent; cf. Examples 21 and 22.

The next step has been made by Noll and Rondepierre [22, Definition 1]. They noticed that, when dealing with alternating projections, the main building block of the method consists of two successive projections:

$$a_1 \in A, \quad b \in P_B(a_1) \quad \text{and} \quad a_2 \in P_A(b) \tag{2}$$

and it is sufficient to consider only the (proximal) normal directions determined by $a_1 - b$ and $b - a_2$ for all a_1, b, a_2 in a neighbourhood of the reference point satisfying (2). In fact, in [22], a more general setting is studied which allows for nonlinear convergence estimates under more subtle nonlinear regularity assumptions.

Another important advancement in this area is considering in [19] of *inexact* alternating projections. Arguing that finding an exact projection of a point on a closed set is in general a difficult problem by itself, Lewis et al. relaxed the requirements to the sequence $\{x_n\}$ in Definition 1 by allowing the points belonging to one of the sets to be “almost” projections. Assuming that the other set is super-regular at the reference point, they established in [19, Theorem 6.1] an inexact version of Theorem 2.

In the next section, we discuss and compare the uniform regularity property of collections of sets and its relaxations mentioned above. Several equivalent characterizations of these properties are provided in a uniform way simplifying the comparison.

The terminology employed in [4, 5, 10, 19, 22] for various regularity properties is not always consistent. We have not found a better way of handling the situation, but to use the terms *BLPW-restricted regularity*, *DIL-restricted regularity*, and *NR-restricted regularity* for the properties introduced in Bauschke, Luke, Phan, and Wang [4, 5], Drusvyatskiy, Ioffe and Lewis [10], and Noll and Rondepierre [22], respectively. The refined version of *BLPW-restricted regularity* due to Drusvyatskiy et al. [10] is referred to in this article as *BLPW-DIL-restricted regularity*.

In Section 3, we study two settings of inexact alternating projections under the assumptions of DIL-restricted regularity and uniform regularity, respectively, and establish and discuss the corresponding convergence estimates.

Our basic notation is standard; cf. [9, 21, 24]. For a normed linear space X , its topological dual is denoted X^* while $\langle \cdot, \cdot \rangle$ denotes the bilinear form defining the pairing between the two spaces. If X is a Hilbert space, X^* is identified with X while $\langle \cdot, \cdot \rangle$ denotes the scalar product. If $\dim X < \infty$, then X is usually assumed equipped with the Euclidean norm. The open and closed unit balls and the unit sphere in a normed space are denoted \mathbb{B} , $\overline{\mathbb{B}}$ and \mathbb{S} , respectively. $\mathbb{B}_\delta(x)$ stands for the open ball with radius $\delta > 0$ and center x . We use the convention $\mathbb{B}_0(x) = \{x\}$.

2 Uniform regularity and related regularity properties

In this section, we discuss and compare the uniform regularity property of collections of sets and its several relaxations which are used in convergence analysis of projection methods.

2.1 Uniform regularity

The uniform regularity property has been studied in [14–18]. Below we consider the case of a collection $\{A, B\}$ of two nonempty closed subsets of a normed linear space.

Definition 3. Suppose X is a normed linear space. The collection $\{A, B\}$ is uniformly regular at $\bar{x} \in A \cap B$ if there exist positive numbers α and δ such that

$$(A - a - x) \cap (B - b - y) \cap (\rho \mathbb{B}) \neq \emptyset$$

for all $\rho \in (0, \delta)$, $a \in A \cap \mathbb{B}_\delta(\bar{x})$, $b \in B \cap \mathbb{B}_\delta(\bar{x})$, and $x, y \in (\alpha\rho)\mathbb{B}$.

The supremum of all α in Definition 3 is denoted $\hat{\theta}[A, B](\bar{x})$ and provides a quantitative characterization of the uniform regularity property, the latter one being equivalent to the inequality $\hat{\theta}[A, B](\bar{x}) > 0$. It is easy to check from the definition that

$$\hat{\theta}[A, B](\bar{x}) = \liminf_{\substack{a \xrightarrow{A} \bar{x}, b \xrightarrow{B} \bar{x}, \rho \downarrow 0}} \frac{\theta_\rho[A - a, B - b](0)}{\rho}, \quad (3)$$

where

$$\theta_\rho[A, B](\bar{x}) := \sup \left\{ r \geq 0 \mid (A - x) \cap (B - y) \cap \mathbb{B}_\rho(\bar{x}) \neq \emptyset, \forall x, y \in r\mathbb{B} \right\}$$

and $a \xrightarrow{A} \bar{x}$ means that $a \rightarrow \bar{x}$ with $a \in A$.

The next proposition contains several characterizations of the uniform regularity property from [14–18]. In its parts (ii) and (iii), $N_A(a)$ stands for the Fréchet normal cone to A at $a \in A$:

$$N_A(a) := \left\{ u \in X^* \mid \limsup_{x \xrightarrow{A} a} \frac{\langle u, x - a \rangle}{\|x - a\|} \leq 0 \right\}. \quad (4)$$

Proposition 4. Let A and B be closed subsets of X .

(i) Suppose X is a normed linear space.

Metric characterization:

$$\hat{\theta}[A, B](\bar{x}) = \liminf_{\substack{z \rightarrow \bar{x}; x, y \rightarrow 0 \\ z \notin (A-x) \cap (B-y)}} \frac{\max \{d(z, A - x), d(z, B - y)\}}{d(z, (A - x) \cap (B - y))}. \quad (5)$$

$\{A, B\}$ is uniformly regular at \bar{x} if and only if there exist positive numbers α and δ such that

$$\alpha d \left(z, (A - x) \cap (B - y) \right) \leq \max \{d(z, A - x), d(z, B - y)\} \quad (6)$$

for all $z \in \mathbb{B}_\delta(\bar{x})$ and $x, y \in \delta\mathbb{B}$.

- (ii) Suppose X is an Asplund space.
Dual characterization.

$$\hat{\theta}[A, B](\bar{x}) = \liminf_{\rho \downarrow 0} \left\{ \|u + v\| \mid u \in N_A(a), v \in N_B(b), \right. \\ \left. a \in A \cap \mathbb{B}_\rho(\bar{x}), b \in B \cap \mathbb{B}_\rho(\bar{x}), \|u\| + \|v\| = 1 \right\}. \quad (7)$$

$\{A, B\}$ is uniformly regular at \bar{x} if and only if there exist positive numbers α and δ such that

$$\alpha(\|u\| + \|v\|) \leq \|u + v\| \quad (8)$$

for all $a \in A \cap \mathbb{B}_\delta(\bar{x})$, $b \in B \cap \mathbb{B}_\delta(\bar{x})$, $u \in N_A(a)$, and $v \in N_B(b)$.

- (iii) Suppose X is a Hilbert space.
Angle characterization. If either $\bar{x} \in \text{bd } A \cap \text{bd } B$ or $\bar{x} \in \text{int}(A \cap B)$, then

$$\hat{\theta}^2[A, B](\bar{x}) = \frac{1}{2} (1 - \hat{c}[A, B](\bar{x})), \quad (9)$$

where

$$\hat{c}[A, B](\bar{x}) := \limsup_{\rho \downarrow 0} \left\{ -\langle u, v \rangle \mid u \in N_A(a) \cap \mathbb{S}, v \in N_B(b) \cap \mathbb{S}, \right. \\ \left. a \in A \cap \mathbb{B}_\rho(\bar{x}), b \in B \cap \mathbb{B}_\rho(\bar{x}) \right\}. \quad (10)$$

Otherwise, $\hat{\theta}[A, B](\bar{x}) = 1$ and $\hat{c}[A, B](\bar{x}) = -\infty$.

$\{A, B\}$ is uniformly regular at \bar{x} if and only if $\hat{c}[A, B](\bar{x}) < 1$, i.e., there exist numbers $\alpha < 1$ and $\delta > 0$ such that

$$-\langle u, v \rangle < \alpha \quad (11)$$

for all $a \in A \cap \mathbb{B}_\delta(\bar{x})$, $b \in B \cap \mathbb{B}_\delta(\bar{x})$, $u \in N_A(a) \cap \mathbb{S}$, and $v \in N_B(b) \cap \mathbb{S}$.

Remark 5. 1. Regularity criteria (6) and (8) are formulated in terms of distances in the primal space and in terms of Fréchet normals, respectively. This explains why we talk about, respectively, the metric and the dual characterizations in parts (i) and (ii) of Proposition 4. The term ‘‘angle characterization’’ in part (iii) comes from the observation that $-\langle u, v \rangle$ in criterion (11) can be interpreted as the cosine of the angle between the unit vectors u and $-v$.

2. Constant (3) is nonnegative while constant (10) can take negative values. It is easy to see from (5) that $\hat{\theta}[A, B](\bar{x}) \leq 1$ if $\bar{x} \notin \text{int}(A \cap B)$ and $\hat{\theta}[A, B](\bar{x}) = \infty$ otherwise. Similarly, $|\hat{c}[A, B](\bar{x})| \leq 1$ if $\bar{x} \in \text{bd } A \cap \text{bd } B$ and $\hat{c}[A, B](\bar{x}) = -\infty$ otherwise.

3. Unlike [17], we assume in (5), (7), and (10) the standard conventions that the infimum and supremum of the empty set in \mathbb{R} equal $+\infty$ and $-\infty$, respectively. As a result, an additional assumption that either $\bar{x} \in \text{bd } A \cap \text{bd } B$ or $\bar{x} \in \text{int}(A \cap B)$ is needed in part (iii) to ensure equality (9).

4. Equality (5) was proved in [13, Theorem 1] while equality (7) was established in [15, Theorem 4(vi)]; see also [13, Theorem 4] for a slightly weaker result containing inequality estimates. Equality (9) is a direct consequence of [17, Theorem 2]. It can be also easily checked directly.

If $\dim X < \infty$, then representations (7) and (10) as well as the corresponding criteria in parts (ii) and (iii) of Proposition 4 can be simplified by using the limiting version of the Fréchet normal cones (4). If, additionally, X is a Euclidean space, then one can also make use of proximal normals.

Recall (cf., e.g., [21]) that, in a Euclidean space, the *limiting (Fréchet) normal cone* to A at \bar{x} and the *proximal normal cone* to A at $a \in A$ are defined as follows:

$$\overline{N}_A(\bar{x}) := \text{Lim sup}_{a \xrightarrow{A} \bar{x}} N_A(a) = \left\{ x^* = \lim x_k^* \mid x_k^* \in N_A(a_k), a_k \xrightarrow{A} \bar{x} \right\}, \quad (12)$$

$$N_A^{\text{prox}}(a) := \text{cone}(P_A^{-1}(a) - a) = \{\lambda(x - a) \mid \lambda \geq 0, a \in P_A(x)\}. \quad (13)$$

Their usage is justified by the following simple observations:

$$N_A^{\text{prox}}(a) \subset N_A(a) \quad \text{and} \quad \overline{N}_A(\bar{x}) = \text{Lim sup}_{a \xrightarrow{A} \bar{x}} N_A^{\text{prox}}(a). \quad (14)$$

Proposition 6. *Let A and B be closed subsets of X .*

(i) *Suppose $\dim X < \infty$.*

Dual characterizations.

$$\begin{aligned} \hat{\theta}[A, B](\bar{x}) &= \inf \{ \|u + v\| \mid u \in \overline{N}_A(\bar{x}), v \in \overline{N}_B(\bar{x}), \|u\| + \|v\| = 1 \} \\ &= \liminf_{\rho \downarrow 0} \left\{ \|u + v\| \mid u \in N_A^{\text{prox}}(a), v \in N_B^{\text{prox}}(b), \right. \\ &\quad \left. a \in A \cap \mathbb{B}_\rho(\bar{x}), b \in B \cap \mathbb{B}_\rho(\bar{x}), \|u\| + \|v\| = 1 \right\}. \end{aligned}$$

$\{A, B\}$ is uniformly regular at \bar{x} if and only if one of the following two equivalent conditions is satisfied:

- (a) $\overline{N}_A(\bar{x}) \cap (-\overline{N}_B(\bar{x})) = \{0\}$;
- (b) there exist positive numbers α and δ such that inequality (8) holds true for all $a \in A \cap \mathbb{B}_\delta(\bar{x})$, $b \in B \cap \mathbb{B}_\delta(\bar{x})$, $u \in N_A^{\text{prox}}(a)$, and $v \in N_B^{\text{prox}}(b)$.

(ii) *Suppose X is a Euclidean space.*

Angle characterizations.

$$\begin{aligned} \hat{c}[A, B](\bar{x}) &= \sup \{ -\langle u, v \rangle \mid u \in \overline{N}_A(\bar{x}) \cap \mathbb{S}, v \in \overline{N}_B(\bar{x}) \cap \mathbb{S} \} \\ &= \limsup_{\rho \downarrow 0} \left\{ -\langle u, v \rangle \mid u \in N_A^{\text{prox}}(a) \cap \mathbb{S}, v \in N_B^{\text{prox}}(b) \cap \mathbb{S}, \right. \\ &\quad \left. a \in A \cap \mathbb{B}_\rho(\bar{x}), b \in B \cap \mathbb{B}_\rho(\bar{x}) \right\}. \end{aligned} \quad (15)$$

$\{A, B\}$ is uniformly regular at \bar{x} if and only if one of the following two equivalent conditions is satisfied:

- (a) $\{(u, v) \in (\overline{N}_A(\bar{x}) \cap \mathbb{S}) \times (\overline{N}_B(\bar{x}) \cap \mathbb{S}) \mid \langle u, v \rangle = -1\} = \emptyset$;
- (b) *there exist numbers $\alpha < 1$ and $\delta > 0$ such that inequality (11) holds true for all $a \in A \cap \mathbb{B}_\delta(\bar{x})$, $b \in B \cap \mathbb{B}_\delta(\bar{x})$, $u \in N_A^{\text{prox}}(a) \cap \mathbb{S}$, and $v \in N_B^{\text{prox}}(b) \cap \mathbb{S}$.*

Remark 7. 1. Condition (a) in part (i) of the above proposition is a ubiquitous qualification condition in optimization and variational analysis; cf. *basic qualification condition* [21] and *transversality condition* [10, 20].

2. If one replaces \mathbb{S} with \mathbb{B} in representation (15), one will get nonnegative constant (1). The relationship between the two constants is straightforward: $\bar{c} = \max\{\hat{c}[A, B](\bar{x}), 0\}$.

2.2 Super-regularity

In the next several subsections, we follow [19] and [4, 5], respectively. Although some definitions and assertions are valid in arbitrary Hilbert spaces, in accordance with the setting of [19] and [4, 5], we assume in these two subsections that X is a finite dimensional Euclidean space.

Unlike the uniform regularity, the super-regularity property is defined for a single set. The next definition contains a list of equivalent characterizations of this property which come from [19, Definition 4.3, Proposition 4.4, and Corollary 4.10], respectively.

Definition 8. A closed subset $A \subset X$ is super-regular at a point $\bar{x} \in A$ if one of the following equivalent conditions is satisfied:

- (i) for any $\gamma > 0$, there exists a $\delta > 0$ such that

$$\langle x - x_A, a - x_A \rangle \leq \gamma \|x - x_A\| \|a - x_A\|$$

for all $x \in \mathbb{B}_\delta(\bar{x})$, $x_A \in P_A(x)$, and $a \in A \cap \mathbb{B}_\delta(\bar{x})$;

- (ii) for any $\gamma > 0$, there exists a $\delta > 0$ such that

$$\langle u, x - a \rangle \leq \gamma \|u\| \|x - a\| \tag{16}$$

for all $x, a \in A \cap \mathbb{B}_\delta(\bar{x})$ and $u \in N_A(a)$;

- (iii) for any $\gamma > 0$, there exists a $\delta > 0$ such that

$$\langle v - u, y - x \rangle \geq -\gamma \|y - x\|$$

for all $x, y \in A \cap \mathbb{B}_\delta(\bar{x})$ and $u \in N_A(x)$ and $v \in N_A(y)$.

Remark 9. 1. Super-regularity is a kind of local “near convexity” property, refining or complementing a number of properties of this kind: *Clarke regularity* [7, 24], *amenability* [24], *prox-regularity* [23, 24], and *subsmoothness* [2] (cf. *first order Shapiro property* [25]). For a detailed discussion and comparing of the properties we refer the reader to [19].

2. Super-regularity of one of the sets is an important ingredient of the convergence analysis of projection methods following the scheme initiated in Lewis et al. [19]; cf. Theorems 2 and 34. In fact, a weaker “quantified” version of this property corresponding to fixing $\gamma > 0$ in Definition 8 (and Definition 10 below), i.e., a kind of γ -*super-regularity* is sufficient for this type of analysis; cf. [5, Definition 8.1] and [22, Definition 2] (The latter definition introduces a more advanced Hölder version of this property.) Of course for alternating projections to converge, γ must be small and the convergence rate depends on γ .

2.3 Restricted normal cones and restricted super-regularity

There have been several successful attempts to relax the discussed above regularity properties by restricting the set of involved (normal) directions to only those relevant for characterizing alternating projections.

The definitions of *restricted normal cones* to a set introduced in [5] take into account another set and generalize proximal and limiting normal cones (13) and (12) in the setting of a Euclidean space:

$$N_A^{B-\text{prox}}(a) := \text{cone}((P_A^{-1}(a) \cap B) - a), \quad (17)$$

$$\overline{N}_A^B(\bar{x}) := \text{Lim sup}_{a \xrightarrow{A} \bar{x}} N_A^{B-\text{prox}}(a). \quad (18)$$

Sets (17) and (18) are called, respectively, the *B-proximal normal cone* to A at $a \in A$ and *B-limiting normal cone* to A at \bar{x} . When B is the whole space, they obviously coincide with (13) and (12) (cf. the representation of the limiting normal cone given by the equality in (14)). Note that cones (17) and (18) can be empty.

Similarly to (17), one can define also the *B-Fréchet normal cone* to A at $a \in A$:

$$N_A^B(a) := N_A(a) \cap \text{cone}(B - a)$$

and the corresponding limiting one. The following inclusions are straightforward:

$$N_A^{B-\text{prox}}(a) \subset N_A^B(a) \subset N_A(a).$$

Definition 10. A closed subset $A \subset X$ is *B-super-regular* at a point $\bar{x} \in A$ if, for any $\gamma > 0$, there exists a $\delta > 0$ such that condition (16) holds true for all $x, a \in A \cap \mathbb{B}_\delta(\bar{x})$ and $u \in N_A^{B-\text{prox}}(a)$.

Remark 11. As observed in [5], *B*-proximal normals in Definition 10 can be replaced with *B*-limiting ones. Similarly, in Definition 8(ii) and (iii), one can replace Fréchet normals with limiting ones.

2.4 BLPW-restricted regularity

The next definition introduces a modification of the property used in the angle characterization of the uniform regularity in Proposition 4(iii). This new property and its subsequent characterizations and application in convergence estimate (Theorem 15) originate in Bauschke, Luke, Phan, and Wang [4, 5]. We are going to use for the regularity property of a collection of two sets discussed below the term *BLPW-restricted regularity*.

Definition 12. A collection of closed sets $\{A, B\}$ is BLPW-restrictedly regular at $\bar{x} \in A \cap B$ if

$$\hat{c}_1[A, B](\bar{x}) := \limsup_{\rho \downarrow 0} \left\{ -\langle u, v \rangle \mid u \in N_A^{B-\text{prox}}(a) \cap \mathbb{S}, v \in N_B^{A-\text{prox}}(b) \cap \mathbb{S}, \right. \\ \left. a \in A \cap \mathbb{B}_\rho(\bar{x}), b \in B \cap \mathbb{B}_\rho(\bar{x}) \right\} < 1, \quad (19)$$

i.e., there exist numbers $\alpha < 1$ and $\delta > 0$ such that condition (11) holds for all $a \in A \cap \mathbb{B}_\delta(\bar{x})$, $b \in B \cap \mathbb{B}_\delta(\bar{x})$, $u \in N_A^{B-\text{prox}}(a) \cap \mathbb{S}$, and $v \in N_B^{A-\text{prox}}(b) \cap \mathbb{S}$.

Proposition 13. (i) *The following representation holds true:*

$$\hat{c}_1[A, B](\bar{x}) = \sup \left\{ -\langle u, v \rangle \mid u \in \overline{N}_A^B(\bar{x}) \cap \mathbb{S}, v \in \overline{N}_B^A(\bar{x}) \cap \mathbb{S} \right\}. \quad (20)$$

(ii) *If either $\overline{N}_A^B(\bar{x}) \cap \mathbb{S} \neq \emptyset$ and $\overline{N}_B^A(\bar{x}) \cap \mathbb{S} \neq \emptyset$, or $\overline{N}_A^B(\bar{x}) \cap \mathbb{S} = \overline{N}_B^A(\bar{x}) \cap \mathbb{S} = \emptyset$, then*

$$\hat{c}_1[A, B](\bar{x}) = 1 - 2\hat{\theta}_1^2[A, B](\bar{x}),$$

where

$$\hat{\theta}_1[A, B](\bar{x}) = \liminf_{\rho \downarrow 0} \left\{ \|u + v\| \mid u \in N_A^{B-\text{prox}}(a), v \in N_B^{A-\text{prox}}(b), \right. \\ \left. a \in A \cap \mathbb{B}_\rho(\bar{x}), b \in B \cap \mathbb{B}_\rho(\bar{x}), \|u\| + \|v\| = 1 \right\} \\ = \inf \left\{ \|u + v\| \mid u \in \overline{N}_A^B(a), v \in \overline{N}_B^A(b), \|u\| + \|v\| = 1 \right\}.$$

(iii) *A collection of closed sets $\{A, B\}$ is BLPW-restrictedly regular at $\bar{x} \in A \cap B$ if and only if one of the following conditions holds true:*

$$\bar{c}_1 := \sup \left\{ -\langle u, v \rangle \mid u \in \overline{N}_A^B(\bar{x}) \cap \overline{\mathbb{B}}, v \in \overline{N}_B^A(\bar{x}) \cap \overline{\mathbb{B}} \right\} < 1, \quad (21)$$

$$\hat{\theta}_1[A, B](\bar{x}) > 0,$$

$$\overline{N}_A^B(\bar{x}) \cap \left(-\overline{N}_B^A(\bar{x}) \right) \subseteq \{0\}. \quad (22)$$

Remark 14. 1. The difference between formula (20) and definition of \bar{c}_1 in (21) is that, in the latter one, closed unit balls are used instead of spheres. As a result, \bar{c}_1 is either nonnegative or equal $-\infty$. (The latter case is possible because restricted normal cones can be empty.) At the same time, conditions

$\hat{c}_1[A, B](\bar{x}) < 1$ and $\bar{c}_1 < 1$ are equivalent and \bar{c}_1 can be used for characterizing BLPW-restricted regularity. The inequality $\bar{c}_1 \leq \bar{c}$, where \bar{c} is given by (1), is obvious. It can be strict; cf. [5, Example 7.1].

2. In [5], a more general setting of four sets $A, B, \tilde{A}, \tilde{B}$ is considered with the A - and B -proximal and limiting normals cones in Definition 12 and Proposition 13 replaced by their \tilde{A} and \tilde{B} versions. As described in [4, Subsection 3.6], this provides additional flexibility in applications when determining regularity properties. To simplify the presentation, in this article we set $\tilde{A} = A$ and $\tilde{B} = B$.

3. Condition (22) is referred to in [5] as (A, B) -qualification condition while constant (19) is called the limiting CQ number.

Theorem 15. *Let X be a finite dimensional Euclidean space. Suppose that*

- (i) $\{A, B\}$ is BLPW-restrictedly regular at $\bar{x} \in A \cap B$;
- (ii) A is B -super-regular at \bar{x} .

Then, for any $c \in (\bar{c}_1, 1)$, a sequence of alternating projections for $\{A, B\}$ with initial point sufficiently close to \bar{x} converges to a point in $A \cap B$ with R -linear rate \sqrt{c} .

2.5 BLPW-DIL-restricted regularity

The concept of BLPW-restricted regularity was further refined in Drusvyatskiy, Ioffe and Lewis [10, Definition 4.4]. We are going to call the amended property *BLPW-DIL-restricted regularity*.

Definition 16. A collection of closed sets $\{A, B\}$ is BLPW-DIL-restrictedly regular at $\bar{x} \in A \cap B$ if

$$\hat{c}_2[A, B](\bar{x}) := \limsup_{\rho \downarrow 0} \left\{ - \frac{\langle a - b_a, b - a_b \rangle}{\|a - b_a\| \|b - a_b\|} \mid b_a \in P_B(a), a_b \in P_A(b), \right. \\ \left. a \in (A \setminus B) \cap \mathbb{B}_\rho(\bar{x}), b \in (B \setminus A) \cap \mathbb{B}_\rho(\bar{x}) \right\} < 1, \quad (23)$$

i.e., there exist numbers $\alpha < 1$ and $\delta > 0$ such that

$$-\langle a - b_a, b - a_b \rangle < \alpha \|a - b_a\| \|b - a_b\|$$

for all $a \in (A \setminus B) \cap \mathbb{B}_\delta(\bar{x})$, $b \in (B \setminus A) \cap \mathbb{B}_\delta(\bar{x})$, $b_a \in P_B(a)$, and $a_b \in P_A(b)$.

Remark 17. The property in Definition 16 is referred to in [10] as *inherent transversality*.

An analogue of Proposition 13 holds true with constant $\hat{\theta}_1[A, B](\bar{x})$ replaced by

$$\hat{\theta}_2[A, B](\bar{x}) = \frac{1}{2} \liminf_{\rho \downarrow 0} \left\{ \left\| \frac{a - b_a}{\|a - b_a\|} + \frac{b - a_b}{\|b - a_b\|} \right\| \mid b_a \in P_B(a), a_b \in P_A(b), \right. \\ \left. a \in (A \setminus B) \cap \mathbb{B}_\rho(\bar{x}), b \in (B \setminus A) \cap \mathbb{B}_\rho(\bar{x}) \right\}$$

and appropriate limiting objects.

It is easy to see that a BLPW-restrictedly regular collection is also BLPW-DIL-restrictedly regular, but the converse is not true in general.

2.6 DIL-restricted regularity

The next definition and its subsequent characterizations originate in Drusvyatskiy, Ioffe and Lewis [10]. We are going to use for the regularity property of a collection of two sets discussed below the term *DIL-restricted regularity*.

Unlike [10], if not specified otherwise, we adopt in this subsection the setting of a general Hilbert space.

Definition 18. A collection of closed sets $\{A, B\}$ is DIL-restrictedly regular at $\bar{x} \in A \cap B$ if

$$\hat{\theta}_4[A, B](\bar{x}) := \lim_{\rho \downarrow 0} \inf_{a \in (A \setminus B) \cap \mathbb{B}_\rho(\bar{x}), b \in (B \setminus A) \cap \mathbb{B}_\rho(\bar{x})} \max \left\{ d \left(\frac{b-a}{\|a-b\|}, N_A(a) \right), d \left(\frac{a-b}{\|a-b\|}, N_B(b) \right) \right\} > 0, \quad (24)$$

i.e., there exist positive numbers γ and $\delta > 0$ such that

$$\max \left\{ d \left(\frac{b-a}{\|a-b\|}, N_A(a) \right), d \left(\frac{a-b}{\|a-b\|}, N_B(b) \right) \right\} > \gamma \quad (25)$$

for all $a \in (A \setminus B) \cap \mathbb{B}_\delta(\bar{x})$ and $b \in (B \setminus A) \cap \mathbb{B}_\delta(\bar{x})$.

Proposition 19. A collection of closed sets $\{A, B\}$ is DIL-restrictedly regular at $\bar{x} \in A \cap B$ if and only if

$$\hat{c}_4[A, B](\bar{x}) := \limsup_{\rho \downarrow 0} \left\{ \frac{\min\{\langle b-a, u \rangle, \langle a-b, v \rangle\}}{\|a-b\|} \mid u \in N_A(a) \cap \mathbb{S}, v \in N_B(b) \cap \mathbb{S}, a \in (A \setminus B) \cap \mathbb{B}_\rho(\bar{x}), b \in (B \setminus A) \cap \mathbb{B}_\rho(\bar{x}) \right\} < 1, \quad (26)$$

i.e., there exist numbers $\alpha < 1$ and $\delta > 0$ such that

$$\min\{\langle b-a, u \rangle, \langle a-b, v \rangle\} < \alpha \|a-b\|$$

for all $a \in (A \setminus B) \cap \mathbb{B}_\delta(\bar{x})$, $b \in (B \setminus A) \cap \mathbb{B}_\delta(\bar{x})$, $u \in N_A(a) \cap \mathbb{S}$, and $v \in N_B(b) \cap \mathbb{S}$.

Moreover, $(\hat{c}_4[A, B](\bar{x}))^2 + (\hat{\theta}_4[A, B](\bar{x}))^2 = 1$.

Remark 20. 1. If $\dim X < \infty$, then, as usual, the Fréchet normals in (24) and (26) can be replaced by the proximal ones:

$$\hat{\theta}_4[A, B](\bar{x}) = \lim_{\rho \downarrow 0} \inf_{a \in (A \setminus B) \cap \mathbb{B}_\rho(\bar{x}), b \in (B \setminus A) \cap \mathbb{B}_\rho(\bar{x})} \max \left\{ d \left(\frac{b-a}{\|a-b\|}, N_A^{\text{prox}}(a) \right), d \left(\frac{a-b}{\|a-b\|}, N_B^{\text{prox}}(b) \right) \right\},$$

$$\hat{c}_4[A, B](\bar{x}) = \limsup_{\rho \downarrow 0} \left\{ \frac{\min\{\langle b-a, u \rangle, \langle a-b, v \rangle\}}{\|a-b\|} \mid u \in N_A^{\text{prox}}(a) \cap \mathbb{S}, \right. \\ \left. v \in N_B^{\text{prox}}(b) \cap \mathbb{S}, a \in (A \setminus B) \cap \mathbb{B}_\rho(\bar{x}), b \in (B \setminus A) \cap \mathbb{B}_\rho(\bar{x}) \right\}.$$

2. In [10], the property in Definition 18 is referred to as *intrinsic transversality*.

The next two examples show that DIL-restricted regularity is in general independent of BLPW-DIL-restricted regularity.

Example 21 (BLPW-DIL-restricted regularity but not DIL-restricted regularity; Figure 1). Define a function $f : [0, 1] \rightarrow \mathbb{R}$ by

$$f(t) := \begin{cases} 0, & \text{if } t = 0, \\ -t + 1/2^{n+1}, & \text{if } t \in (1/2^{n+1}, 3/2^{n+2}], \\ t - 1/2^n, & \text{if } t \in (3/2^{n+2}, 1/2^n], n = 0, 1, \dots \end{cases}$$

and consider the sets: $A = \text{gph } f$ and $B = \{(t, t) : t \in [0, 1]\}$ and the point $\bar{x} = (0, 0) = A \cap B$ in \mathbb{R}^2 . Suppose \mathbb{R}^2 is equipped with the Euclidean norm.




Figure 1: BLPW-DIL-restricted regularity but not DIL-restricted regularity

It is easy to check that f is a continuous function and consequently A is closed; $f(1/2^n) = 0$, $f(3/2^{n+2}) = -1/2^{n+2}$, $n = 0, 1, \dots$

Take any $a \in A \setminus B$, $b \in B \setminus A$, $b_a \in P_B(a)$, and $a_b \in P_A(b)$. Thanks to the properties of the Euclidean distance, we have

$$a_b = (1/2^n, 0), \\ b \in B \cap P_A^{-1}(a_b) = \{(t, t) \mid t \in [3/2^{n+2}, 3/2^{n+1}]\}, \\ a - b_a = k(1, -1)$$

for some $n \in \mathbb{N}$ and $k > 0$. Then,

$$\hat{c}_2[A, B](\bar{x}) = \max_{b \in B \cap P_A^{-1}(a_b)} \left\{ \frac{\langle (-1, 1), b - a_b \rangle}{\sqrt{2}\|b - a_b\|} \right\} = \frac{1}{\sqrt{2}} \left\langle (-1, 1), \frac{\bar{b} - a_b}{\|\bar{b} - a_b\|} \right\rangle,$$

where $\bar{b} := (3/2^{n+2}, 3/2^{n+2})$, and consequently,

$$\hat{c}_2[A, B](\bar{x}) = \frac{\langle (-1, 1), (-1, 3) \rangle}{\sqrt{2} \sqrt{10}} = \frac{2}{\sqrt{5}} > 0.$$

Hence, $\{A, B\}$ is BLPW-DIL-restrictedly regular at \bar{x} .

Given an $n \in \mathbb{N}$, we choose $a := (1/2^n, 0) \in A \setminus B$ and $b := (1/2^{n+1}, 1/2^{n+1}) \in B \setminus A$. Then,

$$\begin{aligned} N_A^{\text{prox}}(a) &= N_A(a) = \{(t_1, t_2) : t_2 \geq |t_1|\}, \\ N_B^{\text{prox}}(b) &= N_B(b) = \mathbb{R}(1, -1), \end{aligned}$$

and consequently,

$$a - b = 1/2^{n+1}(1, -1) \in N_B(b) \cap -N_A(a).$$

It follows that $\hat{c}_4[A, B](\bar{x}) = 1$ and $\{A, B\}$ is not DIL-restrictedly regular at \bar{x} . \triangle

Example 22 (DIL-restricted regularity but not BLPW-DIL-restricted regularity; Figure 2). Consider two sets:

$$\begin{aligned} A &= \{(t, 0) : t \geq 0\} \cup \{(t, -t) : t \geq 0\}, \\ B &= \{(t, 0) : t \geq 0\} \cup \{(t, t) : t \geq 0\} \end{aligned}$$

and the point $\bar{x} = (0, 0) \in A \cap B$ in \mathbb{R}^2 . Suppose \mathbb{R}^2 is equipped with the Euclidean norm.




Figure 2: DIL-restricted regularity but not BLPW-DIL-restricted regularity

For any $a = (t_1, -t_1) \in A \setminus B$ and $b = (t_2, t_2) \in B \setminus A$, we have

$$\begin{aligned} N_A^{\text{prox}}(a) &= N_A(a) = \mathbb{R}(1, 1), \\ N_B^{\text{prox}}(b) &= N_B(b) = \mathbb{R}(1, -1). \end{aligned}$$

and consequently,

$$\hat{c}_4[A, B](\bar{x}) = \sup_{t_1 > 0, t_2 > 0} \frac{\min\{t_1, t_2\}}{\sqrt{t_1^2 + t_2^2}} = \frac{1}{\sqrt{2}} < 1.$$

Hence, $\{A, B\}$ is DIL-restrictedly regular at \bar{x} .

For any $a \in A \setminus B$, $b \in B \setminus A$, $b_a \in P_B(a)$, and $a_b \in P_A(b)$, we have

$$\frac{b - a_b}{\|b - a_b\|} = \frac{b_a - a}{\|a - b_a\|}.$$

It follows that $\hat{c}_2[A, B](\bar{x}) = 1$ and $\{A, B\}$ is not BLPW-DIL-restrictedly regular at \bar{x} . \triangle

The next fact was established in [10, Proposition 4.5].

Proposition 23. *If $\dim X < \infty$, $\{A, B\}$ is BLPW-DIL-restrictedly regular at \bar{x} , and both sets A and B are super-regular at \bar{x} , then $\{A, B\}$ is DIL-restrictedly regular at \bar{x} .*

Remark 24. The assumption of super-regularity of both sets in Proposition 23 is essential. Indeed, in Example 21, $\{A, B\}$ is BLPW-DIL-restrictedly regular and B is super-regular (in fact, convex), while A is not and $\{A, B\}$ is not DIL-restrictedly regular.

2.7 NR-restricted regularity

The next step in relaxing both BLPW- and DIL-restricted regularity properties while preserving the linear convergence of alternating projections has been done in Noll and Rondepierre [22]. In what follows, the resulting property is called *NR-restricted regularity*.

Definition 25. A collection of closed sets $\{A, B\}$ is NR-restrictedly regular at $\bar{x} \in A \cap B$ if

$$\hat{c}_3[A, B](\bar{x}) := \limsup_{\rho \downarrow 0} \left\{ \frac{\langle a_1 - b, a_2 - b \rangle}{\|a_1 - b\| \|a_2 - b\|} \mid a_1 \in A, b \in P_B(a_1), a_2 \in P_A(b), a_1, b, a_2 \in \mathbb{B}_\rho(\bar{x}) \right\} < 1,$$

i.e., there exist numbers $\alpha < 1$ and $\delta > 0$ such that

$$\langle a_1 - b, a_2 - b \rangle \leq \alpha \|a_1 - b\| \|a_2 - b\|$$

for all $a_1 \in A \cap \mathbb{B}_\delta(\bar{x})$, $b \in P_B(a_1) \cap \mathbb{B}_\delta(\bar{x})$, and $a_2 \in P_A(b) \cap \mathbb{B}_\delta(\bar{x})$.

Remark 26. 1. NR-restricted regularity property is not symmetric: NR-restricted regularity of $\{A, B\}$ does not imply that $\{B, A\}$ is NR-restrictedly regular.

2. If $\{A, B\}$ is BLPW- or DIL-restrictedly regular at \bar{x} , then it is NR-restrictedly regular at \bar{x} and the second implication can be strict [22, Propositions 1 and 2 and Example 7.6]. In fact, it is easy to check that NR-restricted regularity is implied by BLPW-DIL-restricted regularity. Example 22 shows that NR-restricted regularity can be strictly weaker.

3. Theorem 15 remains valid if the assumption of BLPW-restricted regularity is replaced by that of NR-restricted regularity and \hat{c}_1 is replaced by $\hat{c}_3[A, B](\bar{x})$.

4. The property in Definition 25 is referred to in [22] as *separable intersection*.
5. In [22], a more general Hölder-type property with exponent $\omega \in [0, 2)$ is considered. Definition 25 corresponds to that property with $\omega = 0$. For the convergence analysis, the authors of [22] introduce also a Hölder version of the superregularity property.

3 Convergence for inexact alternating projections

In this section, we study two settings of inexact alternating projections under the assumptions of DIL-restricted regularity and uniform regularity, respectively, and establish the corresponding convergence estimates.

3.1 Convergence for inexact alternating projections under DIL-restricted regularity

Given a point x and a set A in a Hilbert space and numbers $\tau \in (0, 1]$ and $\sigma \in [0, 1)$, the (τ, σ) -*projection* of x on A is defined as follows:

$$P_A^{\tau, \sigma}(x) := \{a \in A \mid \tau \|x - a\| \leq d(x, A), d(x - a, N_A(a)) \leq \sigma \|x - a\|\}. \quad (27)$$

One obviously has $P_A^{1, \sigma}(x) = P_A(x)$ for any $\sigma \in [0, 1)$. Observe also that the above definition requires a to be an “almost projection” in terms of the distance $\|x - a\|$ being close to $d(x, A)$ and also $x - a$ being an “almost normal” to A .

Definition 27 (Inexact alternating projections). Given $\tau \in (0, 1]$ and $\sigma \in [0, 1)$, $\{x_n\}$ is a sequence of (τ, σ) -alternating projections for $\{A, B\}$ if

$$x_{2n+1} \in P_B^{\tau, \sigma}(x_{2n}) \quad \text{and} \quad x_{2n+2} \in P_A^{\tau, \sigma}(x_{2n+1}) \quad (n = 0, 1, \dots).$$

The next statement is taken from [10, Theorem 5.3] where it is formulated in the setting of a finite dimensional Euclidean space. It is a version of the general metric space *Basic Lemma* from [12]. Recall that the (strong) *slope* [8] of f at a point $u \in X$ with $f(u) < +\infty$ is defined as follows:

$$|\nabla f|(u) := \limsup_{u' \rightarrow u, u' \neq u} \frac{f(u) - f(u')}{d(u', u)}.$$

Lemma 28 (Error bound). *Let X be a complete metric space, $f : X \rightarrow \mathbb{R} \cup \{+\infty\}$ a lower semicontinuous function, $x \in X$ with $f(x) < +\infty$, $\delta > 0$, and $\alpha < f(x)$. Suppose that*

$$\mu := \inf_{u \in \mathbb{B}_\delta(x), \alpha < f(u) \leq f(x)} |\nabla f|(u) > \frac{f(x) - \alpha}{\delta}. \quad (28)$$

Then $S(f, \alpha) := \{u \in X \mid f(u) \leq \alpha\} \neq \emptyset$ and

$$\mu d(x, S(f, \alpha)) \leq f(x) - \alpha.$$

If X is an Asplund space, then a standard argument based on the *subdifferential sum rule* (cf., e.g., [11, Proposition 5(ii)] or [10, Proposition 6.9]) shows that the primal space slopes in the definition of μ in (28) can be replaced by the subdifferential slopes:

$$\mu = \inf_{u \in \mathbb{B}_\delta(x), \alpha < f(u) \leq f(x)} |\partial f|(u). \quad (29)$$

Here

$$|\partial f|(u) := \inf_{x^* \in \partial f(u)} \|x^*\|_*,$$

where $\partial f(u)$ is the Fréchet subdifferential of f at u and $\|\cdot\|_*$ is the norm on X^* dual to the norm on X participating in the definition of the primal space slope. Note that in general $|\nabla f|(u) \leq |\partial f|(u)$.

If X is a finite dimensional Euclidean space, then, instead of the Fréchet subdifferentials, one can use the *proximal* subdifferentials $\partial^{\text{prox}} f(u)$:

$$\mu = \inf_{u \in \mathbb{B}_\delta(x), \alpha < f(u) \leq f(x)} |\partial^{\text{prox}} f|(u), \quad (30)$$

where

$$|\partial^{\text{prox}} f|(u) := \inf_{x^* \in \partial^{\text{prox}} f(u)} \|x^*\|.$$

The next statement is a consequence of Lemma 28. It extends slightly [10, Theorem 3.1].

Proposition 29 (Distance decrease). *Let A be a closed subset of a Hilbert space X , $a \in A$, $b \notin A$, $\delta > 0$, and $\alpha < \|a - b\|$. Suppose that*

$$\mu := \inf_{\substack{u \in A \cap \mathbb{B}_\delta(a) \\ \|u - b\| \leq \|a - b\|}} d\left(\frac{b - u}{\|u - b\|}, N_A(u)\right) > 0. \quad (31)$$

Then $d(b, A) \leq \|a - b\| - \mu\delta$.

If $\dim X < \infty$, then the Fréchet normal cones $N_A(u)$ in (31) can be replaced by the proximal ones $N_A^{\text{prox}}(u)$.

Proof. Consider the lower semicontinuous function $f = d(\cdot, b) + \iota_A$, where ι_A is the indicator function of A : $\iota_A(x) = 0$ if $x \in A$ and $\iota_A(x) = +\infty$ if $x \notin A$. Then $f(a) = \|a - b\|$ and

$$\partial f(u) = \frac{u - b}{\|u - b\|} + N_A(u), \quad |\partial f|(u) = d\left(\frac{b - u}{\|u - b\|}, N_A(u)\right)$$

for any $u \in A$. It follows from the first part of Lemma 28 and representation (29) that $d(b, A) \leq \alpha$ for any $\alpha \in (\|a - b\| - \mu\delta, \|a - b\|)$ and consequently, $d(b, A) \leq \|a - b\| - \mu\delta$.

If $\dim X < \infty$, then instead of representation (29) one can use representation (30). \square

The next statement is essentially [10, Lemma 3.2].

Lemma 30. *Any nonzero vectors x and y in a Hilbert space satisfy*

$$\left\| \frac{x}{\|x\|} - z \right\| \leq \frac{\|x - y\|}{\|y\|},$$

where $z := \left\langle \frac{x}{\|x\|}, \frac{y}{\|y\|} \right\rangle \frac{y}{\|y\|}$ is the projection of $\frac{x}{\|x\|}$ on $\mathbb{R}y$.

Proof.

$$\begin{aligned} \left(\frac{\|x - y\|}{\|y\|} \right)^2 - \left\| \frac{x}{\|x\|} - z \right\|^2 &= \frac{\|x\|^2 - 2\langle x, y \rangle + \|y\|^2}{\|y\|^2} - 1 + \left\langle \frac{x}{\|x\|}, \frac{y}{\|y\|} \right\rangle^2 \\ &= \frac{1}{\|y\|^2} \left(\|x\|^2 - 2\langle x, y \rangle + \frac{\langle x, y \rangle^2}{\|x\|^2} \right) \\ &= \frac{1}{\|y\|^2} \left(\|x\| - \frac{\langle x, y \rangle}{\|x\|} \right)^2 \geq 0. \end{aligned}$$

□

Theorem 31 (Convergence of inexact alternating projections). *Suppose that $\{A, B\}$ is DIL-restrictedly regular at \bar{x} , $0 \leq \sigma < \hat{\theta}_4[A, B](\bar{x})$ and $0 < \tau \leq 1$. Then, for any $\gamma < \hat{\theta}_4[A, B](\bar{x})$ satisfying $0 < \gamma - \sigma \leq \tau$ and*

$$c := \tau^{-1}(1 - \gamma^2 + \gamma\sigma) < 1,$$

any sequence of (τ, σ) -alternating projections for $\{A, B\}$ with initial point sufficiently close to \bar{x} converges to a point in $A \cap B$ with R -linear rate c .

Proof. By Definition 18, there exists a $\rho > 0$ such that condition (25) holds true for all $a \in (A \setminus B) \cap \mathbb{B}_\rho(\bar{x})$ and $b \in (B \setminus A) \cap \mathbb{B}_\rho(\bar{x})$.

Let $a \in A \cap \mathbb{B}_{\rho'}(\bar{x})$ and $b \in P_B^{\tau, \sigma}(a) \cap \mathbb{B}_{\rho'}(\bar{x})$ where $\rho' := \rho/(1 + 2(\gamma - \sigma))$. We are going to show that

$$d(b, A) \leq (1 - \gamma^2 + \gamma\sigma)\|b - a\|.$$

If $b \in A$, the inequality holds true trivially. Suppose $b \notin A$ and denote $\delta := (\gamma - \sigma)\|b - a\|$. Consider any point $u \in A \cap \mathbb{B}_\delta(a)$. Since $\|u - a\| < (\gamma - \sigma)\|b - a\| \leq \tau\|b - a\| \leq d_B(a)$, we see that $u \notin B$; in particular, $a \notin B$ and $u \neq b$. Let z denote the projection of $\frac{u - b}{\|u - b\|}$ on $\mathbb{R}(a - b)$. Then $\|z\| \leq 1$ and, employing Lemma 30,

$$\begin{aligned} d\left(\frac{u - b}{\|u - b\|}, N_B(b)\right) &\leq \left\| \frac{u - b}{\|u - b\|} - z \right\| + d(z, N_B(b)) \\ &\leq \frac{\|u - a\|}{\|b - a\|} + \sigma < (\gamma - \sigma) + \sigma = \gamma. \end{aligned}$$

Since $\|u - \bar{x}\| \leq \|u - a\| + \|a - \bar{x}\| < 2(\gamma - \sigma)\rho' + \rho' = \rho$ and $\|b - \bar{x}\| < \rho' < \rho$, we get from (25) that $d\left(\frac{b-u}{\|u-b\|}, N_A(u)\right) > \gamma$. It follows from Proposition 29 that $d(b, A) \leq \|a - b\| - \gamma\delta = (1 - \gamma^2 + \gamma\sigma)\|a - b\|$. Hence,

$$\|a' - b\| \leq \tau^{-1}d(b, A) \leq c\|a - b\| \quad \text{for all } a' \in P_A^{\tau, \sigma}(b). \quad (32)$$

Now we show that any sequence $\{x_n\}$ of (τ, σ) -alternating projections for $\{A, B\}$ remains in $\mathbb{B}_{\rho'}(\bar{x})$ whenever $x_0 \in \mathbb{B}_{\rho''}(\bar{x})$ where $\rho'' := \left(\frac{\tau^{-1}}{1-c} + 1\right)^{-1} \rho' < \rho'$. Indeed,

$$\|x_1 - x_0\| \leq \tau^{-1}d(x_0, B) \leq \tau^{-1}\|x_0 - \bar{x}\|.$$

Let $n \in \mathbb{N}$ and $x_k \in \mathbb{B}_{\rho'}(\bar{x})$, $k = 0, 1, \dots, n$. It follows from (32) that

$$\|x_{k+1} - x_k\| \leq c^k \|x_1 - x_0\| \quad (k = 0, 1, \dots, n), \quad (33)$$

and consequently,

$$\begin{aligned} \|x_{n+1} - x_0\| &\leq \sum_{k=0}^n c^k \|x_1 - x_0\| \leq \frac{1}{1-c} \|x_1 - x_0\|, \\ \|x_{n+1} - \bar{x}\| &\leq \left(\frac{\tau^{-1}}{1-c} + 1\right) \|x_0 - \bar{x}\| < \rho'. \end{aligned}$$

Thanks to (33), $\{x_k\}$ is a Cauchy sequence containing two subsequences belonging to closed subsets A and B , respectively. Hence, it converges to a point in $A \cap B$ with R -linear rate c . \square

Remark 32. 1. When inexact alternating projections are close to being exact, i.e., τ and σ are close to 1 and 0, respectively (cf. definition (27)), then the assumptions of Theorem 31 are easily satisfied (as long as $\hat{\theta}_4[A, B](\bar{x}) > 0$) while the convergence rate $c = \tau^{-1}(1 - \gamma^2 + \gamma\sigma)$ is mostly determined by the term $1 - \gamma^2$. Recall that γ can be any number in $(0, \hat{\theta}_4[A, B](\bar{x}))$. Thanks to Proposition 18, $1 - \gamma^2 = (\gamma')^2$ where γ' can be any number in $(\hat{c}_4[A, B](\bar{x}), 1)$.

2. When $\dim X < \infty$, the special case $\tau = 1$ and $\sigma = 0$ of Theorem 31 recaptures [10, Theorem 2.3]. The proof given above follows that of [10, Theorem 2.3].

3. It can be of interest to consider a more advanced version of inexact alternating projections than the one given in Definition 27:

$$x_{2n+1} \in P_B^{\tau_1, \sigma_1}(x_{2n}) \quad \text{and} \quad x_{2n+2} \in P_A^{\tau_2, \sigma_2}(x_{2n+1}) \quad (n = 0, 1, \dots),$$

where $\tau_1, \tau_2 \in (0, 1]$ and $\sigma_1, \sigma_2 \in [0, 1)$. For instance, the projections on one of the sets, say, A can be required to be exact, i.e., $\tau_2 = 1$ and $\sigma_2 = 0$. Theorem 31 remains applicable to this situation with $\tau := \min\{\tau_1, \tau_2\}$ and $\sigma := \max\{\sigma_1, \sigma_2\}$. It is possible to obtain a sharper convergence estimate taking into account different ‘‘inexactness’’ parameters for each of the sets. For that, one needs to amend the definition of alternating projections by considering the selection of the pair $\{x_{2n+1}, x_{2n+2}\}$ as a single two-part iteration.

3.2 Convergence for inexact alternating projections under uniform regularity

The motivation for the discussed below version of inexact projections comes from [19, Section 6].

Given a point x and a set A in a Hilbert space and a number $\sigma \in [0, 1)$, the σ -projection of x on A is defined as follows:

$$P_A^\sigma(x) := \{a \in A \mid d(x - a, N_A(a)) \leq \sigma \|x - a\|\}. \quad (34)$$

Observe that

$$P_A^0(x) = \{a \in A \mid x - a \in N_A(a)\} \supset P_A(x)$$

and the inclusion can be strict even in finite dimensions. Furthermore, for any $\sigma \in [0, 1)$, $P_A^\sigma(x)$ can contain points lying arbitrarily far from x .

Definition 33 (Inexact alternating projections). Given a number $\sigma \in [0, 1)$, $\{x_n\}$ is a sequence of σ -alternating projections for $\{A, B\}$ if

$$\begin{aligned} x_{2n+1} &\in P_B^\sigma(x_{2n}) \quad \text{and} \quad x_{2n+2} \in P_A^\sigma(x_{2n+1}), \\ \|x_{n+2} - x_{n+1}\| &\leq \|x_{n+1} - x_n\| \quad (n = 0, 1, \dots). \end{aligned} \quad (35)$$

The role of the monotonicity condition (35) in Definition 33 is to compensate for the lack of good projection properties of the σ -projection operator (34). In the case of standard alternating projections (cf. Definition 1), this condition is satisfied automatically.

Theorem 34 (Convergence of inexact alternating projections under uniform regularity). *Suppose that $\{A, B\}$ is uniformly regular at \bar{x} , A is super-regular at \bar{x} and $\sigma \in [0, 1)$ satisfies*

$$c_0 := \hat{c}[A, B](\bar{x})(1 - \sigma^2) + \sigma^2 + 2\sigma\sqrt{1 - \sigma^2} + \sigma < 1.$$

Then, for any $c \in (c_0, 1)$, any sequence $\{x_k\}$ of σ -alternating projections for $\{A, B\}$ with initial points x_0 and x_1 sufficiently close to \bar{x} converges to a point in $A \cap B$ with R -linear rate \sqrt{c} .

Proof. Let $c \in (c_0, 1)$ and choose a $c_1 > \hat{c}[A, B](\bar{x})$ and a $\gamma > 0$ such that

$$c_1(1 - \sigma^2) + \sigma^2 + (2\sigma + \gamma)\sqrt{1 - \sigma^2} + \sigma < c. \quad (36)$$

By Proposition 4(iii) and Definition 8(ii), there exists a $\delta > 0$ such that

$$-\langle u, v \rangle \leq c_1 \|u\| \|v\|, \quad (37)$$

$$\langle u, x - a \rangle \leq \gamma \|u\| \|x - a\| \quad (38)$$

for all $x, a \in A \cap \mathbb{B}_\delta(\bar{x})$, $b \in B \cap \mathbb{B}_\delta(\bar{x})$, $u \in N_A(a)$ and $v \in N_B(b)$.

Let $a_1 \in A \cap \mathbb{B}_\delta(\bar{x})$, $b \in P_B^\sigma(a_1) \cap \mathbb{B}_\delta(\bar{x})$ and $a_2 \in P_A^\sigma(b) \cap \mathbb{B}_\delta(\bar{x})$. We are going to show that

$$\|a_2 - b\| \leq c\|b - a_1\|. \quad (39)$$

By definition (34), for any $\varepsilon \in (0, 1 - \sigma)$, there exist $u \in N_A(a_2)$ and $v \in N_B(b)$ such that

$$\|b - a_2 - u\| \leq (\sigma + \varepsilon)\|b - a_2\| \quad \text{and} \quad \|a_1 - b - v\| \leq (\sigma + \varepsilon)\|a_1 - b\|. \quad (40)$$

Additionally, one can ensure that

$$\|u\| \leq \sqrt{1 - (\sigma + \varepsilon)^2}\|b - a_2\| \quad \text{and} \quad \|v\| \leq \sqrt{1 - (\sigma + \varepsilon)^2}\|a_1 - b\|. \quad (41)$$

Indeed, take any $u \in N_A(a_2)$ satisfying the first inequality in (40). If $u = 0$, the first inequality in (41) is satisfied too. Suppose $u \neq 0$ and consider $u_1 := \langle b - a_2, u \rangle \frac{u}{\|u\|^2}$ - the projection of $b - a_2$ on $\mathbb{R}u$. Then

$$\|b - a_2\|^2 = \|u_1\|^2 + \|b - a_2 - u_1\|^2, \quad (42)$$

$$\|b - a_2 - u_1\| \leq \|b - a_2 - u\| \leq (\sigma + \varepsilon)\|b - a_2\| \quad (43)$$

and there exists a $t \in (0, 1]$ such that $u_2 := tu_1$ satisfies

$$\|b - a_2 - u_2\| = (\sigma + \varepsilon)\|b - a_2\| \quad (44)$$

(thanks to the continuity of the function $t \mapsto \|b - a_2 - tu_1\|$). Hence, $u_2 \in N_A(a_2)$, vector $u_1 - u_2$ is a projection of $b - a_2 - u_2$ on $\mathbb{R}u$, i.e.,

$$\|b - a_2 - u_2\|^2 = \|u_1 - u_2\|^2 + \|b - a_2 - u_1\|^2, \quad (45)$$

and, using (42), (45), (44), and (43),

$$\begin{aligned} \|u_2\| &= \|u_1\| - \|u_1 - u_2\| \\ &= \sqrt{\|b - a_2\|^2 - \|b - a_2 - u_1\|^2} - \sqrt{(\sigma + \varepsilon)^2\|b - a_2\|^2 - \|b - a_2 - u_1\|^2} \\ &= \frac{(1 - (\sigma + \varepsilon)^2)\|b - a_2\|^2}{\sqrt{\|b - a_2\|^2 - \|b - a_2 - u_1\|^2} + \sqrt{(\sigma + \varepsilon)^2\|b - a_2\|^2 - \|b - a_2 - u_1\|^2}} \\ &\leq \frac{(1 - (\sigma + \varepsilon)^2)\|b - a_2\|^2}{\sqrt{1 - (\sigma + \varepsilon)^2}\|b - a_2\|} = \sqrt{1 - (\sigma + \varepsilon)^2}\|b - a_2\|. \end{aligned}$$

Similarly, given any $v \in N_B(b)$ satisfying the second inequality in (40), one can find a $v_2 \in N_B(b)$ satisfying this inequality and, additionally, the second inequality in (41).

Making use of (37), (40) and (41), we get

$$\begin{aligned} -\langle b - a_2, a_1 - b \rangle &= -\langle u, v \rangle - \langle u, a_1 - b - v \rangle \\ &\quad - \langle b - a_2 - u, v \rangle - \langle b - a_2 - u, a_1 - b - v \rangle \\ &\leq c_1\|u\|\|v\| + \|u\|\|a_1 - b - v\| \\ &\quad + \|b - a_2 - u\|\|v\| + \|b - a_2 - u\|\|a_1 - b - v\| \\ &\leq \left(c_1(1 - (\sigma + \varepsilon)^2) + 2(\sigma + \varepsilon)(\sqrt{1 - (\sigma + \varepsilon)^2}) \right. \\ &\quad \left. + (\sigma + \varepsilon)^2 \right) \|b - a_2\|\|a_1 - b\|. \end{aligned}$$

At the same time, making use of (38) and the first inequalities in (40) and (41), we have

$$\begin{aligned}\langle b - a_2, a_1 - a_2 \rangle &= \langle u, a_1 - a_2 \rangle + \langle b - a_2 - u, a_1 - a_2 \rangle \\ &\leq \gamma \|u\| \|a_1 - a_2\| + (\sigma + \varepsilon) \|b - a_2\| \|a_1 - a_2\| \\ &\leq (\gamma \sqrt{1 - (\sigma + \varepsilon)^2} + (\sigma + \varepsilon)) \|b - a_2\| \|a_1 - a_2\|,\end{aligned}$$

Adding the last two estimates and passing to limit as $\varepsilon \downarrow 0$, we obtain

$$\|b - a_2\|^2 \leq \left(c_1(1 - \sigma^2) + \sigma^2 + (2\sigma + \gamma) \sqrt{1 - \sigma^2} + \sigma \right) \|b - a_2\| \|a_1 - b\|.$$

Thanks to (36), this proves (39).

Now we show that a sequence $\{x_n\}$ of σ -alternating projections for $\{A, B\}$ remains in $\mathbb{B}_\delta(\bar{x})$ if $x_0, x_1 \in \mathbb{B}_\rho(\bar{x})$ where $\rho := \frac{1-c}{5-c}\delta < \delta$. Let $n \in \mathbb{N}$ and $x_k \in \mathbb{B}_\delta(\bar{x})$, $k = 0, 1, \dots, 2n$. It follows from (39) that

$$\|x_{2k} - x_{2k-1}\| \leq c^k \|x_1 - x_0\| \quad (k = 1, 2, \dots, n), \quad (46)$$

and consequently, employing also (35),

$$\begin{aligned}\|x_{2n+2} - x_0\| &\leq \|x_{2n+2} - x_{2n+1}\| + \|x_{2n+1} - x_0\| \leq \|x_{2n+2} - x_{2n+1}\| \\ &\quad + \sum_{k=1}^n (\|x_{2k+1} - x_{2k}\| + \|x_{2k} - x_{2k-1}\|) + \|x_1 - x_0\| \\ &\leq 2 \sum_{k=1}^n \|x_{2k} - x_{2k-1}\| + 2\|x_1 - x_0\| \\ &\leq 2 \sum_{k=0}^n c^k \|x_1 - x_0\| \leq \frac{2}{1-c} \|x_1 - x_0\|.\end{aligned}$$

Thus,

$$\begin{aligned}\max\{\|x_{2n+2} - \bar{x}\|, \|x_{2n+1} - \bar{x}\|\} &\leq \max\{\|x_{2n+2} - x_0\|, \|x_{2n+1} - x_0\|\} \\ &\quad + \|x_0 - \bar{x}\| \leq \frac{2}{1-c} \|x_1 - x_0\| + \|x_0 - \bar{x}\| \\ &\leq \frac{2}{1-c} \|x_1 - \bar{x}\| + \frac{3-c}{1-c} \|x_0 - \bar{x}\| < \frac{5-c}{1-c} \rho = \delta,\end{aligned}$$

i.e., $x_{2n+1}, x_{2n+2} \in \mathbb{B}_\delta(\bar{x})$.

Thanks to (46), $\{x_k\}$ is a Cauchy sequence containing two subsequences belonging to closed subsets A and B , respectively. Hence, it converges to a point in $A \cap B$ with R -linear rate \sqrt{c} . \square

Remark 35. 1. When the ‘‘inexactness’’ parameter σ is small (cf. definition (33)), then the assumptions of Theorem 34 are easily satisfied (as long as $\tilde{c}[A, B](\bar{x}) < 1$ and condition (35) holds) while the convergence rate is close to the one guaranteed by Theorem 2 and [19, Theorem 6.1].

2. One can also consider a more advanced version of inexact alternating projections than the one given in Definition 33:

$$x_{2n+1} \in P_B^{\sigma_1}(x_{2n}) \quad \text{and} \quad x_{2n+2} \in P_A^{\sigma_2}(x_{2n+1}), \quad (n = 0, 1, \dots).$$

where $\sigma_1, \sigma_2 \in [0, 1)$. Theorem 34 remains applicable to this situation with $\sigma := \max\{\sigma_1, \sigma_2\}$ (cf. Remark 32.3).

3. Observe that, thanks to (39), for odd values of n , condition (35) is improved in the proof of Theorem 34:

$$\|x_{n+2} - x_{n+1}\| \leq c\|x_{n+1} - x_n\|,$$

where $c < 1$. However, the assumption is still needed to ensure that x_{n+2} is not too far from \bar{x} and uniform and super-regularity conditions are applicable.

4. Constant c_1 in (37) is an upper estimate of the cosine of the angle φ between vectors u and $-v$ while $\sigma + \varepsilon$ in (40) can be interpreted as an upper estimate of the sine of the angles ψ_1 and ψ_2 between vectors $b - a_2$ and u and $a_1 - b$ and v , respectively. One can use standard trigonometric identities and inequalities (37) and (40) to obtain an upper estimate of the cosine of the angle $\varphi - \psi_1 - \psi_2$ between vectors $b - a_2$ and $b - a_1$ and possibly improve the convergence estimate in the statement of Theorem 34.

5. If both subsets A and B are super-regular, then in the proof of Theorem 34, one can establish an analogue of (39) with subsets A and B interchanged:

$$\|b_2 - a\| \leq c\|a - b_1\|,$$

where $b_1 \in B \cap \mathbb{B}_\delta(\bar{x})$, $a \in P_A^\sigma(b_1) \cap \mathbb{B}_\delta(\bar{x})$ and $b_2 \in P_B^\sigma(a) \cap \mathbb{B}_\delta(\bar{x})$. This guarantees an improvement with rate c on each iteration. As a result, one obtains a better overall R -linear rate c .

6. The conclusion of Theorem 34 remains true if one replaces the assumptions of uniform regularity of $\{A, B\}$ (and the regularity constant $\hat{c}[A, B](\bar{x})$) and super-regularity of A with BLPW-restricted regularity (and the regularity constant $\hat{c}_1[A, B](\bar{x})$) and B -super-regularity, respectively, accompanied by appropriate adjustments in the definition of σ -projections.

Acknowledgements

The authors wish to express their gratitude to the anonymous referee for the careful reading of the manuscript and helpful comments and suggestions.

References

- [1] ATTOUCH, H., BOLTE, J., REDONT, P., AND SOUBEYRAN, A. Proximal alternating minimization and projection methods for nonconvex problems: an approach based on the Kurdyka–Lojasiewicz inequality. *Math. Oper. Res.* 35, 2 (2010), 438–457.

- [2] AUSSEL, D., DANILIDIS, A., AND THIBAUT, L. Subsmooth sets: functional characterizations and related concepts. *Trans. Amer. Math. Soc.* 357, 4 (2005), 1275–1301.
- [3] BAUSCHKE, H. H., AND BORWEIN, J. M. On the convergence of von Neumann’s alternating projection algorithm for two sets. *Set-Valued Anal.* 1, 2 (1993), 185–212.
- [4] BAUSCHKE, H. H., LUKE, D. R., PHAN, H. M., AND WANG, X. Restricted normal cones and the method of alternating projections: applications. *Set-Valued Var. Anal.* 21, 3 (2013), 475–501.
- [5] BAUSCHKE, H. H., LUKE, D. R., PHAN, H. M., AND WANG, X. Restricted normal cones and the method of alternating projections: theory. *Set-Valued Var. Anal.* 21, 3 (2013), 431–473.
- [6] BREGMAN, L. M. The method of successive projection for finding a common point of convex sets. *Sov. Math., Dokl.* 6 (1965), 688–692.
- [7] CLARKE, F. H., STERN, R. J., AND WOLENSKI, P. R. Proximal smoothness and the lower- C^2 property. *J. Conv. Anal.* 2 (1995), 97–116.
- [8] DE GIORGI, E., MARINO, A., AND TOSQUES, M. Problems of evolution in metric spaces and maximal decreasing curve. *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (8)* 68, 3 (1980), 180–187. In Italian.
- [9] DONTCHEV, A. L., AND ROCKAFELLAR, R. T. *Implicit Functions and Solution Mappings. A View from Variational Analysis*. Springer Monographs in Mathematics. Springer, Dordrecht, 2009.
- [10] DRUSVYATSKIY, D., IOFFE, A. D., AND LEWIS, A. S. Alternating projections and coupling slope. *arXiv: 1401.7569* (2014), 1–17.
- [11] FABIAN, M. J., HENRION, R., KRUGER, A. Y., AND OTRATA, J. V. Error bounds: necessary and sufficient conditions. *Set-Valued Var. Anal.* 18, 2 (2010), 121–149.
- [12] IOFFE, A. D. Metric regularity and subdifferential calculus. *Russian Math. Surveys* 55 (2000), 501–558.
- [13] KRUGER, A. Y. Stationarity and regularity of set systems. *Pac. J. Optim.* 1, 1 (2005), 101–126.
- [14] KRUGER, A. Y. About regularity of collections of sets. *Set-Valued Anal.* 14, 2 (2006), 187–206.
- [15] KRUGER, A. Y. About stationarity and regularity in variational analysis. *Taiwanese J. Math.* 13, 6A (2009), 1737–1785.
- [16] KRUGER, A. Y., AND LÓPEZ, M. A. Stationarity and regularity of infinite collections of sets. *J. Optim. Theory Appl.* 154, 2 (2012), 339–369.

- [17] KRUGER, A. Y., AND THAO, N. H. About uniform regularity of collections of sets. *Serdica Math. J.* 39 (2013), 287–312.
- [18] KRUGER, A. Y., AND THAO, N. H. Quantitative characterizations of regularity properties of collections of sets. *J. Optim. Theory Appl.* 164, 1 (2015), 41–67.
- [19] LEWIS, A. S., LUKE, D. R., AND MALICK, J. Local linear convergence for alternating and averaged nonconvex projections. *Found. Comput. Math.* 9, 4 (2009), 485–513.
- [20] LEWIS, A. S., AND MALICK, J. Alternating projection on manifolds. *Math. Oper. Res.* 33, 1 (2008), 216–234.
- [21] MORDUKHOVICH, B. S. *Variational Analysis and Generalized Differentiation. I: Basic Theory*, vol. 330 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer, Berlin, 2006.
- [22] NOLL, D., AND RONDEPIERRE, A. On local convergence of the method of alternating projections. *arXiv: 1312.5681v2* (2014), 1–24.
- [23] POLIQUIN, R. A., ROCKAFELLAR, R. T., AND THIBAUT, L. Local differentiability of distance functions. *Trans. Amer. Math. Soc.* 352, 11 (2000), 5231–5249.
- [24] ROCKAFELLAR, R. T., AND WETS, R. J.-B. *Variational Analysis*. Springer, Berlin, 1998.
- [25] SHAPIRO, A. Existence and differentiability of metric projections in Hilbert spaces. *SIAM J. Optim.* 4, 1 (1994), 130–141.