

Perturbation of error bounds

A. Y. Kruger · M. A. López · M. A. Théra

Dedicated to Professor Terry Rockafellar in honor of his 80th birthday

the date of receipt and acceptance should be inserted later

Abstract Our aim in the current article is to extend the developments in Kruger, Ngai & Théra, *SIAM J. Optim.* **20**(6), 3280–3296 (2010) and, more precisely, to characterize, in the Banach space setting, the stability of the local and global error bound property of inequalities determined by proper lower semicontinuous under data perturbations. We propose new concepts of (arbitrary, convex and linear) perturbations of the given function defining the system under consideration, which turn out to be a useful tool in our analysis. The characterizations of error bounds for families of perturbations can be interpreted as estimates of the ‘radius of error bounds’. The definitions and characterizations are illustrated by examples.

Keywords Error bound · Feasibility problem · Perturbation · Subdifferential · Metric regularity · Metric subregularity

Mathematics Subject Classification (2000) 49J52 · 49J53 · 90C30

The research is supported by the Australian Research Council: project DP160100854; EDF and the Jacques Hadamard Mathematical Foundation: Gaspard Monge Program for Optimization and Operations Research. The research of the second and third authors is also supported by MINECO of Spain and FEDER of EU: grant MTM2014-59179-C2-1-P.

A. Y. Kruger
Centre for Informatics and Applied Optimization, Faculty of Science and Technology, Federation University Australia
E-mail: a.kruger@federation.edu.au

M. A. López
Department of Statistics and Operations Research, University of Alicante, Spain and Federation University Australia
E-mail: marco.antonio@ua.es

M. A. Théra
Laboratoire XLIM, UMR-CNRS 6172, University of Limoges, France and Federation University Australia
E-mail: michel.thera@unilim.fr

1 Introduction and Preliminaries

In this article, we mostly follow the standard terminology and notation used by the optimization and variational analysis community, see, e.g., Rockafellar & Wets [55]. Throughout, X and X^* stand for a real Banach space and its topological dual, respectively. The norms in both spaces are denoted by the same symbol $\|\cdot\|$. For a subset S , we denote its interior and boundary by $\text{int}S$ and $\text{bd}S$, respectively. The distance from a point x to a set S is denoted by $d(x, S) := \inf_{u \in S} \|u - x\|$, and we use the convention $d(x, S) = +\infty$ whenever $S = \emptyset$.

For an extended-real-valued function: $f : X \rightarrow \mathbb{R}_\infty := \mathbb{R} \cup \{+\infty\}$, its domain is the set $\text{dom} f := \{x \in X \mid f(x) < +\infty\}$. A function f is said to be *proper* if $\text{dom} f \neq \emptyset$. We use the symbol $f_+(x)$ to denote $\max(f(x), 0)$. The class of all extended-real-valued proper convex lower semicontinuous functions on X is denoted by $\Gamma_0(X)$. For a convex function $f : X \rightarrow \mathbb{R}_\infty$, its (Moreau) *subdifferential* at $x \in \text{dom} f$ is given by

$$\partial f(x) := \{x^* \in X^* \mid \langle x^*, u - x \rangle \leq f(u) - f(x), \forall u \in X\}.$$

It is a (possibly empty) weak*-closed set in X^* .

The article is concerned with the study of the solution set of a single inequality of the type

$$S_f := \{x \in X \mid f(x) \leq 0\}, \quad (1)$$

where $f : X \rightarrow \mathbb{R}_\infty$. Such sets subsume, e.g., feasible sets in mathematical programming. Indeed, solutions of a finite family of inequalities:

$$\{x \in X \mid f_i(x) \leq 0 \quad \text{for all } i = 1, \dots, n\}$$

can be rewritten as (1) with function f defined by $f(x) = \max\{f_1(x), \dots, f_n(x)\}$.

An important issue when studying systems of the type (1) is to give an upper estimate (error bound) of the distance from a point $x \in X$ to the set S_f in terms of a computable function measuring the violation of the inequality in (1). This can, e.g., be the function f itself.

We say that a function f has (or admits) a *local error bound* at a point $\bar{x} \in S_f$ if there exists a real $\tau > 0$ such that

$$\tau d(x, S_f) \leq f_+(x) \quad (2)$$

for all x near \bar{x} . We similarly say that a function f has a *global error bound* if there exists a real $\tau > 0$ such that inequality (2) is satisfied for all $x \in X$.

The exact upper bound of all such τ (the *error bound modulus*; cf. [22]) equals either

$$\text{Er} f(\bar{x}) := \liminf_{\substack{x \rightarrow \bar{x} \\ f(x) > 0}} \frac{f(x)}{d(x, S(f))} \quad (3)$$

in the local setting, or

$$\text{Er} f := \inf_{f(x) > 0} \frac{f(x)}{d(x, S(f))} \quad (4)$$

in the global case. Constants (3) and (4) provide quantitative estimates of the error bound property.

The starting point of the theory of error bounds goes back to the pioneering work by Hoffman [28] (although some traces of the error bound property can be found in an earlier publication by Rosenbloom [56]¹) who established for a linear function in finite dimensions the following result:

Given an $m \times n$ matrix A and a vector $b \in \mathbb{R}^m$, there exists a positive number $\kappa > 0$ such that the distance from x to the set $S := \{x \in \mathbb{R}^n \mid Ax \leq b\}$ has an upper bound given by $\kappa \|(Ax - b)_+\|$, where for $y := (y_1, \dots, y_m) \in \mathbb{R}^m$, y_+ denotes the vector $(\max(y_1, 0), \dots, \max(y_m, 0))$.

After the work by Hoffmann and its extensions by Robinson [54], Mangasarian [42], Auslender & Crouzeix [1], Pang [52], Lewis and Pang [41], Klatte & Li [34], Jourani [33], there have been significant developments of various aspects of error bounds for convex and nonconvex functions in recent years. The interested reader is referred to the articles by Ng & Zheng [44], Azé [2, 3], Azé & Corvellec [4], Zălinescu [58], Huang & Ng [29], Corvellec & Motreanu [18], Fabian et al [22, 23], Gfrerer [24], Ioffe [30, 31], Ioffe & Outrata [32], Ngai & Théra [46–49], Zheng & Ng [59, 60], Bednarczuk & Kruger [7, 8], Meng & Yang [43], Kruger [35–37] and the references therein.

Many authors have recently studied error bounds in connection with the *metric regularity* and *subregularity* (cf. [19]) as well as *Aubin property* and *calmness* of set-valued mappings: [3, 13, 14, 24, 30–32, 35–37, 46, 48, 50, 51, 59, 60]. The connections between the error bounds and weak sharp minima were studied in [12].

Another typical example where the theory of error bounds plays an important role is the so-called *feasibility problem* [5, 6, 27], which consists in finding a point in the intersection of a finite family of closed (usually convex) sets and has a broad applicability in various areas such as, e.g., image reconstruction [16]. Several iterative methods such as the method of successive orthogonal projections, the cyclic subgradient projections method, etc, are known to solve this problem (see [15]). Error bounds are also used in the convergence analysis of projection algorithms. They allow one to quantify the proximity of an iterate to the solution set of the problem. The reader is referred to the recent survey paper by Ioffe [31] and the references therein and also to some recent contributions by Beck & Teboulle [6], Lewis et al [40], Hesse & Luke [27], Drusvyatskiy et al [20], Bolte et al [10] and Kruger & Thao [39].

In this article, we study stability of local and global error bounds under perturbations. It was observed in Ngai et al [45, Theorem 1] and Kruger et al [38, Theorem 1], that the requirement for the distance from the origin to the subdifferential of a function at a reference point to be strictly positive, while being a conventional sufficient condition for a local error bound for the system (1), is far from being necessary. This condition guarantees the local error bound property not just for the given function, but also for a family of its small perturbations. In Section 2, we exploit the concept of local ε -perturbation from [38], define subfamilies of convex and linear ε -perturbations, and establish in Theorem 2.2 the exact formula for the ‘radius of local error bounds’ for families of arbitrary, convex and linear perturbations in terms of the distance from the origin to the subdifferential.

¹ Private communication by J.-B. Hiriart-Urruty

The same idea applies when considering stability of global error bounds in Section 3. In the global setting, we define families of convex and linear perturbations as well as larger families of weak convex and weak linear perturbations, discuss some limitations of these definitions, and establish in Theorem 3.2 lower and upper estimates for the ‘radius of global error bounds’ for families of such perturbations. The family of convex perturbations considered here is larger than the corresponding one studied in [38]. In particular, the so called *asymptotic qualification condition* is waived.

Some examples are given for the convenience of the reader to illustrate the different concepts introduced along the presentation.

2 Stability of local error bounds

In this section we establish conditions for stability of local error bounds for the constraint system (1). We start with the following statement extracted from [38, Theorem 1].

Theorem 2.1 *Let $f \in \Gamma_0(X)$ and $f(\bar{x}) = 0$. Function f has a local error bound at \bar{x} provided that one of the following two conditions is satisfied:*

- (i) $\tau(f, \bar{x}) := \liminf_{x \rightarrow \bar{x}, f(x) > f(\bar{x})} d(0, \partial f(x)) > 0$;
- (ii) $\zeta(f, \bar{x}) := d(0, \text{bd } \partial f(\bar{x})) > 0$.

Moreover, condition (i) is also necessary for f to have a local error bound at \bar{x} and

$$\zeta(f, \bar{x}) \leq \tau(f, \bar{x}) = \text{Er } f(\bar{x}). \quad (5)$$

Constant $\tau(f, \bar{x})$ is known as the *strict outer subdifferential slope* [22] $|\overline{\partial f}|^>(\bar{x})$ of f at \bar{x} . The sufficient criterion (i) was used in [32, Theorem 2.1 (c)], [49, Corollary 2 (ii)], [53, Theorem 4.12], [57, Theorem 3.1]. Criterion (ii) was used in [25, Corollary 3.4], [26, Theorem 4.2]. The equality in (5) is well known. See also characterizations of linear and nonlinear conditionings in [17, Theorem 5.2].

The inequality in (5) can be strict.

Example 2.1 $f(x) \equiv 0, x \in \mathbb{R}$. Obviously $0 \in \text{bd } \partial f(\bar{x})$, $\zeta(f, \bar{x}) = 0$, while $\tau(f, \bar{x}) = \infty$ for any $\bar{x} \in \mathbb{R}$. △

Example 2.2 $f(x) = 0$ if $x \leq 0$, and $f(x) = x$ if $x > 0$. Then $\partial f(0) = [0, 1]$ and $0 \in \text{bd } \partial f(0)$, $\zeta(f, \bar{x}) = 0$, while $\tau(f, 0) = 1$. △

Thus, condition (ii) in Theorem 2.1 is in general stronger than condition (i). It characterizes a stronger property than just the existence of a local error bound for f at \bar{x} ; namely, it guaranties the local error bound property for the family of functions being small perturbations of f in the sense defined below.

Let $f(\bar{x}) < \infty$ and $\varepsilon \geq 0$. Following [38, Definition 5], we say that $g : X \rightarrow \mathbb{R}_\infty$ is an ε -perturbation of f near \bar{x} if

$$g(\bar{x}) = f(\bar{x}) \quad (6)$$

and

$$\limsup_{x \rightarrow \bar{x}} \frac{|g(x) - f(x)|}{\|x - \bar{x}\|} \leq \varepsilon. \quad (7)$$

If both functions f and g are continuous at \bar{x} , then equality (6) in the above definition is obviously implied by condition (7).

The collection of all ε -perturbations of f near \bar{x} will be denoted $\text{Ptb}(f, \bar{x}, \varepsilon)$. Obviously, if $g \in \text{Ptb}(f, \bar{x}, \varepsilon)$, then $f \in \text{Ptb}(g, \bar{x}, \varepsilon)$, and neither f nor g are required to be convex. If both f and g are convex, then the actual perturbation function $p := g - f$ needs not to be convex; it is in general a d.c. function (difference of convex functions).

The following subsets of $\text{Ptb}(f, \bar{x}, \varepsilon)$ corresponding, respectively, to convex and linear ε -perturbations of f near \bar{x} can be of interest:

$$\begin{aligned} \text{Ptb}_c(f, \bar{x}, \varepsilon) &:= \{g \in \text{Ptb}(f, \bar{x}, \varepsilon) \mid g - f \in \Gamma_0(X)\}, \\ \text{Ptb}_l(f, \bar{x}, \varepsilon) &:= \{g \mid g(u) = f(u) + \langle x^*, u - \bar{x} \rangle (u \in X), x^* \in \varepsilon \mathbb{B}_{X^*}^*\}. \end{aligned} \quad (8)$$

Obviously,

$$\text{Ptb}_l(f, \bar{x}, \varepsilon) \subset \text{Ptb}_c(f, \bar{x}, \varepsilon) \subset \text{Ptb}(f, \bar{x}, \varepsilon). \quad (9)$$

The next proposition provides a sufficient condition for a function g to be a (convex) ε -perturbations of f near \bar{x} .

Proposition 2.1 *Let $f(\bar{x}) < \infty$ and $\varepsilon \geq 0$. Suppose $g = f + p$ where $p : X \rightarrow \mathbb{R}_\infty$ is convex and Lipschitz continuous near \bar{x} with constant ε and $p(\bar{x}) = 0$. Then $g \in \text{Ptb}_c(f, \bar{x}, \varepsilon)$.*

Proof Thanks to the Lipschitz continuity of p , $\partial p(\bar{x}) \neq \emptyset$ and $\|x^*\| \leq \varepsilon$ for all $x^* \in \partial p(\bar{x})$; cf. [11, Proposition 4.1.25 and its proof]. Given an $x^* \in \partial p(\bar{x})$, for any $x \in X \setminus \{\bar{x}\}$, we have:

$$\frac{g(x) - f(x)}{\|x - \bar{x}\|} = \frac{p(x)}{\|x - \bar{x}\|} \geq \frac{\langle x^*, x - \bar{x} \rangle}{\|x - \bar{x}\|} \geq -\varepsilon.$$

On the other hand,

$$\limsup_{x \rightarrow \bar{x}} \frac{g(x) - f(x)}{\|x - \bar{x}\|} = \limsup_{x \rightarrow \bar{x}} \frac{p(x)}{\|x - \bar{x}\|} \leq \varepsilon.$$

Combining the two inequalities proves (7). \square

The assumption of the Lipschitz continuity of p in the above proposition is essential.

Example 2.3 Let $f, p, g : \mathbb{R} \rightarrow \mathbb{R}_\infty$, $f(x) = 0$ for all $x \in \mathbb{R}$ and

$$p(x) = g(x) = \begin{cases} 0 & \text{if } x \leq 0, \\ +\infty & \text{if } x > 0. \end{cases}$$

Then $f, g, p \in \Gamma_0(X)$ and

$$\limsup_{x \rightarrow 0} \frac{|g(x) - f(x)|}{|x|} = +\infty.$$

\triangle

Given a function $f \in \Gamma_0(X)$ with $f(\bar{x}) = 0$ and a number $\varepsilon \geq 0$, denote

$$\begin{aligned} \text{Er}\{\text{Ptb}(f, \bar{x}, \varepsilon)\}(\bar{x}) &:= \inf_{g \in \Gamma_0(X) \cap \text{Ptb}(f, \bar{x}, \varepsilon)} \text{Er} g(\bar{x}), \\ \text{Er}\{\text{Ptb}_c(f, \bar{x}, \varepsilon)\}(\bar{x}) &:= \inf_{g \in \text{Ptb}_c(f, \bar{x}, \varepsilon)} \text{Er} g(\bar{x}), \\ \text{Er}\{\text{Ptb}_l(f, \bar{x}, \varepsilon)\}(\bar{x}) &:= \inf_{g \in \text{Ptb}_l(f, \bar{x}, \varepsilon)} \text{Er} g(\bar{x}). \end{aligned} \quad (10)$$

These numbers characterize the error bound property for families of ε -perturbations of f near \bar{x} . Thanks to (9), it holds

$$\begin{aligned} \text{Er}\{\text{Ptb}(f, \bar{x}, \varepsilon)\}(\bar{x}) &\leq \text{Er}\{\text{Ptb}_c(f, \bar{x}, \varepsilon)\}(\bar{x}) \\ &\leq \text{Er}\{\text{Ptb}_l(f, \bar{x}, \varepsilon)\}(\bar{x}) \leq \text{Er} f(\bar{x}) \end{aligned} \quad (11)$$

for any $\varepsilon \geq 0$.

Theorem 2.2 *Let $f \in \Gamma_0(X)$, $f(\bar{x}) = 0$. Then*

$$\begin{aligned} \zeta(f, \bar{x}) &= \inf\{\varepsilon > 0 \mid \text{Er}\{\text{Ptb}(f, \bar{x}, \varepsilon)\}(\bar{x}) = 0\} \\ &= \inf\{\varepsilon > 0 \mid \text{Er}\{\text{Ptb}_c(f, \bar{x}, \varepsilon)\}(\bar{x}) = 0\} \\ &= \inf\{\varepsilon > 0 \mid \text{Er}\{\text{Ptb}_l(f, \bar{x}, \varepsilon)\}(\bar{x}) = 0\}. \end{aligned} \quad (12)$$

Proof Thanks to the first two inequalities in (11), we always have

$$\begin{aligned} &\inf\{\varepsilon > 0 \mid \text{Er}\{\text{Ptb}(f, \bar{x}, \varepsilon)\}(\bar{x}) = 0\} \\ &\leq \inf\{\varepsilon > 0 \mid \text{Er}\{\text{Ptb}_c(f, \bar{x}, \varepsilon)\}(\bar{x}) = 0\} \\ &\leq \inf\{\varepsilon > 0 \mid \text{Er}\{\text{Ptb}_l(f, \bar{x}, \varepsilon)\}(\bar{x}) = 0\}. \end{aligned} \quad (13)$$

We are going to show that

$$\zeta(f, \bar{x}) \leq \inf\{\varepsilon > 0 \mid \text{Er}\{\text{Ptb}(f, \bar{x}, \varepsilon)\}(\bar{x}) = 0\} \quad (14)$$

and

$$\inf\{\varepsilon > 0 \mid \text{Er}\{\text{Ptb}_l(f, \bar{x}, \varepsilon)\}(\bar{x}) = 0\} \leq \zeta(f, \bar{x}). \quad (15)$$

By [38, Theorem 8 (i)], $\text{Er} g(\bar{x}) \geq \zeta(f, \bar{x}) - \varepsilon$ for any $\varepsilon > 0$ and any $g \in \Gamma_0(X) \cap \text{Ptb}(f, \bar{x}, \varepsilon)$. By definition (10), we have

$$\zeta(f, \bar{x}) \leq \text{Er}\{\text{Ptb}(f, \bar{x}, \varepsilon)\}(\bar{x}) + \varepsilon$$

for any $\varepsilon > 0$, and consequently $\zeta(f, \bar{x}) \leq \varepsilon$ if $\text{Er}\{\text{Ptb}(f, \bar{x}, \varepsilon)\}(\bar{x}) = 0$, which yields inequality (14).

If $\zeta(f, \bar{x}) = \infty$, inequality (15) holds trivially. Let $\zeta(f, \bar{x}) < \varepsilon < \infty$. We are going to show that $\text{Er}\{\text{Ptb}_l(f, \bar{x}, \varepsilon)\}(\bar{x}) = 0$. By the definition of $\zeta(f, \bar{x})$, there exists an $x^* \in \text{bd} \partial f(\bar{x})$ such that $\|x^*\| < \varepsilon$. Choose a positive number $\xi < \varepsilon - \|x^*\|$. Thus,

$$f(u) \geq \langle x^*, u - \bar{x} \rangle \quad \text{for all } u \in X, \quad (16)$$

and there exists an $\hat{x}^* \in X^*$ such that $\|\hat{x}^* - x^*\| < \xi$ and $\hat{x}^* \notin \partial f(\bar{x})$, which means that there exists a point $\hat{x} \in X \setminus \{\bar{x}\}$ satisfying

$$f(\hat{x}) = f(\hat{x}) - f(\bar{x}) < \langle \hat{x}^*, \hat{x} - \bar{x} \rangle.$$

Thanks to the convexity of f , one also has

$$f(x_k) < \langle \hat{x}^*, x_k - \bar{x} \rangle, \quad k = 1, 2, \dots, \quad (17)$$

where $x_k := \bar{x} + \frac{1}{k}(\hat{x} - \bar{x})$. Now select a $z^* \in X^*$ such that $\|z^*\| = 1$ and $\langle z^*, \hat{x} - \bar{x} \rangle = \|\hat{x} - \bar{x}\|$, and define

$$g(u) := f(u) + \langle -x^* + \xi z^*, u - \bar{x} \rangle, \quad u \in X. \quad (18)$$

One has $\| -x^* + \xi z^* \| \leq \|x^*\| + \xi < \varepsilon$, and consequently, $g \in \text{Ptb}_l(f, \bar{x}, \varepsilon)$. It follows from (16) and (18) that

$$g(u) \geq \xi \langle z^*, u - \bar{x} \rangle \quad \text{for all } u \in X. \quad (19)$$

In particular,

$$g(x_k) \geq \xi \langle z^*, x_k - \bar{x} \rangle = \frac{\xi}{k} \|\hat{x} - \bar{x}\| > 0, \quad k = 1, 2, \dots \quad (20)$$

At the same time, by (17) and (18),

$$g(x_k) < \langle \hat{x}^* - x^* + \xi z^*, x_k - \bar{x} \rangle, \quad k = 1, 2, \dots \quad (21)$$

For a fixed k , choose a positive $t < 1$ such that $g(x_k) < t \langle \hat{x}^* - x^* + \xi z^*, x_k - \bar{x} \rangle$ and denote

$$\eta := t \langle \hat{x}^* - x^* + \xi z^*, x_k - \bar{x} \rangle > 0, \quad \lambda := \frac{t}{k} \|\hat{x} - \bar{x}\| > 0.$$

Thus, $g(x_k) < \eta$. By virtue of the Ekeland variational principle [21] (see also [11, Theorem 4.3.1], [19, Theorem 4B.5]) applied to the function $u \mapsto g_+(u) := \max\{g(u), 0\}$, there exists a point $u_k \in X$ such that

$$\|u_k - x_k\| \leq \lambda, \quad (22)$$

$$g_+(u) + \frac{\eta}{\lambda} \|u - u_k\| \geq g_+(u_k) \quad \text{for all } u \in X. \quad (23)$$

By (22) and the definitions of λ and x_k , we have

$$\|u_k - \bar{x}\| \leq \|u_k - x_k\| + \|x_k - \bar{x}\| \leq \lambda + \|x_k - \bar{x}\| < \frac{2}{k} \|\hat{x} - \bar{x}\|. \quad (24)$$

By (19) and (22) and the definitions of x_k , λ and z^* ,

$$\begin{aligned} g(u_k) &\geq \xi \langle z^*, u_k - \bar{x} \rangle = \xi \langle z^*, x_k - \bar{x} \rangle + \xi \langle z^*, u_k - x_k \rangle \\ &\geq \frac{\xi}{k} \|\hat{x} - \bar{x}\| - \xi \lambda = (1-t) \frac{\xi}{k} \|\hat{x} - \bar{x}\| > 0, \end{aligned} \quad (25)$$

and consequently, $g_+(u_k) = g(u_k)$. Thanks to this observation and the fact that $g_+(u) \geq g(u)$ for all $u \in X$, it follows from (23) that u_k is a global minimum of the convex function

$$u \mapsto h(u) := g(u) + \frac{\eta}{\lambda} \|u - u_k\|,$$

and so, $0 \in \partial h(u_k) = \partial g(u_k) + \frac{\eta}{\lambda} \mathbb{B}^*$, i.e.,

$$\begin{aligned} d(0, \partial g(u_k)) &\leq \frac{\eta}{\lambda} = \frac{\langle \hat{x}^* - x^* + \xi z^*, x_k - \bar{x} \rangle}{\frac{1}{k} \|\hat{x} - \bar{x}\|} \\ &= \frac{k \langle \hat{x}^* - x^*, x_k - \bar{x} \rangle}{\|\hat{x} - \bar{x}\|} + \xi \\ &\leq \|\hat{x}^* - x^*\| + \xi < 2\xi. \end{aligned} \quad (26)$$

By (24), we have $u_k \rightarrow \bar{x}$ as $k \rightarrow \infty$. Thanks to (25) and (26), it follows from Theorem 2.1 that $\text{Er} g(\bar{x}) \leq 2\xi$. As $\xi > 0$ can be chosen arbitrarily small, we conclude that $\text{Er}\{\text{Ptb}_l(f, \bar{x}, \varepsilon)\}(\bar{x}) = 0$, which proves (15). \square

Corollary 2.1 *Let $f \in \Gamma_0(X)$, $f(\bar{x}) = 0$, $\varepsilon \geq 0$. Then*

$$\begin{aligned} \varepsilon < \zeta(f, \bar{x}) &\Rightarrow \text{Er}\{\text{Ptb}(f, \bar{x}, \varepsilon)\}(\bar{x}) > 0 \\ &\Rightarrow \text{Er}\{\text{Ptb}_c(f, \bar{x}, \varepsilon)\}(\bar{x}) > 0 \\ &\Rightarrow \text{Er}\{\text{Ptb}_l(f, \bar{x}, \varepsilon)\}(\bar{x}) > 0 \Rightarrow \varepsilon \leq \zeta(f, \bar{x}). \end{aligned}$$

Remark 2.1 Theorem 2.2 strengthens [38, Theorem 8 and Corollary 9] which establishes inequality (14) for any $\zeta(f, \bar{x})$ as well as the first equality in (12) in the case $\zeta(f, \bar{x}) = 0$. The above proof of inequality (14) is more straightforward than that of the corresponding one in [38, Theorem 8 (ii)].

Remark 2.2 Thanks to Theorem 2.2, $\zeta(f, \bar{x})$ can be interpreted as the *radius of error bounds* for a family of (arbitrary or convex or linear) perturbations of f at \bar{x} .

Remark 2.3 The given above proof of the inequality (15) is constructive: for an $\varepsilon > \zeta(f, \bar{x})$ and an arbitrarily small $\xi > 0$, the linear ε -perturbation g of f near \bar{x} satisfying $\text{Er} g(\bar{x}) \leq 2\xi$ is given by (18) where x^* is any element from $\text{bd} \partial f(\bar{x})$ satisfying $\|x^*\| < \varepsilon$ while element \hat{z}^* is fully determined by an arbitrary $\hat{x}^* \notin \partial f(\bar{x})$ with $\|\hat{x}^* - x^*\| < \xi$.

3 Stability of global error bounds

This section deals with the global error bound property for the constraint system (1). The next theorem extracted from [38, Theorem 22] represents a nonlocal analogue of Theorem 2.1.

Theorem 3.1 *Let $f \in \Gamma_0(X)$ and $S_f \neq \emptyset$. Function f has a global error bound provided that one of the following two conditions is satisfied:*

- (i) $\tau(f) := \inf_{f(x) > 0} d(0, \partial f(x)) > 0$;
- (ii) $\zeta(f) := \inf_{f(x) = 0} d(0, \text{bd} \partial f(x)) > 0$.

Moreover, condition (i) is also necessary for f to have a global error bound and

$$\zeta(f) \leq \tau(f) = \operatorname{Er} f. \quad (27)$$

Obviously $\tau(f) \leq \tau(f, x)$ and

$$\zeta(f) = \inf_{x \in S_f^-} \zeta(f, x). \quad (28)$$

The inequality in (27) guarantees, in particular, that $\|x^*\| \geq \zeta(f)$ for any $x^* \in \partial f(x)$ with $f(x) > 0$. Examples 2.1 and 2.2 in Section 2 are also applicable to global error bounds to show that this inequality can be strict. We shall prove in this section, that condition (ii) in Theorem 3.1 corresponds to the existence of a global error bound for a family of functions being small perturbations of f .

Alongside $\zeta(f)$ and $\tau(f)$, we are going to consider a “localized” quantity associated with a point $x \in S_f^-$ (and depending also on numbers $\varepsilon \geq 0$ and $\delta \geq 0$):

$$\tau(f, x, \varepsilon, \delta) := \begin{cases} \inf_{f(u) \geq -\varepsilon \|u-x\| - \delta} d(0, \partial f(u)) & \text{if } 0 \notin \operatorname{int} \partial f(x), \\ d(0, \operatorname{bd} \partial f(x)) & \text{if } 0 \in \operatorname{int} \partial f(x). \end{cases} \quad (29)$$

The two cases in definition (29) are mutually exclusive in the sense that they cannot happen simultaneously for the same function even at different points: if $0 \in \operatorname{int} \partial f(x)$ for some $x \in S_f^-$, then $S_f^- = \{x\}$ and there are no points $x \in S_f^-$ where $0 \notin \operatorname{int} \partial f(x)$. In the second case, we are actually dealing with local error bounds. This case has been added for completeness to ensure that $\tau(f, x, \varepsilon, \delta)$ is defined for all $x \in S_f^-$. For the purposes of the current paper, only the first case is important. Unlike $\zeta(f)$ and $\tau(f)$, when $0 \notin \operatorname{int} \partial f(x)$ for all $x \in S_f^-$ and either $\varepsilon > 0$ or $\delta > 0$, this definition takes into account also certain points u with $f(u) < 0$. Obviously,

$$\tau(f, x, \varepsilon, \delta) \leq \zeta(f)$$

for all $x \in S_f^-$, $\varepsilon \geq 0$ and $\delta \geq 0$, and the equality holds when $\varepsilon = \delta = 0$ or $0 \in \partial f(x)$. Observe that $\tau(f, x, 0, \delta)$ does not depend on x , and $\tau(f, x, \varepsilon, \delta)$ does not depend on ε and δ when $0 \in \partial f(x)$. The function $(\varepsilon, \delta) \mapsto \tau(f, x, \varepsilon, \delta)$ is nonincreasing:

$$\tau(f, x, \varepsilon_2, \delta_2) \leq \tau(f, x, \varepsilon_1, \delta_1)$$

if $0 \leq \varepsilon_1 \leq \varepsilon_2$ and $0 \leq \delta_1 \leq \delta_2$.

The next example illustrates the computation and properties of the localized constant (29).

Example 3.1 Consider the piecewise linear function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by

$$f(x) := \max\{-2x + 2, -x + 1, 2x - 5\} = \begin{cases} -2x + 2 & \text{if } x \leq 1, \\ -x + 1 & \text{if } 1 < x \leq 2, \\ 2x - 5 & \text{if } x > 2. \end{cases}$$

It can be easily verified that $S_f^- = \{1, 2.5\}$, $\zeta(f) = 1$, $\tau(f) = 2$, and the only point u where $0 \in \partial f(u)$ is $u = 2$. When computing (29) with $x = 1$ (and nonnegative ε and δ), we see that $2 \in \{u \mid f(u) \geq -\varepsilon|u-1| - \delta\}$ if and only if $\varepsilon + \delta \geq 1$. Hence,

$$\tau(f, 1, \varepsilon, \delta) = \inf_{f(u) \geq -\varepsilon|u-1| - \delta} d(0, \partial f(u)) = \begin{cases} 1 & \text{if } \varepsilon + \delta < 1, \\ 0 & \text{if } \varepsilon + \delta \geq 1. \end{cases} \quad (30)$$

For small ε and δ ($\varepsilon + \delta < 1$), we have $\tau(f, 1, \varepsilon, \delta) = \zeta(f)$. \triangle

When examining stability issues of global error bounds, estimates of the difference $\zeta(f) - \tau(f, x, \varepsilon, \delta)$ are needed.

Let $S_f \neq \emptyset$ and $\varepsilon \geq 0$. We say that $g : X \rightarrow \mathbb{R}_\infty$ is an ε -perturbation of f if $S_g \neq \emptyset$, $g = f + p$ where $p : X \rightarrow \mathbb{R}$ is convex, and there exist a point $x \in S_f^-$ and a number $\xi \geq 0$ such that

$$\xi + \zeta(f) - \tau(f, x, \xi, |p(x)|) \leq \varepsilon \quad (31)$$

and

$$|p(u) - p(x)| \leq \xi \|u - x\| \quad \text{for all } u \in X. \quad (32)$$

The collection of all ε -perturbations of f will be denoted $\text{Ptb}(f, \varepsilon)$.

Condition (31) implies $\xi \leq \varepsilon$. Conditions (31) and (32) are obviously satisfied with $g \equiv f$, any $x \in S_f^-$ and $\xi = 0$. Hence, $f \in \text{Ptb}(f, \varepsilon)$ for any $\varepsilon \geq 0$. If $\tau(f, x, \varepsilon, \delta) = \zeta(f)$ for some $x \in S_f^-$ and all $\delta \geq 0$ (for instance, if $0 \in \partial f(x)$), then any function p satisfying condition (32) with $\xi = \varepsilon$ defines an ε -perturbation $f + p$ of f .

Function p in (32) enjoys some nice properties. The next lemma is unlikely to be new, cf. e.g., [11, Exercise 4.1.28]. We provide the proof for completeness.

Lemma 3.1 *Suppose $p : X \rightarrow \mathbb{R}$ is convex and satisfies condition (32) with some $\xi \geq 0$. Then p is Lipschitz continuous with constant ξ and, for any $u \in X$ and any $x^* \in \partial p(u)$, it holds $\|x^*\| \leq \xi$.*

Proof Let $u_1, u_2 \in X$, $u_1 \neq u_2$. Denote $\lambda := \|u_1 - u_2\|$ and take an arbitrary $t > 0$. Since,

$$u_1 = \frac{t}{t+\lambda} u_2 + \frac{\lambda}{t+\lambda} \left(u_1 + \frac{t}{\lambda} (u_1 - u_2) \right),$$

using the convexity of p and condition (32), we obtain

$$\begin{aligned} p(u_1) &\leq \frac{t}{t+\lambda} p(u_2) + \frac{\lambda}{t+\lambda} p \left(u_1 + \frac{t}{\lambda} (u_1 - u_2) \right) \\ &\leq \frac{t}{t+\lambda} p(u_2) + \frac{\lambda}{t+\lambda} \left(\xi \left\| u_1 + \frac{t}{\lambda} (u_1 - u_2) - x \right\| + p(x) \right) \\ &\leq \frac{t}{t+\lambda} p(u_2) + \frac{\lambda}{t+\lambda} (\xi (\|u_1\| + t + \|x\|) + p(x)) \\ &= \frac{t}{t+\lambda} p(u_2) + \frac{\lambda}{t+\lambda} (\xi (\|u_1\| + \|x\|) + p(x)) + \frac{\lambda \xi t}{t+\lambda}, \end{aligned}$$

and after passing to the limit as $t \rightarrow \infty$, we conclude that $p(u_1) \leq p(u_2) + \lambda \xi$. The same inequality obviously holds true with u_1 and u_2 reversed. Hence,

$$|p(u_1) - p(u_2)| \leq \lambda \xi = \xi \|u_1 - u_2\|,$$

and consequently, p is Lipschitz continuous with constant ξ . If $u \in X$ and $x^* \in \partial p(u)$, then

$$\langle x^*, x \rangle \leq p(u+x) - p(u) \leq \xi \|x\|$$

for any $x \in X$. Hence, $\|x^*\| \leq \xi$. \square

Observe that the above ‘‘global’’ definition of an ε -perturbation contains a local element: it requires the existence of a point $x \in S_f^-$, and the perturbation is defined by (31) and (32) relative to this point and depends on the number ξ . If $\xi = 0$ in (32), then $p(u) = p(x)$ for all $u \in X$.

Condition (31) imposes a strong restriction on the choice of such a point and a number: for all points $u \in X$ with $f(u) \geq -\xi \|u - x\| - |p(x)|$, the norms of subgradients of f must not be much smaller than $\zeta(f)$. This is also a restriction on the class of functions which admit nontrivial ε -perturbations.

Proposition 3.1 *Let $f \in \Gamma_0(X)$ and $S_f^- \neq \emptyset$. If there exist sequences $\{x_k\} \subset X$ and $\{x_k^*\} \subset X^*$ such that $x_k^* \in \partial f(x_k)$ ($k = 1, 2, \dots$), $\|x_k\| \rightarrow \infty$, $\|x_k^*\| \rightarrow 0$ as $k \rightarrow \infty$ and the sequence $\{f(x_k)\}$ is bounded below, then, for any $\varepsilon \in [0, \zeta(f)]$ and $g \in \text{Ptb}(f, \varepsilon)$, it holds $g = f + \text{const}$.*

Proof Let $\varepsilon \in [0, \zeta(f)]$, $g \in \text{Ptb}(f, \varepsilon)$, and sequences $\{x_k\} \subset X$ and $\{x_k^*\} \subset X^*$ satisfy the conditions of the proposition. Since $g \in \text{Ptb}(f, \varepsilon)$, there are a convex function p , an $x \in S_f^-$ and a $\xi \geq 0$ satisfying (31) and (32). If $\xi > 0$, then, because $\{f(x_k)\}$ is bounded below, it holds $f(x_k) \geq -\xi \|x_k - x\| - |p(x)|$ for all sufficiently large k , and consequently, $\tau(f, x, \xi, |p(x)|) = 0$. This contradicts (31) in view of the assumption that $\varepsilon \leq \zeta(f)$. Hence, $\xi = 0$, and it follows from (32) that $p(u) = p(x)$ for all $u \in X$. \square

Remark 3.1 The conclusion of Proposition 3.1 remains valid if condition $\|x_k^*\| \rightarrow 0$ as $k \rightarrow \infty$ is replaced by a weaker one: $\lim_{k \rightarrow \infty} \|x_k^*\| \leq \zeta(f) - \varepsilon$. The assumption that the sequence $\{f(x_k)\}$ is bounded below can be relaxed too: it is sufficient to assume that $\min\{f(x_k), 0\}/\|x_k\| \rightarrow 0$ as $k \rightarrow \infty$.

The conditions of Proposition 3.1 are satisfied, for instance, when the set $S := \text{argmin} f$ is nonempty and unbounded. Then, obviously, $0 \in \partial f(x)$ for all $x \in S$, and any sequence $\{x_k\} \subset S$ with $\|x_k\| \rightarrow \infty$ as $k \rightarrow \infty$ will do the job. Below are some specific examples.

Example 3.2 1. $f(x) \equiv 0$, $x \in \mathbb{R}$ (cf. Example 2.1). Then $S_f = S_f^- = \text{argmin} f = \mathbb{R}$, $\tau(f, x, \varepsilon, \delta) = \zeta(f) = 0$ for all $x \in \mathbb{R}$, $\varepsilon \geq 0$, $\delta \geq 0$. Thus, $\text{Ptb}(f, 0) = \{g \equiv c \mid c \leq 0\}$. If $\varepsilon > 0$, then $g \in \text{Ptb}(f, \varepsilon)$ if and only if $S_g \neq \emptyset$ and

$$|g(u) - g(x)| \leq \varepsilon |u - x| \quad \text{for some } x \in \mathbb{R} \text{ and all } u \in \mathbb{R}.$$

2. $f(x) := \max\{x, -1\}$, $x \in \mathbb{R}$. Then $S_f =]-\infty, 0]$, $S_f^- = \{0\}$, $\operatorname{argmin} f =]-\infty, -1]$, $\zeta(f) = 1$, and

$$\tau(f, 0, \xi, \delta) = \begin{cases} 0 & \text{if } \xi > 0, \delta \geq 0 \text{ or } \xi = 0, \delta \geq 1, \\ 1 & \text{if } \xi = 0, 0 \leq \delta < 1. \end{cases}$$

When $0 \leq \varepsilon < 1$, then condition (31) with $x = 0$ is satisfied only when $\xi = 0$ and $|g(0)| < 1$. Hence, $\operatorname{Ptb}(f, \varepsilon) = \{f + c \mid |c| < 1\}$.

3. $f : \mathbb{R} \rightarrow \mathbb{R}$ is defined by

$$f(x) := \begin{cases} -2x & \text{if } x \leq 0, \\ -x & \text{if } x \in]0, 1], \\ -1 - \frac{1}{2}(x-1) & \text{if } x \in]1, 2], \\ \dots & \dots \\ -1 - \frac{1}{2} - \dots - \frac{1}{2^{n-1}} - \frac{1}{2^n}(x-n) & \text{if } x \in]n, n+1], \\ \dots & \dots \end{cases}$$

Then $S_f = [0, \infty[$, $S_f^- = \{0\}$, $\zeta(f) = 1$, and $\inf f = -2$. Observe that in this example $\operatorname{argmin} f = \emptyset$ and the graph has no horizontal parts. Nevertheless, it is easy to construct sequences satisfying the assumptions of Proposition 3.1. To determine the exact representation of the collection of ε -perturbations, similar to the previous example, $\tau(f, 0, \xi, \delta)$ needs to be computed:

$$\tau(f, 0, \xi, \delta) = \begin{cases} 0 & \text{if } \xi > 0, \delta \geq 0 \text{ or } \xi = 0, \delta \geq 2, \\ 1 & \text{if } \xi = 0, 0 \leq \delta < 1, \\ \frac{1}{2} & \text{if } \xi = 0, 1 \leq \delta < \frac{3}{2}, \\ \dots & \dots \\ \frac{1}{2^n} & \text{if } \xi = 0, 2 - \frac{1}{2^{n-1}} \leq \delta < 2 - \frac{1}{2^n}, \\ \dots & \dots \end{cases}$$

When $0 \leq \varepsilon < 1$, then condition (31) with $x = 0$ is satisfied only when $\xi = 0$ and $\tau(f, 0, 0, |g(0)|) \geq 1 - \varepsilon$. In particular, if $0 \leq \varepsilon < 1/2$, then it must hold $|g(0)| < 1$. Hence, in this case, $\operatorname{Ptb}(f, \varepsilon) = \{f + c \mid |c| < 1\}$. \triangle

In [38], when investigating stability of global error bounds, a special *asymptotic qualification condition* (\mathcal{AQL}) (generalizing the corresponding finite dimensional condition from [45]) was imposed on function f :

- (\mathcal{AQL}) $\liminf_{k \rightarrow \infty} \|x_k^*\| \geq \zeta(f)$ for any sequences $x_k \in X$ with $f(x_k) < 0$ and $x_k^* \in \partial f(x_k)$, $k = 1, 2, \dots$, satisfying the following:
- (a) either the sequence $\{x_k\}$ is bounded and $\lim_{k \rightarrow \infty} f(x_k) = 0$,
 - (b) or $\lim_{k \rightarrow \infty} \|x_k\| = \infty$ and $\lim_{k \rightarrow \infty} f(x_k)/\|x_k\| = 0$.

If $\inf_{x \in X} f(x) < 0$, then, under this condition, thanks to [38, Proposition 23], for any point $x \in S_f^-$ and any $\varepsilon > 0$, one can find a positive number ξ such that condition (31) is satisfied as long as $|p(x)| \leq \xi$. Due to this observation and Lemma 3.1, under the (\mathcal{AQL}) the collection $\operatorname{Ptb}(f, \varepsilon)$ of ε -perturbations defined above is larger than the corresponding one considered in [38].

The next example illustrates the computation of ε -perturbations.

Example 3.3 Consider again the piecewise linear function $f : \mathbb{R} \rightarrow \mathbb{R}$ examined in Example 3.1. We target small perturbations of this function related to the point $x = 1 \in S_f^-$. Let $\varepsilon \in [0, 1[$. Since $\zeta(f) = 1$ and $\tau(f, 1, \varepsilon, \delta)$ equals either 0 or 1 (cf. (30)), condition (31) can only be satisfied when $\tau(f, 1, \xi, |g(1)|) = 1$, i.e., when $0 \leq \xi \leq \varepsilon$ and $\xi + |g(1)| < 1$. Condition (32) takes the following form:

$$|p(u) - p(1)| \leq \xi |u - 1| \quad \text{for all } u \in \mathbb{R}.$$

In particular, when $\varepsilon = 0$, all 0-perturbations are of the form $f + c$ with $|c| < 1$. \triangle

If condition (31) in the definition of ε -perturbation is dropped we will talk about weak ε -perturbations: $g : X \rightarrow \mathbb{R}_\infty$ is a *weak ε -perturbation* of f if $S_g \neq \emptyset$, $g = f + p$ where $p : X \rightarrow \mathbb{R}$ is convex, and there exists a point $x \in S_f^-$ such that

$$|p(u) - p(x)| \leq \varepsilon \|u - x\| \quad \text{for all } u \in X.$$

In this case, one obviously has $g - p(x) \in \text{Ptb}_c(f, x, \varepsilon)$. The collection of all weak ε -perturbations of f will be denoted $\text{Ptb}^w(f, \varepsilon)$. Obviously $\text{Ptb}(f, \varepsilon) \subset \text{Ptb}^w(f, \varepsilon)$.

The following subsets of $\text{Ptb}(f, \varepsilon)$ and $\text{Ptb}^w(f, \varepsilon)$ corresponding to linear ε -perturbations of f can be of interest:

$$\begin{aligned} \text{Ptb}_l(f, \varepsilon) &:= \{g \mid g(u) = f(u) + \langle x^*, u - x \rangle \ (u \in X), \\ &\quad x \in S_f^-, \xi \geq 0, x^* \in \xi \mathbb{B}_{X^*}^*, \xi + \zeta(f) - \tau(f, x, \xi, 0) \leq \varepsilon\}, \\ \text{Ptb}_l^w(f, \varepsilon) &:= \{g \mid g(u) = f(u) + \langle x^*, u - x \rangle \ (u \in X), x \in S_f^-, x^* \in \varepsilon \mathbb{B}_{X^*}^*\}. \end{aligned}$$

Obviously,

$$\text{Ptb}_l(f, \varepsilon) \subset \text{Ptb}(f, \varepsilon), \quad \text{Ptb}_l^w(f, \varepsilon) \subset \text{Ptb}^w(f, \varepsilon), \quad (33)$$

$$\text{Ptb}_l^w(f, \varepsilon) = \bigcup_{x \in S_f^-} \text{Ptb}_l(f, x, \varepsilon). \quad (34)$$

Given a function $f \in I_0(X)$ and a number $\varepsilon \geq 0$, denote

$$\begin{aligned} \text{Er}\{\text{Ptb}(f, \varepsilon)\} &:= \inf_{g \in \text{Ptb}(f, \varepsilon)} \text{Er } g, \\ \text{Er}\{\text{Ptb}_l(f, \varepsilon)\} &:= \inf_{g \in \text{Ptb}_l(f, \varepsilon)} \text{Er } g, \\ \text{Er}\{\text{Ptb}^w(f, \varepsilon)\} &:= \inf_{g \in \text{Ptb}^w(f, \varepsilon)} \text{Er } g, \\ \text{Er}\{\text{Ptb}_l^w(f, \varepsilon)\} &:= \inf_{g \in \text{Ptb}_l^w(f, \varepsilon)} \text{Er } g. \end{aligned}$$

These numbers characterize the error bound property for families of ε -perturbations of f . Thanks to (33) and (34), it holds

$$\text{Er}\{\text{Ptb}(f, \varepsilon)\} \leq \text{Er}\{\text{Ptb}_l(f, \varepsilon)\} \leq \text{Er } f, \quad (35)$$

$$\text{Er}\{\text{Ptb}^w(f, \varepsilon)\} \leq \text{Er}\{\text{Ptb}_l^w(f, \varepsilon)\} = \inf_{x \in S_f^-} \text{Er}\{\text{Ptb}_l(f, x, \varepsilon)\}(x) \quad (36)$$

for any $\varepsilon \geq 0$.

Theorem 3.2 *Let $f \in \Gamma_0(X)$, $S_f \neq \emptyset$. Then*

$$\begin{aligned} & \inf\{\varepsilon > 0 \mid \text{Er}\{\text{Ptb}^w(f, \varepsilon)\} = 0\} \\ & \leq \inf\{\varepsilon > 0 \mid \text{Er}\{\text{Ptb}_l^w(f, \varepsilon)\} = 0\} \leq \zeta(f) \leq \inf\{\varepsilon > 0 \mid \text{Er}\{\text{Ptb}(f, \varepsilon)\} = 0\} \\ & \leq \inf\{\varepsilon > 0 \mid \text{Er}\{\text{Ptb}_l(f, \varepsilon)\} = 0\}. \end{aligned} \quad (37)$$

Proof Thanks to (35) and (36), we always have

$$\inf\{\varepsilon > 0 \mid \text{Er}\{\text{Ptb}(f, \varepsilon)\} = 0\} \leq \inf\{\varepsilon > 0 \mid \text{Er}\{\text{Ptb}_l(f, \varepsilon)\} = 0\}, \quad (38)$$

$$\inf\{\varepsilon > 0 \mid \text{Er}\{\text{Ptb}^w(f, \varepsilon)\} = 0\} \leq \inf\{\varepsilon > 0 \mid \text{Er}\{\text{Ptb}_l^w(f, \varepsilon)\} = 0\}. \quad (39)$$

We first show that

$$\zeta(f) \leq \inf\{\varepsilon > 0 \mid \text{Er}\{\text{Ptb}(f, \varepsilon)\} = 0\}. \quad (40)$$

If $\zeta(f) = 0$, the inequality holds true trivially. Let $0 < \varepsilon < \zeta(f)$ and $g \in \text{Ptb}(f, \varepsilon)$, i.e. conditions (31) and (32) are satisfied for some convex function $p : X \rightarrow \mathbb{R}$, a point $x \in S_f^-$ and a number $\xi \geq 0$. Two cases are possible.

If $0 \in \text{int } \partial f(\bar{x})$ for some $\bar{x} \in S_f^-$, then $S_f = \{\bar{x}\}$, $x = \bar{x}$, $\zeta(f) = \zeta(f, \bar{x})$, and

$$\zeta(f)\mathbb{B}^* \subset \partial f(\bar{x}). \quad (41)$$

We need to show that

$$\partial f(\bar{x}) \subset \partial g(\bar{x}) + \xi\mathbb{B}^*. \quad (42)$$

Let $x^* \in \partial f(\bar{x})$, i.e.,

$$f(u) - f(\bar{x}) - \langle x^*, u - \bar{x} \rangle \geq 0 \quad \text{for all } u \in X.$$

At the same time, by (32),

$$p(u) - p(\bar{x}) \geq -\xi\|u - \bar{x}\| \quad \text{for all } u \in X. \quad (43)$$

Adding the last two inequalities, we obtain

$$g(u) - g(\bar{x}) + \xi\|u - \bar{x}\| - \langle x^*, u - \bar{x} \rangle \geq 0 \quad \text{for all } u \in X,$$

i.e., x^* is a subgradient at \bar{x} of the sum of two convex functions: g and $u \mapsto \varphi(u) := \xi\|u - \bar{x}\|$. Function φ is Lipschitz continuous and $\partial\varphi(\bar{x}) = \xi\mathbb{B}^*$. Hence, $x^* \in \partial g(\bar{x}) + \xi\mathbb{B}^*$, which proves (42). Since $\xi \leq \varepsilon$, it follows from (41) and (42) that

$$\zeta(f)\mathbb{B}^* \subset \partial g(\bar{x}) + \varepsilon\mathbb{B}^*.$$

Then, thanks to the *Rådström cancellation principle* (cf. Beer [9, 7.4.1]),

$$(\zeta(f) - \varepsilon)\mathbb{B}^* \subset \partial g(\bar{x}),$$

and consequently, \bar{x} is a minimum point of g . Since $S_g \neq \emptyset$, we have $\bar{x} \in S_g = S_g^-$. We next show that $\|x^*\| \geq \zeta(f) - \varepsilon$ as long as $x^* \in \partial g(u)$ for some $u \neq \bar{x}$. Indeed, for any $u \neq \bar{x}$, $x^* \in \partial g(u)$ and $u^* \in \partial g(x)$, one has

$$g(u) \geq g(\bar{x}) + \langle u^*, u - \bar{x} \rangle, \quad g(\bar{x}) \geq g(u) + \langle x^*, \bar{x} - u \rangle.$$

Hence,

$$\langle x^*, u - \bar{x} \rangle \geq \langle u^*, u - \bar{x} \rangle,$$

and consequently, taking supremum in the right-hand side over all $u^* \in (\zeta(f) - \varepsilon)\mathbb{B}^*$,

$$\langle x^*, u - \bar{x} \rangle \geq (\zeta(f) - \varepsilon)\|u - \bar{x}\|.$$

Since $u \neq \bar{x}$, it follows immediately from the last inequality that $\|x^*\| \geq \zeta(f) - \varepsilon$. Hence, $\text{Er}\{\text{Ptb}(f, \varepsilon)\} \geq \zeta(f) - \varepsilon > 0$ which yields (40).

$0 \notin \text{int } \partial f(x)$ for all $x \in S_f^-$. If $g(u) > 0$, then, in view of (32), $f(u) > -\xi\|u - x\| - |p(x)|$ and, by (31), $\|u^*\| \geq \zeta(f) + \xi - \varepsilon$ for all $u^* \in \partial f(u)$. If $x^* \in \partial g(u)$, then, in view of Lemma 3.1, $x^* \in \partial f(u) + \partial p(u) \subset \partial f(u) + \xi\mathbb{B}^*$, and consequently, $\|x^*\| \geq \zeta(f) - \varepsilon$. Hence, $\text{Er}\{\text{Ptb}(f, \varepsilon)\} \geq \zeta(f) - \varepsilon > 0$, and consequently, (40) holds true. Together with (38), this proves the second group of inequalities in (37).

Next we show that

$$\inf\{\varepsilon > 0 \mid \text{Er}\{\text{Ptb}_l^w(f, \varepsilon)\} = 0\} \leq \zeta(f). \quad (44)$$

If $\zeta(f) = \infty$, the inequality holds trivially. Let $\zeta(f) < \varepsilon < \infty$. By (28), there is an $x \in S_f^-$ such that $\zeta(f, x) < \varepsilon$. By Theorem 2.2, $\text{Er}\{\text{Ptb}_l(f, x, \varepsilon)\}(x) = 0$. Finally, by the equality in (36), $\text{Er}\{\text{Ptb}_l^w(f, \varepsilon)\} = 0$, and consequently, (44) holds true. Together with (39), this proves the first group of inequalities in (37). \square

Corollary 3.1 *Let $f \in \Gamma_0(X)$, $S_f \neq \emptyset$, $\varepsilon \geq 0$. Then*

$$\begin{aligned} \text{Er}\{\text{Ptb}^w(f, \varepsilon)\} > 0 &\Rightarrow \text{Er}\{\text{Ptb}_l^w(f, \varepsilon)\} > 0 \Rightarrow \varepsilon \leq \zeta(f), \\ \varepsilon < \zeta(f) &\Rightarrow \text{Er}\{\text{Ptb}(f, \varepsilon)\} > 0 \Rightarrow \text{Er}\{\text{Ptb}_l(f, \varepsilon)\} > 0. \end{aligned}$$

Remark 3.2 Theorem 3.2 strengthens [38, Theorem 25]. Nonnegative number $\zeta(f)$ provides an estimate for the *radius of error bounds* for a family of (arbitrary or convex and in finite dimensions also linear) perturbations of f .

Remark 3.3 On different stages of the above proof of Theorem 3.2, different features of the definition (31), (32) of an ε -perturbation were used. Accordingly, the statement of the theorem can be strengthened by cutting it into pieces and, for each piece, assuming exactly those features of the definition of an ε -perturbation, that are needed there.

In the proof of the inequality (40) when $0 \in \text{int } \partial f(\bar{x})$ for some $\bar{x} \in S_f^-$, the one-sided estimate (43) was used instead of the ‘full’ inequality (32) in the definition of an ε -perturbation while the inequality (31) was not needed at all. Similarly, in the proof of the same inequality (40) in the case when $0 \notin \text{int } \partial f(x)$ for all $x \in S_f^-$, the opposite one-sided estimate

$$p(u) - p(\bar{x}) \leq \xi\|u - \bar{x}\| \quad \text{for all } u \in X \quad (45)$$

was used.

In the proof of the inequality (44) we worked with ‘weak’ ε -perturbations and, again, only the one-sided estimate (45) was used.

4 Concluding remarks and perspectives

The main results of this article establish the exact formula for the radius of local error bounds (Theorem 2.2) and estimates of global error bounds (Theorem 3.2) for families of (arbitrary, convex and linear) perturbations of a convex inequality system. The next natural step would be to attack the stability of error bounds for inequality systems defined by non-convex functions. The recent article by Zheng & Wei [61] suggests some important classes of such systems to be studied as a starting point.

References

1. Auslender, A., Crouzeix, J.P.: Global regularity theorems. *Math. Oper. Res.* **13**(2), 243–253 (1988). DOI 10.1287/moor.13.2.243
2. Azé, D.: A survey on error bounds for lower semicontinuous functions. In: Proceedings of 2003 MODE-SMAI Conference, *ESAIM Proc.*, vol. 13, pp. 1–17. EDP Sci., Les Ulis (2003)
3. Azé, D.: A unified theory for metric regularity of multifunctions. *J. Convex Anal.* **13**(2), 225–252 (2006)
4. Azé, D., Corvellec, J.N.: Characterizations of error bounds for lower semicontinuous functions on metric spaces. *ESAIM Control Optim. Calc. Var.* **10**(3), 409–425 (2004)
5. Bauschke, H.H., Borwein, J.M.: On projection algorithms for solving convex feasibility problems. *SIAM Rev.* **38**(3), 367–426 (1996). DOI 10.1137/S0036144593251710
6. Beck, A., Teboulle, M.: Convergence rate analysis and error bounds for projection algorithms in convex feasibility problems. *Optim. Methods Softw.* **18**(4), 377–394 (2003). DOI 10.1080/10556780310001604977. The Second Japanese-Sino Optimization Meeting, Part II (Kyoto, 2002)
7. Bednarczuk, E.M., Kruger, A.Y.: Error bounds for vector-valued functions: necessary and sufficient conditions. *Nonlinear Anal.* **75**(3), 1124–1140 (2012). DOI 10.1016/j.na.2011.05.098
8. Bednarczuk, E.M., Kruger, A.Y.: Error bounds for vector-valued functions on metric spaces. *Vietnam J. Math.* **40**(2-3), 165–180 (2012)
9. Beer, G.: Topologies on Closed and Closed Convex Sets, *Mathematics and its Applications*, vol. 268. Kluwer Academic Publishers Group, Dordrecht (1993). DOI 10.1007/978-94-015-8149-3
10. Bolte, J., Nguyen, T.P., Peypouquet, J., Suter, B.W.: From error bounds to the complexity of first order descent methods for convex functions. Preprint, ArXiv **1510.08234** (2015)
11. Borwein, J.M., Vanderwerff, J.D.: Convex Functions: Constructions, Characterizations and Counterexamples, *Encyclopedia of Mathematics and its Applications*, vol. 109. Cambridge University Press, Cambridge (2010). DOI 10.1017/CBO9781139087322
12. Burke, J.V., Deng, S.: Weak sharp minima revisited. II. Application to linear regularity and error bounds. *Math. Program., Ser. B* **104**(2-3), 235–261 (2005)
13. Cánovas, M.J., Henrion, R., López, M.A., Parra, J.: Outer limit of subdifferentials and calmness moduli in linear and nonlinear programming. *J. Optim. Theory Appl.* pp. 1–28 (2015). DOI 10.1007/s10957-015-0793-x
14. Cánovas, M.J., Kruger, A.Y., López, M.A., Parra, J., Théra, M.A.: Calmness modulus of linear semi-infinite programs. *SIAM J. Optim.* **24**(1), 29–48 (2014)
15. Censor, Y.: Iterative methods for the convex feasibility problem. In: Convexity and Graph Theory (Jerusalem, 1981), *North-Holland Math. Stud.*, vol. 87, pp. 83–91. North-Holland, Amsterdam (1984). DOI 10.1016/S0304-0208(08)72812-3
16. Combettes, P.L.: The convex feasibility problem in image recovery. In: P. Hawkes (ed.) *Advances in Imaging and Electron Physics*, vol. 95, pp. 155–270. Academic Press, New York (1996)
17. Cornejo, O., Jourani, A., Zălinescu, C.: Conditioning and upper-Lipschitz inverse subdifferentials in nonsmooth optimization problems. *J. Optim. Theory Appl.* **95**(1), 127–148 (1997). DOI 10.1023/A:1022687412779
18. Corvellec, J.N., Motreanu, V.V.: Nonlinear error bounds for lower semicontinuous functions on metric spaces. *Math. Program., Ser. A* **114**(2), 291–319 (2008)
19. Dontchev, A.L., Rockafellar, R.T.: *Implicit Functions and Solution Mappings. A View from Variational Analysis*, 2 edn. Springer Series in Operations Research and Financial Engineering. Springer, New York (2014)

20. Drusvyatskiy, D., Ioffe, A.D., Lewis, A.S.: Transversality and alternating projections for nonconvex sets. *Found. Comput. Math.* **15**(6), 1637–1651 (2015). DOI 10.1007/s10208-015-9279-3
21. Ekeland, I.: On the variational principle. *J. Math. Anal. Appl.* **47**, 324–353 (1974)
22. Fabian, M.J., Henrion, R., Kruger, A.Y., Outrata, J.V.: Error bounds: necessary and sufficient conditions. *Set-Valued Var. Anal.* **18**(2), 121–149 (2010)
23. Fabian, M.J., Henrion, R., Kruger, A.Y., Outrata, J.V.: About error bounds in metric spaces. In: D. Klatte, H.J. Lüthi, K. Schmedders (eds.) *Operations Research Proceedings 2011. Selected papers of the Int. Conf. Operations Research (OR 2011), August 30 – September 2, 2011, Zurich, Switzerland*, pp. 33–38. Springer-Verlag, Berlin (2012)
24. Gfrerer, H.: First order and second order characterizations of metric subregularity and calmness of constraint set mappings. *SIAM J. Optim.* **21**(4), 1439–1474 (2011)
25. Henrion, R., Jourani, A.: Subdifferential conditions for calmness of convex constraints. *SIAM J. Optim.* **13**(2), 520–534 (2002)
26. Henrion, R., Outrata, J.V.: A subdifferential condition for calmness of multifunctions. *J. Math. Anal. Appl.* **258**(1), 110–130 (2001)
27. Hesse, R., Luke, D.R.: Nonconvex notions of regularity and convergence of fundamental algorithms for feasibility problems. *SIAM J. Optim.* **23**(4), 2397–2419 (2013). DOI 10.1137/120902653
28. Hoffman, A.J.: On approximate solutions of systems of linear inequalities. *J. Research Nat. Bur. Standards* **49**, 263–265 (1952)
29. Huang, L.R., Ng, K.F.: On first- and second-order conditions for error bounds. *SIAM J. Optim.* **14**(4), 1057–1073 (2004). DOI 10.1137/S1052623401390549
30. Ioffe, A.D.: Metric regularity and subdifferential calculus. *Russian Math. Surveys* **55**, 501–558 (2000)
31. Ioffe, A.D.: Metric regularity. Theory and applications – a survey. Preprint, arXiv: **1505.07920** (2015)
32. Ioffe, A.D., Outrata, J.V.: On metric and calmness qualification conditions in subdifferential calculus. *Set-Valued Anal.* **16**(2-3), 199–227 (2008)
33. Jourani, A.: Hoffman’s error bound, local controllability, and sensitivity analysis. *SIAM J. Control Optim.* **38**(3), 947–970 (2000)
34. Klatte, D., Li, W.: Asymptotic constraint qualifications and global error bounds for convex inequalities. *Math. Program., Ser. A* **84**(1), 137–160 (1999)
35. Kruger, A.Y.: Error bounds and Hölder metric subregularity. *Set-Valued Var. Anal.* **23**(4), 705–736 (2015). DOI 10.1007/s11228-015-0330-y
36. Kruger, A.Y.: Error bounds and metric subregularity. *Optimization* **64**(1), 49–79 (2015). DOI 10.1080/02331934.2014.938074
37. Kruger, A.Y.: Nonlinear metric subregularity. *J. Optim. Theory Appl.* pp. 1–36 (2015). DOI 10.1007/s10957-015-0807-8
38. Kruger, A.Y., Ngai, H.V., Théra, M.: Stability of error bounds for convex constraint systems in Banach spaces. *SIAM J. Optim.* **20**(6), 3280–3296 (2010). DOI 10.1137/100782206
39. Kruger, Y., Thao, N.H.: Regularity of collections of sets and convergence of inexact alternating projections. *J. Convex Anal.* **23**(1) (2016)
40. Lewis, A.S., Luke, D.R., Malick, J.: Local linear convergence for alternating and averaged nonconvex projections. *Found. Comput. Math.* **9**(4), 485–513 (2009). DOI 10.1007/s10208-008-9036-y
41. Lewis, A.S., Pang, J.S.: Error bounds for convex inequality systems. In: *Generalized Convexity, Generalized Monotonicity: Recent Results* (Luminy, 1996), pp. 75–110. Kluwer Acad. Publ., Dordrecht (1998)
42. Mangasarian, O.L.: A condition number for differentiable convex inequalities. *Math. Oper. Res.* **10**(2), 175–179 (1985). DOI 10.1287/moor.10.2.175
43. Meng, K.W., Yang, X.Q.: Equivalent conditions for local error bounds. *Set-Valued Var. Anal.* **20**(4), 617–636 (2012)
44. Ng, K.F., Zheng, X.Y.: Error bounds for lower semicontinuous functions in normed spaces. *SIAM J. Optim.* **12**(1), 1–17 (2001)
45. Ngai, H.V., Kruger, A.Y., Théra, M.: Stability of error bounds for semi-infinite convex constraint systems. *SIAM J. Optim.* **20**(4), 2080–2096 (2010). DOI 10.1137/090767819
46. Ngai, H.V., Théra, M.: Error bounds and implicit multifunction theorem in smooth Banach spaces and applications to optimization. *Set-Valued Anal.* **12**(1-2), 195–223 (2004). DOI 10.1023/B:SVAN.0000023396.58424.98
47. Ngai, H.V., Théra, M.: Error bounds for convex differentiable inequality systems in Banach spaces. *Math. Program., Ser. B* **104**(2-3), 465–482 (2005)
48. Ngai, H.V., Théra, M.: Error bounds in metric spaces and application to the perturbation stability of metric regularity. *SIAM J. Optim.* **19**(1), 1–20 (2008). DOI 10.1137/060675721

49. Ngai, H.V., Théra, M.: Error bounds for systems of lower semicontinuous functions in Asplund spaces. *Math. Program., Ser. B* **116**(1-2), 397–427 (2009)
50. Ngai, H.V., Tron, N.H., Théra, M.: Implicit multifunction theorems in complete metric spaces. *Math. Program.* **139**(1-2, Ser. B), 301–326 (2013). DOI 10.1007/s10107-013-0673-9
51. Ngai, H.V., Tron, N.H., Théra, M.: Metric regularity of the sum of multifunctions and applications. *J. Optim. Theory Appl.* **160**(2), 355–390 (2014). DOI 10.1007/s10957-013-0385-6
52. Pang, J.S.: Error bounds in mathematical programming. *Math. Programming, Ser. B* **79**(1-3), 299–332 (1997)
53. Penot, J.P.: Error bounds, calmness and their applications in nonsmooth analysis. In: *Nonlinear analysis and optimization II. Optimization, Contemp. Math.*, vol. 514, pp. 225–247. Amer. Math. Soc., Providence, RI (2010). DOI 10.1090/conm/514/10110
54. Robinson, S.M.: An application of error bounds for convex programming in a linear space. *SIAM J. Control* **13**, 271–273 (1975)
55. Rockafellar, R.T., Wets, R.J.B.: *Variational Analysis*. Springer, Berlin (1998)
56. Rosenbloom, P.C.: Quelques classes de problèmes extrémaux. *Bulletin de la Société Mathématique de France* **79**, 1–58 (1951). URL <http://eudml.org/doc/86848>
57. Wu, Z., Ye, J.J.: Sufficient conditions for error bounds. *SIAM J. Optim.* **12**(2), 421–435 (2001/02)
58. Zălinescu, C.: Sharp estimates for Hoffman’s constant for systems of linear inequalities and equalities. *SIAM J. Optim.* **14**(2), 517–533 (2003). DOI 10.1137/S1052623402403505
59. Zheng, X.Y., Ng, K.F.: Metric subregularity and calmness for nonconvex generalized equations in Banach spaces. *SIAM J. Optim.* **20**(5), 2119–2136 (2010). DOI 10.1137/090772174
60. Zheng, X.Y., Ng, K.F.: Metric subregularity for proximal generalized equations in Hilbert spaces. *Nonlinear Anal.* **75**(3), 1686–1699 (2012). DOI 10.1016/j.na.2011.07.004
61. Zheng, X.Y., Wei, Z.: Perturbation analysis of error bounds for quasi-subsmooth inequalities and semi-infinite constraint systems. *SIAM J. Optim.* **22**(1), 41–65 (2012). DOI 10.1137/100806199