

# Matroid Optimisation Problems with Nested Non-linear Monomials in the Objective Function

Anja Fischer\*      Frank Fischer†      S. Thomas McCormick‡

14th March 2016

## Abstract

Recently, Buchheim and Klein [4] suggested to study polynomial-time solvable optimisation problems with linear objective functions combined with exactly one additional quadratic monomial. They concentrated on special quadratic spanning tree or forest problems. We extend their results to general *matroid* optimisation problems with a *set of nested monomials* in the objective function. The monomials are linearised and we study the corresponding polytope with the aim to better understand the structure of polytopes arising from such linearisations and to provide strengthened cutting planes as well as separation algorithms for linearisations of matroid optimisation problems with general polynomial objective function. Extending results by Edmonds [7] for the matroid polytope we present a complete description for the linearised polytope. Indeed, apart from the standard linearisation one needs appropriately strengthened rank inequalities satisfying certain non-separability conditions. The separation problem of these extended rank inequalities reduces to a submodular function minimisation problem. In the case of exactly one additional non-linear monomial we completely characterise the facetial structure of the associated polytope.

**Keywords:** polyhedral combinatorics, matroid, matroid polytope, linearisation, polynomial optimisation, separation

**MSC2010:** 90C57, 90C27

## 1 Introduction

Some combinatorial optimisation problems are well-understood in the sense that they can be solved in polynomial time and the complete description of the associated polytope is known. Examples are the spanning tree problem, the optimisation over matroids or over the intersection of two matroids as well as the matching problem, see, e. g., [7, 8, 23]. Optimisation problems with general polynomial objective functions are often solved by introducing new variables for each monomial, see, e. g., [10, 12, 13, 16, 25] and considering the associated linear problem, for the quadratic unconstrained case see, e. g., [21]. Approaches for solving some kinds of non-linear matroid optimization problem can be found, e. g., in [1, 17, 19]. In this paper we study a simpler but related problem with the aim to better understand the structure of polytopes arising from linearisations. Instead of considering an arbitrary polynomial objective function, we restrict to linear terms and a set of nested monomials or in a special case one additional monomial of arbitrary degree. We will show that studying the polynomially-solvable matroid optimisation problem, the corresponding problem with this particular polynomial objective function is solvable in polynomial time, too (see Remark 2). But although the well-known “separation equals

---

\*University of Göttingen, Institute for Numerical and Applied Mathematics, [anja.fischer@mathematik.uni-goettingen.de](mailto:anja.fischer@mathematik.uni-goettingen.de)

†University of Kassel, Institute for Mathematics, [frank.fischer@uni-kassel.de](mailto:frank.fischer@uni-kassel.de)

‡Sauder School of Business, University of British Columbia, [tom.mccormick@sauder.ubc.ca](mailto:tom.mccormick@sauder.ubc.ca)

optimisation” result [15] then implies that the separation problem for our problem can be solved in polynomial time, we want to find the exact complete description of the associated linearised polytope and combinatorial separation algorithms for the newly derived cutting planes.

Our main motivation is that this study can also be helpful for (matroid) problems with general linearised polynomial objective functions because the strengthened inequalities for the case with linearised nested monomials usually remain valid for the general case and the respective separation algorithms can be used as well (but in most cases the facet defining inequalities of the case with few linearised monomials will not remain facet defining for the general case). One advantage of our approach in comparison to general linearisation techniques, like the procedure of Sherali-Adams [24, 25], is that we strengthen the constraints of the original linear problem without introducing new variables for non-linear monomials that are not explicitly contained in the objective function.

Focusing on exactly one additional non-linear monomial has first been suggested by Buchheim and Klein [3, 4] for the quadratic case. Results for the spanning tree and forest problem with exactly one additional quadratic monomial can be found in [3, 4, 9]. Buchheim and Klein also showed that the use of strengthened cutting planes can reduce the number of nodes of a branch-and-cut tree in the solution of general quadratic spanning tree problems significantly. In this paper we extend the results for the quadratic forest problems to general *matroid* optimisation problems with one monomial of *arbitrary* degree and to a *set of nested monomials* to be described below.

In the following we will describe our setting in more detail. We start with a short review of matroids and their associated polytopes [20, 26]. In 1970 Edmonds [7] presented a complete description of the polytope  $P_M$  associated with a matroid  $M = (E, \mathcal{J})$  (the convex hull over all incidence vectors of independent sets) with finite set  $E$  and a family  $\mathcal{J}$  of subsets of  $E$  satisfying the three matroid properties

$$(M1) \quad \emptyset \in \mathcal{J},$$

$$(M2) \quad T \in \mathcal{J}, S \subset T \Rightarrow S \in \mathcal{J},$$

$$(M3) \quad S, T \in \mathcal{J}, |S| < |T|: \exists e \in T \setminus S \text{ with } S \cup \{e\} \in \mathcal{J}.$$

A pair  $M = (E, \mathcal{J})$  satisfying only (M1) and (M2) is called *independence system*. We denote by  $\mathcal{P}(X)$  the power set of a finite set  $X$ . Then the rank function  $r: \mathcal{P}(E) \rightarrow \mathbb{N}_0$  of an independence system  $M = (E, \mathcal{J})$  is defined as  $r(T) = \max\{|B|: B \subseteq T, B \in \mathcal{J}\}$ . For some  $T \subseteq E$  a set  $B$  with  $B \subseteq T, B \in \mathcal{J}$  and  $r(T) = |B|$  is called *basis* (of  $T$ ). It is well known that the rank function of a matroid has the following properties (see, e. g., [20, 26]):

$$(R1) \quad \textit{subcardinality}: \forall X \subseteq E: r(X) \leq |X|,$$

$$(R2) \quad \textit{monotonicity}: \forall X \subseteq Y \subseteq E: r(X) \leq r(Y),$$

$$(R3) \quad \textit{submodularity}: \forall X, Y \subseteq E: r(X \cap Y) + r(X \cup Y) \leq r(X) + r(Y).$$

Furthermore, we will often make use of the following well-known theorem.

**Theorem 1 (Brualdi [2], also Strong Basis Exchange Theorem).** *Let  $S, S'$  be two different bases of a matroid  $M$ . Then for any  $e \in S \setminus S'$  there is an  $f \in S' \setminus S$  so that  $(S \setminus \{e\}) \cup \{f\}$ ,  $(S' \setminus \{f\}) \cup \{e\}$  are both bases of  $M$ .*

For further results on matroid theory we refer the reader to the excellent books [20, 26].

Let  $M = (E, \mathcal{J})$  be a matroid with rank function  $r: \mathcal{P}(E) \rightarrow \mathbb{N}_0$  and  $\bar{E} := \{e_1, \dots, e_k\} \subseteq E$ ,  $|\bar{E}| \geq 2$ , be a set with cardinality  $k$ . We set  $\bar{E}_j := \{e_1, \dots, e_j\}$  for all  $j \in \{1, \dots, k\}$  and  $\bar{E}_0 := \emptyset$ . Throughout the paper we assume that the following conditions are satisfied:

$$\forall e \in E: \{e\} \in \mathcal{J} \quad \text{and} \quad \bar{E} \in \mathcal{J}, \quad (1)$$

as well as that  $M$  is given via a rank oracle. Furthermore, we use the following notation. Let  $S, T \subseteq E$  and let  $e \in E$ . If the meaning is clear from the context, we will often simply write  $S + T$  and  $S - T$  instead of  $S \cup T$  and  $S \setminus T$  as well as  $S + e$  and  $S - e$  instead of  $S \cup \{e\}$  and  $S \setminus \{e\}$ .

We can now state our optimisation problem. Given an index set  $K := \{k_1, \dots, k_l\} \subseteq \{2, \dots, k\}$  for the (non-linear) monomials and cost functions  $c: E \rightarrow \mathbb{R}$  and  $\bar{c}: K \rightarrow \mathbb{R}$  we consider the maximisation problem

$$\text{maximise } \left\{ \sum_{e \in E} c(e) \cdot x(e) + \sum_{j=1}^l \bar{c}(k_j) \cdot \left( \prod_{i=1}^{k_j} x(e_i) \right), x \in P_M \cap \{0, 1\}^E \right\} \quad (\text{OPT-}K)$$

where  $P_M$  is the polytope that is associated with the matroid  $M$  (note that, w.l.o.g., we will always assume that  $k_l = k$ ). Furthermore we set  $k_0 := 0$ . An important special case of (OPT- $K$ ) is the problem

$$\text{maximise } \left\{ \sum_{e \in E} c(e) \cdot x(e) + \bar{c}(k) \cdot \left( \prod_{i=1}^k x(e_i) \right), x \in P_M \cap \{0, 1\}^E \right\} \quad (\text{OPT-}k)$$

where  $K = \{k\}$  in (OPT- $K$ ) with exactly one non-linear monomial. Although we are mainly interested in the structure of the associated polytopes, we first note that these problems can be solved in polynomial time. This motivates us by the ‘‘separation equals optimisation’’ result to detect the complete descriptions of the associated linearised polytopes.

**Remark 2.** The optimisation problem (OPT- $K$ ) can be solved in polynomial time.

**Proof.** We solve (OPT- $K$ ) with  $K = \{k_1, \dots, k_l\}$  in two main steps. First, we determine for each  $i \in \{1, \dots, l\}$  a cost maximal solution  $X_i^* \in \mathcal{J}, \bar{E}_{k_{i-1}} \subseteq X_i^*$  such that at most  $k_i - k_{i-1} - 1$  elements of  $\bar{E}_{k_i} \setminus \bar{E}_{k_{i-1}}$  are contained in  $X_i^*$ . This can be done by solving a matroid intersection problem of matroid  $M$  and a matroid  $M'_i = (E, \mathcal{J}'_i)$  with  $\mathcal{J}'_i = \{X \subseteq E: |X \cap (\bar{E}_{k_i} \setminus \bar{E}_{k_{i-1}})| \leq k_i - k_{i-1} - 1\}$ . The objective value of problem (OPT- $K$ ) evaluated for  $X_i^*$  equals  $\sum_{e \in X_i^*} c(e) + \sum_{j=1}^{i-1} \bar{c}(k_j)$  in this case. Furthermore we solve one additional matroid optimisation problem in order to determine a set  $X_{i+1}^* \in \mathcal{J}, \bar{E}_{k_i} \subseteq X_{i+1}^*$  such that  $\sum_{e \in X_{i+1}^*} c(e)$  is maximised. The objective value of problem (OPT- $K$ ) evaluated for  $X_{i+1}^*$  is then  $\sum_{e \in X_{i+1}^*} c(e) + \sum_{j=1}^l \bar{c}(k_j)$ . At the end we take a set  $X_i^*$ ,  $i = 1, \dots, l + 1$ , with maximal total objective value.  $\square$

Because (OPT- $k$ ) is a special case of (OPT- $K$ ) the problem (OPT- $k$ ) can be solved by one matroid optimisation and one matroid intersection problem. Alternatively one can solve (OPT- $K$ ) by comparing the optimal values of  $k + 1$  matroid optimisation problems.

Our aim is to present a complete description of the linearised polytopes

$$P_M^{\bar{E}, K} := \text{conv} \left\{ (x, y) \in \{0, 1\}^{E+K} : x \in P_M, y_{k_j} = \prod_{i=1}^{k_j} x(e_i), j = 1, \dots, l \right\},$$

$$P_M^{\bar{E}, k} := \text{conv} \left\{ (x, y) \in \{0, 1\}^{E+\{k\}} : x \in P_M, y = \prod_{i=1}^k x(e_i) \right\}.$$

Note, if  $K = \{k\}$ , we often simply write  $y$  instead of  $y_k$  or  $y_{k_1}$ .

Starting point of our work is a result of Edmonds on the polytope  $P_M$  associated with a matroid  $M$ . Before repeating this result we need some notation. For any  $S \subseteq E$  we define the characteristic vector  $\chi_S \in \{0, 1\}^E$  by

$$\chi_S(e) = \begin{cases} 1, & e \in S, \\ 0, & e \notin S. \end{cases}$$

(In slight abuse of notation, we often do not distinguish between the characteristic vector of an independent set  $J \in \mathcal{J}$  and the set  $J$ .) Then the polytope associated with the matroid  $M$  is

$$P_M = \text{conv}\{\chi_S \in \{0, 1\}^E : S \in \mathcal{J}\}.$$

**Theorem 3 (Edmonds, [7]).** *Let  $M$  be a matroid with rank function  $r: \mathcal{P}(E) \rightarrow \mathbb{N}_0$ , then*

$$P_M = \left\{ x \in \mathbb{R}_+^E : \sum_{e \in T} x(e) \leq r(T), T \subseteq E \right\}.$$

The inequalities

$$x(T) \leq r(T), \quad T \subseteq E \quad (2)$$

with  $x(T) := \sum_{e \in T} x(e)$  are often called *rank constraints/inequalities*. Together with the non-negativity constraints  $x(e) \geq 0, e \in E$ , they form a complete description of  $P_M$ . Edmonds also characterised which sets  $T$  lead to facet defining inequalities of  $P_M$ .

The paper is structured as follows. In Section 2 we present complete descriptions of  $P_M^{\bar{E},k}$  and  $P_M^{\bar{E},K}$ . A main part of the proof consists of showing that apart from the standard linearisation, i. e., all constraints that are needed in the unconstrained case ( $\mathcal{J} = \mathcal{P}(E)$ ) as well, all coefficients of facet defining inequalities  $a^T x + \alpha^T y \leq b$  of the polytopes are non-negative. We start with proving this result for  $P_M^{\bar{E},k}$  and extend this to  $P_M^{\bar{E},K}$  with  $K = \{2, \dots, k\}$ . Based on the complete description in this case, we show that it is possible to derive a complete description of  $P_M^{\bar{E},K}$  for arbitrary sets  $K \subseteq \{2, \dots, k\}$ . In Section 3 we study the facetial structure of  $P_M^{\bar{E},K}$ . So we have a closer look at the so called *trivial facets* and give some necessary conditions for some specially lifted rank inequalities to be facet defining for  $P_M^{\bar{E},K}$ . For  $P_M^{\bar{E},k}$  we even fully characterise all facets. In Section 4 we show that the separation problem for  $P_M^{\bar{E},K}$  can be solved in strongly polynomial time by complete enumeration of the trivial facets and by submodular function minimisation. Finally, in Section 5, we describe further possible extensions of our results and give suggestions for future work. Some rather technical proofs are deferred to the Appendix.

## 2 Complete Description of $P_M^{\bar{E},K}$

In this section we provide a complete description of  $P_M^{\bar{E},K}$  for arbitrary sets  $K \subseteq \{2, \dots, k\}$ . Before we present a formulation of (OPT- $K$ ) we need some definitions.

**Definition 4.** Let  $T \subseteq E$ . The *closure* of set  $T$  is the set

$$\text{cl}(T) := T \cup \{e \in E : r(T) = r(T + e)\}.$$

The set  $T$  is called *closed* if  $T = \text{cl}(T)$ .

Later we will need the following simple result that follows directly from the last definition and the submodularity of the rank function:

**Observation 5.** *Let  $T \subseteq E$  and let  $e \in \text{cl}(T)$ . Then  $r(T + S) = r(T + S + e)$  for all  $S \subseteq E$ .*

We will present strengthened versions of the rank inequalities (2). In order to express the coefficients of the  $y$ -variables corresponding to the linearised monomials we introduce the following objects:




Figure 1: Illustration of  $\alpha_i(\cdot)$ . We consider the graphical matroid on a complete graph  $G = (V, E)$  with six nodes and  $\bar{E} = \{e_1, e_2, e_3, e_4\}$ . The edges of  $T \subseteq E$  are drawn solid and the edges of  $\bar{E}$  are drawn with dotted lines. All other edges are omitted in the picture for the sake of clarity.

**Definition 6.** Let  $T \subseteq E$ . We define  $\alpha_0(T) := 0$  as well as

$$\alpha_i(T) := |\{e_i\} \setminus T| + r(T + \bar{E}_{i-1}) - r(T + \bar{E}_i), \quad i = 1, \dots, k.$$

Furthermore we write  $\alpha_{i,j}(T) := \sum_{m=i}^j \alpha_m(T)$  for  $i, j \in \{1, \dots, k\}, i \leq j$ .

Fig. 1 illustrates  $\alpha_i(T)$  for some sets  $T \subseteq E$  for the graphical matroid on a complete graph  $G = (V, E)$  with six nodes and  $\bar{E} = \{e_1, e_2, e_3, e_4\}$ . The edges of  $T \subseteq E$  are drawn solid and the edges of  $\bar{E}$  are drawn with dotted lines.

The following results follow directly from the definition of  $\alpha_i(\cdot)$ . For details we refer to the Appendix.

**Observation 7.** Let  $T \subseteq E$ .

- (A1) If  $e_i \in T, i \in \{1, \dots, k\}$ , then  $\alpha_i(T) = 0$ .
- (A2) We have  $\alpha_i(T) \in \{0, 1\}$  for all  $i \in \{1, \dots, k\}$ .
- (A3) Let  $i \in \{1, \dots, k\}$ . If  $j \in \mathbb{N}, j > i$ , then  $\alpha_j(T) = \alpha_j(T + e_i)$ .
- (A4) We have  $\alpha_{1,j}(T) = |\bar{E}_j \setminus T| + r(T) - r(T + \bar{E}_j)$  for all  $j \in \{1, \dots, k\}$ .
- (A5) Let  $i \in \{1, \dots, k\}$  and  $e \in \text{cl}(T) \setminus \{e_i\}$ . Then  $\alpha_i(T) = \alpha_i(T + e)$  for all  $i \in \{1, \dots, k\}$ .
- (A6) Let  $i \in \{1, \dots, k\}$  and  $e_i \in \text{cl}(T) \setminus T$ . Then  $\alpha_i(T) = 1 = \alpha_i(T + e_i) + 1$ .

Studying (OPT- $K$ ) we define for each set  $S \subseteq E$  the  $j$ -th (linearised) monomial term of  $S$  by

$$y_{k_j}(S) = \begin{cases} 1, & \bar{E}_{k_j} \subseteq S, \\ 0, & \bar{E}_{k_j} \not\subseteq S, \end{cases}$$

for  $j \in \{1, \dots, l\}$ . So  $P_M^{\bar{E}, K}$  can also be written as

$$P_M^{\bar{E}, K} = \text{conv} \left\{ (\chi_S, y_{k_1}(S), \dots, y_{k_l}(S)) \in \{0, 1\}^{E+K} : S \in \mathcal{J} \right\}.$$

With these definitions we are able to give a formulation of (OPT- $K$ ):

**Lemma 8.** A formulation of (OPT-K) is given by:

$$\text{maximise } \sum_{e \in E} c(e)x(e) + \sum_{j=1}^l \bar{c}(k_j)y_{k_j} \quad (3)$$

$$-y_{k_l} \leq 0, \quad (3)$$

$$y_{k_j} - y_{k_{j-1}} \leq 0, \quad j \in \{2, \dots, l\}, \quad (4)$$

$$\sum_{j=1}^{k_1} x(e_j) - y_{k_1} \leq k_1 - 1, \quad (5)$$

$$\sum_{i=k_{j-1}+1}^{k_j} x(e_i) + y_{k_{j-1}} - y_{k_j} \leq k_j - k_{j-1}, \quad j \in \{2, \dots, l\}, \quad (6)$$

$$-x(e_i) + y_{k_j} \leq 0, \quad j \in \{1, \dots, l\}, i \in \{k_{j-1} + 1, \dots, k_j\}, \quad (7)$$

$$-x(e) \leq 0, \quad e \in E \setminus \bar{E}, \quad (8)$$

$$x(T) + \sum_{j=1}^l \alpha_{k_{j-1}+1, k_j}(T)y_{k_j} \leq r(T), \quad T \subseteq E, \quad (9)$$

$$(x, y) \in \{0, 1\}^{E+K}. \quad (10)$$

Note, we often denote  $(y_{k_1}, \dots, y_{k_l})$  by  $y$ .

**Proof.** Constraints (3)–(7) are the standard linearisation of the non-linear monomials in this case and (8) are the non-negativity conditions for the  $x$ -variables. By the integrality conditions (10) and the complete description of the matroid polytope  $P_M$  in Theorem 3 it remains to show the validity of the strengthened rank inequalities (9). Let  $T \subseteq E$  be an arbitrary set and let  $S \in \mathcal{J}$ . If  $\bar{E}_{k_1} \not\subseteq S$ , then the validity follows from the validity of (2) for  $P_M$ . So let  $\bar{E}_{k_i} \subseteq S, i \in \{1, \dots, l\}$ , and  $\bar{E}_{k_{i+1}} \not\subseteq S$  in the case  $i < l$ . Then by (A4) we have

$$\begin{aligned} \sum_{e \in T} \chi_S(e) + \sum_{j=1}^l \alpha_{k_{j-1}, k_j}(T)y_{k_j}(S) &= |S \cap T| + \underbrace{|\bar{E}_{k_i} \setminus T| + r(T) - r(T \cup \bar{E}_{k_i})}_{=\alpha_{1, k_i}(T)} \\ &\text{(by } (S \cap T) \cap (\bar{E}_{k_i} \setminus T) = \emptyset) = |(S \cap T) \cup (\bar{E}_{k_i} \setminus T)| - r(T \cup \bar{E}_{k_i}) + r(T) \\ &\text{(by } S \in \mathcal{J}, \bar{E}_{k_i} \subseteq S) = r((S \cap T) \cup (\bar{E}_{k_i} \setminus T)) - r(T \cup \bar{E}_{k_i}) + r(T) \\ &\text{(by } ((S \cap T) \cup (\bar{E}_{k_i} \setminus T)) \subseteq T \cup \bar{E}_{k_i}) \leq r(T \cup \bar{E}_{k_i}) - r(T \cup \bar{E}_{k_i}) + r(T) = r(T). \quad \square \end{aligned}$$

A visualisation of the extended rank inequalities (9) on the example of a graphical matroid is given in Fig. 2.

**Remark 9.** With the last result the formulation of (OPT- $k$ ) reduces to

$$\max \sum_{e \in E} c(e)x(e) + \bar{c}(k)y \quad (11)$$


$$x(\bar{E}) - y \leq |\bar{E}| - 1, \quad (11)$$

$$-x(e) + y \leq 0, \quad e \in \bar{E}, \quad (12)$$

$$x \geq 0, y \geq 0, \quad (13)$$

$$x(T) + \alpha_{1, k}(T)y \leq r(T), \quad T \subseteq E, \quad (14)$$

$$x, y \text{ binary.}$$



$T = \{f_1, f_2, f_3\}, \bar{E} = \{e_1, e_2, e_3, e_4\}$
case $K = \{4\}$ :
$x(f_1) + x(f_2) + x(f_3) + 2y_4 \leq 2$
case $K = \{3, 4\}$ :
$x(f_1) + x(f_2) + x(f_3) + y_3 + y_4 \leq 2$

$T = \{f_1, f_2\}, \bar{E} = \{e_1, e_2, e_3\}$
case $K = \{3\}$ :
$x(f_1) + x(f_2) + 2y_3 \leq 2$
case $K = \{2, 3\}$ :
$x(f_1) + x(f_2) + y_2 + y_3 \leq 2$

$T = \{f_1, f_2, f_3\}, \bar{E} = \{e_1, e_2, e_3\}$
case $K = \{2\}$ :
$x(f_1) + x(f_2) + x(f_3) \leq 3$
case $K = \{2, 3\}$ :
$x(f_1) + x(f_2) + x(f_3) + y_3 \leq 3$

Figure 2: Visualisation of the extended rank inequalities (9). We consider the graphical matroid on an undirected graph with six nodes with different sets  $K$  for a given  $\bar{E}$ . The set  $T$  is visualised with bold lines, the edges associated with  $\bar{E}$  with dotted lines. All other edges are omitted in the pictures for the sake of clarity. Note, all sets  $T$  are closed, see also Remark 10.

**Remark 10.** In the formulation and the complete description of  $P_M^{\bar{E}, K}$  and  $P_M^{\bar{E}, k}$  it is sufficient to restrict to closed sets  $T \subseteq E$  for the extended rank inequalities (9) and (14), respectively, i. e.,

$$x(T) + \sum_{j=1}^l \alpha_{k_{j-1}+1, k_j}(T) y_{k_j} \leq r(T), \quad T \subseteq E, \text{cl}(T) = T, \quad (15)$$

for arbitrary sets  $K$  and in the case  $K = \{k\}$  we can restrict to

$$x(T) + \alpha_{1, k}(T) y \leq r(T), \quad T \subseteq E, \text{cl}(T) = T. \quad (16)$$

For proving this results, let  $T \subseteq E$  be an arbitrary set. Then we define  $\bar{T} := \text{cl}(T) \setminus T$ . By **(A5)** and **(A6)** we get  $\alpha_i(T) = \alpha_i(\text{cl}(T)) + |\{e_i\} \setminus \bar{T}|$ ,  $i = 1, \dots, k$ , because  $\alpha_i(T) = \alpha_i(T + e)$  for some  $e \in E$  as long as  $e \neq e_i$  and  $\alpha_i(T) = 1$  if  $e_i \in \bar{T}$ . Let  $(\bar{x}, \bar{y})$  be a point that satisfies (3)–(8) as well as (15), then

$$\begin{aligned} & \bar{x}(T) + \sum_{j=1}^l \alpha_{k_{j-1}+1, k_j}(T) \bar{y}_{k_j} \\ & \leq \underbrace{\bar{x}(T) + \bar{x}(\bar{T} \cap \bar{E}) - \sum_{j=1}^l |\{e_{k_{j-1}+1}, \dots, e_{k_j}\} \setminus \bar{T}| \bar{y}_{k_j}}_{\geq 0 \text{ by (7)}} + \sum_{j=1}^l \alpha_{k_{j-1}+1, k_j}(T) \bar{y}_{k_j} \\ & \leq \bar{x}(T) + \bar{x}(\bar{T} \cap \bar{E}) + \underbrace{\bar{x}(\bar{T} \setminus \bar{E})}_{\geq 0 \text{ by (8)}} + \sum_{j=1}^l \underbrace{\alpha_{k_{j-1}+1, k_j}(\text{cl}(T))}_{=\alpha_{k_{j-1}+1, k_j}(T)} \bar{y}_{k_j} \\ & \quad - |\{e_{k_{j-1}+1}, \dots, e_{k_j}\} \setminus \bar{T}| \\ & \leq \bar{x}(\text{cl}(T)) + \sum_{j=1}^l \alpha_{k_{j-1}+1, k_j}(\text{cl}(T)) \bar{y}_{k_j} \stackrel{(15)}{\leq} r(\text{cl}(T)) = r(T). \end{aligned}$$

So all extended rank inequalities (9) for non-closed sets  $T \subseteq E$  are implied by (3)–(8) and (15).

In the remaining section we will prove that (3)–(8) and (15) are a complete description of  $P_M^{\bar{E}, K}$  and therefore that (11)–(13) and (16) are a complete description of  $P_M^{\bar{E}, k}$ . The proof for

$P_M^{\bar{E},K}$  consists of three steps. First, we consider the special case  $K = \{2, \dots, k\}$ . Indeed one can show that then each facet defining inequality of  $P_M^{\bar{E},K}$  that is not positive multiple of one of (3)–(8) is an inequality of the type  $a^T x + \alpha^T y \leq b$  with  $a \geq 0, \alpha \geq 0$ . This result then allows to derive the complete description of  $P_M^{\bar{E},K}$  in this special case. Finally, we consider the complete description of  $P_M^{\bar{E},K}$  for arbitrary sets  $K \subseteq \{2, \dots, k\}$ .

We start with an observation about the structure of facet defining inequalities  $a^T x + \alpha^T y \leq b$  of  $P_M^{\bar{E},K}$  that are not a positive multiple of one of (3)–(8).

**Observation 11.** *Let  $a^T x + \alpha^T y \leq b$  be a facet defining inequality of  $P_M^{\bar{E},K}$  that is not a positive multiple of the standard-linearisation and the non-negativity constraints (3)–(8). Then we know  $a \geq 0$ .*

**Proof.** Let  $a^T x + \alpha^T y \leq b$  be a facet defining inequality of  $P_M^{\bar{E},K}$  that is not a positive multiple of one of (3)–(8). We consider two cases.

1.  $a(e) < 0$  for some  $e \in E \setminus \bar{E}$ : Because  $a^T x + \alpha^T y \leq b$  is not a positive multiple of  $x(e) \geq 0$  (8), there exists a set  $S \in \mathcal{J}$  with  $e \in S$  and  $a^T \chi_S + \alpha^T y(S) = b$ . Assumption  $e \notin \bar{E}$  implies  $y_{k_i}(S) = y_{k_i}(S - e), i = 1, \dots, l$ , thus  $b = a^T \chi_S + \alpha^T y(S) < a^T \chi_{S-e} + \alpha^T y(S - e)$ . This contradicts the validity of  $a^T x + \alpha^T y \leq b$ .
2.  $a(e) < 0$  for some  $e = e_i \in \bar{E}$ : Because  $a^T x + \alpha^T y \leq b$  is not a positive multiple of  $y_{k_m} \leq x(e_i)$  for  $m \in \{1, \dots, l\}$  such that  $i \in \{k_{m-1} + 1, \dots, k_m\}$ , there exists some  $S \in \mathcal{J}$  with  $e \in S$  and  $y_{k_m}(S) = 0$ . With  $y_{k_j}(S) = y_{k_j}(S - e), j \in \{1, \dots, l\}$ , this implies  $b = a^T \chi_S + \alpha^T y(S) < a^T \chi_{S-e} + \alpha^T y(S - e)$ , again a contraction to validity.  $\square$

Next, we consider the coefficient of the  $y$ -variables of facet defining inequalities  $a^T x + \alpha^T y \leq b$ . Although the result for  $P_M^{\bar{E},k}$  will be implied by the results for  $P_M^{\bar{E},K}$  we will present next the proof in detail because it is much shorter than in the general case and already shows the main ideas.

**Lemma 12.** *Each facet defining inequality  $a^T x + \alpha y \leq b$  of  $P_M^{\bar{E},k}$  that is not a positive multiple of one of (11)–(13) satisfies  $\alpha \geq 0$ .*

**Proof.** Let  $a^T x + \alpha y \leq b$  be a facet defining inequality of  $P_M^{\bar{E},k}$  that is not a positive multiple of one of (11)–(13). By Observation 11 we may assume  $a \geq 0$ . Because  $a^T x + \alpha y \leq b$  is not a positive multiple of  $x(\bar{E}) - y \leq |\bar{E}| - 1$  there exists an  $S \in \mathcal{J}$  with  $|\bar{E} \setminus S| \geq 2$  and  $a^T \chi_S = b$ . Similarly, because  $a^T x + \alpha y \leq b$  is not a positive multiple of  $y \geq 0$ , there exists an  $S' \in \mathcal{J}$  with  $\bar{E} \subseteq S'$  and  $a^T \chi_{S'} + \alpha = b$ . Set  $T := \{e \in E \setminus \bar{E} : a(e) > 0\} \cup (S \cap \bar{E})$ . We may assume  $S \subseteq T$  (otherwise use  $S \cap T$  because  $a \geq 0$  implies  $a(e) = 0$  for all  $e \notin T$ ) and  $r(S) = r(T)$  (otherwise there would be an  $e \in T \setminus S$  with  $S + e \in \mathcal{J}$ , implying  $a^T \chi_{S+e} = a^T \chi_S + a(e) > a^T \chi_S = b$ ). Analogously we may assume  $S' \subseteq T \cup \bar{E}$  as well as  $r(S') = r(T \cup \bar{E})$ . We distinguish two cases.

1.  $r(T) < r(T \cup \bar{E})$ . Then there exists an  $e \in \bar{E} \setminus T$  with  $r(T + e) > r(T) = r(S)$ , so  $S + e \in \mathcal{J}$ . Therefore  $0 = b - b \geq a^T \chi_{S+e} - a^T \chi_S = a(e)$ , so  $a(e) = 0$ . But since  $y(S') = 1$  and  $e \in \bar{E} \subseteq S'$ , this implies  $y(S' - e) = 0$  and so  $a^T \chi_{S'} = a^T \chi_{S'-e} \leq b = a^T \chi_{S'} + \alpha$ . This proves  $\alpha \geq 0$ .
2.  $r(T) = r(T \cup \bar{E})$ . By assumption  $S, S'$  are both bases of  $T \cup \bar{E}$ . We may assume that  $|S \cap S'|$  is maximal. By Theorem 1 there exists for each  $f \in S' \setminus S$  an  $e \in S \setminus S'$  so that  $S - e + f, S' + e - f \in \mathcal{J}$ . Because  $a^T \chi_{S-e+f} \leq b = a^T \chi_S$  we get  $a(e) \geq a(f)$ . We choose  $f \in (S' \setminus S) \cap \bar{E} \neq \emptyset$ , then  $a^T \chi_{S'+e-f} \leq b = a^T \chi_{S'} + \alpha$  implying  $a(e) \leq a(f) + \alpha$ . This proves  $\alpha \geq 0$ .  $\square$



Now, we extend the previous result to  $P_M^{\bar{E},K}$  in the case  $K = \{2, \dots, k\}$ . This is done in three main steps, which make use of the following observation. For further details and the complete, rather technical proofs we refer to the Appendix.

**Observation 13.** *Let  $a^T x + \alpha^T y \leq b$  be a facet defining inequality of  $P_M^{\bar{E},K}$  with  $K = \{2, \dots, k\}$  and let  $S$  be a root of it. Let  $X \subseteq E$ . We define*

$$\text{lhs}(X) := a^T \chi_X + \alpha^T y(X) \quad \text{and} \quad \tilde{a}(X, e) := \text{lhs}(X) - \text{lhs}(X - e), e \in X.$$

Then the following conditions are satisfied:

- (i) for all  $e \in S$ :  $\tilde{a}(S, e) \geq 0$ ,
- (ii) for all  $e \notin S$  with  $S + e \in \mathcal{J}$ :  $\tilde{a}(S + e, e) \leq 0$ ,
- (iii) for all  $e \in S, f \notin S$  with  $S - e + f \in \mathcal{J}$ :  $\tilde{a}(S, e) \geq \tilde{a}(S - e + f, f)$ .

**Proof.** Immediately clear because otherwise  $a^T x + \alpha^T y \leq b$  would not be valid for the sets  $S - e$ ,  $S + e$  or  $S - e + f$ , respectively.  $\square$

First one can show that for each facet defining inequality of  $P_M^{\bar{E},K}$  ( $K = \{2, \dots, k\}$ ) there exists a root with a certain structure.

**Lemma 14.** *Let  $a^T x + \alpha^T y \leq b$  be a facet defining inequality of  $P_M^{\bar{E},K}$  with  $K = \{2, \dots, k\}$  that is not a positive multiple of one of (3)–(8) and let  $m \in \{2, \dots, k\}$ . Then there exists a root  $S$  of  $a^T x + \alpha^T y \leq b$  with the structure*

$$\bar{E}_m \setminus \{e_j, e_m\} = S \cap \bar{E}_m \text{ for some } j \in \{1, \dots, m-1\}.$$

This implies the non-negativity of particular sums of coefficients of facet defining inequalities satisfying certain structures.

**Lemma 15.** *Let  $a^T x + \alpha^T y \leq b$  be a facet defining inequality of  $P_M^{\bar{E},K}$  with  $K = \{2, \dots, k\}$  that is not a positive multiple of one of (3)–(8). Let  $m \in \{2, \dots, k\}$  so that there is a root  $S$  of  $a^T x + \alpha^T y \leq b$  satisfying  $S \cap \bar{E}_m = \bar{E}_m \setminus \{e_j, e_m\}$  for some  $j \in \{1, \dots, m-1\}$ . Then  $\alpha_m + \dots + \alpha_h \geq 0$  for all  $h \in \{m, \dots, k\}$  with  $h = k \vee e_{h+1} \notin S$ .*

This leads to the following result.

**Lemma 16.** *Let  $a^T x + \alpha^T y \leq b$  be a facet defining inequality of  $P_M^{\bar{E},K}$  with  $K = \{2, \dots, k\}$  that is not a positive multiple of one of (3)–(8). Then  $\alpha_m + \dots + \alpha_h \geq 0$  for all  $m, h \in \{2, \dots, k\}$ ,  $m \leq h$ .*

The following corollary summarises the previous lemmas.

**Corollary 17.** *Let  $a^T x + \alpha^T y \leq b$  be a facet defining inequality of  $P_M^{\bar{E},K}$  with  $K = \{2, \dots, k\}$ , that is not a positive multiple of one of (3)–(8). Then  $a \geq 0$  and  $\alpha \geq 0$ .*

**Proof.** Let  $a^T x + \alpha^T y \leq b$  be a facet defining inequality of  $P_M^{\bar{E},K}$  with  $K = \{2, \dots, k\}$  satisfying the requirements. Then  $a \geq 0$  follows from Observation 11 and  $\alpha \geq 0$  follows from Lemma 16 for  $m = h$ .  $\square$

This allows us to prove our main result for  $P_M^{\bar{E},K}$  in the case  $K = \{2, \dots, k\}$ .

**Theorem 18.** *The inequalities (3)–(8) and (15) are a complete description of  $P_M^{\bar{E},K}$ ,  $K = \{2, \dots, k\}$ .*

**Proof.** Let  $a^T x + \alpha^T y \leq b$  be a facet defining inequality of  $P_M^{\bar{E}, K}$ ,  $K = \{2, \dots, k\}$ , that is not a positive multiple of one of (3)–(8) and (15). Then Corollary 17 implies  $a \geq 0$  and  $\alpha \geq 0$ . Let  $T' := \{e \in E : a(e) > 0\}$  and  $T := \text{cl}(T')$ .

If we can prove for some  $S \in \mathcal{J}$  with  $a^T \chi_S + \alpha^T y(S) = b$  that there exists an  $e \in T' \setminus S$  so that  $S + e \in \mathcal{J}$ , then  $a^T \chi_{S+e} + \alpha^T y(S+e) > a^T \chi_S + \alpha^T y(S) = b$ , hence the inequality would not be valid, a contradiction. We distinguish two cases.

We consider the constraint (15) associated with  $T$ . Because  $a^T x + \alpha^T y \leq b$  is not a positive multiple of this constraint, there exists an  $S \in \mathcal{J}$  with  $\sum_{e \in T} \chi_S(e) + \alpha(T)^T y(S) < r(T)$  and  $a^T \chi_S + \alpha^T y(S) = b$ . In particular,  $r(S \cap T) = \sum_{e \in T} \chi_S(e)$  implies

$$r(S \cap T) + \alpha(T)^T y(S) < r(T). \quad (17)$$

Let  $\hat{i} \in \{1, \dots, k\}$  be maximal so that  $y_i = 1 \Leftrightarrow i \leq \hat{i}$  ( $\hat{i} = 1$  means that  $\bar{E}_2 \not\subseteq S$ ). Note that  $\hat{i} > 1$  implies  $\bar{E}_{\hat{i}} \subseteq S$  and  $\hat{i} < n$  implies  $\bar{E}_{\hat{i}+1} \not\subseteq S$  as well. We first consider the case  $\hat{i} > 1$ . Then we may assume  $S \subseteq T' \cup \bar{E}_{\hat{i}}$  (otherwise use  $\bar{S} := S \cap (T' \cup \bar{E}_{\hat{i}})$  with  $a^T \chi_{\bar{S}} + \alpha^T y(\bar{S}) = a^T \chi_S + \alpha^T y(S) = b$ ). Note that  $\alpha^T y(S) = \alpha_{1, \hat{i}}(T)$ , so (17) and **(A4)** imply  $r(S \cap T) < r(T \cup \bar{E}_{\hat{i}}) - |\bar{E}_{\hat{i}} \setminus T|$ . Therefore

$$r(S) = r(S \cap T) + |S \setminus T| < r(T \cup \bar{E}_{\hat{i}}) - \underbrace{|\bar{E}_{\hat{i}} \setminus T| + |S \setminus T|}_{=0 \text{ by } \bar{E}_{\hat{i}} \subseteq S \subseteq T \cup \bar{E}_{\hat{i}}} = r(T \cup \bar{E}_{\hat{i}}) = r(T' \cup \bar{E}_{\hat{i}}).$$

Here the last equation follows from Observation 5. So there exists an  $e \in (T' \cup \bar{E}_{\hat{i}}) \setminus S = T' \setminus S$  with  $S + e \in \mathcal{J}$ , which contradicts the validity of  $a^T x + \alpha^T y \leq b$ .

It remains to consider the case  $\hat{i} = 1$ . But then  $y_i(S) = 0, i = 2, \dots, k$ , and we know with  $S' := S \cap T'$  that  $r(S') \leq r(S \cap T) < r(T) = r(T')$  by (17). Hence there exists an  $e \in T' \setminus S'$  with  $S' + e \in \mathcal{J}$ , a contradiction.  $\square$

**Remark 19.** If we consider (OPT- $k$ ), i. e., (OPT- $K$ ) with  $K = \{k\}$ , the proof of Theorem 18 in combination with Lemma 12 also shows that (11)–(13) and (16) define a complete description of  $P_M^{\bar{E}, k}$  because setting  $\hat{i} = k$  implies  $\alpha_{1, k}(T) = \sum_{j=1}^{\hat{i}} \alpha_j(T)$ .

In our last step we extend the previous results to the complete description of  $P_M^{\bar{E}, K}$  with arbitrary  $K = \{k_1, \dots, k_l\} \subseteq \{2, \dots, k\}$ .

**Theorem 20.** *Inequalities (3)–(8) and (15) are a complete description for  $P_M^{\bar{E}, K}$  with arbitrary set  $K = \{k_1, \dots, k_l\} \subseteq \{2, \dots, k\}$ .*

**Proof.** We have already proved that (3)–(8) and (15) are a complete description for  $P_M^{\bar{E}, K}$  if  $K = \{2, \dots, k\}$ .

Let  $j \in \{1, \dots, l\}$  so that either  $j \geq 2$  and  $k_j - k_{j-1} \geq 2$  or  $j = 1$  and  $k_1 > 2$ . Set

$$\hat{k} := \begin{cases} k_{j-1} + 1, & j \geq 2, \\ 2, & j = 1. \end{cases}$$

It will be convenient throughout the proof to identify formally  $\bar{y}_1 := x(e_1)$ . We show that the complete description for  $K$  can be derived from the complete description for  $P_M^{\bar{E}, K'}$  with  $K' = K \cup \{\hat{k}\}$ .

Let  $(\bar{x}, \bar{y})$  be a point that satisfies (3)–(8) as well as (15) for (OPT- $K$ ). We consider two cases:

**Case 1** If setting  $\bar{y}_{\hat{k}} := \bar{y}_{k_j}$ , the point  $(\bar{x}, \bar{y}_{k_1}, \dots, \bar{y}_{k_{j-1}}, \bar{y}_{\hat{k}}, \bar{y}_{k_j}, \dots, \bar{y}_{k_l})$  is feasible for  $P_M^{\bar{E}, K'}$ , the statement immediately follows from the complete description of the known case. The only inequality that might be violated by this definition of  $\bar{y}_{\hat{k}}$  is  $\bar{y}_{\hat{k}-1} + \bar{x}(e_{\hat{k}}) \leq 1 + \bar{y}_{\hat{k}}$ . The validity of all other constraints follows directly from the feasibility of  $(\bar{x}, \bar{y})$  and the fact that  $\bar{x}(e) \leq 1, e \in E$  (by (15)). For (15) with  $T \subseteq E$ ,  $T$  closed, we just split  $\alpha_{\hat{k}, k_j}(T) \bar{y}_{k_j}$  into  $\alpha_{\hat{k}, \hat{k}}(T) \bar{y}_{\hat{k}} + \alpha_{\hat{k}+1, k_j}(T) \bar{y}_{k_j}$  (note, in the case  $\hat{k} = 2$  the coefficient of  $\bar{y}_{\hat{k}}$  is  $\alpha_{1, \hat{k}}(T)$ , but  $T$  closed implies  $\alpha_1(T) = 0$  and so  $\alpha_{1, \hat{k}}(T) = \alpha_{\hat{k}, \hat{k}}(T)$ ).

**Case 2** Otherwise, we set  $\bar{y}_{\hat{k}} := \bar{y}_{\hat{k}-1} + \bar{x}(e_{\hat{k}}) - 1$  and so  $\bar{y}_{\hat{k}} > \bar{y}_{k_j}$ . But then  $(\bar{x}, \hat{y}) := (\bar{x}, \bar{y}_{k_1}, \dots, \bar{y}_{\hat{k}-1}, \bar{y}_{\hat{k}}, \bar{y}_{k_j}, \dots, \bar{y}_{k_l})$  might possibly violate (15) of the formulation of  $P_M^{\bar{E}, K'}$ . All other constraints of the formulation of  $P_M^{\bar{E}, K'}$  ((3)–(8)) remain satisfied by this definition of  $\bar{y}_{\hat{k}}$ , either trivially or because

$$\sum_{i=\hat{k}+1}^{k_j} \bar{x}(e_i) + \underbrace{\bar{x}(e_{\hat{k}}) + \bar{y}_{\hat{k}-1}}_{=\bar{y}_{\hat{k}+1}} \leq \bar{y}_{k_j} + k_j - k_{j-1} \Leftrightarrow \sum_{i=\hat{k}+1}^{k_j} \bar{x}(e_i) + \bar{y}_{\hat{k}} \leq \bar{y}_{k_j} + (k_j - \hat{k}).$$

Let  $T \subseteq E$ ,  $\text{cl}(T) = T$ , (see also Remark 10) be a closed set and consider the associated rank constraint (15). If  $\alpha_{\hat{k}}(T) = 0$ , then the validity of this rank constraint for  $P_M^{\bar{E}, K'}$  follows directly from the validity of the associated rank constraint for  $P_M^{\bar{E}, K}$ . So we may assume  $\alpha_{\hat{k}}(T) = 1$ . This implies  $r(T + \bar{E}_{\hat{k}-1}) = r(T + \bar{E}_{\hat{k}})$  and with  $T' := T + e_{\hat{k}}$

$$\begin{aligned} \alpha_{1,m}(T') - \alpha_{1,m}(T) &= 1 - r(T' + \bar{E}_m) + r(T + \bar{E}_m) \geq 0, \quad m = 1, \dots, \hat{k} - 1, \\ \alpha_{1,\hat{k}-1}(T') - \alpha_{1,\hat{k}-1}(T) &= 1 - r(T + \bar{E}_{\hat{k}}) + r(T + \bar{E}_{\hat{k}-1}) = 1. \end{aligned} \quad (18)$$

Let  $\gamma_i := \bar{y}_{k_i} - \bar{y}_{k_{i+1}} \geq 0, i = 1, \dots, j-2$ . Then we can write  $\bar{y}_{k_i} = \bar{y}_{\hat{k}-1} + \sum_{m=i}^{j-2} \gamma_m$ . Using these relations we get for the rank constraint associated with  $T$  for  $K'$

$$\begin{aligned} \bar{x}(T) + \sum_{i=1}^{j-1} \alpha_{k_{i-1}+1, k_i}(T) \bar{y}_{k_i} + \underbrace{\alpha_{\hat{k}}(T)}_{=1} \bar{y}_{\hat{k}} + \underbrace{\alpha_{\hat{k}+1, k_j}(T)}_{=\alpha_{\hat{k}+1, k_j}(T')} \bar{y}_{k_j} + \sum_{i=j+1}^l \underbrace{\alpha_{k_{i-1}+1, k_i}(T)}_{=\alpha_{k_{i-1}+1, k_i}(T')} \bar{y}_{k_i} \\ = \bar{x}(T) + \underbrace{\alpha_{1, \hat{k}-1}(T)}_{=\alpha_{1, \hat{k}-1}(T')-1} \bar{y}_{\hat{k}-1} + \sum_{i=1}^{j-2} \underbrace{\alpha_{1, k_i}(T)}_{\leq \alpha_{1, k_i}(T')} \gamma_i + \bar{y}_{\hat{k}} + \underbrace{\alpha_{\hat{k}+1, k_j}(T')}_{=\alpha_{\hat{k}, k_j}(T')} \bar{y}_{k_j} \\ + \sum_{i=j+1}^l \alpha_{k_{i-1}+1, k_i}(T') \bar{y}_{k_i} \end{aligned}$$

and using  $-\bar{y}_{\hat{k}-1} = -y_{\hat{k}} + \bar{x}(e_{\hat{k}}) - 1$  and the definition of the  $\gamma_{\bullet}$

$$\begin{aligned} &\leq \bar{x}(T') + \sum_{i=1}^l \alpha_{k_{i-1}+1, k_i}(T') \bar{y}_{k_i} - 1 \\ &\leq \bar{x}(\text{cl}(T')) + \sum_{i=1}^l \alpha_{k_{i-1}, k_i}(\text{cl}(T')) \bar{y}_{k_i} - 1 \leq r(\text{cl}(T')) - 1 = r(T). \end{aligned}$$

For the last step we used that each rank constraint associated with  $T'$  is always dominated by the rank constraint associated with  $\text{cl}(T')$  by Remark 10. So the result follows again by the known complete description of  $P_M^{\bar{E}, K'}$ .  $\square$

So the complete description of  $P_M^{\bar{E}, K}$  for arbitrary sets  $K \subseteq \{2, \dots, k\}$  can easily be derived from the complete description of  $P_M^{\bar{E}, K'}$  with  $K' = \{2, \dots, k\}$  by adapting the constraints of the standard linearisation and by adjusting the extended rank inequalities. Indeed, the coefficient of  $y_{k_j}, k_j \in K$ , equals the sum of the coefficients of variables  $y_i, i = k_{j-1} + 1, \dots, k_j$ , in the problem with  $K'$ . Although feasibility of the extended rank inequalities is trivially preserved by this approach, it is remarkable that we even get the complete description of  $P_M^{\bar{E}, K}$  in this way.

### 3 Facetial Structure of $P_M^{\bar{E},K}$ and $P_M^{\bar{E},k}$

The polytope  $P_M$  of some matroid  $M$  is well-understood. In 1970 Edmonds [7] presented a complete description of  $P_M$  and characterised its facetial structure, see also [14]. We will extend these results to the polytopes  $P_M^{\bar{E},k}$  and  $P_M^{\bar{E},K}$  and study their facetial structures in more detail. This includes the dimension of the polytopes as well as results on the facetness of the constraints of the standard linearisation. Furthermore, we have a closer look at the extended rank inequalities (9). If a closed set  $T$  does not satisfy certain non-separability conditions (to be defined below) the associated rank inequality cannot be facet defining for  $P_M^{\bar{E},K}$ . For  $P_M^{\bar{E},k}$  we even exactly characterise the facet defining inequalities and provide a minimal description.

We start with studying the dimension and the trivial facets of  $P_M^{\bar{E},K}$ . One can easily prove the following result:

**Observation 21.** *Under the assumptions (1) the polytope  $P_M^{\bar{E},K}$  is full-dimensional. The inequalities (3)–(8) define facets of  $P_M^{\bar{E},K}$ .*

In the next step we consider the extended rank inequalities (14). These are extensions of the rank inequalities (2). Before repeating results on the facetness of (2) for  $P_M$  we need a definition, which will be extended to our setting later on.

**Definition 22.** Let  $T \subseteq E$ . A set  $T$  is called *separable* if there exists a partition  $T_1, T_2 \subsetneq T, T_1 \cup T_2 = T, T_1 \cap T_2 = \emptyset$  such that  $r(T_1) + r(T_2) = r(T)$ , otherwise it is called *non-separable*.

Indeed, all closed and non-separable sets  $T \subseteq E$  lead to facet defining rank inequalities (2) of  $P_M$  [7, 14]. They also lead to facet defining inequalities (9) of  $P_M^{\bar{E},K}$  whenever  $\alpha_{k_{j-1}+1, k_j}(T) > 0$  for all  $j \in \{1, \dots, l\}$ .

**Proposition 23.** *Let  $T \subseteq E$  be a closed and non-separable set associated with a facet defining inequality (2) of  $P_M$ . If  $\alpha_{k_{j-1}+1, k_j}(T) > 0$  for all  $j \in \{1, \dots, l\}$ , then the inequality (9) is facet defining for  $P_M^{\bar{E},K}$ , too.*

**Proof.** Because  $x(T) \leq r(T)$  is facet defining for  $P_M$  there exist pairwise different  $S_i \in \mathcal{J}$ ,  $S_i \subseteq T$ ,  $r(S_i) = r(T)$ ,  $i = 1, \dots, d$ ,  $d := \dim P_M$ . By  $\alpha_{k_{j-1}+1, k_j}(T) > 0, j \in \{1, \dots, l\}$ , we know  $(\bar{E}_{k_j} \setminus \bar{E}_{k_{j-1}}) \not\subseteq T, j \in \{1, \dots, l\}$ , hence  $(\bar{E}_{k_j} \setminus \bar{E}_{k_{j-1}}) \not\subseteq S_i$  for all  $i \in \{1, \dots, d\}, j \in \{1, \dots, l\}$ . If we choose for each  $j \in \{1, \dots, l\}$  a set  $\hat{S}_j \in \mathcal{J}$  with  $\bar{E}_{k_j} \subseteq \hat{S}_j \subseteq T \cup \bar{E}_{k_j}$  and  $|\hat{S}_j| = r(\hat{S}_j) = r(T \cup \bar{E}_{k_j})$ , the left-hand side of  $x(\hat{S}_j \cap T) + \sum_{j=1}^l \alpha_{k_{j-1}+1, k_j}(T) y_{k_j}(\hat{S}_j) \leq r(T)$  computes to

$$\begin{aligned} r(\hat{S}_j \cap T) + \alpha_{1, k_j}(T) &= r(T \cup \bar{E}_{k_j}) - |\hat{S}_j \setminus T| + \alpha_{1, k_j}(T) \\ &= r(T \cup \bar{E}_{k_j}) - |\bar{E}_{k_j} \setminus T| + \alpha_{1, k_j}(T) = r(T) \end{aligned}$$

by (A4). So there exist  $d + l$  affinely independent sets that are roots of (9), which proves the statement.  $\square$

Considering the first type of extended subtour elimination constraints (constraints (5) in [4]) for the spanning forest problem with one linearised quadratic monomial in [4], the facetness of these inequalities follows directly from the previous result. But as already observed by Buchheim and Klein [4] for this problem (see constraints (6) in [4]), not all facet defining inequalities of  $P_M^{\bar{E},K}$  are directly related to facet defining inequalities of  $P_M$ . So we need further properties that allow us to describe the facetial structure of  $P_M^{\bar{E},K}$ .

**Definition 24.** Let  $T \subseteq E$  be closed and  $K = \{k_1, \dots, k_l\} \subset \{2, \dots, k\}$ . A set  $T$  is called  $(\bar{E}, K)$ -*separable* if there exists a partition  $T_1, T_2 \subsetneq T, T_1 \cap T_2 = \emptyset, T_1 \cup T_2 = T$  with

$$r(T) = r(T_1) + r(T_2), \quad \text{and} \quad \alpha_{k_{j-1}+1, k_j}(T) = \alpha_{k_{j-1}+1, k_j}(T_1) + \alpha_{k_{j-1}+1, k_j}(T_2), \quad j = 1, \dots, l, \quad (19)$$

otherwise it is called  $(\bar{E}, K)$ -*non-separable*.




Figure 3: Visualisation of  $(\bar{E}, K)$ -(non)-separability. We consider the graphical matroid on an undirected graph with four nodes. Although set  $T$  (bold lines) is the same on both sides, depending on  $\bar{E}$  (dotted lines), it is  $(\bar{E}, \{2\})$ -separable or not. Note,  $T$  is separable with  $T_1 = \{e_1\}, T_2 = \{e_2\}$  by Definition 22.

Note, the definition of separability by Edmonds [7] only included the first equation of the last definition (see Definition 22). A visualisation of the new setting is given in Fig. 3.

A closed set  $T \subseteq E$  that is  $(\bar{E}, K)$ -separable does not lead to a facet defining rank inequality (9) of  $P_M^{\bar{E}, K}$ .

**Observation 25.** *Let (9) be a facet defining inequality of  $P_M^{\bar{E}, K}$  associated with a closed set  $T \subseteq E$ . Then  $T$  is  $(\bar{E}, K)$ -non-separable.*

**Proof.** Let  $T \subseteq E$  be closed and assume that  $T$  is  $(\bar{E}, K)$ -separable with partition  $T_1, T_2 \subsetneq T, T_1 \cap T_2 = \emptyset, T_1 \cup T_2 = T$  satisfying (19). Then we can derive (9) by adding the two rank inequalities (9) for  $T_1, T_2$ :

$$\underbrace{x(T_1) + x(T_2)}_{=x(T)} + \sum_{j=1}^l \underbrace{(\alpha_{k_{j-1}+1, k_j}(T_1) + \alpha_{k_{j-1}+1, k_j}(T_2))}_{=\alpha_{k_{j-1}+1, k_j}(T)} y_{k_j} \leq r(T).$$

So the extended rank inequality associated with  $T$  does not define a facet of  $P_M^{\bar{E}, K}$ .  $\square$

However, not all  $(\bar{E}, K)$ -non-separable sets  $T \subseteq E$  lead to facet defining rank inequalities (9). This may depend on the structure of  $K$  as well. If, for instance,  $k_{j-1}, k_j \in K$  with  $k_{j-1} + 1 = k_j$ , then the inequality  $x(e_{k_j}) \leq 1$  can be derived by adding  $-y_{k_{j-1}} + y_{k_j} \leq 0$  and  $y_{k_{j-1}} + x(e_{k_j}) - y_{k_j} \leq 1$  (see also property **(P1)<sup>k</sup>** to be defined below for the case  $K = \{k\}$ ).

In the following we will fully characterise the facetial structure of  $P_M^{\bar{E}, k}$ . We start with some easy results that follow directly from the definition of  $(\bar{E}, \{k\})$ -separability.

**Observation 26.** *Let  $T \subseteq E$  be a closed set with  $|T| \geq 2$ .*

1. *If  $T \subseteq \bar{E}$ , then  $T$  is  $(\bar{E}, \{k\})$ -separable.*
2. *If  $\alpha_{1, k}(T \setminus \bar{E}) = 0$  and  $\bar{E} \cap T \neq \emptyset$ , then  $T$  is  $(\bar{E}, \{k\})$ -separable.*

In addition to closedness and  $(\bar{E}, \{k\})$ -non-separability it will turn out that a set  $T \subseteq E$  leading to facet defining extended rank inequalities (14) has to satisfy the following properties:

**Definition 27.** Let  $T \subseteq E$  be a closed and  $(\bar{E}, \{k\})$ -non-separable set. We say it satisfies properties

- (P1)<sup>k</sup>** if  $\bar{E} \not\subseteq T$  and if  $T \neq \{e\}, e \in \bar{E}$ , in the case  $|\bar{E}| = 2$ ,
- (P2)<sup>k</sup>** if  $\bar{E} \subseteq T$  and  $\alpha_{1, k}(T \setminus \bar{E}) \geq 2$ .

These properties are necessary conditions that a set  $T \subseteq E$  leads to a facet defining rank inequality (14) of  $P_M^{\bar{E}, k}$ .

**Observation 28.** Let  $T \subseteq E$  be a closed and  $(\bar{E}, \{k\})$ -non-separable set. If the rank inequality associated with  $T$  is facet defining for  $P_M^{\bar{E},k}$ , then it satisfies either  $(\mathbf{P1}^k)$  or  $(\mathbf{P2}^k)$ .

Next we will prove that for a set  $T \subseteq E$  closedness,  $(\bar{E}, \{k\})$ -non-separability and properties  $(\mathbf{P1}^k)$ ,  $(\mathbf{P2}^k)$  lead to facetness of the associated inequality (14). In the proof of this result we need the following observation:

**Observation 29.** Let  $T \subseteq E$  be a closed set with  $\bar{E} \subseteq T$ . If  $\alpha_{1,k}(T \setminus \bar{E}) \geq 2$ , then there exists a basis  $B$  of  $T$  such that  $|B \cap \bar{E}| \leq |\bar{E}| - 2$ .

**Lemma 30.** Let  $T \subseteq E$  be a closed and  $(\bar{E}, \{k\})$ -non-separable set satisfying properties  $(\mathbf{P1}^k)$  or  $(\mathbf{P2}^k)$ . Then the extended rank inequality (14) defines a facet of  $P_M^{\bar{E},k}$ .

**Proof.** Let  $T \subseteq E$  be a closed,  $(\bar{E}, \{k\})$ -non-separable set satisfying  $(\mathbf{P1}^k)$  or  $(\mathbf{P2}^k)$ . Further we assume that  $a^T x + \bar{a}y \leq b$  is a facet defining inequality of  $P_M^{\bar{E},k}$  so that  $a^T \chi_S + \bar{a}y(S) = b$  whenever  $S$  is a root of  $x(T) + \alpha_{1,k}(T)y \leq r(T)$ , the extended rank constraint associated with  $T$ . We will show that in this case  $a^T x + \bar{a}y \leq b$  is a positive multiple of  $x(T) + \alpha_{1,k}(T)y \leq r(T)$ . Our proof consists of four main steps.

**Claim 1.** Under the given assumptions the constraints (14) are not implied by the constraints of the standard linearisation (see Observation 21).

*Proof of Claim 1.* In order to prove that  $x(T) + \alpha_{1,k}(T)y \leq r(T)$  is not implied by (11)–(13) we can easily determine in each of the cases an independent set  $J \in \mathcal{J}$  such that  $J$  is a root of the rank inequality (14), but  $J$  is not a root of one of (11)–(13). Technical details can be found in the Appendix.

So by Lemma 12 we know that  $a \geq 0, \bar{a} \geq 0$ . Next we prove the following:

**Claim 2.** A coefficient  $a(e), e \in \bar{E}$ , or  $\bar{a}$  is zero if the corresponding coefficient in  $x(T) + \alpha_{1,k}(T)y \leq r(T)$  is zero.

*Proof of Claim 2.* Assume, for a contradiction,  $a(e) > 0$  for some  $e \notin T$ . Let  $S$  be a basis of  $T$ . Then  $S$  is a root of  $x(T) + \alpha_{1,k}(T)y \leq r(T)$  and consequently of  $a^T x + \bar{a}y \leq b$ , too. Because  $T$  is closed  $S + e \in \mathcal{J}$ , but  $S + e$  violates  $a^T x + \bar{a}y \leq b$ , a contradiction.

Now assume the coefficient  $\alpha_{1,k}(T)$  of  $y$  is zero in  $x(T) + \alpha_{1,k}(T)y \leq r(T)$ , i. e.,

$$r(T) + |\bar{E} \setminus T| = r(T \cup \bar{E}) \quad (20)$$

by **(A4)**. First, if  $\bar{E} \not\subseteq T$ , we choose a basis  $B$  of  $T$  with  $(T \cap \bar{E}) \subseteq B$ . Then by (20) we know  $B + (\bar{E} \setminus T) \in \mathcal{J}$  and this independent set is also a root of  $x(T) + \alpha_{1,k}(T)y \leq r(T)$ . However, because  $y(B + (\bar{E} \setminus T)) = 1$  it follows  $\bar{a} = 0$ . Second, let  $\bar{E} \subseteq T$ . Let  $\bar{B}$  be a basis of  $T$  with  $\bar{E} \subseteq \bar{B}$ . Furthermore, by assumption  $(\mathbf{P2}^k)$  we may apply Observation 29, and find a basis  $B$  of  $T$  with  $|B \cap \bar{E}| \leq |\bar{E}| - 2$ . So there exist two elements  $e_i, e_j \in (\bar{B} \setminus B) \cap \bar{E}, i \neq j$ . Because  $M$  is a matroid there exists an  $f_i \in B \setminus \bar{B} \subseteq B \setminus \bar{E}$  with  $\bar{B} - e_i + f_i \in \mathcal{J}$  and an  $f_j \in B \setminus (\bar{B} - e_i + f_i) \subseteq B \setminus \bar{E}$  with  $\bar{B} - e_i - e_j + f_i + f_j \in \mathcal{J}$ . Note that all three sets  $\bar{B}, \bar{B} - e_i + f_i, \bar{B} - e_i - e_j + f_i + f_j$  are bases of  $T$  and define roots of  $x(T) + \alpha_{1,k}(T)y \leq r(T)$ . In particular, this implies  $a(e_i) + \bar{a} = a(f_i)$  and  $a(e_j) = a(f_j)$ . Submodularity gives

$$\underbrace{r(\bar{B} - e_j)}_{=r(T)-1} + \underbrace{r(\bar{B} - e_j + f_i + f_j)}_{=r(T)} \leq r(\bar{B} - e_j + f_i) + r(\bar{B} - e_j + f_j),$$

so at least one of  $\bar{B} - e_j + f_i, \bar{B} - e_j + f_j$  is a basis of  $T$ . In the second case we conclude  $a(e_i) = a(f_i)$ , hence  $\bar{a} = 0$ . Otherwise, if  $\bar{B} - e_j + f_j$  is not a basis,  $a(e_i) = a(f_j) = a(e_j)$  and again by submodularity

$$\underbrace{r(\bar{B} - e_i - e_j + f_j)}_{=r(T)-1} + \underbrace{r(\bar{B} + f_j)}_{=r(T)} \leq r(\bar{B} - e_i + f_j) + \underbrace{r(\bar{B} - e_j + f_j)}_{=r(T)-1},$$

hence  $\bar{B} - e_i + f_j$  is a basis and we get  $a(f_i) = a(f_j) = a(e_i)$  and again  $\bar{a} = 0$ . This proves Claim 2.

It remains to prove that all coefficients  $a(e), e \in T$ , have the same value and that the coefficient  $\bar{a}$  of  $y$  has the correct value. We start with the first statement.

**Claim 3.** All coefficients  $a(e), e \in T$ , have the same value.

*Proof of Claim 3.* We define the simple undirected graph  $G = (T, A)$  with set of nodes  $T$  and set of edges

$$A := \{ \{e, f\} : \exists B \subseteq T, e, f \in T \setminus B, B + e, B + f \text{ bases of } T \} \\ \cup \left\{ \{e, f\} : \exists B \subseteq T \cup \bar{E}, e, f \in T \setminus B, \bar{E} \subseteq B, B + e, B + f \text{ bases of } T \cup \bar{E} \right\}.$$

We prove the statement in two steps.

**Claim 3.1.** If  $\{e, f\} \in A$ , then we have  $a(e) = a(f)$ .

*Proof of Claim 3.1.* Let  $\{e, f\} \in A$ . If, on the one hand, there exists  $B \subseteq T$ ,  $e, f \in T \setminus B$ , with  $B + e, B + f$  bases of  $T$ , then both bases define roots of  $x(T) + \alpha_{1,k}(T)y \leq r(T)$  and consequently  $\sum_{e' \in B+e} a(e') + \bar{a}y(B+e) = \sum_{e' \in B+f} a(e') + \bar{a}y(B+f)$ , hence  $a(e) + \bar{a}y(B+e) = a(f) + \bar{a}y(B+f)$ . If  $\bar{E} \not\subseteq T$ , then  $y(B+e) = y(B+f) = 0$ , otherwise  $\alpha_{1,k}(T) = r(T) + |\bar{E} \setminus T| - r(T \cup \bar{E}) = r(T) + 0 - r(T) = 0$  by **(A4)** implying  $\bar{a} = 0$ , so  $a(e) = a(f)$  follows. On the other hand, if there exists  $B \subseteq T \cup \bar{E}$ ,  $e, f \in T \setminus B$ , with  $\bar{E} \subseteq B$  and  $B + e, B + f$  bases of  $T \cup \bar{E}$ , then  $y(B+e) = y(B+f) = 1$ . Furthermore

$$\begin{aligned} |(B+e) \cap T| + \alpha_{1,k}(T)y(B+e) &= |(B+e) \cap T| + r(T) + |\bar{E} \setminus T| - r(T \cup \bar{E}) \\ &= |(B+e) \cap T| + r(T) + |(B+e) \setminus T| - r(T \cup \bar{E}) \\ &= r(T \cup \bar{E}) + r(T) - r(T \cup \bar{E}) = r(T) \end{aligned}$$

by **(A4)**, so  $B + e$  defines a root of  $x(T) + \alpha_{1,k}(T)y \leq r(T)$  (and, analogously,  $B + f$ ). As above we conclude  $a(e) = a(f)$  and Claim 3.1 follows. Therefore, if we can prove that  $G$  is connected, all coefficients  $a(e), e \in T$ , must be equal.

**Claim 3.2.** Graph  $G$  is connected.

*Proof of Claim 3.2.* Assume, for a contradiction, that  $G$  is not connected, i.e., there exist  $T_1, T_2 \subseteq T$ ,  $T_1, T_2 \neq \emptyset$ ,  $T_1 \cap T_2 = \emptyset$  and  $T_1 \cup T_2 = T$  such that there do not exist edges between  $T_1$  and  $T_2$ . First assume  $r(T) < r(T_1) + r(T_2)$  and let  $B'$  be a basis of  $T$  with  $r(T_1) = r(B' \cap T_1)$ . Then  $r(T_2) > r(B' \cap T_2)$ , so there exists an  $e \in T_2 \setminus B'$  so that  $(B' \cap T_2) + e \in \mathcal{J}$ . However,  $B' + e \notin \mathcal{J}$  ( $B'$  basis), but because  $M$  is a matroid there exists an  $f \in (T_1 \cap B')$  so that  $B' + e - f$  is a basis of  $T$ . Consequently, choosing  $B := B' - f$  proves  $\{e, f\} \in A$ , a contradiction.

Secondly, if  $r(T) = r(T_1) + r(T_2)$  then we know that  $|\bar{E}| + r(T \cup \bar{E}) < r(T_1 \cup \bar{E}) + r(T_2 \cup \bar{E})$  by **(A4)** and because  $T$  is  $(\bar{E}, \{k\})$ -non-separable. Let  $B'$  be a basis of  $T \cup \bar{E}$  with  $\bar{E} \subseteq B'$  and  $r(B' \cap (T_1 \cup \bar{E})) = r(T_1 \cup \bar{E})$ . Consequently,

$$\begin{aligned} r(B' \cap (T_2 \cup \bar{E})) &= |\bar{E}| + |B' \cap (T_2 \setminus \bar{E})| \\ &< r(T_1 \cup \bar{E}) + r(T_2 \cup \bar{E}) - r(T \cup \bar{E}) + |B' \cap (T_2 \setminus \bar{E})| \\ &= |B' \cap (T_1 \cup \bar{E})| + r(T_2 \cup \bar{E}) - r(T \cup \bar{E}) + |B' \cap (T_2 \setminus \bar{E})| \\ &= |B'| + r(T_2 \cup \bar{E}) - r(T \cup \bar{E}) = r(T_2 \cup \bar{E}). \end{aligned}$$

Hence there exists an  $e \in T_2 \setminus B'$  so that  $(B' + e) \cap (T_2 \cup \bar{E}) \in \mathcal{J}$ , but  $B' + e \notin \mathcal{J}$ . Because  $M$  is a matroid there exists an  $f \in T_1$  such that  $B' + e - f$  is a basis of  $T$  (otherwise  $(B' + e) \cap (T_2 \cup \bar{E}) \notin \mathcal{J}$ ), too. Consequently, choosing  $B := B' - f$  proves  $\{e, f\} \in A$ , a contradiction. So  $G$  is connected and this proves Claim 3.2 and so Claim 3. Finally, it remains to prove:

**Claim 4.** The coefficient  $\bar{a}$  of  $y$  has the correct value.

*Proof of Claim 4.* By Claim 2 we know that  $\bar{a} = 0$  if  $\alpha_{1,k}(T) = 0$ . So, e.g., in the cases  $T \subseteq \bar{E}$  or  $\bar{E} \subseteq T$  the statement follows. So assume  $T \not\subseteq \bar{E}$ ,  $\bar{E} \not\subseteq T$  and  $\alpha_{1,k}(T) > 0$ . Let  $B$  be a

basis of  $T \cup \bar{E}$  with  $r(B \cap (T \cap \bar{E})) = r(T \cap \bar{E})$  and  $r(B \cap T) = r(T)$ . Then there exists a set  $S \subseteq B \setminus \bar{E} \neq \emptyset$  so that  $B' := (B \setminus S) \cup (\bar{E} \setminus B)$  is also a basis of  $T \cup \bar{E}$ . For this note that

$$\begin{aligned} |S| &= |\bar{E} \setminus B| = |\bar{E} \setminus T| - |B \setminus T| \\ &= |B \cap T| + |\bar{E} \setminus T| - |B| = r(T) + |\bar{E} \setminus T| - r(T \cup \bar{E}) = \alpha_{1,k}(T) > 0. \end{aligned}$$

In particular,  $y(B) = 0$  because  $B$  does not contain all elements of  $\bar{E}$ . Thus, with  $\tilde{a} := a(e)$  being the value of all coefficients  $a(e)$ ,  $e \in T$ , we get that  $B, B'$  are roots of  $x(T) + \alpha_{1,k}(T)y \leq r(T)$  (thus also of  $a^T x + \bar{a}y \leq b$ ),

$$\sum_{e \in B \cap T} a(e) + \underbrace{\bar{a}y(B)}_{=0} = \sum_{e \in B' \cap T} a(e) + \underbrace{\bar{a}y(B')}_{=1}$$

so  $\bar{a} = |S| \cdot \tilde{a} = \alpha_{1,k}(T)\tilde{a}$ . Consequently Claim 4 follows. In total, this proves that  $a^T x + \bar{a}y \leq b$  is a positive multiple (with factor  $\tilde{a}$ ) of  $x(T) + \alpha_{1,k}(T)y \leq r(T)$ .  $\square$

The following theorem summarises the results on the facetial structure of  $P_M^{\bar{E},k}$ .

**Theorem 31.** *All facet defining inequalities of  $P_M^{\bar{E},k}$  are given by (11) and (12),  $x(e) \geq 0, e \in E \setminus \bar{E}, y \geq 0$  and all extended rank inequalities (14)*

$$\begin{aligned} x(T) + \alpha_{1,k}(T)y \leq r(T), \quad T \subseteq E, \quad T \text{ closed, } (\bar{E}, \{k\})\text{-non-separable and} \\ T \text{ satisfies property } (\mathbf{P1}^k) \text{ or } (\mathbf{P2}^k). \end{aligned}$$

## 4 Separation

By the well-known ‘‘optimisation equals separation’’ result [15] we know that the separation problem for the polytope  $P_M^{\bar{E},K}$  can be solved in polynomial time because the corresponding optimisation problem can be solved in polynomial time. But the proof of the next result shows how to separate the extended rank inequalities more directly. Note, all inequalities of the standard linearisation (3)-(7) as well as the non-negativity constraints (8) can be separated in polynomial time by complete enumeration.

**Theorem 32.** *Let  $K = \{k_1, \dots, k_l\} \subset \{2, \dots, k\}$ . Given a point  $(\bar{x}, \bar{y})$  the separation problem for the extended rank inequalities (9) of  $P_M^{\bar{E},K}$  can be solved in (strongly) polynomial time if  $\bar{y}_{k_j} \geq \bar{y}_{k_{j+1}}, j = 1, \dots, l-1$ , as well as  $\bar{y}_{k_1} \leq 1, \bar{y}_{k_l} \geq 0$ .*

**Proof.** Let  $(\bar{x}, \bar{y})$  be a point that satisfies all requirements. Then we want to find a set  $T \subseteq E$  such that (9) evaluated for  $(\bar{x}, \bar{y})$  is violated or we want to prove that there does not exist such a set  $T$ . Therefore we define the function  $d: \mathcal{P}(E) \rightarrow \mathbb{R}$ ,

$$d(T) = r(T) - \bar{x}(T) - \sum_{j=1}^l \alpha_{k_{j-1}+1, k_j}(T) \bar{y}_{k_j}.$$

Our separation problem is then equivalent to the problem to decide whether the optimal value of

$$\min_{T \subseteq E} d(T)$$

is negative or not. Using **(A4)** we can write  $d(T)$  as:

$$\begin{aligned} d(T) &= r(T) \cdot \underbrace{(1 - \bar{y}_{k_1})}_{\geq 0} + \sum_{j=1}^{l-1} (r(T \cup \bar{E}_{k_j}) \cdot \underbrace{(\bar{y}_{k_j} - \bar{y}_{k_{j+1}})}_{\geq 0}) + r(T \cup \bar{E}_{k_l}) \cdot \underbrace{\bar{y}_{k_l}}_{\geq 0} \\ &\quad - \bar{x}(T) - |\bar{E}_{k_1} \setminus T| \bar{y}_{k_1} - \sum_{j=2}^l (|\{e_m : m \in \{k_{j-1} + 1, \dots, k_j\}\} \setminus T| \cdot \bar{y}_{k_j}). \end{aligned}$$






Figure 4: Visualisation of the four different graphs in Examples 33–35.

So,  $d(T)$  is submodular by the submodularity of the rank function  $r(\cdot)$  and our separation problem reduces to minimising the submodular function  $d(T)$ , which can be done in strongly polynomial time [11, 18], e. g., using a combinatorial algorithm [22, 23].  $\square$

## 5 Extensions and Future Work

A natural question is if it is possible to extend the previous results to related problems. For instance, Edmonds [7] also provided a complete description for the polytope that is associated with the intersection of two matroids. However, the following example shows that even in the case with exactly one additional quadratic monomial there exist facet defining inequalities  $a^T x + \bar{a}y \leq b$  of the associated linearised polytope that are not a positive multiple of the standard linearisation but with  $a \geq 0$  and  $\bar{a} < 0$ . Thus, it might be necessary to extend the line of arguments in Section 2 significantly.

**Example 33.** We consider the directed graph  $G = (V, E)$  in Fig. 4 (I) with four nodes and four arcs  $E = \{e_1, e_2, f_1, f_2\}$  and the branching problem [6] (intersection of two matroids) with  $\bar{E} = \{e_1, e_2\}$ . Then one can check by direct computations that the inequality  $x(f_1) + x(f_2) + x(e_1) + x(e_2) - y \leq 2$  is facet defining for the associated linearised polytope although the coefficient of  $y_{e_1 e_2}$  is  $-1$ . Note, the inequality above is of the form

$$x(T) + x(e_1) + x(e_2) - y_{e_1 e_2} \leq r(T),$$

with  $T \subseteq E, r(T) = r(T + e_1) = r(T + e_2) = r(T + e_1 + e_2) - 1$  ( $r$  denotes the minimum of the rank of  $T$  for  $M_1$  and  $M_2$ ). Furthermore, considering the branching problem in Fig. 4 (II) the inequality  $2x(e_1) + x(e_2) + x(f_1) + x(f_2) - y_{e_1 e_2} \leq 2$  is facet defining. So, also coefficients of the  $x$ -variables larger than one may appear. This is a large contrast to the results by Edmonds for the polytope associated with the intersection of two matroids [7]. Indeed, given two matroids  $M_1 = (E, \mathcal{J}_1), M_2 = (E, \mathcal{J}_2)$  with rank functions  $r_i: E \rightarrow \mathbb{N}_0, i = 1, 2$ , the convex hull of all sets  $I \subseteq E$  with  $I \in \mathcal{J}_1, I \in \mathcal{J}_2$  can be described by the non-negativity constraints and the rank inequalities for the two single matroids [7].

Furthermore, it is interesting to extend the results to two or more monomials that are not necessarily nested. But again, this might lead to facet defining inequalities containing variables with negative coefficients apart from the standard linearisation. We start with an example where the two monomials are in pairwise conflict.

**Example 34.** We consider the undirected graph  $G = (V, E)$  in Fig. 4 (III) with four nodes and five edges  $E = \{f_1, e_1, e_2, e_3, e_4\}$  and the optimisation problem over the graphical matroid with the two additional monomials  $x(e_1)x(e_2), x(e_3)x(e_4)$ . Then the inequality  $x(f_1) + y_{e_1 e_2} + x(e_3) + x(e_4) - y_{e_3 e_4} \leq 2$  is facet defining. So one of the coefficients of the linearised variables is negative although the inequality does not belong to the standard linearisation in this case.

One might think that the problem in Example 34 arises because  $\{e_i: i = 1, \dots, 4\} \notin \mathcal{J}$ . But even if the union of all elements contained in the additional monomials is an independent set and none of the elements  $e \in E$  appears in more than one of these non-linear monomials, facet

defining inequalities can contain variables with negative coefficients apart from the standard linearisation.

**Example 35.** We consider the undirected graph  $G = (V, E)$  in Fig. 4 (IV) with six nodes and set of edges  $E = \{f_1, f_2, f_3, e_1, e_2, e_3, e_4\}$  and the optimisation problem over the graphical matroid with the two additional monomials  $x(e_1)x(e_2)$ ,  $x(e_3)x(e_4)$  and  $\{e_1, \dots, e_4\} \in \mathcal{J}$ . Then the inequality  $x(f_1) + x(f_2) + x(f_3) + y_{e_1e_2} + x(e_3) + x(e_4) - y_{e_3e_4} \leq 4$  is facet defining for the associated polytope.

Let, in general, a matroid  $M = (E, \mathcal{J})$  be given and let  $x(e_1)x(e_2)$ ,  $x(e_3)x(e_4)$ ,  $\{e_1, e_2\} \cap \{e_3, e_4\} = \emptyset$ , denote the two additional quadratic monomials with associated variables  $y_{e_1e_2}, y_{e_3e_4}$ . Then inequalities

$$x(T) + y_{e_1e_2} + x(e_3) + x(e_4) - y_{e_3e_4} \leq r(T) + 1,$$

$T \subset E \setminus \{e_1, e_2, e_3, e_4\}, r(T + e_3 + e_4) > r(T) + 1, (T + e_1 + e_2 + e_i) \notin \mathcal{J}, i = 3, 4$ , are valid for the associated linearised polytope. The facet defining inequalities in Examples 34 and 35 are exactly of this type.

Extending our results to problems with objective functions with more than one non-linear monomial, a good starting point might be to consider the three quadratic monomials  $x(e_1)x(e_2)$ ,  $x(e_1)x(e_3)$ ,  $x(e_2)x(e_3)$  on the three elements  $e_1, e_2, e_3 \in E$  with  $\{e_1, e_2, e_3\} \in \mathcal{J}$  or  $\{e_1, e_2, e_3\}$  forming a circuit. In both mentioned cases the monomials then fulfil some kind of up- and downwards compatibility, see [5]. Furthermore, it remains for future work to extend the results to (integral) polymatroids [7], see also [11, 23]. Here, the setting might be quite different because the variables are not necessarily binary.

## Appendix

In the following we present several detailed, rather technical proofs of the results stated in the main part of this paper.

**Proof (of Observation 5).** Let  $e \in \text{cl}(T)$ , then  $r(T + e) = r(T)$ .

$$\begin{aligned} r(T + S) + r(T) &\stackrel{(R2)}{\leq} r(T + S + e) + r((T + S) \cap (T + e)) \stackrel{(R3)}{\leq} r(T + S) + r(T + e) \\ &= r(T + S) + r(T). \end{aligned} \quad \square$$

**Proof (of Observation 7).** Let  $T \subseteq E$ .

**(A1):** If  $e_i \in T, i \in \{1, \dots, k\}$ , then  $\alpha_i(T) = 0 + r(T + \bar{E}_i) - r(T + \bar{E}_i) = 0$ .

**(A2):** Let  $i \in \{1, \dots, k\}$ . If  $r(T + \bar{E}_i) = r(T + \bar{E}_{i-1})$ , the statement is clear by  $|\{e_i\} \setminus T| \in \{0, 1\}$ . If  $r(T + \bar{E}_i) = r(T + \bar{E}_{i-1}) + 1$ , this implies  $e_i \notin T$  and so  $\alpha_i(T) = 0$ .

**(A3):** Let  $i \in \{1, \dots, k\}$  and  $j \in \mathbb{N}, j > i$ . Then  $\alpha_j(T) = |\{e_j\} \setminus T| + r(T + \bar{E}_{j-1}) - r(T + \bar{E}_j) = |\{e_j\} \setminus (T + e_i)| + r(T + e_i + \bar{E}_{j-1}) - r(T + e_i + \bar{E}_j) = \alpha_j(T + e_i)$ .

**(A4):** Let  $j \in \{1, \dots, k\}$ . Then  $\alpha_{1,j}(T) = \sum_{i=1}^j (|\{e_i\} \setminus T| + r(T + \bar{E}_{i-1}) - r(T + \bar{E}_i)) = |\bar{E}_j \setminus T| + r(T + \bar{E}_0) - r(T + \bar{E}_j) = |\bar{E}_j \setminus T| + r(T) - r(T + \bar{E}_j)$ .

**(A5):** Fix  $i \in \{1, \dots, k\}$ . If  $e \in T$ , the statement is clear. So let  $e \in (\text{cl}(T) \setminus T) \setminus \{e_i\}$ . Then  $\alpha_i(T) = |\{e_i\} \setminus T| + r(T + \bar{E}_{i-1}) - r(T + \bar{E}_i) = |\{e_i\} \setminus (T + e)| + r(T + e + \bar{E}_{i-1}) - r(T + e + \bar{E}_i) = \alpha_i(T + e)$  by Observation 5.

**(A6):** Let  $i \in \{1, \dots, k\}$  and  $e_i \in \text{cl}(T) \setminus T$ . Then Observation 5 and **(A1)** show  $\alpha_i(T) = |\{e_i\} \setminus T| + r(T + \bar{E}_{i-1}) - r(T + \bar{E}_i) = 1 + r(T + e_i + \bar{E}_{i-1}) - r(T + \bar{E}_i) = 1 = \alpha_i(T + e_i) + 1$ .

□

**Proof (of Lemma 14).** Let  $a^T x + \alpha^T y \leq b$  be a facet defining inequality of  $P_M^{\bar{E}, K}$ ,  $K = \{2, \dots, k\}$ , satisfying the requirements. Let  $S_i$  denote a root of  $a^T x + \alpha^T y \leq b$  such that  $\bar{E}_i \subseteq S_i$  and  $(i = k \vee e_{i+1} \notin S_i)$ . For  $m = 2$  the desired statement follows directly from the fact that  $a^T x + \alpha^T y \leq b$  is not a positive multiple of (5). For  $m > 2$  we know by (6) that there exists a root  $S$  of  $a^T x + \alpha^T y \leq b$  with  $e_m \notin S$  and  $\bar{E}_{m-1} \not\subseteq S$ . So we may assume that

$$\mu := |S \cap \bar{E}_{m-1}| \text{ is maximal.} \quad (21)$$

If  $|S \cap \bar{E}_m| = |\bar{E}_m| - 2$ , then  $S$  satisfies the requirements. Otherwise there must exist two elements  $e_i, e_j \notin S$  with  $i < j < m$  and  $\bar{E}_{j-1} \setminus \{e_i\} \subseteq S$ . Let  $S_{j-1}$  be a root of  $a^T x + \alpha^T y \leq b$  with  $\bar{E}_{j-1} \subseteq S_{j-1}$  and  $e_j \notin S_{j-1}$  (exists by (4) or by (7) in the case  $j = 2$ ).

Now assume in addition that  $S$  and  $S_{j-1}$  are chosen so that

$$|S \cap \bar{E}_{m-1}| = \mu \quad \text{and} \quad |S \cap S_{j-1}| \text{ is maximal.} \quad (22)$$

1. If  $r(S) < r(S_{j-1})$ , then there exists an  $e \in S_{j-1} \setminus S$  so that  $e + S \in \mathcal{J}$ . By Observation 13 we get  $\tilde{a}(S + e, e) \leq 0 \leq \tilde{a}(S_{j-1}, e)$ . If  $e \neq e_i$ , then  $e \notin \bar{E}_j$  and so  $\tilde{a}(S_{j-1}, e) = a(e) = \tilde{a}(S + e, e)$  proving  $a(e) = 0$ . Thus  $S + e$  is also a root of  $a^T x + \alpha^T y \leq b$  contradicting (22). Otherwise we have  $\tilde{a}(S_{j-1}, e) = a(e) + \alpha_i + \dots + \alpha_{j-1} = \tilde{a}(S + e, e) = 0$ . As before  $S + e$  is also root of  $a^T x + \alpha^T y \leq b$ , a contradiction to (21).
2. If  $r(S) > r(S_{j-1})$ , then there exists an  $e \in S$  so that  $S_{j-1} + e \in \mathcal{J}$ . Certainly  $e \notin \bar{E}_j$ , so as above it follows  $a(e) = 0$ . So  $S, (S_{j-1} + e)$  contradict to assumption (22).
3. If  $r(S) = r(S_{j-1})$ , then we know by Theorem 1 that there exists an  $e \in S \setminus S_{j-1}$  so that  $S - e + e_i, S_{j-1} + e - e_i \in \mathcal{J}$ . Note,  $e \notin \bar{E}_j$ , which implies  $\tilde{a}(S, e) = a(e) = \tilde{a}(S_{j-1} + e - e_i, e)$  as well as  $\tilde{a}(S_{j-1}, e_i) = a(e_i) + \alpha_i + \dots + \alpha_{j-1} = \tilde{a}(S - e + e_i, e_i)$ . Applying Observation 13 twice then shows  $\tilde{a}(S - e + e_i, e_i) \leq \tilde{a}(S, e) = \tilde{a}(S_{j-1} + e - e_i, e) \leq \tilde{a}(S_{j-1}, e_i) = \tilde{a}(S - e + e_i, e_i)$  and so  $S - e + e_i$  is also a root of  $a^T x + \alpha^T y \leq b$ , contradicting (21). □

**Proof (of Lemma 15).** Let  $a^T x + \alpha^T y \leq b$  be a facet defining inequality of  $P_M^{\bar{E}, K}$  with  $K = \{2, \dots, k\}$  that is not a positive multiple of one of (3)–(8). Let  $S_i$  be defined as in the proof of Lemma 14. Assume, for a contradiction, the assertion is false. Then there must exist an  $m \in \{2, \dots, k\}$  and a minimal  $h \in \{m, \dots, k\}$  so that  $\alpha_m + \dots + \alpha_h < 0$  and there exists a root  $S$  of  $a^T x + \alpha^T y \leq b$  satisfying  $\bar{E}_m \setminus \{e_j, e_m\} = S \cap \bar{E}_m$ ,  $j < m$ , and  $h = k$  or  $e_{h+1} \notin S$ . Furthermore, by (4) there exists a root  $S_h$  with  $\bar{E}_h \subseteq S_h$  and  $h = k$  or  $e_{h+1} \notin S$ . We may assume that

$$|S \cap S_h| \text{ is maximal.} \quad (23)$$

1. If  $r(S) < r(S_h)$ , then there exists an  $e \in S_h \setminus S$  so that  $S + e \in \mathcal{J}$ . We consider four cases depending on  $e$  and apply Observation 13.
  - If  $e \notin \bar{E}_h$ , then  $\tilde{a}(S + e, e) = a(e)$  and  $\tilde{a}(S_h, e) = a(e)$ . Observation 13 shows that  $a(e) = 0$ , so  $S + e$  is also a root of  $a^T x + \alpha^T y \leq b$ , contradicting (23).
  - If  $e = e_m$ , then  $0 \geq \tilde{a}(S + e_m, e_m) = a(e_m)$  and  $0 \leq \tilde{a}(S_h, e_m) = a(e_m) + \alpha_m + \dots + \alpha_h$ , hence  $\alpha_m + \dots + \alpha_h \geq 0$ , a contradiction.
  - If  $e = e_j$ , then we similarly get  $0 \geq \tilde{a}(S + e_j, e_j) = a(e_j) + \alpha_j + \dots + \alpha_{m-1}$  and  $0 \leq \tilde{a}(S_h, e_j) = a(e_j) + \alpha_j + \dots + \alpha_h$ , hence  $\alpha_m + \dots + \alpha_h \geq 0$ , a contradiction.
  - If  $e = e_i$ ,  $m < i \leq h$ , then  $0 \geq \tilde{a}(S + e_i, e_i) = a(e_i)$  and  $0 \leq \tilde{a}(S_h, e_i) = a(e_i) + \alpha_i + \dots + \alpha_h$ , hence we derive  $\alpha_i + \dots + \alpha_h \geq 0$ . Because  $e_i \notin S$  we know by minimality of  $h$  that  $\alpha_m + \dots + \alpha_{i-1} \geq 0$ . Together we have  $\alpha_m + \dots + \alpha_h \geq 0$ , a contradiction.

2. If  $r(S) > r(S_h)$ , then there exists an  $e \in S \setminus S_h$  so that  $S_h + e \in \mathcal{J}$ . Note that  $e \notin \bar{E}_{h+1}$  (if  $h < k$ ), so by Observation 13  $0 \geq \tilde{a}(S_h + e, e) = a(e) \geq 0$ . Therefore  $S_h + e$  is also a root of  $a^T x + \alpha^T y \leq b$ , contradicting (23).
3. If  $r(S) = r(S_h)$ , then by Theorem 1 there exists an  $e \in S \setminus S_h$  so that  $S - e + e_m, S_h + e - e_m \in \mathcal{J}$ . By assumption  $h = k \vee e \notin \bar{E}_{h+1}$ , thus we have  $\tilde{a}(S, e) = a(e) = \tilde{a}(S_h + e - e_m, e)$  as well as  $\tilde{a}(S_h, e_m) = a(e_m) + \alpha_m + \dots + \alpha_h$  and  $\tilde{a}(S - e + e_m, e_m) = a(e_m)$ . We know by Observation 13 that  $a(e_m) = \tilde{a}(S - e + e_m, e_m) \leq \tilde{a}(S, e) = \tilde{a}(S_h - e_m + e, e) \leq \tilde{a}(S_h, e_m) = a(e_m) + \alpha_m + \dots + \alpha_h$ , so  $\alpha_m + \dots + \alpha_h \geq 0$ , a contradiction.  $\square$

**Proof (of Lemma 16).** Let  $a^T x + \alpha^T y \leq b$  be a facet defining inequality of  $P_M^{\bar{E}, K}$ ,  $K = \{2, \dots, k\}$ , satisfying the requirements defined above. Let  $S_i$  be defined as in the proof of Lemma 14. Fix some  $m \in \{2, \dots, k\}$  and suppose that the claim is false. Then there exists a maximal  $h \in \{m, \dots, k\}$  so that  $\alpha_m + \dots + \alpha_h < 0$ . Choose a root  $S$  with  $\bar{E}_m \setminus \{e_j, e_m\} = S \cap \bar{E}_m$  (exists by Lemma 14) and a root  $S_h$  with  $\bar{E}_h \subseteq S_h$ ,  $h = k \vee e_{h+1} \notin S_h$  (exists by (3), (4)) so that  $|S \cap S_h|$  is maximal. If  $e_{h+1} \notin S$  or  $h = k$ , then the claim follows from Lemma 15, so we may assume  $e_{h+1} \in S$ . We use similar arguments as in the previous proof.

1. If  $r(S) < r(S_h)$ , then there is an  $e \in S_h \setminus S$  so that  $S + e \in \mathcal{J}$ . If additionally  $e \neq e_i$ ,  $m < i \leq h$ , then all arguments of the previous proof apply (in particular, we do not require the assumption of  $h$  being minimal). We only have to consider the case  $e = e_i$ ,  $m < i \leq h$ . Then we get  $\tilde{a}(S + e_i, e_i) = a(e_i)$  and  $\tilde{a}(S_h, e_i) = a(e_i) + \alpha_i + \dots + \alpha_h$ . So by Observation 13 we have  $\alpha_i + \dots + \alpha_h \geq 0$ . Note that  $e_i \notin S$  by construction, so we may apply Lemma 15 with this  $S$ ,  $j$  and  $m$  and  $h = i - 1$  to derive  $\alpha_m + \dots + \alpha_{i-1} \geq 0$ . Hence, we conclude that  $\alpha_m + \dots + \alpha_h \geq 0$ , a contradiction.
2. If  $r(S) > r(S_h)$ , then there is an  $e \in S \setminus S_h$  so that  $S_h + e \in \mathcal{J}$ . The case  $e \neq e_{h+1}$  works as in the previous proof, so it remains  $e = e_{h+1}$ . Then  $\tilde{a}(S_h + e_{h+1}, e_{h+1}) = a(e_{h+1}) + \alpha_{h+1} + \dots + \alpha_o$  for some  $o \in \{h + 1, \dots, k\}$  and  $\tilde{a}(S, e_{h+1}) = a(e_{h+1})$ . Therefore, applying Observation 13 yields  $\alpha_{h+1} + \dots + \alpha_o \leq 0$ . By maximality of  $h$  we know  $\alpha_m + \dots + \alpha_o \geq 0$  and can conclude  $\alpha_m + \dots + \alpha_h \geq 0$ , a contradiction.
3. If  $r(S) = r(S_h)$ , then by Theorem 1 there exists an  $e \in S \setminus S_h$  so that  $S - e + e_m, S_h + e - e_m \in \mathcal{J}$ . Note that  $e \notin \bar{E}_h$ . We have  $\tilde{a}(S, e) = a(e) = \tilde{a}(S_h - e_m + e, e)$ ,  $\tilde{a}(S_h, e_m) = a(e_m) + \alpha_m + \dots + \alpha_h$  and  $\tilde{a}(S - e + e_m, e_m) = a(e_m)$ . Observation 13 shows  $a(e_m) = \tilde{a}(S - e + e_m, e_m) \leq \tilde{a}(S, e) = \tilde{a}(S_h + e - e_m, e) \leq \tilde{a}(S_h, e_m) = a(e_m) + \alpha_m + \dots + \alpha_h$ . This implies  $\alpha_m + \dots + \alpha_h \geq 0$ , a contradiction.  $\square$

**Proof (of Observation 21).** The polytope  $P_M^{\bar{E}, K}$  is full-dimensional with dimension  $|E| + l$ . We prove this by explicitly constructing  $|E| + l + 1$  independent sets whose incidence vectors are affinely independent. We use  $\emptyset \in \mathcal{J}$ ,  $\{e\} \in \mathcal{J}$  for all  $e \in E$  and  $\{e_1, \dots, e_{k_j}\} \in \mathcal{J}$ ,  $j = \{1, \dots, l\}$ . Similarly for the facet defining inequalities, we explicitly present the  $|E| + l$  respective sets.

- $y_{k_l} \geq 0$  (3): We can use the independent sets of the dimension proof except for  $\{e_1, \dots, e_{k_l}\}$ .
- $y_{k_j} - y_{k_{j-1}} \leq 0$ ,  $j = 2, \dots, l$  (4): We can use the independent sets of the dimension proof except for  $\{e_1, \dots, e_{k_{j-1}}\}$ .
- $\sum_{i=1}^{k_1} x(e_i) - y_{k_1} \leq k_1 - 1$  (5): We use  $\bar{E}_{k_j}$ ,  $j = 1, \dots, l$ , and  $\bar{E}_{k_1} \setminus \{e\}$  for all  $e \in \bar{E}_{k_1}$ . By assumptions (1) we know that for each  $e \in E \setminus \bar{E}_{k_1}$  there exists an  $f \in \bar{E}_{k_1}$  such that  $(\bar{E}_{k_1} + e) - f \in \mathcal{J}$ . So we use one such set for each  $e \in \bar{E} \setminus \bar{E}_{k_1}$ .

- $\sum_{i=k_{j-1}+1}^{k_j} x(e_i) + y_{k_{j-1}} - y_{k_j} \leq k_j - k_{j-1}, j = 2, \dots, l$ , (6): We use the independent sets  $\bar{E}_{k_j} \setminus \bar{E}_{k_{j-1}}, (\bar{E}_{k_j} \setminus \bar{E}_{k_{j-1}}) + e, e \in \bar{E}_{k_{j-1}}$ , and  $(\bar{E}_{k_j} \setminus \bar{E}_{k_{j-1}}) + \bar{E}_{k_m}, m = 1, \dots, j-2$ , as well as  $\bar{E}_{k_j} - e, e \in \bar{E}_{k_j} \setminus \bar{E}_{k_{j-1}}$ . For each  $e \in E \setminus \bar{E}_{k_j}$  there exists an  $f \in \bar{E}_{k_j}$  such that  $\bar{E}_{k_j} + e - f \in \mathcal{J}$ . We use one such set for each  $e \in E \setminus \bar{E}_{k_j}$ . Furthermore we take the sets  $\bar{E}_{k_m}, m = j, \dots, l$ .
- $-x(e_i) + y_{k_j} \leq 0, j = 1, \dots, l, i = k_{j-1} + 1, \dots, k_j$ , (7): We can use the independent sets of the dimension proof except for  $\{e_i\}$ .
- $x(e) \geq 0, e \in E \setminus \bar{E}$  (8): We can use the independent sets of the dimension proof except for  $\{e\}$ .

Further note, constraints  $-x(e) \leq 0, e \in \bar{E}$ , are implied by (3), (4) and (7).  $\square$

**Proof (of Observation 26). 1:** Let  $T \subseteq \bar{E}$ . Then  $T$  is  $(\bar{E}, \{k\})$ -separable with  $T_1 = T \setminus \{e\} \neq \emptyset$  and  $T_2 = \{e\}$  for each  $e \in T$ .

**2:** Let  $T \subseteq E, \bar{E} \cap T \neq \emptyset$  and  $\alpha_{1,k}(T \setminus \bar{E}) = 0$ . If  $T \subseteq \bar{E}$ , we are in case 1. Otherwise  $\alpha_{1,k}(T \setminus \bar{E}) = 0$  and **(A4)** imply  $r(T \setminus \bar{E}) + |\bar{E}| = r(T \cup \bar{E})$ , hence  $r(T \setminus \bar{E}) + |T \cap \bar{E}| = r(T)$  (by  $r(T) \leq r(T \setminus \bar{E}) + r(T \cap \bar{E}) = r(T \cup \bar{E}) + r(T \cap \bar{E}) - |\bar{E}| \leq r(T)$ ). Furthermore, this implies  $\alpha_{1,k}(T) = 0$  as well. Consequently  $T$  is  $(\bar{E}, \{k\})$ -separable with  $T_1 = T \setminus \bar{E} \neq \emptyset$  and  $T_2 = T \cap \bar{E}$ .  $\square$

**Proof (of Observation 28).** First we consider a rank inequality with set  $T = \{e\}, e \in \bar{E}$ , in the case  $|\bar{E}| = 2$ .  $T$  is closed and  $(\bar{E}, \{k\})$ -non-separable but it can be derived by adding the two constraints  $x(\bar{E}) - y \leq 1$  and  $-x(\bar{e}) + y \leq 0$  for  $\bar{e} = \bar{E} \setminus \{e\}$ .

Now assume that  $T$  is closed and  $(\bar{E}, \{k\})$ -non-separable with  $\bar{E} \subseteq T$  and  $\alpha_{1,k}(T \setminus \bar{E}) \leq 1$ . By Observation 26 there remains to consider the case  $\alpha_{1,k}(T \setminus \bar{E}) = 1$ . Then we can derive (14) for  $T$  by adding (11) and (14) for  $T' := T \setminus \bar{E}$ .  $\square$

**Proof (of Observation 29).** Let  $T \subseteq E$  be a closed set with  $\bar{E} \subseteq T$  and  $\alpha_{1,k}(T \setminus \bar{E}) = r(T \setminus \bar{E}) + |\bar{E}| - r(T) \geq 2$  (by **(A4)**). Then we take a basis  $B$  of  $T \setminus \bar{E}$  and extend  $B$  to a basis of  $T$  by adding elements of  $\bar{E}$ . By the assumption on  $\alpha_{1,k}(T \setminus \bar{E})$  at most  $r(T) - r(T \setminus \bar{E}) \leq |\bar{E}| - 2$  elements are added to  $B$ .  $\square$

**Proof (of Claim 1 in the proof of Lemma 30).** We want to prove that inequalities  $x(T) + \alpha_{1,k}(T)y \leq r(T)$  for a closed and  $(\bar{E}, \{k\})$ -non-separable set  $T \subseteq E$  satisfying **(P1<sup>k</sup>)** or **(P2<sup>k</sup>)** are not implied by (11)–(13). For this, we determine in each of the cases an independent set  $J \in \mathcal{J}$  such that  $J$  is a root of (14), but  $J$  is not a root of the considered other constraint out of (11)–(13).

- $x(\bar{E}) - y \leq |\bar{E}| - 1$  (11): We consider three cases.
  - $|T \cap \bar{E}| < |\bar{E}| - 1$ : Let  $B$  be a basis of  $T$ . Then we set  $J := B$ .
  - $|T \cap \bar{E}| = |\bar{E}| - 1$ : Observation 26 and **(P1<sup>k</sup>)** imply  $T \not\subseteq \bar{E}$  and  $|T| \geq 2$ . Choose  $f \in T \cap \bar{E}$ , then  $r(T - f) = r(T)$  because otherwise  $T$  would be  $(\bar{E}, \{k\})$ -separable with  $T_1 = \{f\}$  and  $T_2 = T - f$ . Let  $B$  be a basis of  $T - f$ , then  $B$  is also a basis of  $T$  with  $|B \cap \bar{E}| \leq |\bar{E}| - 2$ . So we set  $J := B$ .
  - $\bar{E} \subseteq T$ : In this case, by **(P2<sup>k</sup>)** and Observation 29, there exists a basis  $B$  of  $T$  such that  $|B \cap \bar{E}| \leq |\bar{E}| - 2$ . Then we set  $J := B$ .
- $y - x(e) \leq 0, e \in \bar{E}$  (12): We consider four cases:

$e \in T, \bar{E} \not\subseteq T$ : Let  $B$  be a basis of  $T$  with  $e \in T$ . Then we have that  $y(B) = 0, \chi_B(e) = 1$  and  $\sum_{e \in T} \chi_B(e) + 0 = r(T)$ . So we use  $J := B$ .

$e \in T, \bar{E} \subseteq T$ : In this case we know by **(P2<sup>k</sup>)** and Observation 29 that there exists a basis  $B$  of  $T$  with  $|B \cap \bar{E}| \leq |\bar{E}| - 2$ . If  $e \in B$ , we set  $J := B$ , otherwise there exists an  $f \in B$  such that  $B + e - f \in \mathcal{J}$  is a basis of  $T$  and so we use  $J := B + e - f$  (note  $y(B + e - f) = 0$ ).

$e \notin T, \bar{E} \not\subseteq T + e$ : Let  $B$  be a basis of  $T$ . Then  $B + e =: J \in \mathcal{J}$  because  $T$  is closed.

$e \notin T, \bar{E} \subseteq T + e$ : Observation 26 and **(P1<sup>k</sup>)** imply  $T \not\subseteq \bar{E}$ , in particular  $|T| \geq 2$ . Choose  $f \in T \cap \bar{E}$ , then  $r(T - f) = r(T)$  because otherwise  $T$  would be  $(\bar{E}, \{k\})$ -separable with  $T_1 = \{f\}$  and  $T_2 = T - f$ . Let  $B$  be a basis of  $T - f$ , then  $B$  is also a basis of  $T$ , and because  $T$  is closed we can use  $J := B + e$ , which is a basis of  $T + e$  with  $\bar{E} \not\subseteq J$  by the choice of  $f$ .

- $-x(e) \leq 0, e \in E \setminus \bar{E}$  (13): First, we consider the case  $e \notin T$ . Let  $B$  be a basis of  $T$ . By  $T$  closed we know  $B + e \in \mathcal{J}$  and set  $J := B + e$ . Second, if  $e \in T$ , there exists a basis  $B$  of  $T$  with  $e \in B$  and we can use  $J := B$ .
- $-y \leq 0$  (13): We consider two cases. If  $\bar{E} \subseteq T$ , there exists a basis  $B$  of  $T$  with  $\bar{E} \subseteq B$  (note  $\bar{E} \in \mathcal{J}$ ) and so we use  $J := B$ . If, otherwise,  $\bar{E} \not\subseteq T$ , let  $B$  be a basis of  $T \cup \bar{E}$  with  $\bar{E} \subset B$ . Then by  $\sum_{e \in T} \chi_B(e) + |\bar{E} \setminus T| = |B \cap T| + |\bar{E} \setminus T| = |((B \setminus \bar{E}) \cap T) \cup \bar{E}| = r(B) = r(T \cup \bar{E})$  we get  $\sum_{e \in T} \chi_B(e) + r(T) + |\bar{E} \setminus T| - r(T \cup \bar{E}) = r(T)$ . So  $B$  is a root of  $x(T) + \alpha_{1,k}(T)y \leq r(T)$  and we can set  $J := B$ .

So  $x(T) + \alpha_{1,k}(T)y \leq r(T)$  is not a positive multiple of one of (11)–(13). This proves Claim 1 in Lemma 30.  $\square$

## References

- [1] Yael Berstein, Jon Lee, Hugo Maruri-Aguilar, Shmuel Onn, Eva Riccomagno, Robert Weismantel, and Henry Wynn. Nonlinear matroid optimization and experimental design. *SIAM Journal on Discrete Mathematics*, 22(3):901–919, 2008.
- [2] Richard A. Brualdi. Comments on bases in dependence structures. *Bulletin of the Australian Mathematical Society*, 1(2):161–167, 8 1969.
- [3] Christoph Buchheim and Laura Klein. The spanning tree problem with one quadratic term. In Kamiel Cornelissen, Ruben Hoeksma, Johann Hurink, and Bodo Manthey, editors, *CTW*, volume WP 13-01 of *CTIT Workshop Proceedings*, pages 31–34, 2013.
- [4] Christoph Buchheim and Laura Klein. Combinatorial optimization with one quadratic term: Spanning trees and forests. *Discrete Applied Mathematics*, 177:34–52, 2014.
- [5] Christoph Buchheim and Giovanni Rinaldi. Efficient reduction of polynomial zero-one optimization to the quadratic case. *SIAM Journal on Optimization*, 18(4):1398–1413, 2008.
- [6] Jack Edmonds. Optimum Branchings. *Journal of Research of the National Bureau of Standards*, 71B:233–240, 1967.
- [7] Jack Edmonds. Submodular functions, matroids, and certain polyhedra. *Combinatorial Structures and Their Applications*, pages 69–87, 1970.
- [8] Jack Edmonds. Matroids and the greedy algorithm. *Mathematical Programming*, 1(1):127–136, 1971.

- [9] Anja Fischer and Frank Fischer. Complete description for the spanning tree problem with one linearised quadratic term. *Operations Research Letters*, 41(6):701–705, 2013.
- [10] R. Fortet. L’Algèbre de Boole et ses applications en recherche opérationnelle. *Cahiers du Centre d’Etudes de Recherche Opérationnelle*, 4:5–36, 1959.
- [11] Satoru Fujishige. *Submodular Functions and Optimization*. Elsevier, 2005.
- [12] Fred Glover and Eugene Woolsey. Further reduction of zero-one polynomial programming problems to zero-one linear programming problems. *Operations Research*, 21(1):156–161, 1973.
- [13] Fred Glover and Eugene Woolsey. Converting the 0-1 polynomial programming problem to a 0-1 linear program. *Operations Research*, 22(1):180–182, 1974.
- [14] Martin Grötschel. *Polyedrische Charakterisierungen kombinatorischer Optimierungsprobleme*, volume 36 of *Mathematical Systems in Economics*. Verlag Anton Hain, Meisenheim am Glan, 1977.
- [15] Martin Grötschel, László Lovász, and Alexander Schrijver. *Geometric Algorithms and Combinatorial Optimization*, volume 2 of *Algorithms and Combinatorics*. Springer, second corrected edition, 1993.
- [16] Jon Lee and Janny Leung. On the boolean quadric forest polytope. *INFOR*, 42(2):125–141, 2004.
- [17] Jesús A. De Loera, David C. Haws, Jon Lee, and Allison O’Hair. Computation in multicriteria matroid optimization. *Journal of Experimental Algorithmics*, 14:8:1.8–8:1.33, January 2010.
- [18] S. Thomas McCormick. *Handbook on Discrete Optimization*, chapter Submodular Function Minimization, pages 321–391. Elsevier, 2006.
- [19] Shmuel Onn. Convex matroid optimization. *SIAM Journal on Discrete Mathematics*, 17(2):249–253, February 2004.
- [20] James G. Oxley. *Matroid theory*. Oxford University Press, 1992.
- [21] Manfred Padberg. The boolean quadric polytope: Some characteristics, facets and relatives. *Mathematical Programming*, 45(1-3):139–172, 1989.
- [22] Alexander Schrijver. A combinatorial algorithm minimizing submodular functions in strongly polynomial time. *Journal of Combinatorial Theory Ser. B*, 80(2):346–355, November 2000.
- [23] Alexander Schrijver. *Combinatorial Optimization – Polyhedra and Efficiency*. Springer, 2003.
- [24] Hanif D. Sherali and Warren P. Adams. A hierarchy of relaxations between the continuous and convex hull representations for zero-one programming problems. *SIAM Journal on Discrete Mathematics*, 3(3):411–430, 1990.
- [25] Hanif D. Sherali and Warren P. Adams. *Reformulation-Linearization Technique for Solving Discrete and Continuous Nonconvex Problems*. Springer, 1998.
- [26] Dominic J. A. Welsh. *Matroid theory*. Academic Press, London, 1976.