

On the generalized ϑ -number and related problems for highly symmetric graphs

Lennart Sinjorgo ^{*} Renata Sotirov [†]

Abstract

This paper is an in-depth analysis of the generalized ϑ -number of a graph. The generalized ϑ -number, $\vartheta_k(G)$, serves as a bound for both the k -multichromatic number of a graph and the maximum k -colorable subgraph problem. We present various properties of $\vartheta_k(G)$, such as that the series $(\vartheta_k(G))_k$ is increasing and bounded above by the order of the graph G . We study $\vartheta_k(G)$ when G is the graph strong, disjunction and Cartesian product of two graphs. We provide closed form expressions for the generalized ϑ -number on several classes of graphs including the Kneser graphs, cycle graphs, strongly regular graphs and orthogonality graphs. Our paper provides bounds on the product and sum of the k -multichromatic number of a graph and its complement graph, as well as lower bounds for the k -multichromatic number on several graph classes including the Hamming and Johnson graphs.

Keywords k -multicoloring, k -colorable subgraph problem, generalized ϑ -number, Johnson graphs, Hamming graphs, strongly regular graphs.

AMS subject classifications. 90C22, 05C15, 90C35

1 Introduction

The k -multicoloring of a graph is to assign k distinct colors to each vertex in the graph such that two adjacent vertices are assigned disjoint sets of colors. The k -multicoloring is also known as k -fold coloring, n -tuple coloring or simply multicoloring. We denote by $\chi_k(G)$ the minimum number of colors needed for a valid k -multicoloring of a graph G , and refer to it as the k -th chromatic number of G or the multichromatic number of G . Multicoloring seems to have been independently introduced by Hilton et al. [40] and Stahl [75]. The k -multicoloring is a generalization of the well known standard graph coloring. Namely, $\chi(G) := \chi_1(G)$ is known as the chromatic number of a graph G . Not surprisingly, multicoloring finds applications in comparable areas, such as job scheduling [27, 38], channel assignment in cellular networks [63] and register allocation in computers [13]. There exist several results on $\chi_k(G)$ for specific classes of graphs. In particular, Lin [50] and Lin et al. [51] consider multicoloring the Mycielskian of graphs, Ren and Bu [68] study multicoloring of planar graphs while Marx [58] proves that the multicoloring problem is strongly NP-hard in binary trees. Cranston and Rabern [18] show that for any planar graph G , $\chi_2(G) \leq 9$, which can be seen as an extension of the famous four-color theorem by Appel and Haken [4].

The maximum k -colorable subgraph ($MkCS$) problem is to find the largest induced subgraph in a given graph that can be colored with k colors so that no two adjacent vertices have the same color. When $k = 1$, the $MkCS$ problem reduces to the well known maximum stable set problem. The $MkCS$ problem is one of the NP-complete problems considered by Lewis and Yannakakis [49]. We denote by $\alpha_k(G)$ the number of vertices in the maximum k -colorable subgraph of G , and by $\omega_k(G)$ the size of the largest induced subgraph that can be covered with k cliques. When $k = 1$, the graph parameter $\omega(G) := \omega_1(G)$ is known as the clique number of a graph, and $\alpha(G) := \alpha_1(G)$ as the independence number of a graph. We note that $\alpha_k(G) = \omega_k(\bar{G})$, where \bar{G} denotes the complement of G . The $MkCS$ problem has a number of applications such as channel assignment in spectrum sharing networks [47, 76], VLSI design [57] and human genetic research [22, 52]. There exist several results on $\alpha_k(G)$ for specific classes of graphs. The size of the maximum k -colorable subgraph for the Kneser graph $K(v, 2)$ is provided by Füredi and Frankl [24]. Yannakakis and Gavril [79] consider the $MkCS$ problem for chordal graphs,

^{*}CentER, Tilburg University, The Netherlands, l.m.sinjorgo@tilburguniversity.edu

[†]Department of Econometrics and OR, Tilburg University, The Netherlands, r.sotirov@uvt.nl

Addario-Berry et al. [2] study the problem for an i -triangulated graph, and Narasimhan [61] computes $\alpha_k(G)$ for circular-arc graphs and tolerance graphs.

Narasimhan and Manber [62] introduce a graph parameter $\vartheta_k(G)$ that serves as a bound for both the minimum number of colors needed for a k -multicoloring of a graph G and the number of vertices in the maximum k -colorable subgraph of G . The parameter $\vartheta_k(G)$ generalizes the concept of the famous ϑ -number that was introduced by Lovász [54] for bounding the Shannon capacity of a graph [71]. The Lovász theta number is a widely studied graph parameter see e.g., [14, 33, 35, 46, 55, 59]. The Lovász theta number provides bounds for both the clique number and the chromatic number of a graph, both of which are NP-hard to compute. The well known result that establishes the following relation $\alpha_1(G) \leq \vartheta_1(G) \leq \chi_1(\overline{G})$ or equivalently $\omega_1(G) \leq \vartheta_1(\overline{G}) \leq \chi_1(G)$ is known as the sandwich theorem [55]. The Lovász theta number can be computed in polynomial time as an semidefinite programming (SDP) problem by using interior point methods. Thus, when the clique number and chromatic number of a graph coincide such as for weakly perfect graphs, the Lovász theta function provides those quantities in polynomial time.

Despite the popularity of the Lovász theta number, the function $\vartheta_k(G)$ has received little attention in the literature. Narasimhan and Manber [62] show that $\alpha_k(G) \leq \vartheta_k(G) \leq \chi_k(\overline{G})$ or equivalently $\omega_k(G) \leq \vartheta_k(\overline{G}) \leq \chi_k(G)$. These inequalities can be seen as a generalization of the the Lovász sandwich theorem. Alizadeh [3] formulate the generalized ϑ -number using semidefinite programming. Kuryatnikova et al. [48] introduce the generalized ϑ' -number that is obtained by adding non-negativity constraints to the SDP formulation of the ϑ_k -number. The generalized ϑ -number and ϑ' -number are evaluated numerically as upper bounds for the $MkCS$ problem in [48]. The authors of [48] characterise a family of graphs for which $\vartheta_k(G)$ and $\vartheta'_k(G)$ provide tight bounds for $\alpha_k(G)$. Here, we study also a relation between $\vartheta_k(G)$ and $\chi_k(G)$, and extend many known results for the Lovász ϑ -number to the generalized ϑ -number. This paper is based on the thesis of Sinjorgo [74].

Main results and outline

This paper provides various theoretical results for $\alpha_k(G)$, $\vartheta_k(G)$ and $\chi_k(G)$. We provide numerous properties of $\vartheta_k(G)$ including results on different graph products of two graphs such as the Cartesian product, strong product and disjunction product. We show that the series $(\vartheta_k(G))_k$ is increasing towards the number of vertices in G , and that the increments of the series can be arbitrarily small. The latter result is proven by constructing a particular graph that satisfies the desired property. We also provide a closed form expression on the generalized ϑ -number for several graph classes including complete graphs, cycle graphs, complete multipartite graphs, strongly regular graphs, orthogonality graphs, the Kneser graphs and some Johnson graphs. We compute $\vartheta_k(G)$ for circulant graphs and the Johnson graphs. Our results show that $\vartheta_k(G) = k\vartheta(G)$ for the Kneser graphs and more general Johnson graphs, strongly regular graphs, cycle graphs and circulant graphs. Our paper presents lower bounds on the k -th chromatic number for all regular graphs, but also specialized bounds for the Hamming, Johnson and orthogonality graphs. We also provide bounds on the product and sum of $\chi_k(G)$ and $\chi_k(\overline{G})$, and present graphs for which those bounds are attained. Those results generalize well known results of Nordhaus and Gaddum [65] for $\chi(G)$ and $\chi(\overline{G})$.

This paper is organized as follows. Notation and definitions of several graphs and graph products are given in section 1.1. In section 2 we formally introduce $\vartheta_k(G)$ and $\chi_k(G)$ and show how those graph parameters relate. In section 3 we study the series $(\vartheta_k(G))_k$. Section 4 provides bounds for $\vartheta_k(G)$ when G is the strong graph product of two graphs and the disjunction product of two graphs. In section 5 one can find values of the generalized ϑ -number for complete graphs, cycle graphs, circulant graphs and complete multipartite graphs. In section 5.1 we provide a closed form expression for the generalized ϑ -function on the Kneser graphs, as well as for the Johnson graphs. Section 5.2 relates $\vartheta(K_k \square G)$ and $\vartheta_k(G)$. We provide a closed form expression for the generalized ϑ -function for strongly regular graphs in section 6. In the same section we also relate the Schrijver's number $\vartheta'(K_k \square G)$ with $\vartheta_k(G)$ when G is a strongly regular graph. In section 7 we study a relation between the orthogonality graphs and here considered graph parameters. Section 8 provides new lower bounds on the k -th chromatic number for regular graphs and triangular graphs. We present several results for the multichromatic number on the Hamming graphs in section 8.1.

1.1 Notation and definitions

Let \mathbb{S}^n be the space of symmetric n by n matrices. For matrix $X \in \mathbb{S}^n$, we write $X \succeq 0$ when X is positive semidefinite. Entries of matrix X are given by X_{ij} . The trace inner product for symmetric matrices is denoted $\langle X, Y \rangle = \text{Tr}(XY)$. The Kronecker product of matrices is denoted by $X \otimes Y$. By abuse of notation, we use the same symbol for the tensor product of graphs. The matrix of all ones is denoted by J , while the identity matrix is denoted by I . We sometimes use subscripts to indicate the size of a matrix. Denote by $\mathbf{0}$ and $\mathbf{1}$ the vector of all zeroes and ones respectively.

For any graph $G = (V(G), E(G))$, we denote its adjacency matrix by A_G , or simply A when the context is clear. Similarly, we use V and E to denote the vertex and edge set of G when it is clear from the context. We assume that $|V| = n$, unless stated differently. The Laplacian matrix of a graph G is denoted by L_G . The complement graph of G , denoted by \overline{G} , is defined as the graph such that $A_{\overline{G}} + A_G = J - I$.

For the eigenvalues of $X \in \mathbb{S}^n$, we follow $\lambda_1(X) \geq \lambda_2(X) \geq \dots \geq \lambda_n(X)$, and denote by $\sigma(A)$ the spectrum of matrix A . That is, $\sigma(A) = \{\lambda_1(A), \dots, \lambda_n(A)\}$. We denote the set $\{1, \dots, n\}$ by $[n]$.

In the rest of this section we provide several definitions. The first definition introduces several graph products, and the remaining ones different classes of graphs.

Definition 1 (Graph products). *An arbitrary graph product of graphs $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ is denoted by $G_1 * G_2$, having as vertex set the Cartesian product $V_1 \times V_2$. Table 1 shows when vertices (v_1, v_2) and (u_1, u_2) are adjacent in $G_1 * G_2$, for the lexicographic, tensor, Cartesian, strong and disjunction (Abdo and Dimitrov [1]) graph products.*

Graph product	$G_1 * G_2$	Condition for $((v_1, v_2), (u_1, u_2)) \in E(G_1 * G_2)$
Lexicographic	$G_1 \circ G_2$	$(v_1, u_1) \in E_1$ or $[v_1 = u_1 \text{ and } (v_2, u_2) \in E_2]$
Tensor	$G_1 \otimes G_2$	$(v_1, u_1) \in E_1$ and $(v_2, u_2) \in E_2$
Cartesian	$G_1 \square G_2$	$[v_1 = u_1 \text{ and } (v_2, u_2) \in E_2]$ or $[v_2 = u_2 \text{ and } (v_1, u_1) \in E_1]$
Strong	$G_1 \boxtimes G_2$	$((v_1, v_2), (u_1, u_2)) \in E(G_1 \square G_2) \cup E(G_1 \otimes G_2)$
Disjunction	$G_1 \vee G_2$	$(v_1, u_1) \in E_1$ or $(v_2, u_2) \in E_2$

Table 1: Graph products

In order to define the Hamming graphs, we first state the definition of the Hamming distance.

Definition 2 (Hamming distance). *For two integer valued vectors \mathbf{u} and \mathbf{v} , the Hamming distance between them, denoted by $d(\mathbf{u}, \mathbf{v})$, is the number of positions in which their entries differ.*

Definition 3 (Hamming graph). *The Hamming graph $H(n, q, F)$ for $n, q \in \mathbb{N}$ and $F \subset \mathbb{N}$ has as vertices all the unique elements in $(\mathbb{Z}/q\mathbb{Z})^n$. In the Hamming graph, vertices u and v are adjacent if their Hamming distance $d(u, v) \in F$.*

Many authors define the Hamming graphs only for $F = \{1\}$.

Definition 4 (Johnson graph). *Let $n, m \in \mathbb{N}$, $1 \leq m \leq n/2$, $f \in \{0, 1, \dots, m\}$ and $N = \{1, 2, \dots, n\}$. The Johnson graph $J(n, m, f)$ has as vertices all the possible m -sized subsets of N . Denote the subset corresponding to a vertex u by $s(u)$. Then $|s(u)| = m$ and vertices u and v are adjacent if and only if $|s(u) \cap s(v)| = f$.*

Many authors define the Johnson graph only for $f = m - 1$. When $f = 0$, the Johnson graph is better known as the Kneser graph.

Definition 5 (Kneser graph). *Let $n, m \in \mathbb{N}$ and $1 \leq m \leq n/2$. Then the Kneser graph $K(n, m)$ is the Johnson graph $J(n, m, 0)$.*

Definition 6 (Strongly regular graph). *A d -regular graph G of order n is called strongly regular with parameters (n, d, λ, μ) if any two adjacent vertices share λ common neighbors and any two non-adjacent vertices share μ common neighbors.*

2 $\vartheta_k(G)$ and $\chi_k(G)$ formulations and their relation

In this section, we formally introduce the multichromatic number and the generalized ϑ -number of a graph. We also show a relationship between these two graph parameters.

Let $G = (V, E)$ be a simple undirected graph with n vertices. A valid k -multicoloring of G that uses R colors is a mapping $f : V \rightarrow 2^R$, such that $|f(i)| = k$ for all vertices $i \in V$ and $|f(i) \cap f(j)| = 0$ for all edges $(i, j) \in E$. The multichromatic number $\chi_k(G)$ is defined to be the smallest R such that a valid k -multicoloring of G exists. Here we consider only valid k -multicoloring and refer to it as k -multicoloring.

Multicoloring can be reduced to standard graph coloring by use of the lexicographic product of graphs, see definition 1. Namely, Stahl [75] showed that for any graph H such that $\chi(H) = k$, we have $\chi_k(G) = \chi(G \circ H)$. For clarity purposes, the simplest choice for H is K_k , the complete graph of order k . This results in

$$\chi_k(G) = \chi(G \circ K_k). \quad (1)$$

For bounds on the chromatic number of (lexicographic) graph products, we refer readers to Klavžar [44] and Geller and Stahl [29]. By the lexicographic product, any bound on $\chi(G)$ can also be transformed to a bound on $\chi_k(G)$. In particular:

$$\chi_k(G) = \chi(G \circ K_k) \geq \omega(G \circ K_k) = \omega(G)\omega(K_k) = k\omega(G). \quad (2)$$

Here we use that $\omega(G \circ H) = \omega(G)\omega(H)$ for general graphs G and H . We also mention the following result

$$\alpha(G \circ H) = \alpha(G)\alpha(H). \quad (3)$$

Both results are proven by Geller and Stahl [29]. Let us also state the following known result:

$$\chi(G \circ H) \leq \chi(G)\chi(H). \quad (4)$$

This result can be explained as follows. Denote the vertex sets of G and H by $V(G)$ and $V(H)$, respectively. For an optimal coloring of G and H , define $c(u)$ as the color of some vertex u . Graph $G \circ H$ has vertices (g_i, h_i) . Every vertex in $G \circ H$ can then be assigned a 2-color combination $(c(g_i), c(h_i))$. Note that by interpreting these 2-color combinations as simply colors, this constitutes a valid coloring of $G \circ H$ by using $\chi(G)\chi(H)$ colors. Combining inequalities (2) and (4) results in

$$k\omega(G) \leq \chi_k(G) = \chi(G \circ K_k) \leq k\chi(G). \quad (5)$$

Note that by (5), any upper bound on $\chi(G)$ can be transformed into an upper bound on $\chi_k(G)$. To compute (or approximate) $\chi_k(G)$ one can consult the wide range of existing literature on standard graph coloring by using $\chi_k(G) = \chi(G \circ K_k)$. Next to that, more specific literature on multicoloring can also be examined. Malaguti and Toth [56] use a combination of tabu search and population management procedures as a metaheuristic to solve (slightly generalized) multicoloring problems. Mehrotra and Trick [60] apply branch and price to generate independent sets for solving the multicoloring problem.

Narasimhan and Manber [62] generalize $\vartheta(G)$ by introducing $\vartheta_k(G)$ as follows:

$$\vartheta_k(G) = \underset{A \in \mathcal{A}(G)}{\text{Minimize}} \sum_{i=1}^k \lambda_i(A), \quad (6)$$

where

$$\mathcal{A}(G) := \{A \in \mathbb{S}^n \mid A_{ij} = 1 \ \forall (i, j) \notin E(G)\}. \quad (7)$$

Narasimhan and Manber prove that $\vartheta_k(G)$ satisfies the following inequality

$$\alpha_k(\overline{G}) \leq \vartheta_k(\overline{G}) \leq \chi_k(G) \quad (8)$$

and thus also $\omega_k(G) \leq \vartheta_k(\overline{G}) \leq \chi_k(G)$. Here, $\alpha_k(G)$ is the cardinality of largest subset $C \subseteq V$ such that the subgraph induced in G by C , denoted $G[C]$, satisfies $\chi(G[C]) \leq k$. Inequality (8) generalizes the Lovász's sandwich theorem [55].

Alizadeh [3] derived the following SDP formulation of $\vartheta_k(G)$, see also [48]:

$$\begin{aligned} \vartheta_k(G) = \underset{\mu \in \mathbb{R}, X, Y \in \mathbb{S}^n}{\text{Minimize}} \quad & \langle I, Y \rangle + \mu k \\ \text{subject to} \quad & X_{ij} = 0 \quad \forall (i, j) \notin E(G) \\ & \mu I + X - J + Y \succeq 0, \ Y \succeq 0. \end{aligned} \quad (\vartheta_k\text{-SDP})$$

The dual problem for ϑ_k -SDP is:

$$\begin{aligned} \vartheta_k(G) = \text{Maximize} \quad & \langle J, Y \rangle \\ \text{subject to} \quad & Y_{ij} = 0 \quad \forall (i, j) \in E(G) \\ & \langle I, Y \rangle = k, \quad 0 \preceq Y \preceq I. \end{aligned} \quad (\vartheta_k\text{-SDP2})$$

Note that for $k = 1$ constraint $Y \preceq I$ is redundant. We show below that $\vartheta_k(\overline{G}) \leq \chi_k(G)$. To prove the result we use different arguments than the arguments used in [62]. In an optimal k -multicoloring of G , define for each of the $\chi_k(G)$ colors used a vector $\mathbf{y}^j \in \{0, 1, k\}^{n+1}$, $1 \leq j \leq \chi_k(G)$. For the entries of \mathbf{y}^j , we have $\mathbf{y}_0^j = k$ and $\mathbf{y}_i^j = 1$ if vertex i has color j , 0 otherwise. Then

$$\frac{1}{k^2} \sum_{j=1}^{\chi_k(G)} \mathbf{y}^j (\mathbf{y}^j)^\top = \begin{bmatrix} \chi_k(G) & \mathbf{1}^\top \\ \mathbf{1} & \frac{1}{k} I + \frac{1}{\chi_k(G)} X \end{bmatrix}.$$

For some $X \in \mathbb{S}^n$ satisfying $X_{ij} = 0$ for all $(i, j) \in E(G)$. By the Schur complement we find $\frac{\chi_k(G)}{k} I + X - J \succeq 0$. Simply set $Y = 0 \in \mathbb{S}^n$. Then the triple $(\frac{\chi_k(G)}{k}, X, Y)$ is feasible for ϑ_k -SDP (for \overline{G}) with objective value $\chi_k(G)$.

To conclude this section we state the following result:

$$\vartheta_k(\overline{G}) \leq k\vartheta(\overline{G}) \leq \chi_k(G). \quad (9)$$

Narasimhan and Manber [62] prove the first inequality in (9). To show this, let $\tilde{A} \in \mathcal{A}(\overline{G})$ such that $\lambda_1(\tilde{A}) = \vartheta(\overline{G})$. Then $\vartheta_k(\overline{G}) \leq \sum_{i=1}^k \lambda_i(\tilde{A}) \leq k\lambda_1(\tilde{A})$ and the proof follows. The second inequality follows from $\vartheta(\overline{G \circ K_k}) = k\vartheta(\overline{G})$ and $\vartheta(\overline{G \circ K_k}) \leq \chi(G \circ K_k) = \chi_k(G)$. The second inequality in (9) also follows from the following known results $\vartheta(\overline{G}) \leq \chi_f(G)$ and $k\chi_f(G) \leq \chi_k(G)$ where $\chi_f(G)$ is the fractional chromatic number of a graph, see e.g., [12]. In this paper we show that $\vartheta_k(G) = k\vartheta(G)$ for many highly symmetric graphs.

3 The series $(\vartheta_k(G))_k$

In this section we consider the series $\vartheta_1(G), \vartheta_2(G), \dots, \vartheta_n(G)$ where G is a graph of order n . We first prove that this series is bounded from above (proposition 1) and increasing (proposition 2). Then, we prove that the increments of the series i.e., $\vartheta_k(G) - \vartheta_{k-1}(G)$ are decreasing in k , see theorem 7. We also show that this increment can be arbitrarily small for a particular graph, see theorem 8.

Let us first establish a relation between $\vartheta_k(G)$ and $\chi(G)$.

Proposition 1. *For $k \geq \chi(G)$, $G = (V, E)$, we have $\vartheta_k(G) = |V|$. Furthermore, $\vartheta_k(G) \leq |V|$ for all k .*

Proof. Let $k = \chi(G)$. Then $\alpha_k(G) = |V|$, where we take the k independent sets to be the color classes in an optimal coloring of G . Thus, it follows from (8) that $|V| \leq \vartheta_k(G)$.

Furthermore, note that for any graph G , matrix $J \in \mathcal{A}(G)$ is feasible for (6). Since matrix J has eigenvalue $|V|$ with multiplicity one and the other eigenvalues equal 0, we have $\vartheta_k(G) \leq |V|$ for any graph G . However, when $k \geq \chi(G)$ we have $\vartheta_k(G) = |V|$. \square

Part of proposition 1 can be more succinctly stated as $\vartheta_{\chi(G)}(G) = |V|$. The parameter $\vartheta_k(G)$ induces a series of parameters for a graph, given by $\vartheta_1(G), \vartheta_2(G), \dots, \vartheta_n(G) = |V|$. Proposition 1 shows that this series is bounded from above by $|V|$. The next proposition shows that this series is non-decreasing in k .

Proposition 2. *For any graph G , $\vartheta_k(G) \leq \vartheta_{k+1}(G)$, with equality if and only if $\vartheta_k(G) = |V|$.*

Proof. Consider graph G of order n and let Y be optimal for ϑ_k -SDP2. We have $\text{Tr}(Y) = k$ and $0 \preceq Y \preceq I$. Define matrix Z as follows: $Z := \left(1 - \frac{1}{n-k}\right)Y + \frac{1}{n-k}I$. It follows that matrix Z is feasible for ϑ_{k+1} -SDP2 and thus

$$\vartheta_{k+1}(G) \geq \langle J, Z \rangle = \vartheta_k(G) + \frac{n - \vartheta_k(G)}{n - k} \geq \vartheta_k(G).$$

\square

Proposition 2 allows us to further restrict ϑ_k -SDP.

Proposition 3. *Let (X^*, Y^*, μ^*) be an optimal solution to ϑ_k -SDP for an arbitrary graph G . Then $\mu^* \geq 0$.*

Proof. We prove the statement by contradiction. Assume that the triple (X^*, Y^*, μ^*) is optimal for ϑ_k -SDP and $\mu^* < 0$. Note that the triple (X^*, Y^*, μ^*) is then also feasible for ϑ_{k+1} -SDP. Since $\mu^* < 0$, this would imply that $\vartheta_k(G) > \vartheta_{k+1}(G)$, which contradicts proposition 2. Thus $\mu^* \geq 0$. \square

Next, we investigate the increments of the series $(\vartheta_k(G))_k$. For that purpose, we define for any graph G and $k \geq 2$ the increment of $(\vartheta_k(G))_k$ as follows:

$$\Delta_k(G) := \vartheta_k(G) - \vartheta_{k-1}(G), \quad (10)$$

and set $\Delta_1(G) = \vartheta_1(G)$.

Theorem 7. *For any graph G and $k \geq 1$, $\Delta_k(G) \geq \Delta_{k+1}(G)$.*

Proof. Let $k \geq 1$ and matrix $A_k \in \mathcal{A}(G)$, where $\mathcal{A}(G)$ is defined in (7), satisfies

$$\sum_{i=1}^k \lambda_i(A_k) = \vartheta_k(G). \quad (11)$$

Stated differently, matrix A_k is the optimal solution to (6) for computing $\vartheta_k(G)$. Since (6) is a minimization problem,

$$\vartheta_k(G) \leq \sum_{i=1}^k \lambda_i(A_{k'}), \quad k' \neq k. \quad (12)$$

By substituting (11) and (12) in the definition of $\Delta_k(G)$ for $k \geq 2$, see (10), we obtain:

$$\Delta_k(G) \leq \sum_{i=1}^k \lambda_i(A_{k-1}) - \sum_{i=1}^{k-1} \lambda_i(A_{k-1}) = \lambda_k(A_{k-1}). \quad (13)$$

Similarly,

$$\Delta_k(G) \geq \sum_{i=1}^k \lambda_i(A_k) - \sum_{i=1}^{k-1} \lambda_i(A_k) = \lambda_k(A_k). \quad (14)$$

Combining (13) and (14) yields $\Delta_k(G) \geq \lambda_k(A_k) \geq \lambda_{k+1}(A_k) \geq \Delta_{k+1}(G)$, $k \geq 2$. The inequality $\Delta_1(G) \geq \Delta_2(G)$ follows from (9). \square

Let us summarize the implications of propositions 2 and theorem 7. Proposition 2 proves that

$$\Delta_k(G) = 0 \iff \vartheta_{k-1}(G) = |V|. \quad (15)$$

For complete graphs we have $\Delta_k(K_n) = 1$, see theorem 11. There exist however graphs for which $\Delta_k(G) < 1$. We investigate the limiting behaviour of $\Delta_k(G)$ in section 3.1.

When we consider the series induced by $\vartheta_k(G)$ as a function of k , we know that this series is increasing towards $|V(G)|$. Theorem 7 shows that the increments in this series decrease in k . Loosely speaking, one might say the second derivative of $f(k) = \vartheta_k(G)$ is negative.

3.1 Limiting behaviour of $\Delta_k(G)$

In this section we show that for any real number $\varepsilon > 0$, there exists a graph G and a number $k \geq 1$ such that $0 < \Delta_k(G) < \varepsilon$. For this purpose, define graph $\mathcal{G}_n = (V(\mathcal{G}_n), E(\mathcal{G}_n))$ as follows:

$$V(\mathcal{G}_n) := \{1, 2, \dots, n\} \text{ and } E(\mathcal{G}_n) := \{(i, j) \mid i < j \leq n-1\} \cup \{(n-1, n)\}. \quad (16)$$

Graph \mathcal{G}_n is thus a complete graph on $n-1$ vertices plus one additional vertex. This additional vertex is connected to the complete graph K_{n-1} by a single edge.

Theorem 8. *For $n \geq 5$, we have*

$$\vartheta_{n-2}(\mathcal{G}_n) = n - 2 + \frac{2}{n-3} \sqrt{(n-2)(n-4)}.$$

Proof. We prove the theorem by finding a lower and upper bound on $\vartheta_{n-2}(\mathcal{G}_n)$, both of which equal the expression stated in theorem 8. Let $p = \sqrt{\frac{n-4}{(n-2)(n-3)^2}}$. Define matrix $Y \in \mathbb{S}^n$ as follows:

$$Y = \begin{bmatrix} \frac{n-4}{n-3}I_{n-2} & \mathbf{0}_{n-2} & p\mathbf{1}_{n-2} \\ \mathbf{0}_{n-2}^\top & 1 & 0 \\ p\mathbf{1}_{n-2}^\top & 0 & \frac{1}{n-3} \end{bmatrix}.$$

Matrix Y is feasible for ϑ_{n-2} -SDP2 if $0 \preceq Y \preceq I$. Therefore we derive

$$I - Y = \begin{bmatrix} \frac{1}{n-3}I_{n-2} & \mathbf{0}_{n-2} & -p\mathbf{1}_{n-2} \\ \mathbf{0}_{n-2}^\top & 0 & 0 \\ -p\mathbf{1}_{n-2}^\top & 0 & \frac{n-4}{n-3} \end{bmatrix},$$

and take the Schur complement of the block $\frac{1}{n-3}I_{n-2}$ of $I - Y$:

$$\begin{bmatrix} 0 & 0 \\ 0 & \frac{n-4}{n-3} \end{bmatrix} - \begin{bmatrix} \mathbf{0}_{n-2}^\top \\ p\mathbf{1}_{n-2}^\top \end{bmatrix} (n-3)I_{n-2} \begin{bmatrix} \mathbf{0}_{n-2} & p\mathbf{1}_{n-2} \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \succeq 0.$$

Thus $Y \preceq I$. Similarly, by taking the Schur complement of the upper left $(n-1) \times (n-1)$ block matrix of Y , we find that $Y \succeq 0$. We omit the details of this computation. This implies that Y is feasible for ϑ_{n-2} -SDP2 and

$$\vartheta_{n-2}(\mathcal{G}_n) \geq \langle J, Y \rangle = n - 2 + \frac{2}{n-3} \sqrt{(n-2)(n-4)}. \quad (17)$$

Finding the (equal) upper bound on $\vartheta_{n-2}(\mathcal{G}_n)$ is a bit more involved. Let $\alpha = \frac{n-5}{n-3} \sqrt{\frac{n-2}{n-4}}$ and set

$$A = \begin{bmatrix} \alpha J_{n-2} + (1-\alpha)I_{n-2} & \mathbf{0}_{n-2} & \mathbf{1}_{n-2} \\ \mathbf{0}_{n-2}^\top & 1 & 0 \\ \mathbf{1}_{n-2}^\top & 0 & 1 \end{bmatrix}.$$

Note that $A \in \mathcal{A}(\mathcal{G}_n)$ (see (7)). We show that for

$$\beta_{1,2} = \frac{-\alpha(n-3) \pm \sqrt{\alpha^2(n-3)^2 - 4(2-n)}}{2}, \quad (18)$$

the vectors $v_i = [\mathbf{1}_{n-2}^\top, 0, \beta_i]^\top$, $i \in \{1, 2\}$ are two eigenvectors of matrix A . Consider

$$Av_i = \begin{bmatrix} (\alpha(n-3) + 1 + \beta_i)\mathbf{1}_{n-2} \\ 0 \\ (\frac{n-2}{\beta_i} + 1)\beta_i \end{bmatrix}. \quad (19)$$

The right-hand side of (19) equals the scaled vector v_i , since by (18),

$$\alpha(n-3) + 1 + \beta_i = \frac{n-2}{\beta_i} + 1. \quad (20)$$

For $i \in \{1, 2\}$, (20) gives the two eigenvalues corresponding to v_i . Also $u = [\mathbf{0}_{n-2}^\top, 1, 0]^\top$ is an eigenvector of A with corresponding eigenvalue one (and multiplicity one). Since A is a real symmetric matrix, its eigenvectors are orthogonal. The remaining eigenvectors are thus $w_i = [c_i^\top, 0, 0]^\top$ where $c_i \in \mathbb{R}^{n-2}$ is a vector whose entries sum to 0. The eigenvectors w_i correspond to eigenvalues of $1 - \alpha$. We have described all the eigenvectors of A . By substituting (18) in (20) one can verify that the four unique eigenvalues of A are ordered as follows:

$$\sqrt{(n-2)(n-4)} + 1 > 1 > 1 - \alpha > 1 - \sqrt{\frac{n-2}{n-4}},$$

with corresponding multiplicities 1, 1, $n-3$, 1, respectively. The sum of the largest $n-2$ eigenvalues of A serves as upper bound on $\vartheta_{n-2}(\mathcal{G}_n)$, see (6). That is,

$$\vartheta_{n-2}(\mathcal{G}_n) \leq \sum_{i=1}^{n-2} \lambda_i(A) = n - 2 + \frac{2}{n-3} \sqrt{(n-2)(n-4)}.$$

This upper bound on $\vartheta_{n-2}(\mathcal{G}_n)$ coincides with the lower bound (17), which proves the theorem. \square

Using theorem 8 we can show that $\Delta_{n-1}(\mathcal{G}_n)$ ($n \geq 5$) converges to zero. Namely,

$$\Delta_{n-1}(\mathcal{G}_n) = \vartheta_{n-1}(\mathcal{G}_n) - \vartheta_{n-2}(\mathcal{G}_n) \leq 2 \left(1 - \sqrt{\frac{(n-2)(n-4)}{(n-3)^2}} \right),$$

from where it follows that $\Delta_{n-1}(\mathcal{G}_n)$ ($n \geq 5$) converges to zero. To conclude, strictly positive values of $\Delta_k(G)$ can be arbitrarily small. It is unclear whether lower bounds exist on $\Delta_k(G)$ for fixed k . One example of such a bound is simple for $k = 1$ i.e., $\Delta_1(G) = \vartheta_1(G) \geq \alpha(G) \geq 1$.

4 Graph products and the generalized ϑ -number

In this section we present bounds for $\vartheta_k(G)$ when G is the graph strong product of two graphs (theorem 9) and the graph disjunction product of two graphs (theorem 10).

In [54], Lovász proved the following result:

$$\vartheta(G_1 \boxtimes G_2) = \vartheta(G_1)\vartheta(G_2), \quad (21)$$

where $G_1 \boxtimes G_2$ is the graph strong product of G_1 and G_2 , see definition 1. Since $G_1 \boxtimes K_k$ is isomorphic to $G_1 \circ K_k$ and $\vartheta(K_k) = 1$ we have that

$$\vartheta(G \circ K_k) = \vartheta(G \boxtimes K_k) = \vartheta(G) \leq \vartheta_k(G),$$

see also proposition 2. Below, we generalize the result for $\vartheta(G_1 \boxtimes G_2)$ to $\vartheta_k(G_1 \boxtimes G_2)$. For that purpose we need the following well known result.

Lemma 1. *For square matrices A and B with eigenvalues λ_i and μ_i respectively, the eigenvalues of $A \otimes B$ equal $\lambda_i \mu_i$.*

For a reference of lemma 1 one can confer Horn and Johnson [42] for example.

Theorem 9. *For any graphs G_1 and G_2*

$$\frac{1}{k} \vartheta_k(G_1) \vartheta_k(G_2) \leq \vartheta_k(G_1 \boxtimes G_2) \leq k \vartheta(G_1) \vartheta(G_2).$$

Proof. Let X_1^* and X_2^* be optimal for ϑ_k -SDP2 of G_1 and G_2 respectively. The adjacency matrix of $G_1 \boxtimes G_2$ is given by

$$A_{G_1 \boxtimes G_2} = (A_{G_1} + I) \otimes (A_{G_2} + I) - I,$$

see e.g., Sayama [70]. Here \otimes denotes the Kronecker product. Consider $Y = \frac{1}{k} X_1^* \otimes X_2^*$. From the adjacency matrix of $G_1 \boxtimes G_2$ it can be verified that $Y_{ij} = 0, \forall (i, j) \in E(G_1 \boxtimes G_2)$. By lemma 1, the eigenvalues of Y lie between 0 and 1 and thus $0 \preceq Y \preceq I$. More specifically, the eigenvalues of Y lie between 0 and $\frac{1}{k}$. It follows that matrix Y is feasible for ϑ_k -SDP2 of $G_1 \boxtimes G_2$ and attains the following objective value:

$$\langle J, Y \rangle = \frac{1}{k} \langle J, X_1^* \otimes X_2^* \rangle = \frac{1}{k} \vartheta_k(G_1) \vartheta_k(G_2).$$

This proves the lower bound. The upper bound follows from (9) and (21). \square

The bounds from theorem 9 are attained for some graphs, e.g., complete graphs. In general, the bounds are more crude for larger values of k . We now focus on the disjunction graph product (see definition 1). For graphs G_1 and G_2 of order n_1 and n_2 respectively, we have

$$A_{G_1 \vee G_2} = J_{n_1} \otimes A_{G_2} + A_{G_1} \otimes (A_{\overline{G_2}} + I_{n_2}).$$

Equivalently, $A_{G_1 \vee G_2} = \min \left(J_{n_1} \otimes A_{G_2} + A_{G_1} \otimes J_{n_2}, 1 \right)$. Our next result provides an upper bound on the generalized ϑ -number for the graph disjunction product.

Theorem 10. *For graphs G_1 and G_2 of orders n_1 and n_2 respectively, we have*

$$\vartheta_k(G_1 \vee G_2) \leq \min \{ n_1 \vartheta_k(G_2), n_2 \vartheta_k(G_1) \}.$$

Proof. Consider the SDP problem ϑ_k -SDP2 for $G_1 \vee G_2$. This maximization problem is least constrained when $G_1 = \overline{K}_{n_1}$. Thus

$$\vartheta_k(G_1 \vee G_2) \leq \vartheta_k(\overline{K}_{n_1} \vee G_2). \quad (22)$$

We will show that $\vartheta_k(\overline{K}_{n_1} \vee G_2) = n_1 \vartheta_k(G_2)$. Let X^* be an optimal solution to ϑ_k -SDP2 for G_2 . Matrix $J_{n_1} \otimes \frac{1}{n_1} X^*$ is a feasible solution to ϑ_k -SDP2 for $\overline{K}_{n_1} \vee G_2$. The objective value of this solution equals

$$\langle J, J_{n_1} \otimes \frac{1}{n_1} X^* \rangle = n_1 \langle J, X^* \rangle = n_1 \vartheta_k(G_2) \implies \vartheta_k(\overline{K}_{n_1} \vee G_2) \geq n_1 \vartheta_k(G_2). \quad (23)$$

Let (Y^*, X^*, μ^*) be an optimal solution to ϑ_k -SDP for G_2 . Then $J_{n_1} \otimes Y^*$, $J_{n_1} \otimes X^*$ and $n_1 \mu^*$ form a feasible solution to ϑ_k -SDP for $\overline{K}_{n_1} \vee G_2$. Namely, by lemma 1 we have that $J_{n_1} \otimes Y^* \succeq 0$. Also

$$n_1 \mu^* I + J_{n_1} \otimes X^* - J + J_{n_1} \otimes Y^* = \mu^* (n_1 I_{n_1} - J_{n_1}) \otimes I_{n_2} + J_{n_1} \otimes (\mu^* I_{n_2} + X^* - J_{n_2} + Y^*) \succeq 0,$$

where we use that $\mu^* \geq 0$, see proposition 3. Lastly, this feasible solution to the minimization problem obtains an objective value of

$$\langle I, J_{n_1} \otimes Y^* \rangle + n_1 \mu^* k = n_1 \left(\langle I, Y^* \rangle + \mu^* k \right) = n_1 \vartheta_k(G_2) \implies \vartheta_k(\overline{K}_{n_1} \vee G_2) \leq n_1 \vartheta_k(G_2). \quad (24)$$

Now (23) and (24) imply that $\vartheta_k(\overline{K}_{n_1} \vee G_2) = n_1 \vartheta_k(G_2)$. This result combined with (22) proves that

$$\vartheta_k(G_1 \vee G_2) \leq n_1 \vartheta_k(G_2). \quad (25)$$

From the definition of the disjunction graph product (see definition 1), it follows that the disjunction graph product is commutative and thus

$$\vartheta_k(G_1 \vee G_2) = \vartheta_k(G_2 \vee G_1) \leq n_2 \vartheta_k(G_1). \quad (26)$$

Combining equations (25) and (26) proves the theorem. \square

The proof shows that when either G_1 or G_2 is the complement of a complete graph, graph $G_1 \vee G_2$ attains the bound of theorem 10.

5 Value of ϑ_k for some graphs

In [54], Lovász derived an explicit expression for the ϑ -number of cycle graphs and the Kneser graphs. In this section, we derive the generalized ϑ -number for those graphs, as well as for circulant, complete, complete multipartite graphs, and the Johnson graphs. In section 5.1 we present bounds for $\vartheta_k(G)$ when G is a regular graph and show that the bound is tight for edge-transitive graphs. Section 5.2 provides an analysis of $\vartheta(K_k \square G)$, which is an upper bound on the number of vertices in the maximum k -colorable subgraph of G .

We denote cycle graphs of order n by C_n , complete graphs of order n by K_n , and complete multipartite graph by K_{m_1, \dots, m_p} . Note that K_{m_1, \dots, m_p} is a graph on $n = \sum_{i=1}^p m_i$ vertices.

Theorem 11. For $k \leq n$, $\vartheta_k(K_n) = k$.

Proof. Consider the SDP problem ϑ_k -SDP2. For the complete graph, the only matrices feasible for ϑ_k -SDP2 are diagonal matrices with trace equal to k . Set for example $Y = \frac{k}{n} I$. Then Y is feasible for ϑ_k -SDP2 and has objective value k . \square

Stahl [75] determined $\chi_k(C_n)$. For odd cycles, he showed that $\chi_k(C_{2n+1}) = 2k + 1 + \left\lfloor \frac{k-1}{n} \right\rfloor$, and for even cycles that $\chi_k(C_{2n}) = 2k$. The latter result follows trivially from (60).

Since C_n is bipartite when n is even, it follows from proposition 1 that $\vartheta_2(C_n) = n$. To compute $\vartheta_2(C_n)$ for odd cycle graphs, we require the following lemma.

Lemma 2. For n odd, $n \geq 5$, we have $0.447 \approx \frac{\sqrt{5}}{5} \leq \frac{\vartheta(C_n)}{n} < \frac{\vartheta(C_{n+2})}{n+2} < \frac{1}{2}$.

Proof. By Lovász [54], we have

$$\vartheta(C_n) = \frac{n \cos(\pi/n)}{1 + \cos(\pi/n)}, \quad n \text{ odd.} \quad (27)$$

Define $f(n) := \vartheta(C_n)/n$. Then $f'(n) = \frac{\pi \sin(\frac{\pi}{n})}{(1 + \cos(\frac{\pi}{n}))^2 n^2}$. For $n \geq 5$, $f'(n) > 0$. Moreover, for $n \geq 5$, we have $\cos(\pi/n) < 1$. This results in $f(n) < 1/2$ and since $f(5) = \sqrt{5}/5$, this proves the lemma. \square

We also require the definition of a *circulant matrix*. Each row of a circulant matrix equals the preceding row in the matrix rotated one element to the right. Circulant matrices thus have a constant row sum. This constant row sum is also one of the eigenvalues with $\mathbf{1}$ as its corresponding eigenvector.

Theorem 12. *Let n be odd and $n > 1$. Then $\vartheta_2(C_n) = 2\vartheta(C_n)$.*

Proof. For $n = 3$, $C_3 = K_3$ and the result follows from theorem 11. Thus let $n \geq 5$. Let $\Gamma \subset \mathbb{S}^n$ be the set of optimal feasible solutions to ϑ_1 -SDP2 for C_n and let $Y \in \Gamma$. Note Γ is convex. Let $p(Y)$ denote an optimal solution to ϑ_1 -SDP2 obtained by permuting the vertices of C_n by automorphism p . Matrix $p(Y) \in \Gamma$. Denote the average over all automorphisms p by \bar{Y} . Then $\bar{Y} \in \Gamma$ by convexity of Γ and since C_n is edge transitive, \bar{Y} is a circulant matrix, like the adjacency matrix of C_n .

As $\bar{Y} \in \Gamma$, we find

$$\langle J, \bar{Y} \rangle = \text{Tr}(\mathbf{1}\mathbf{1}^\top \bar{Y}) = \text{Tr}(\mathbf{1}^\top \bar{Y}\mathbf{1}) = \mathbf{1}^\top \bar{Y}\mathbf{1} = \vartheta_1(C_n). \quad (28)$$

As \bar{Y} is also circulant, it has eigenvector $\mathbf{1}$. By (28), its corresponding eigenvalue equals $\bar{\lambda} = \vartheta(C_n)/n$.

We will prove that the largest eigenvalue of \bar{Y} equals $\bar{\lambda}$. Assume that the largest eigenvalue of \bar{Y} does not equal $\bar{\lambda}$. Then \bar{Y} has eigenvalue Λ , for some $\Lambda > \bar{\lambda}$. Since \bar{Y} is a symmetric circulant matrix of odd dimension, \bar{Y} has only one eigenvalue with odd multiplicity (Tee [77]). Thus Λ or $\bar{\lambda}$ have multiplicity greater than one. Note that since \bar{Y} is feasible for ϑ_1 -SDP2, it has non-negative eigenvalues that sum to one. However, both terms $\Lambda + 2\bar{\lambda}$ and $2\Lambda + \bar{\lambda}$ are strictly greater than one by lemma 2, and hence, the assumption that $\bar{\lambda}$ is not the largest eigenvalue of \bar{Y} leads to a contradiction.

The largest eigenvalue of \bar{Y} is thus smaller than $1/2$. Then $2\bar{Y} \preceq I$. Clearly, $2\bar{Y}$ satisfies the other feasibility conditions of ϑ_2 -SDP2. Thus $2\bar{Y}$ is feasible for ϑ_2 -SDP2 and $\vartheta_2(C_n) \geq 2\vartheta(C_n)$. Combined with (9), the theorem follows. \square

Since $\chi(C_n) = 3$ for odd cycles, $\vartheta_3(C_n) = n$ follows trivially from proposition 1.

Graphs for which the adjacency matrix is a circulant matrix are called *circulant graphs*, like the cycle graphs and some Paley graphs. There has been research done on computing $\vartheta(G)$ for circulant graphs [5, 9, 10, 19]. In particular, Crespi [19] computes the Lovász theta function for the circulant graphs of degree four having even displacement, while Brimkov et al. [10] consider $\vartheta(C_{n,j})$, where $V(C_{n,j}) = \{1, 2, \dots, n\}$ and $E(C_{n,j}) = E(C_n) \cup \{(i, i') \mid i - i' = j \pmod n\}$.

Let H_n be a connected circulant graph on n vertices. Then H_n contains a Hamiltonian cycle (Boesch and Tindell [7]). Equivalently, the cycle graph C_n is a minor of H_n . Maximization problem ϑ_k -SDP2 is then more restricted for H_n than it is for C_n . Thus

$$\vartheta_1(H_n) \leq \vartheta_1(C_n) \leq \frac{n}{2}.$$

Consider ϑ_1 -SDP2 for H_n . Graph H_n has a circulant adjacency matrix, meaning we can restrict optimization of ϑ_1 -SDP2 over the Lee scheme, the association scheme of symmetric circulant matrices, without loss of generality [28]. As (28) shows, ϑ_1 -SDP2 is now equivalent to maximizing the largest (scaled) eigenvalue over feasible matrices. Let M be a matrix optimal for ϑ_1 -SDP2 for graph H_n . Then $\lambda_1(M) = \vartheta(H_n)/n \leq 1/2$. Then $2M$ is also optimal for ϑ_2 -SDP2 for graph H_n . More generally, if $k \leq n/\vartheta(H_n)$, then $\lambda_1(kM) \leq 1$ and kM is then feasible for ϑ_k -SDP2, attaining the objective value $\min\{k\vartheta(H_n), n\}$. In case $k > n/\vartheta(H_n)$, we have $\vartheta_k(H_n) = n$. Thus, in general

$$\vartheta_k(H_n) = \min\{k\vartheta(H_n), n\}.$$

For any k , there exists a graph P on n vertices such that $\vartheta_k(P) < n$. Specifically, if P is the Paley graph of order n , then $\vartheta(P) = \sqrt{n}$ (cf. [34]). For fixed k and n large enough, $k\sqrt{n} < n$.

Theorem 13. *For $m_1 \geq m_2 \geq \dots \geq m_p$, $\vartheta_k(K_{m_1, \dots, m_p}) = \sum_{i=1}^k m_i$.*

Proof. Let $n = \sum_{i=1}^p m_i$. For notational convenience, we write $K = K_{m_1, \dots, m_p}$, with corresponding adjacency matrix A_K . Since K is a graph on n vertices, A_K can be written as $A_K = J_n - \text{Diag}(J_{m_1}, \dots, J_{m_p})$. Note that $X := \text{Diag}(J_{m_1}, \dots, J_{m_p}) \in \mathcal{A}(K)$, see (7). Therefore X is feasible for (6). The eigenvalues of X are the eigenvalues of the block matrices J , i.e., $\lambda_i(X) = m_i$ for $i \leq p$. Thus, we have $\vartheta_k(K) \leq \sum_{i=1}^k \lambda_i(X) = \sum_{i=1}^k m_i$. Note that $\alpha_k(K) = \sum_{i=1}^k m_i$, and the proof follows from (8) and the above inequality. \square

Recall again the definition of $\Delta_k(G)$, given in (10). In section 3.1, we show that strictly positive values $\Delta_k(G)$ can be arbitrarily small. We show now, by use of theorem 13, that the ratio between strictly positive successive values of $\Delta_k(G)$ can be arbitrarily small. More formally, for any $\varepsilon > 0$ and any $k \geq 1$, there exists a graph G such that

$$0 < \frac{\Delta_{k+1}(G)}{\Delta_k(G)} < \varepsilon. \quad (29)$$

We again ignore the case $\Delta_k(G) = 0$, see (15). In view of theorem 13 we have

$$\frac{\Delta_2(K_{n,1})}{\Delta_1(K_{n,1})} = \frac{1}{n} < \varepsilon,$$

for some integer n sufficiently large. Thus for sufficiently large n , graph $K_{n,1}$ satisfies (29) for $k = 1$. Graph $K_{n,n,1}$ satisfies (29) for $k = 2$. Graph $K_{n,n,n,1}$ satisfies (29) for $k = 3$, and so on.

5.1 Regular graphs

In this section we present an upper bound on the ϑ_k -function for regular graphs, see theorem 15. This result can be seen as a generalization of the Lovász upper bound on the ϑ -function for regular graphs. We exploit the result of theorem 15 to derive an explicit expression for the generalized theta function for the Kneser graph, see theorem 16. Moreover, we prove that $\vartheta_k(G) = k\vartheta(G)$ when G is the Johnson graph, see theorem 17.

Let us first state the following well known result.

Theorem 14 (Lovász [54]). *For a regular graph G of order n , having adjacency matrix A_G and $\lambda_1(A_G) \geq \lambda_2(A_G) \geq \dots \geq \lambda_n(A_G)$, we have*

$$\vartheta(G) \leq \frac{n\lambda_n(A_G)}{\lambda_n(A_G) - \lambda_1(A_G)}.$$

If G is an edge-transitive graph, this inequality holds with equality.

For a finite set of real numbers P , we denote by $\mathbf{S}_k(P)$ the sum of the largest k elements in P . Now, we state our result.

Theorem 15. *For any regular graph G of order n , we have*

$$\vartheta_k(G) \leq \min_x \mathbf{S}_k(\sigma(J + xA)) \leq n + \frac{n}{\lambda_n(A_G) - \lambda_1(A_G)} \left(\lambda_1(A_G) + \sum_{i=0}^{k-2} \lambda_{n-i}(A_G) \right). \quad (30)$$

Where we set the summation equal to 0 when $k = 1$ and $\sigma(\cdot)$ denotes the spectrum of a matrix. The first inequality holds with equality if G is also edge-transitive.

Proof. The proof is an extension of Lovász' [54] proof of theorem 14. Let G be a regular graph of order n . For notational convenience, we write $A_G = A$ and $\lambda_i(A_G) = \lambda_i$. Since G is a regular graph, vector $\mathbf{1}$ is an eigenvector of A . Let $v \neq \mathbf{1}$ be an eigenvector of A . As A is symmetric, its eigenvectors are orthogonal. Thus $\mathbf{1}^\top v = 0$, which implies that $Jv = 0$. Thus the eigenvectors of A are also eigenvectors of $J + xA$. In particular, we have $\sigma(J + xA) = \{n + x\lambda_1, x\lambda_2, \dots, x\lambda_n\}$, for some $x \in \mathbb{R}$. Note that $J + xA \in \mathcal{A}(G)$, see (7). Therefore, it follows from (6) that

$$\vartheta_k(G) \leq \min_x \mathbf{S}_k(\sigma(J + xA)).$$

Minimizing $\mathbf{S}_k(\sigma(J + xA))$ can be done analytically when $k = 1$. Lovász [54] showed that setting $x = \frac{n}{\lambda_n - \lambda_1}$ minimizes $\mathbf{S}_k(\cdot)$ when $k = 1$. Setting x to this value provides the second upper bound in the theorem. Note that $x = \frac{n}{\lambda_n - \lambda_1}$ implies that $n + x\lambda_1 = x\lambda_n$.

We now prove that $\vartheta_k(G) = \min_x \mathbf{S}_k(\sigma(J + xA))$ when G is edge-transitive. Assume that G is edge-transitive. It is known that the sum of the k largest eigenvalues of a matrix is a convex function (Overton and Womersley [67]). Thus the average over all optimal solutions to (6) of all automorphisms of G is also optimal. Since G is edge-transitive, this average is of the form $J + xA$, which proves the equality claim. \square

We remark that that theorem 12 can also be proven by applying theorem 15.

In order to obtain sharper bounds for $\vartheta_k(G)$, one can minimize $\mathbf{S}_k(\sigma(J + xA))$, or compute $\vartheta_k(G)$ directly. Note that computing $\vartheta_k(G)$ by interior point methods is computationally demanding already for some graphs with 200 vertices, see [48]. In general, $\mathbf{S}_k(\sigma(J + xA))$ is the sum of the k largest linear functions given by $\sigma(J + xA)$. Ogryczak and Tamir [66] consider this problem which they show is solvable in linear time, but unfortunately, obtaining a general solution is not possible. When we consider specific graphs, and $\sigma(J + xA)$ is thus explicit, minimizing $\mathbf{S}_k(\cdot)$ can be done analytically, as we show for the Kneser graph.

Lovász [54] proved that $\vartheta(K(n, m)) = \binom{n-1}{m-1}$, where $K(n, m)$ is the Kneser graph, see definition 5. The Kneser graph $K(n, m)$ is regular of valency $\binom{n-m}{m}$. We provide an explicit expression for $\vartheta_k(K(n, m))$.

Theorem 16. *For $k \leq \lfloor \frac{n}{m} \rfloor$, $k \leq n - 2m + 1$ and $2m \leq n$, we have*

$$\vartheta_k(K(n, m)) = k\vartheta(K(n, m)) = k\binom{n-1}{m-1}.$$

When $k > \frac{n}{m}$ or $k > n - 2m + 1$

$$\vartheta_k(K(n, m)) = \binom{n}{m}.$$

Proof. Let $k \leq \lfloor \frac{n}{m} \rfloor$, $k \leq n - 2m + 1$ and $2m \leq n$. Note that n does not refer to the number of vertices but to a parameter of the Kneser graph $K(n, m)$. We will use v to denote the number of vertices of $K(n, m)$, i.e., $v = \binom{n}{m}$. Let A be the adjacency matrix of $K(n, m)$, having eigenvalues $\lambda_1 \geq \dots \geq \lambda_v$. We compute the minimum of $\mathbf{S}_k(\sigma(J + xA))$, see theorem 15. Note that $\sigma(\cdot)$ denotes the spectrum of a matrix, and $\mathbf{S}_k(\sigma(\cdot))$ the sum of k largest eigenvalues in a matrix. Define $f(x) := \mathbf{S}_k(\sigma(J + xA))$. For $x^* = \frac{v}{\lambda_v - \lambda_1}$ we have

$$f(x^*) = v + x^* \left(\lambda_1 + \sum_{i=0}^{k-2} \lambda_{v-i} \right). \quad (31)$$

The eigenvalues of A are as follows $(-1)^i \binom{n-m-i}{m-i}$, $i = 0, 1, \dots, m$, see [54]. Substituting the eigenvalues in (31) yields

$$f(x^*) = v + x^* \left[\binom{n-m}{m} - (k-1) \binom{n-m-1}{m-1} \right] = v + x^* (\lambda_1 + (k-1)\lambda_v). \quad (32)$$

Let us consider the case $k < \frac{n}{m}$. For any $\varepsilon > 0$ we have

$$f(x^* + \varepsilon) = f(x^*) + \varepsilon \left[\binom{n-m}{m} - (k-1) \binom{n-m-1}{m-1} \right],$$

and thus $k < \frac{n}{m} \implies \binom{n-m}{m} - (k-1) \binom{n-m-1}{m-1} > 0 \implies f(x^* + \varepsilon) > f(x^*)$. Similarly, for any sufficiently small $\varepsilon > 0$, we have

$$f(x^* - \varepsilon) = k(x^* - \varepsilon)\lambda_v = kx^*\lambda_v - k\varepsilon\lambda_v > x^*\lambda_v + (k-1)x^*\lambda_v = f(x^*). \quad (33)$$

Here we used the fact that $x^*\lambda_v = v + x^*\lambda_1$. From the previous discussion it follows that for any sufficiently small positive ε we have

$$f(x^*) < f(x^* \pm \varepsilon).$$

By convexity of f , x^* is the global minimizer of f . From $f(x^*) = k\vartheta(K(n, m))$ the theorem follows for the case $k < \frac{n}{m}$.

Now we consider the case $k = \frac{n}{m}$. From (32) it follows $f(x^*) = v$. In fact, for any β satisfying $x^* \leq \beta \leq 0$, $f(\beta) = v$. For any $\varepsilon > 0$,

$$f(\varepsilon) = v + \varepsilon \left(\sum_{i=1}^k \lambda_i \right). \quad (34)$$

As the λ_i sum to zero, the sum of the k largest λ_i must be strictly positive. Thus $f(\varepsilon) > v$. The derivation from (33) is also valid for the case $k = \frac{n}{m}$. Invoking again the convexity of f proves that v is the minimum value of f . Thus

$$\vartheta_{n/m}(K(n, m)) = \frac{n}{m} \binom{n-1}{m-1} = \binom{n}{m} = v.$$

When $k > \frac{n}{m}$, similar derivations show that the minimum of f equals v . Lastly, Kneser's conjecture (Kneser [45]), which was proved by Lovász [53], states that $\chi(K(n, m)) = n - 2m + 2$. The inequality $k > n - 2m + 1$ is thus equivalent to $k \geq \chi(K(n, m))$. Therefore, we can apply proposition 1 to prove the last claim. \square

Since the Johnson graphs (definition 4) are edge-transitive (see e.g., Chen and Lih [16]) and we can apply theorem 15 to compute the corresponding generalized ϑ -number. The next theorem generalizes theorem 16.

Theorem 17. *For $0 \leq f < m$, $k \leq n$ and the Johnson graph $J(n, m, f)$, it follows*

$$\vartheta_k(J(n, m, f)) = \min \left\{ k\vartheta(J(n, m, f)), \binom{n}{m} \right\}.$$

Proof. Let v denote the order of $J(n, m, f)$, i.e., $v = \binom{n}{m}$. The multiplicities μ_i of the (not necessarily distinct) $m + 1$ eigenvalues λ_i of the adjacency matrix A of $J(n, m, f)$ are

$$\mu_i = \binom{n}{i} - \binom{n}{i-1}, \quad 0 \leq i \leq m,$$

see e.g., Brouwer et al. [11]. We set $\mu_0 = 1$, corresponding to the multiplicity of λ_1 . The multiplicities are unordered, that is, the multiplicity of λ_i does not necessarily equal μ_i . When $n \leq 4$, the theorem can be verified numerically. We will now assume that $n > 4$. Then $\mu_i \geq n - 1$, and in particular, the multiplicity of λ_v is at least $n - 1$. From theorem 15 it follows that

$$\vartheta_k(J(n, m, f)) = \min \mathbf{S}_k(\sigma(J + xA)).$$

Define $f(x) := \mathbf{S}_k(\sigma(J + xA))$. Because the multiplicity of λ_v is at least $n - 1$, for $k \leq n$, we have that $x^* = n/(\lambda_v - \lambda_1)$ minimizes $f(x)$, and $f(x^*) = v + x^*(\lambda_1 + (k - 1)\lambda_v)$. If $f(x^*) > n$, the minimum will occur at $f(0) = n$. \square

We can explicitly compute $\vartheta_k(J(n, m, m - 1))$. Taking the eigenvalues of this graph from [11], chapter 9, we find

$$\vartheta_k(J(n, m, m - 1)) = \frac{k}{n+1} \binom{n+1}{m}, \quad k \leq n - m + 1.$$

For $k^* = n - m + 1$, we see that $\vartheta_{k^*}(J(n, m, m - 1))$ equals the number of vertices in $J(n, m, m - 1)$.

5.2 Relation between $\vartheta(K_k \square G)$ and $\vartheta_k(G)$

Gvozdenović and Laurent [35] show how to exploit an upper bound on the independence number of a graph to obtain a lower bound for the chromatic number of the graph. They do not consider the generalized ϑ -number in the bounding procedure. Kuryatnikova et al. [48] exploit the generalized ϑ -number to compute bounds on the chromatic number of a graph. For some graphs, the lower bounds on $\chi(G)$ from [48] coincide with the bounds obtained by using the theta function as suggested by [35]. Here we explain that finding by analyzing $\vartheta(K_k \square G)$ for symmetric graphs. We also show that the gap between $\vartheta_k(G)$ and $\vartheta(K_k \square G)$ can be arbitrarily large.

Chvátal [17] noted that

$$\alpha_k(G) = |V(G)| \iff \chi(G) \leq k.$$

Stated differently, $\chi(G) = \min\{k \mid k \in \mathbb{N}, \alpha_k(G) = |V(G)|\}$, or in plain words, the k independent sets of $\alpha_k(G)$ correspond to the color classes of G in an optimal coloring. Analogue to $\chi_k(G) = \chi(G \circ K_k)$, it is known (cf. [48]) that

$$\alpha_k(G) = \alpha(K_k \square G), \quad (35)$$

where $K_k \square G$ is the graph Cartesian product, see definition 1. For a graph parameter $\beta(G)$ that satisfies

$$\alpha(G) \leq \beta(G) \leq \chi(\overline{G}),$$

Gvozdenović and Laurent [35] define $\Psi_\beta(G)$ as follows:

$$\Psi_\beta(G) := \min\{k \mid k \in \mathbb{N}, \beta(K_k \square G) = |V(G)|\}.$$

Then $\Psi_\alpha(G) = \chi(G)$. The operator $\Psi_\beta(\cdot)$ can be applied to a variety of graph parameters $\beta(G)$, and enables obtaining a hierarchy of bounds for $\chi(G)$ from a hierarchy of bounds for $\alpha(G)$. For example, when $\beta(G) = \vartheta(G)$ Gvozdenović and Laurent [35] show that $\Psi_\vartheta(G) = \lceil \vartheta(\overline{G}) \rceil$. It follows from (35) that parameters $\vartheta_k(G)$ and $\vartheta(K_k \square G)$ both provide upper bounds on $\alpha_k(G)$. Therefore, it is natural to compare $\Psi_\vartheta(G)$ with

$$\Psi_{\vartheta_k}(G) = \min\{k \mid k \in \mathbb{N}, \vartheta_k(G) = |V(G)|\}. \quad (36)$$

This comparison boils down to the comparison of $\vartheta(K_k \square G)$ and $\vartheta_k(G)$. Numerical results in [48] suggest the following conjecture.

Conjecture 1. *For any graph G and any natural number k , $\vartheta(K_k \square G) \leq \vartheta_k(G)$. Equality holds when $\vartheta_k(G) = k\vartheta(G)$.*

We show below that the gap between $\vartheta_k(G)$ and $\vartheta(K_k \square G)$ can be made arbitrarily large. We first state the following lemma that is needed in the rest of this section.

Lemma 3 (Gvozdenović and Laurent [35]). *Given $A, B \in \mathbb{S}^n$ and $Y = I_k \otimes A + (J_k - I_k) \otimes B$, then $Y \succeq 0$ if and only if $A - B \succeq 0$ and $A + (k-1)B \succeq 0$. Furthermore, $\sigma(Y) = \sigma(A + (k-1)B) \cup \sigma(A - B)^{\{k-1\}}$.*

Now, we are ready to present our result.

Proposition 4. *For any positive number $M \geq 0$, there exists a graph and integer k such that*

$$\vartheta_k(G) - \vartheta(K_k \square G) \geq M.$$

Proof. Consider again graph \mathcal{G}_n , as defined in (16) for even n and set $k = n/2$. We will show that $\vartheta_{n/2}(\mathcal{G}_n) - \vartheta(K_{n/2} \square \mathcal{G}_n)$ is increasing in n . Let $p = 1/(2\sqrt{n-2})$ and consider first

$$X = \begin{bmatrix} \frac{1}{2}I_{n-2} & \mathbf{0}_{n-2} & p\mathbf{1}_{n-2} \\ \mathbf{0}_{n-2}^\top & \frac{1}{2} & 0 \\ p\mathbf{1}_{n-2}^\top & 0 & \frac{1}{2} \end{bmatrix}.$$

Taking the Schur complement of the bottom right 2×2 block of X shows that $0 \preceq X \preceq I$ (see the proof of theorem 8 for more details). Combined with the fact that $\langle I, X \rangle = k$, it follows that X is feasible for ϑ_k -SDP2. Hence,

$$\vartheta_{n/2}(\mathcal{G}_n) \geq \langle J, X \rangle = n/2 + \sqrt{n-2}. \quad (37)$$

As for $\vartheta(K_{n/2} \square \mathcal{G}_n)$, let

$$A = \begin{bmatrix} -kJ_{n-1} + (k+1)I_{n-1} & \mathbf{1}_{n-1} \\ \mathbf{1}_{n-1}^\top & 1 \end{bmatrix}, \quad B = \begin{bmatrix} J_{n-1} & \mathbf{1}_{n-1} \\ \mathbf{1}_{n-1}^\top & -k \end{bmatrix},$$

and set $Y := I \otimes A + (J - I) \otimes B$. Then matrix $Y \in \mathcal{A}(K_{n/2} \square \mathcal{G}_n)$, see (7). Furthermore, matrix Y is of the form described in lemma 3. Then the largest eigenvalue of Y satisfies $\lambda_1(Y) = \max\{\lambda_1(A - B), \lambda_1(A + (k-1)B)\}$. Similar to the methods used in the proof of theorem 8, it can be shown that $\lambda_1(Y) = k + 1$. Thus,

$$\vartheta(K_{n/2} \square \mathcal{G}_n) \leq \lambda_1(Y) = n/2 + 1. \quad (38)$$

Combining (37) and (38) for fixed M and large enough (even) n , gives $\vartheta_{n/2}(\mathcal{G}_n) - \vartheta(K_{n/2} \square \mathcal{G}_n) \geq \sqrt{n-2} - 1 \geq M$. \square

We prove conjecture 1 only for a particular class of graphs. Let us first show the following result.

Theorem 18. *Let G be graph of order n that is both edge-transitive and vertex-transitive. Then*

$$\vartheta(K_k \square G) = \min\{k\vartheta(G), n\}.$$

Proof. For notational convenience we denote $A = A_G$. Because G is regular, edge-transitive and vertex-transitive, we may assume without loss of generality that $\mathcal{A}(K_k \square G)$, see (7), contains matrices of the form $X = I_k \otimes (J_n + xA) + (J_k - I_k) \otimes (J_n + yI_n)$. In order to minimize the largest eigenvalue of X we apply lemma 3 and find

$$\begin{aligned} \lambda_1(X) = f(x, y) &= \max \left\{ \lambda_1(xA - yI_n), \lambda_1(kJ_n + xA + (k-1)yI_n) \right\} \\ &= \max \begin{cases} f_1 = x\lambda_1 - y, \\ f_2 = x\lambda_n - y, \\ f_3 = kn + x\lambda_1 + (k-1)y, \\ f_4 = x\lambda_n + (k-1)y. \end{cases} \end{aligned}$$

We minimize $\lambda_1(X)$ by considering different intervals of x . In case $x \geq 0$, $f(0, -n) = n$ is the minimum. Furthermore,

$$\frac{kn}{\lambda_n - \lambda_1} \leq x \leq 0 \implies f(x, y) = \max\{f_2, f_3\}.$$

The minimum here is attained when

$$f_2 = f_3 \implies y = x \left(\frac{\lambda_n - \lambda_1}{k} \right) - n \implies f_2 = n + \frac{1}{k}((k-1)\lambda_n + \lambda_1)x.$$

Depending on the sign of $(k-1)\lambda_n + \lambda_1$ we find either $f(0, -n) = n$ or $f(\frac{kn}{\lambda_n - \lambda_1}, 0) = k\frac{n\lambda_n}{\lambda_n - \lambda_1} = k\vartheta(G)$, by theorem 14. Lastly, the case $x < \frac{kn}{\lambda_n - \lambda_1}$ leads again to $k\vartheta(G)$. The minimum value of $\lambda_1(X)$, equivalently, the value $\vartheta(K_k \square G)$, thus equals $\min\{k\vartheta(G), n\}$. \square

As a corollary of theorem 18, conjecture 1 holds when G is both edge-transitive and vertex-transitive (every vertex-transitive graph is regular) and moreover $\vartheta_k(G) = k\vartheta(G)$. A graph that is both edge-transitive and vertex-transitive is also known as a symmetric graph. Many Johnson graphs (definition 4) satisfy these properties.

Kuryatnikova et al. [48] compute $\vartheta_k(G)$ for several highly symmetric graphs (table 13 in [48]). They remark that for those graphs, $\Psi_{\vartheta_k}(G) = \lceil \vartheta(\overline{G}) \rceil$. We explain this result for all the graphs present in table 13 except for the graph $H(12, 2, \{i \mid 1 \leq i \leq 7\})$ (see section 8.1 for the notation). All the other graphs evaluated by Kuryatnikova et al. in table 13 satisfy the assumption of theorem 18. The graphs then satisfy the inequality of theorem 15 with equality. Applying theorem 15 with the explicit eigenvalues of the adjacency matrices of the graphs in table 13 shows that $\vartheta_k(G) = k\vartheta(G)$. Consequently, $\vartheta_k(G) = \vartheta(K_k \square G)$ for those graphs. Therefore,

$$\Psi_{\vartheta_k}(G) = \Psi_{\vartheta}(G) = \lceil \vartheta(\overline{G}) \rceil. \quad (39)$$

Note that the Johnson graph $J(n, m, \{m-1\})$ is regular, vertex-transitive and edge-transitive. Therefore, equation (39) holds and $\lceil \vartheta(J(n, m, m-1)) \rceil = n - m + 1$.

6 Strongly regular graphs

In the previous section we show that certain classes of graphs allow an analytical computation of $\vartheta_k(G)$. This section expands on the considered classes with strongly regular graphs, see definition 6. We also derive analogous of theorem 18 for strongly regular graphs, see theorem 20.

It is known that strongly regular graphs attain Lovász bound of theorem 14, see e.g., Haemers [37]. In particular, for a strongly regular graph G we have

$$\vartheta(G) = \frac{n\lambda_n(A_G)}{\lambda_n(A_G) - \lambda_1(A_G)}.$$

In the following theorem we derive an explicit expression for $\vartheta_k(G)$ for strongly regular graphs.

Theorem 19. For any strongly regular graph G of order n , we have

$$\vartheta_k(G) = \min\{k\vartheta(G), n\} = \min\left\{k\frac{n\lambda_n(A_G)}{\lambda_n(A_G) - \lambda_1(A_G)}, n\right\}.$$

Proof. Let G be a strongly regular graph with parameters (n, d, λ, μ) . We prove the result by showing that the lower and upper bound on $\vartheta_k(G)$ coincide. Consider ϑ_k -SDP2, and set $Y = \frac{k}{n}I + xA_{\overline{G}}$. When $0 \preceq Y \preceq I$, Y is feasible for ϑ_k -SDP2. These SDP constraints on Y can be rewritten in terms of x . As ϑ_k -SDP2 is a maximization problem we may assume w.l.g. $x \geq 0$. Thus, for all $i \leq n$,

$$\lambda_i(Y) = k/n + x\lambda_i(A_{\overline{G}}). \quad (40)$$

It is known that adjacency matrices of (connected) strongly regular graphs have three distinct eigenvalues, which we denote by $d \geq r > s$. Accordingly,

$$\sigma(A_{\overline{G}}) = (n - d - 1, -1 - s, -1 - r). \quad (41)$$

Substituting (41) in (40) and exploiting the fact that $n - d - 1 > -(s + 1) \geq -(1 + r)$ we have:

$$0 \preceq Y \preceq I \implies 0 \leq \lambda_i(Y) \leq 1 \implies \begin{cases} k/n + x(-1 - r) \geq 0 \\ k/n + x(n - d - 1) \leq 1. \end{cases}$$

The last two inequalities provide upper bounds on x , i.e.,

$$x \leq \min\left\{\frac{k}{n(1+r)}, \frac{n-k}{n(n-d-1)}\right\}. \quad (42)$$

When x satisfies (42), Y is thus feasible for ϑ_k -SDP2 and $\langle J, Y \rangle$ will provide a lower bound for $\vartheta_k(G)$. In particular,

$$\langle J, Y \rangle = k + n(n - d - 1)x = \min\left\{k\left(\frac{r + n - d}{1 + r}\right), n\right\}. \quad (43)$$

Equation (43) implies

$$\vartheta_k(G) \geq \min\left\{k\left(\frac{r + n - d}{1 + r}\right), n\right\}. \quad (44)$$

By (9) and proposition 1, we have $\vartheta_k(G) \leq \min\{k\vartheta(G), n\}$. It remains only to show that $k(r + n - d)/(1 + r) = k\vartheta(G)$. The eigenvalues of A_G can be written in terms of the parameters of G , i.e.,

$$rs = \mu - d, \quad r + s = \lambda - \mu. \quad (45)$$

Furthermore, the parameters of any strongly regular graph satisfy

$$(n - d - 1)\mu = d(d - \lambda - 1). \quad (46)$$

Let us now rewrite the term:

$$\frac{r + n - d}{1 + r} = \frac{ns}{s - d} \frac{(r + n - d)(s - d)}{ns(1 + r)} = \frac{ns}{s - d} \frac{ns + nrs + [d^2 - nd - (n - 1)sr - d(r + s)]}{ns + nrs}, \quad (47)$$

and evaluate the expression between the square brackets by using (45) and (46), i.e.,

$$d^2 - nd - (n - 1)rs - d(r + s) = d\lambda + d + (n - d - 1)\mu - nd - (n - 1)(\mu - d) - d(\lambda - \mu) = 0.$$

Thus (47) equals $ns/(s - d)$ and $ns/(s - d) = \vartheta(G)$, which proves the theorem. \square

Recall that in section 5.2 we consider symmetric graphs (graphs that are both edge-transitive and vertex-transitive). Although many graphs belong to both symmetric and strongly regular classes, note that neither one is a subset of the other. The graph C_6 is an example of a graph that is symmetric, but not strongly regular. The strongly regular Chang graphs (Chang [15]) provide an example of a strongly regular graph which is not symmetric.

In section 5.2 we prove that $\vartheta(K_k \square G) = \min\{k\vartheta(G), n\}$ holds for symmetric graphs, see theorem 18. We show below that a similar relation holds also for strongly regular graphs. In fact we prove a result for the generalized ϑ' -number, denoted by $\vartheta'_k(G)$, that is the optimal value of the SDP relaxation ϑ_k -SDP2 strengthened by adding non-negativity constraints on the matrix variable. The generalized ϑ' -number for $k = 1$ is also known as the Schrijver's number.

To prove our result we first present an SDP relaxation that relates $\vartheta'(K_k \square G)$ and $\vartheta_k(G)$. Kuryatnikova et al. [48] introduce the following SDP relaxation

$$\begin{aligned} \theta_k^3(G) = & \text{Maximize} && \langle I, Y \rangle \\ & \text{subject to} && Y_{ij} = 0 \quad \forall (i, j) \in E(G) \\ & && Y_{ii} \leq 1 \quad \forall i \in [n] \\ & && \begin{bmatrix} k & \text{diag}(Y)^\top \\ \text{diag}(Y) & Y \end{bmatrix} \succeq 0, Y \geq 0, \end{aligned}$$

that provides an upper bound for $\alpha_k(G)$, the optimal value for the MkCS problem. The above relaxation can be simplified when G is a highly symmetric graph. In particular, if G is a strongly regular graph one can restrict optimization of the above SDP relaxation to feasible points in the coherent algebra spanned by $\{I, A, J - I - A\}$. By applying symmetry reduction, the above SDP relaxation reduces to the following convex optimization problem:

$$\begin{aligned} \theta_k^3(G) := & \text{Maximize} && ny_1 && (48a) \\ & \text{subject to} && y_1 + (n - d - 1)y_2 - \frac{n}{k}y_1^2 \geq 0 && (48b) \\ & && y_1 - (r + 1)y_2 \geq 0 && (48c) \\ & && y_1 - (s + 1)y_2 \geq 0 && (48d) \\ & && y_1 \leq 1 && (48e) \\ & && y_1, y_2 \geq 0. && (48f) \end{aligned}$$

For details on symmetry reduction see e.g., [28, 48] and references therein.

Lemma 4. *Let G be a strongly regular graph with parameters (n, d, λ, μ) and restricted eigenvalues $r \geq 0$ and $s \leq -1$. Then*

$$\theta_k^3(G) = \min \left\{ k \left(\frac{n + r - d}{r + 1} \right), n \right\},$$

which equals $\vartheta_k(G)$.

Proof. Note that for $s \leq -1$ constraint (48d) is trivially satisfied. Points in which constraints (48b) and (48c) intersect are $(0, 0)$ and $(\frac{k(n+r-d)}{n(r+1)}, \frac{k(n+r-d)}{n(r+1)^2})$. The result follows by combining the latter point and constraint (48e). \square

In [48] the authors conjecture that $\theta_k^3(G) \leq \vartheta'_k(G)$ for any graph G . Here we show that $\theta_k^3(G) = \vartheta'_k(G) = \vartheta_k(G)$ for strongly regular graphs.

Further, it is known that $\vartheta'(K_k \square G) \leq \theta_k^3(G)$, see section 5.1 in [48]. We show below that the relaxations corresponding to $\vartheta'(K_k \square G)$ and $\theta_k^3(G)$ are equivalent when G is a strongly regular graph.

The SDP relaxation for $\vartheta'(K_k \square G)$, see also ϑ_k -SDP2, is invariant under permutations of k colors when the graph under the consideration is $K_k \square G$. This was exploited in [48] to derive the following symmetry reduced relaxation:

$$\begin{aligned} \vartheta'(K_k \square G) = & \text{Maximize} && \langle I, X \rangle \\ & \text{subject to} && X_{ij} = 0 \quad \forall (i, j) \in E(G) \\ & && Z_{ii} = 0 \quad \forall i \in [n] \\ & && X \geq 0, Z \geq 0, X - Z \succeq 0 \\ & && \begin{bmatrix} 1 & \text{diag}(X)^\top \\ \text{diag}(X) & X + (k - 1)Z \end{bmatrix} \succeq 0. \end{aligned}$$

The above relaxation can be further simplified when G is a strongly regular graph. One can restrict optimization to the corresponding coherent algebra. By applying symmetry reduction, the above SDP relaxation reduces to the following optimization problem:

$$\vartheta'(K_k \square G) := \text{Maximize} \quad nx_1 \quad (49a)$$

$$\text{subject to} \quad x_1 + (n-d-1)x_2 - (dz_1 + (n-d-1)z_2) \geq 0 \quad (49b)$$

$$x_1 - (r+1)x_2 - (rz_1 - (r+1)z_2) \geq 0 \quad (49c)$$

$$x_1 - (s+1)x_2 - (sz_1 - (s+1)z_2) \geq 0 \quad (49d)$$

$$x_1 + (n-d-1)x_2 + (k-1)(dz_1 + (n-d-1)z_2) - nx_1^2 \geq 0 \quad (49e)$$

$$x_1 - (r+1)x_2 + (k-1)(rz_1 - (r+1)z_2) \geq 0 \quad (49f)$$

$$x_1 - (s+1)x_2 + (k-1)(sz_1 - (s+1)z_2) \geq 0 \quad (49g)$$

$$x_1 \leq 1 \quad (49h)$$

$$x_1, x_2, z_1, z_2 \geq 0. \quad (49i)$$

Our next results relates optimization problems (48) and (49).

Proposition 5. *Let G be a strongly regular graph with parameters (n, d, λ, μ) and restricted eigenvalues $r \geq 0$ and $s \leq -1$. Then the optimization problems (48) and (49) are equivalent.*

Proof. Let (x_1, x_2, z_1, z_2) be feasible for (49). We show that (y_1, y_2) where $y_1 := x_1$ and $y_2 := x_2$ is feasible for (48).

From (49b) and (49e) we have

$$\begin{cases} x_1 + (n-d-1)x_2 \geq (dz_1 + (n-d-1)z_2) \\ x_1 + (n-d-1)x_2 \geq nx_1^2 - (k-1)(dz_1 + (n-d-1)z_2), \end{cases}$$

from where it follows

$$x_1 + (n-d-1)x_2 - \frac{n}{k}x_1^2 \geq \max \left\{ (dz_1 + (n-d-1)z_2) - \frac{n}{k}x_1^2, (k-1)\left(\frac{n}{k}x_1^2 - (dz_1 + (n-d-1)z_2)\right) \right\}.$$

To verify that the right hand side above is non-negative, note that either $dz_1 + (n-d-1)z_2 \geq \frac{n}{k}x_1^2$ or $dz_1 + (n-d-1)z_2 < \frac{n}{k}x_1^2$. Therefore $x_1 + (n-d-1)x_2 - \frac{n}{k}x_1^2 \geq 0$ and constraint (48b) is satisfied.

Similarly, from (49c) and (49f) it follows that constraint (48c) is satisfied. Constraint (48d) is trivially satisfied.

Conversely, let (y_1, y_2) be feasible for (48). Define $x_1 := y_1$ and $x_2 := y_2$. Let z_1 and z_2 be the solutions of the following system of equations:

$$rz_1 = (r+1)z_2, \quad dz_1 + (n-d-1)z_2 = \frac{n}{k}x_1^2.$$

Thus, $z_1 = \frac{n(r+1)}{k(d+r(n-1))}x_1^2$, $z_2 = z_1 \frac{r}{r+1}$. Therefore, constraint (49b) follows from (48b) and the construction of z_1 and z_2 . Similar arguments can be used to verify that (49c) and (49f) are satisfied. To verify (49e) we rewrite the constraint as follows

$$\begin{aligned} x_1 + (n-d-1)x_2 + (k-1)(dz_1 + (n-d-1)z_2) - nx_1^2 = \\ x_1 + (n-d-1)x_2 - \frac{n}{k}x_1^2 + (k-1)(dz_1 + (n-d-1)z_2 - \frac{n}{k}x_1^2) \geq 0. \end{aligned}$$

To verify constraint (49d) we exploit the construction of z_1 and z_2 to obtain: $-(sz_1 - (s+1)z_2) = \frac{r-s}{r+1}z_1 \geq 0$. It remains to show that constraint (49g) is redundant. We first rewrite the constraint as follows

$$x_1 - (s+1)x_2 + (k-1)(sz_1 - (s+1)z_2) = x_1 - (s+1)x_2 - n \frac{(k-1)(r-s)}{k(d+r(n-1))}x_1^2 \geq 0.$$

Now, one can verify that the above constraint is implied by the constraint $x_1 + (n-d-1)x_2 - \frac{n}{k}x_1^2 \geq 0$.

It follows trivially that the objective values coincide for feasible solutions of two models that are related as described. \square

Now, from the previous discussion it follows the next result.

Theorem 20. *Let G be a strongly regular graph of order n . Then $\vartheta'(K_k \square G) = \min\{k\vartheta'(G), n\}$.*

Proof. The proof follows from lemma 4 and proposition 5. \square

7 Orthogonality graphs

In this section we compute the generalized ϑ -number for the orthogonality graphs.

We motivate the study of orthogonality graphs by a scenario, taken from Galliard et al. [26]. Let $n = 2^r$ for some $r \geq 1$. Consider a game where two players, Alice and Bob, each receive an n -dimensional binary vector as input. These vectors are either equal or their Hamming distance (see definition 2) equals $n/2$ that is they differ in exactly 2^{r-1} positions. Given these inputs, Alice and Bob must each return a r -dimensional binary vector as output. To win the game, Alice and Bob must return equal outputs if and only if their inputs were equal. Alice and Bob are not permitted to communicate once they receive their inputs. The players are however, allowed to coordinate a strategy beforehand. One such strategy results in the definition of an orthogonality graph.

Vertices of the orthogonality graph Ω_n are represented by all the unique n -dimensional binary vectors. Vertices (equivalently vectors) are adjacent if their Hamming distance equals $n/2$ and thus, $\Omega_n = H(n, 2, \{n/2\})$. Here $H(n, 2, \{n/2\})$ denotes the Hamming graph, see definition 3.

The strategy of Alice and Bob then comprises graph coloring for Ω_n before the game starts. After being given their input vector, Alice and Bob should respond the color of their vector, encoded as r -dimensional binary vector. With this r -dimensional vector, Alice and Bob can indicate $2^r = n$ distinct colors. Disregarding any luck in guessing, the game can always be won if and only if $\chi(\Omega_n) \leq n$.

The orthogonality graph got its name from another description of the graph, that is when the vectors have $\{\pm 1\}$ entries. The Hamming distance between two binary vectors of $n/2$ then corresponds to those $\{\pm 1\}$ vectors being orthogonal to each other.

Godsil and Newman [30] prove that $\chi(\Omega_{2^r}) = 2^r$ for $r \in \{1, 2, 3\}$ and $\chi(\Omega_{2^r}) > 2^r$ otherwise. This means that the game can only be won for $r \leq 3$.

Clearly, for odd n , Ω_n is edgeless. We therefore restrict the analysis of Ω_n to the case when n is a multiple of 4. When that is the case, Ω_n consists of two isomorphic components, for vectors of even and odd Hamming weights respectively.

Next to $\chi(\Omega_n)$, the independence number $\alpha(\Omega_n)$ has been studied in multiple papers. The two graph properties are related by $|V| \leq \chi(G)\alpha(G)$, for any graph $G = (V, E)$. Galliard [25] showed a construction to find stable sets in Ω_n of size $\underline{\alpha}(n)$ for $n \equiv 0 \pmod{4}$. In particular,

$$\alpha(\Omega_n) \geq \underline{\alpha}(n) = 4 \sum_{i=0}^{n/4-1} \binom{n-1}{i}. \quad (50)$$

On the other hand de Klerk and Pasechnik [20] used an SDP relaxation to find $\alpha(\Omega_{16}) = \underline{\alpha}(16) = 2306$. More recently, Ihringer and Tanaka [43] have proven $\alpha(\Omega_{2^r}) = \underline{\alpha}(2^r)$ for $r \geq 2$. The conjecture by Godsil and Newman [30] whether $\alpha(\Omega_{4m}) = \underline{\alpha}(4m)$ for $m \geq 1$ remains an open problem.

We proceed by computing $\vartheta_k(\Omega_n)$ when n is a multiple of four. From Newman [64], the (unordered) eigenvalues of Ω_n are then given by

$$\lambda_r = \frac{2^{n/2}}{(n/2)!} \prod_{i=1}^{n/2} (2i - 1 - r), \quad 1 \leq r \leq n,$$

and $\lambda_0 = \binom{n}{n/2}$ (since Ω_n is regular with degree λ_0). The smallest eigenvalue is obtained for $r = 2$ and thus

$$\lambda_2 = \frac{1}{1-n} \binom{n}{n/2}.$$

Since $\Omega(n)$ is isomorphic to a binary Hamming graph, $\Omega(n)$ is a symmetric graph. The bound of theorem 15 thus holds with equality. Newman [64] also shows that the multiplicity of λ_2 equals $n^2 - n$. This multiplicity exceeds n since n is a multiple of 4. Then it is not hard to show (by a method comparable to the one used in the proof of theorem 16) that

$$\vartheta_k(\Omega(n)) = k \frac{2^n}{n}, \quad k \leq n.$$

When $k = 1$, $\vartheta_k(\Omega(n))$ coincides here with the so called ratio bound. This bound refers to (59) for regular graphs and was also computed for $\Omega(n)$ in [64].

Let S be the stable set of size $\underline{\alpha}(n)$ that contains no vectors that have their Hamming weight contained in $W = \{n/4 + 1, n/4 + 3, \dots, 3n/4 - 1\}$. Furthermore, note that the Johnson graphs (see definition

4) appear as induced subgraphs of Ω_n . Let $w \in W$ and consider the subgraph of Ω_n induced by $J(n, w, w - n/4)$. This subgraph contains no vertices in S and thus

$$\alpha_2(\Omega_n) \geq \underline{\alpha}(n) + 4 \max_{w \in W} \{\alpha(J(n, w, w - n/4))\}. \quad (51)$$

We may multiply the independence number of $J(n, w, w - n/4)$ by 4 since we can take bitwise complements and find an isomorphic stable set in the isomorphic second component of Ω_n .

In section 8.1 we prove that that $\chi_k(\Omega_{4n+2}) = 2k$.

8 New bounds on $\chi_k(G)$

In this section we first we derive bounds on the product and sum of $\chi_k(G)$ and $\chi_k(\overline{G})$. Then, we provide graphs for which the bounds are sharp. Lastly, we derive spectral lower bounds on the multichromatic number of a graph.

A famous result by Nordhaus and Gaddum [65] states that

$$n \leq \chi(G)\chi(\overline{G}) \leq \left(\frac{n+1}{2}\right)^2, \quad (52)$$

$$2\sqrt{n} \leq \chi(G) + \chi(\overline{G}) \leq n + 1. \quad (53)$$

Various papers have been published on determining Nordhaus–Gaddum inequalities for other graph parameters, such as the independence and edge-independence number. We provide Nordhaus–Gaddum inequalities for k -multicoloring.

Theorem 21. *For any graph $G = (V, E)$, $|V| = n$, we have*

$$k^2 n \leq \chi_k(G)\chi_k(\overline{G}) \leq k^2 \left(\frac{n+1}{2}\right)^2,$$

$$2k\sqrt{n} \leq \chi_k(G) + \chi_k(\overline{G}) \leq k(n+1).$$

Proof. We follow the original proof as given by Nordhaus and Gaddum [65], extended to the k -multicoloring case. Consider an optimal k -multicoloring of G , using $\chi_k(G)$ colors. Then for $i = 1, 2, \dots, \chi_k(G)$, define n_i as the set of vertices that are colored with color i . We have $\sum_{i=1}^{\chi_k(G)} |n_i| = nk$. Furthermore,

$$\max |n_i| \geq \frac{nk}{\chi_k(G)}. \quad (54)$$

Consider the largest set n_i . Since the vertices in this set share a color, they form a stable set in G . Thus they form a clique in \overline{G} . Accordingly,

$$\chi_k(\overline{G}) \geq k\omega(\overline{G}) \geq k \max |n_i|. \quad (55)$$

Combining (54) and (55) proves the lower bound on the product of $\chi_k(G)$ and $\chi_k(\overline{G})$.

The lower bound on the sum of $\chi_k(G)$ and $\chi_k(\overline{G})$ can be proven by algebraic manipulation:

$$\begin{aligned} (\chi_k(G) - \chi_k(\overline{G}))^2 \geq 0 &\implies \chi_k(G)^2 + \chi_k(\overline{G})^2 + 2\chi_k(G)\chi_k(\overline{G}) \geq 4\chi_k(G)\chi_k(\overline{G}) \\ \implies \chi_k(G) + \chi_k(\overline{G}) &\geq 2\sqrt{\chi_k(G)\chi_k(\overline{G})} \geq 2k\sqrt{n}. \end{aligned}$$

The two upper bounds can be proven by combining (5), (52) and (53) as follows:

$$\begin{aligned} \chi_k(G)\chi_k(\overline{G}) &\leq k^2\chi(G)\chi(\overline{G}) \leq k^2\left(\frac{n+1}{2}\right)^2, \\ \chi_k(G) + \chi_k(\overline{G}) &\leq k(\chi(G) + \chi(\overline{G})) \leq k(n+1). \end{aligned}$$

□

The second upper bound in theorem 21 can also be found in [8] (in a slightly generalized form). We present below graphs for which the bounds in theorem 21 are attained. For that purpose we define the

graph sum of two graphs. The graph sum of graphs G_1 and G_2 is the graph, denoted by $G_1 + G_2$, whose vertices and edges are defined as follows:

$$V(G_1 + G_2) := V(G_1) \cup V(G_2), \quad E(G_1 + G_2) := E(G_1) \cup E(G_2).$$

Nordhaus and Gaddum [65] show that the upper bounds in their theorem are attained by graph $G = K_p + \overline{K}_{p-1}$. Graph G has $n = 2p - 1$ vertices. It is clear that $\chi(G) = \chi(\overline{G}) = p = \frac{n+1}{2}$. Thus G attains both upper bounds simultaneously. As both G and \overline{G} are weakly perfect graphs, we can apply (60) to find $\chi_k(G) = \chi_k(\overline{G}) = kp = k\frac{n+1}{2}$. This implies that graph G also attains the upper bounds in theorem 21. Nordhaus and Gaddum [65] also provide an example of a graph which attains the lower bounds in their theorem. This example extends to the multichromatic variant as well. Let $m_1 = m_2 = \dots = m_p = p$ and consider the complete multipartite graph $G = K_{m_1, \dots, m_p}$. Then $\chi_k(G) = \chi_k(\overline{G}) = kp = k\sqrt{n}$. Thus this graph G attains the lower bounds in theorem 21. In fact, for each graph G such that $\chi(G)\chi(\overline{G}) = |V(G)|$ we have that $k^2|V(G)| = \chi_k(G)\chi_k(\overline{G})$. The set of vertex-transitive graphs provides a number of examples for which this bound is attained, such as the Johnson graph $J(n, 2, 1)$ when n is even.

The chromatic number of a graph is bounded by the spectrum of matrices related to its adjacency matrix. This well known result is given below.

Theorem 22 (Hoffman [41]). *If G has at least one edge, then*

$$\chi(G) \geq 1 - \frac{\lambda_1(A_G)}{\lambda_n(A_G)}.$$

Since each color class has size at most $\alpha(G)$, we have that

$$\chi(G) \geq \frac{n}{\alpha(G)}, \quad (56)$$

where n is the number of vertices in G . Therefore one can use upper bounds for $\alpha(G)$ to derive lower bounds for $\chi(G)$. From (3) it follows that $\alpha(G \circ K_k) = \alpha(G)$. Thus we can establish the multicoloring variant of (56):

$$\chi_k(G) = \chi(G \circ K_k) \geq \frac{|V(G \circ K_k)|}{\alpha(G \circ K_k)} = k \frac{n}{\alpha(G)}. \quad (57)$$

Note that the above result also follows from (54). The bound (57) is also given in [12], where the authors show that the lower bound is tight for webs and antiwebs. Note that for a graph G such that $\alpha_k(G) = k\vartheta(G)$ we have that $\alpha_k(G) = k\alpha(G)$, see lemma 5 in [48], and thus $\chi_k(G) \geq \frac{k^2 n}{\alpha_k(G)}$. The above inequality is satisfied for example for the Johnson graph $J(n, 2, 1)$ when n is even, and for $J(n, 3, 2)$ when $v \equiv 1$ or $3 \pmod{6}$.

Let us now present known upper bounds for the independence number of a graph.

Theorem 23 (Hoffman [41]). *For any d -regular graph G of order n , we have*

$$\alpha(G) \leq n \frac{\lambda_n(A_G)}{\lambda_n(A_G) - d}.$$

The result of theorem 23 applies only to regular graphs with no loops. Haemers generalizes the Hoffman bound as follows.

Theorem 24 (Haemers [36]). *Let G have minimum vertex degree δ . Then*

$$\alpha(G) \leq n \frac{\lambda_1(A_G)\lambda_n(A_G)}{\lambda_1(A_G)\lambda_n(A_G) - \delta^2}. \quad (58)$$

If G is regular then the result of theorem 24 reduces to Hoffman's bound. Another extension of the bound of Hoffman is given by Godsil and Newman.

Theorem 25 (Godsil and Newman [31]). *Let G be a loopless graph and L_G its Laplacian matrix. Then*

$$\alpha(G) \leq n \frac{\lambda_1(L_G) - \overline{d}_G}{\lambda_1(L_G)}, \quad (59)$$

where \overline{d}_G denotes the average degree of the vertices of G .

Now we are ready to present our results.

Lemma 5. *Let G have minimum vertex degree δ . Then*

$$\chi_k(G) \geq k \frac{\lambda_1(A_G)\lambda_n(A_G) - \delta^2}{\lambda_1(A_G)\lambda_n(A_G)}.$$

Proof. The result follows by combining (57) and (58). \square

Lemma 6. *For any loopless graph G , we have*

$$\chi_k(G) \geq k \frac{\lambda_1(L_G)}{\lambda_1(L_G) - \bar{d}_G},$$

where \bar{d}_G denotes the average degree of its vertices, and L_G the Laplacian matrix of G .

Proof. The result follows by combining (57) and (59). \square

When G is a regular graph, (59) is equivalent to the result of theorem 23, and therefore the result of lemma 6 is equivalent to:

$$\chi_k(G) \geq k \left(1 - \frac{\lambda_1(A_G)}{\lambda_n(A_G)} \right).$$

It is not difficult to verify that complete graphs attain the bound of lemma 5 and lemma 6.

We end this section by presenting bounds on the multichromatic number of the Johnson graphs, see definition 4. We study the simple case $J(n, 2, 1)$, $n \geq 4$. Graph $J(n, 2, 1)$ is sometimes referred to as the triangular graph. The graph $J(n, 2, 1)$ is the complement graph of the Kneser graph $K(n, 2)$, and both are known to be strongly regular. Every vertex of $J(n, 2, 1)$ corresponds to a set of two elements. These two elements can be thought of as two vertices of the complete graph K_n , with the vertex in $J(n, 2, 1)$ representing the edge between these two vertices of K_n . Graph $J(n, 2, 1)$ is thus the line graph of the complete graph K_n . For any graph G , its line graph is denoted $L(G)$.

Proposition 6. *For the triangular graph $J(n, 2, 1)$ and $n \geq 4$, we have*

$$k(n-1) \leq \chi_k(J(n, 2, 1)) \leq k(2 \lfloor \frac{n-1}{2} \rfloor + 1).$$

Proof. As $J(n, 2, 1)$ is isomorphic to $L(K_n)$, a coloring of $J(n, 2, 1)$ is equivalent to an edge coloring of K_n . It is not hard to see that $\omega(L(G))$ equals the maximum degree of a vertex of G . Thus $\omega(L(K_n)) = n-1$. For even n , $\chi(L(K_n)) = n-1$, see Baranyai [6]. Therefore, for even n we have

$$\chi(L(K_n)) = \omega(L(K_n)) \implies \chi_k(L(K_n)) = k\chi(L(K_n)) = k(n-1).$$

Where the implication follows from (5). For odd n , $\chi(L(K_n)) = n$, see Vizing [78]. By (5), the proposition follows. \square

Note that in the proof of the previous proposition we could also exploit the following well known result: $\alpha(K(n, 2)) = n-1$.

8.1 Hamming graphs

In this section we present results for the (multi)chromatic number of the Hamming graphs (definition 3). We also provide sufficient and necessary conditions for the Hamming graph to be perfect.

In the Hamming graph $H(n, q, F)$, the vertex set is the set of n -tuples of letters from an alphabet of size q , and vertices u and v are adjacent if their Hamming distance satisfies $d(u, v) \in F$. Note that $|V(H(n, q, F))| = q^n$. By slight abuse of notation, we will use the terms vectors and vertices interchangeably, as they permit a one-to-one correspondence in Hamming graphs. Many authors refer to $H(n, q, \{1\})$ as the Hamming graph. The graph $Q_n := H(n, 2, \{1\})$ is also known as the binary Hamming graph or hypercube graph.

We first list several known results for $H(n, q, \{1\})$. Graph $H(n, q, \{1\})$ equals the Cartesian product of n copies of K_q . Thus $H(n, q, \{1\}) = \square^n K_q$, see definition 1. Furthermore, it holds $\chi(G_1 \square G_2) = \max\{\chi(G_1), \chi(G_2)\}$, see Sabidussi [69]. Therefore, $\chi(H(n, q, \{1\})) = q$. To derive the independence

number of $H(n, q, \{1\})$, we proceed as follows. Let $S \subset V$ be a stable set of $H(n, q, \{1\})$. Then

$\min_{u, v \in S, u \neq v} d(u, v) \geq 2$. From coding theory, the Singleton bound (Singleton [73]) is an upper bound on the maximum number of codes of length n , using an alphabet of size q , such that a Hamming distance between any two codes is at least two. In particular, from the Singleton bound we have $\alpha(H(n, q, \{1\})) \leq q^{n-1}$.

To show that $\alpha(H(n, q, \{1\})) \geq q^{n-1}$, we construct an independent set in the Hamming graph of size q^{n-1} , by a construction employed in [73]. Consider all the vectors in $(\mathbb{Z}/q\mathbb{Z})^n$ for which the coordinates sum to some $x \in \mathbb{Z}/q\mathbb{Z}$. By symmetry, there exist q^{n-1} vectors satisfying this condition. Note that any two different vectors satisfying this condition must differ in at least two positions, which implies they are not adjacent. Thus, $\alpha(H(n, q, \{1\})) \geq q^{n-1}$ and combined with the Singleton bound, this gives $\alpha(H(n, q, \{1\})) = q^{n-1}$.

To the best of our knowledge, the following results are not known in the literature.

Lemma 7. For $k \leq q^n$, $\chi_k(H(n, q, \{1\})) = kq$.

Proof. Let us denote $H = H(n, q, \{1\})$. Consider the vectors in H for which the first entry ranges from 0 up to and including $q - 1$, while the other entries equal 0. This gives a clique of size q and since $\omega(H) \leq \chi(H) = q$, we have $\omega(H) = q$. From (5), it follows the result. \square

The proof of lemma 7 relies on the fact that $\omega(H) = \chi(H)$, or equivalently, that $H(n, q, \{1\})$ is a weakly perfect graph. In general, for any weakly perfect graph G

$$\omega(G) = \chi(G) \implies \chi_k(G) = k\chi(G). \quad (60)$$

This gives rise to the question for which values of q and n the graph $H(n, q, \{1\})$ is perfect. The strong perfect graph theorem states that a graph is perfect if and only if it does not contain C_{2n+1} or \overline{C}_{2n+1} as induced subgraphs, for all $n > 1$.

Proposition 7. The Hamming graph $H(n, q, \{1\})$ is a perfect graph if and only if $n \leq 2$ or $q \leq 2$.

Proof. Denote $H(n, q) = H(n, q, \{1\})$. Graph $H(1, q)$ is K_q , which is clearly a perfect graph. Graph $H(2, q)$ is a lattice graph, or Rook's graph, which is also a perfect graph. Graph $H(n, 1)$ is a single vertex and thus also perfect. Lastly, graph $H(n, 2)$ is bipartite and thus perfect. For $q \geq 3$, the following vectors from $H(3, q)$ form C_7 :

$$\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}.$$

Then by the strong perfect graph theorem, $H(3, q)$ is not perfect. An odd cycle in $H(n, q)$ for general $n, q \geq 3$ is obtained by simply adjoining zeros to the above vectors such that they become n -dimensional. \square

As $H(n, q, \{f\})$ is edgeless for $f > n$, we consider the extremal case $H(n, q, \{n\})$ for $n > 1$. Note that $H(1, q, \{1\}) = K_q$. Graph $H(n, q, \{n\})$ can be described by use of the tensor product of graphs (see definition 1). In particular, we have that $H(n, q, \{n\}) = \otimes^n K_q$.

Since all the edges of $G_1 \otimes G_2$ also appear in $G_1 \circ G_2$, it follows that $\chi(G_1 \otimes G_2) \leq \chi(G_1 \circ G_2)$. Moreover, by Hedetniemi [39], we have

$$\chi(G_1 \otimes G_2) \leq \min\{\chi(G_1), \chi(G_2)\}. \quad (61)$$

Hedetniemi's conjecture states that (61) holds with equality. The conjecture was recently disproved by Shitov [72]. Inequality (61) implies that $\chi(\otimes^n K_q) \leq q$.

The vectors $i \cdot \mathbf{1}$ for $0 \leq i \leq q-1$ form a clique of size q in graph $H(n, q, \{n\})$. Thus $q \leq \omega(H(n, q, \{n\}))$. Now, from this inequality and $\chi(\otimes^n K_q) \leq q$ it follows that $\chi(H(n, q, \{n\})) = q$. Using (5) or (60), we find $\chi_k(H(n, q, \{n\})) = kq$. The coloring of these tensor products of graphs has been previously considered by Greenwell and Lovász [32], where they also proved this result.

Let us now define $f^+ := \{i \in \mathbb{N} \mid f \leq i\}$. The Hamming graph $H(n, q, f^+)$ has been studied by El Rouayheb et al. [21] among others. They show that, under some condition on the parameters n, q and f , $\chi(H(n, q, f^+)) = q^{n-f+1}$. We extend this result to multicoloring in the following proposition.

Proposition 8. For $q \geq n - f + 2$ and $1 \leq f \leq n$, we have $\chi_k(H(n, q, f^+)) = kq^{n-f+1}$.

Proof. For parameters n, q and f satisfying the conditions of the proposition, it is known (cf. [23]) that $\alpha(H(n, q, f^+)) = q^{f-1}$. By (5) and (57), the proposition follows. \square

Binary Hamming graphs $H(n, 2, \{f\})$, $f \leq d$ form another interesting case. Recall that the Hamming weight of a vector is its Hamming distance to the zero vector, and that the Hamming graphs are vertex-transitive.

Theorem 26. *For all $d \in \mathbb{N}$, f odd and $f \leq d$, $\chi_k(H(n, 2, \{f\})) = 2k$.*

Proof. Let $d, f \in \mathbb{N}$, f odd and $f \leq d$. Consider the zero vector in $H(n, 2, \{f\})$. Note that every vector adjacent (orthogonal) to the zero vector has an odd Hamming weight. By vertex transitivity, all vectors of even Hamming weight only have vectors of odd Hamming weight as neighbors. Similarly, vectors of odd Hamming weight only have vectors of even Hamming weight as neighbors. Graph $H(n, 2, \{f\})$ is thus bipartite, which, combined with (5), proves the theorem. \square

It readily follows the following results for orthogonality graphs.

Corollary 26.1. *Let Ω_{4n+2} ($n \in \mathbb{N}$) be the orthogonality graph. Then, $\chi_k(\Omega_{4n+2}) = 2k$.*

Proof. Graph Ω_{4n+2} is isomorphic to $H(4n+2, 2, \{2n+1\})$. This corollary is thus a special case of theorem 26. \square

9 Conclusion

In this paper we study the generalized ϑ -number for highly symmetric graphs and beyond. The parameter $\vartheta_k(G)$ generalizes the concept of the famous ϑ -number that was introduced by Lovász [54]. Since $\vartheta_k(G)$ is sandwiched between the $\alpha_k(G)$ and $\chi_k(\overline{G})$ it serves as a bound for both graph parameters.

Several results in this paper are not restricted to highly symmetric graphs. In particular, the results in section 2, section 3 and section 4. In section 2 we present in an elegant way a known result that $\vartheta_k(G)$ is a lower bound for $\chi_k(\overline{G})$. Another lower bound for $\chi_k(\overline{G})$ is $k\vartheta(G)$, see (9). The inequality (9) is rather counter-intuitive since it is more difficult to compute $\vartheta_k(G)$ than $\vartheta(G)$, while $k\vartheta(G)$ provides a better bound for the k -th chromatic number. However, the generalized ϑ -number can also be used to compute lower bounds for the (classical) chromatic number of a graph, see section 5.2.

In section 3 we show that the series $(\vartheta_k(G))_k$ is increasing and bounded above by the order of G (proposition 2 and theorem 7), and that the increments of the series can be arbitrarily small (theorem 8). Section 4 provides bounds for $\vartheta_k(G)$ where G is the graph strong graph product of two graphs (theorem 9) and the graph disjunction product of two graphs (theorem 10).

Sections 5, 6 and 7 consider highly symmetric graphs. We derive closed from expressions for the generalized ϑ -number on cycles (theorem 12), Kneser graphs (theorem 16), Johnson graphs (theorem 17), strongly regular graphs (theorem 19), among other results. It is known that $\vartheta(K_k \square G)$ and $\vartheta_k(G)$ provide upper bounds on $\alpha_k(G)$. However, it is more computationally demanding to compute $\vartheta(K_k \square G)$ than $\vartheta_k(G)$. We show that for graphs that are both edge-transitive and vertex-transitive as well as for strongly regular graphs it suffices to solve $\vartheta_k(G)$, see theorem 18 and theorem 20. However, the gap between $\vartheta_k(G)$ and $\vartheta(K_k \square G)$ can be arbitrarily large (proposition 4). Section 7 presents results for $\vartheta_k(G)$ and $\chi_k(G)$ on orthogonality graphs.

Bounds on the k -th chromatic number of various graphs are given in section 8. In particular, bounds on the product and sum of $\chi_k(G)$ and $\chi_k(\overline{G})$ are presented in theorem 21. Lemma 7, proposition 8 and theorem 26 provide the multichromatic number for several Hamming graphs, while proposition 6 provides bounds for the multichromatic number on triangular graphs.

References

- [1] H Abdo and D Dimitrov. The total irregularity of graphs under graph operations. *Miskolc Mathematical Notes*, 15(1):3–17, 2014.
- [2] L. Addario-Berry, W.S. Kennedy, A. D. King, Z. Li, and B. Reed. Finding a maximum-weight induced k -partite subgraph of an i -triangulated graph. *Discrete Applied Mathematics*, 158(7):765–770, 2010.

- [3] F. Alizadeh. Interior point methods in semidefinite programming with applications to combinatorial optimization. *SIAM Journal on Optimization*, 5(1):13–51, 1995.
- [4] K Appel and W Haken. Every planar map is four colorable. Part I: Discharging. *Illinois Journal of Mathematics*, 21(3):429–490, 1977.
- [5] Christine Bachoc, Arnaud Pêcher, and Alain Thiéry. On the theta number of powers of cycle graphs. *Combinatorica*, 33(3):297–317, 2013.
- [6] Zsolt Baranyai. On the factorization of the complete uniform hypergraphs. In *Infinite and finite sets (Proc. Intern. Coll. Keszthely, A. Hajnal, R. Rado, V.T. Sós, eds.)*, pages 91–108. Bolyai J. Mat. Társulat, Budapest North-Holland, Amsterdam, 1975.
- [7] Frank Boesch and Ralph Tindell. Circulants and their connectivities. *Journal of Graph Theory*, 8(4):487–499, 1984.
- [8] Robert C Brigham and Ronald D Dutton. Generalized k -tuple colorings of cycles and other graphs. *Journal of Combinatorial Theory, Series B*, 32(1):90–94, 1982.
- [9] Valentin Brimkov. Algorithmic and explicit determination of the Lovász number for certain circulant graphs. *Discrete Applied Mathematics*, 155(14):1812–1825, 2007.
- [10] Valentin E Brimkov, Bruno Codenotti, Valentino Crespi, and Mauro Leoncini. On the Lovász number of certain circulant graphs. In Gian Carlo Bongiovanni, Giorgio Gambosi, and Rossella Petreschi, editors, *Algorithms and Complexity, 4th Italian Conference, CIAC 2000, Rome, Italy, March 2000, Proceedings*, volume 1767 of *Lecture Notes in Computer Science*, pages 291–305. Springer, 2000.
- [11] Andries E Brouwer, Arjeh M Cohen, and Arnold Neumaier. *Distance-regular graphs*. Berlin; New York: Springer-Verlag, 1989.
- [12] Manoel Campêlo, Ricardo C Corrêa, Phablo FS Moura, and Marcio C Santos. On optimal k -fold colorings of webs and antiwebs. *Discrete Applied Mathematics*, 161(1-2):60–70, 2013.
- [13] Gregory J Chaitin. Register allocation & spilling via graph coloring. *ACM Sigplan Notices*, 17(6):98–101, 1982.
- [14] TH Hubert Chan, Kevin L Chang, and Rajiv Raman. An SDP primal-dual algorithm for approximating the Lovász-theta function. In *2009 IEEE International Symposium on Information Theory, Korea (South)*, pages 2808–2812. IEEE, 2009.
- [15] Li-Chien Chang. The uniqueness and nonuniqueness of the triangular association scheme. *Science Record*, 3:604–613, 1959.
- [16] Bor-Liang Chen and Ko-Wei Lih. Hamiltonian uniform subset graphs. *Journal of Combinatorial Theory, Series B*, 42(3):257–263, 1987.
- [17] Vásek Chvátal. Edmonds polytopes and a hierarchy of combinatorial problems. *Discrete Mathematics*, 4(4):305–337, 1973.
- [18] Daniel W Cranston and Landon Rabern. Planar graphs are $9/2$ -colorable. *Journal of Combinatorial Theory, Series B*, 133:32–45, 2018.
- [19] Valentino Crespi. Exact formulae for the Lovász theta function of sparse circulant graphs. *SIAM Journal on Discrete Mathematics*, 17(4):670–674, 2004.
- [20] Etienne de Klerk and Dmitrii V Pasechnik. A note on the stability number of an orthogonality graph. *European Journal of Combinatorics*, 28(7):1971–1979, 2007.
- [21] Salim Y El Rouayheb, Costas N Georgiades, Emina Soljanin, and Alex Sprintson. Bounds on codes based on graph theory. In *2007 IEEE International Symposium on Information Theory, Nice, France*, pages 1876–1879. IEEE, June, 2007.
- [22] P. Fouilhoux and A. R. Mahjoub. Solving VLSI design and DNA sequencing problems using bipartization of graphs. *Computational Optimization and Applications*, 51(2):749–781, 2012.

- [23] Peter Frankl and Norihide Tokushige. The Erdős–Ko–Rado theorem for integer sequences. *Combinatorica*, 19(1):55–63, 1999.
- [24] Z. Füredi and P. Frankl. Extremal problems concerning Kneser graphs. *Journal of Combinatorial Theory, Series B*, 40(3):270–284, 1986.
- [25] Viktor Galliard. *Classical pseudo telepathy and coloring graphs*. Diploma Thesis, ETH Zurich, 2001. Available at <http://math.galliard.ch/Cryptography/Papers/PseudoTelepathy/SimulationOfEntanglement.pdf>.
- [26] Viktor Galliard, Alain Tapp, and Stefan Wolf. The impossibility of pseudotelepathy without quantum entanglement. In *IEEE International Symposium on Information Theory, 2003. Proceedings, Yokohama, Japan*, page 457. IEEE, 29 June - 4 July, 2003.
- [27] Rajiv Gandhi, Magnús M Halldórsson, Guy Kortsarz, and Hadas Shachnai. Improved bounds for sum multicoloring and scheduling dependent jobs with minsum criteria. In *International Workshop on Approximation and Online Algorithms*, pages 68–82. Springer, 2004.
- [28] K. Gatermann and P.A. Parrilo. Symmetry groups, semidefinite programs, and sums of squares. *Journal of Pure and Applied Algebra*, 192:95–128, 2004.
- [29] Dennis Geller and Saul Stahl. The chromatic number and other functions of the lexicographic product. *Journal of Combinatorial Theory, Series B*, 19(1):87–95, 1975.
- [30] Chris D Godsil and Michael W Newman. Coloring an orthogonality graph. *SIAM Journal on Discrete Mathematics*, 22(2):683–692, 2008.
- [31] Chris D Godsil and Mike W Newman. Eigenvalue bounds for independent sets. *Journal of Combinatorial Theory, Series B*, 98(4):721–734, 2008.
- [32] Don Greenwell and László Lovász. Applications of product colouring. *Acta Mathematica Academiae Scientiarum Hungarica*, 25(3-4):335–340, 1974.
- [33] Martin Grötschel, László Lovász, and Alexander Schrijver. The ellipsoid method and its consequences in combinatorial optimization. *Combinatorica*, 1(2):169–197, 1981.
- [34] Nebojša Gvozdenović. *Approximating the stability number and the chromatic number of a graph via semidefinite programming*. PhD thesis, Universiteit van Amsterdam, 2008.
- [35] Nebojša Gvozdenović and Monique Laurent. The operator Ψ for the chromatic number of a graph. *SIAM Journal on Optimization*, 19(2):572–591, 2008.
- [36] W. H. Haemers. *Eigenvalue techniques in design and graph theory*. PhD thesis, (T.H. Eindhoven, 1979) = Math. Centr. Tract 121 (Amsterdam, 1980), Reidel, Dordrecht, 1980.
- [37] Willem Haemers. An upper bound for the Shannon capacity of a graph. In *Colloq. Math. Soc. János Bolyai*, volume 25, pages 267–272, 1978.
- [38] Magnús M Halldórsson and Guy Kortsarz. Multicoloring: Problems and techniques. In Jiří Fiala, Václav Koubek, and Jan Kratochvíl, editors, *Mathematical Foundations of Computer Science 2004*, pages 25–41, Berlin, Heidelberg, 2004. Springer Berlin Heidelberg.
- [39] Stephen T Hedetniemi. Homomorphisms of graphs and automata. Technical report, Michigan univ Ann Arbor Communication Sciences Program, 1966.
- [40] AJW Hilton, R Rado, and SH Scott. A (< 5)-colour theorem for planar graphs. *Bulletin of the London Mathematical Society*, 5(3):302–306, 1973.
- [41] A.J. Hoffman. On eigenvalues and colorings of graphs. In B. Harris, editor, *Graph Theory and its Applications*, pages 79–91. Academic Press, New York, 1970.
- [42] Roger A Horn and Charles R Johnson. *Topics in Matrix Analysis*. Cambridge University Press, 1994.
- [43] Ferdinand Ihringer and Hajime Tanaka. The independence number of the orthogonality graph in dimension 2^k . *Combinatorica*, 39(6):1425–1428, 2019.

- [44] Sandi Klavžar. Coloring graph products—a survey. *Discrete Mathematics*, 155(1-3):135–145, 1996.
- [45] M Kneser. Aufgabe 300, jahresbericht. *Deutschen Math. Ver.*, 58(2), 1955.
- [46] Donald E Knuth. The sandwich theorem. *The Electronic Journal of Combinatorics*, 1(1):A1, 1994.
- [47] A.M.C.A. Koster and M. Scheffel. A routing and network dimensioning strategy to reduce wave-length continuity conflicts in all-optical networks. In *Proceedings of INOC 2007, International Network Optimization Conference, Spa, Belgium, April 22-25, 2007*.
- [48] Olga Kuryatnikova, Renata Sotirov, and Juan Vera. The maximum k -colorable sub-graph problem and related problems. *INFORMS Journal on Computing (accepted)*, 2021. <https://arxiv.org/abs/2001.09644v3>.
- [49] J.M. Lewis and M. Yannakakis. The node-deletion problem for hereditary properties is NP-complete. *Journal of Computer and System Sciences*, 20:219–230, 1980.
- [50] Wensong Lin. Multicoloring and Mycielski construction. *Discrete Mathematics*, 308(16):3565–3573, 2008.
- [51] Wensong Lin, Daphne Der-Fen Liu, and Xuding Zhu. Multi-coloring the Mycielskian of graphs. *Journal of Graph Theory*, 63(4):311–323, 2010.
- [52] R. Lippert, R. Schwartz, G. Lancia, and S. Istrail. Algorithmic strategies for the single nucleotide polymorphism haplotype assembly problem. *Briefings in Bioinformatics*, 3(1):23–31, 2002.
- [53] László Lovász. Kneser’s conjecture, chromatic number, and homotopy. *Journal of Combinatorial Theory, Series A*, 25(3):319–324, 1978.
- [54] László Lovász. On the Shannon capacity of a graph. *IEEE Transactions on Information theory*, 25(1):1–7, 1979.
- [55] László Lovász. *An algorithmic theory of numbers, graphs, and convexity*. SIAM, Philadelphia, PA, 1986.
- [56] Enrico Malaguti and Paolo Toth. An evolutionary approach for bandwidth multicoloring problems. *European Journal of Operational Research*, 189(3):638–651, 2008.
- [57] M. Marek-Sadowska. An unconstrained topological via minimization problem for two-layer routing. *IEEE Transactions on Computer-Aided Design of Integrated Circuits and Systems*, 3(3):184–190, 1984.
- [58] Dániel Marx. The complexity of tree multicolorings. In Rytter W. Diks K., editor, *Mathematical Foundations of Computer Science, MFCS*, volume 2420, pages 532–542. Lecture Notes in Computer Science, Springer, Berlin, Heidelberg, 2002.
- [59] Robert J McEliece, Eugene R Rodemich, and Howard C Rumsey Jr. The Lovász bound and some generalizations. *J. Combin. Inform. System Sci*, 3(3):134–152, 1978.
- [60] Anuj Mehrotra and Michael A Trick. A branch-and-price approach for graph multi-coloring. In Edward K. Baker, Anito Joseph, Anuj Mehrotra, and Michael A. Trick, editors, *Extending the Horizons: Advances in Computing, Optimization, and Decision Technologies*, pages 15–29. Springer, Boston, MA, 2007.
- [61] G. Narasimhan. *The maximum k -colorable subgraph problem*. PhD thesis, University of Wisconsin-Madison, 1989.
- [62] Giri Narasimhan and Rachel Manber. A generalization of Lovász sandwich theorem. Technical report, University of Wisconsin-Madison Department of Computer Sciences, 1988.
- [63] Lata Narayanan. Channel assignment and graph multicoloring. In Ivan Stojmenović, editor, *Handbook of Wireless Networks and Mobile Computing*, chapter 4, pages 71–94. Wiley Online Library, 2002.
- [64] Michael William Newman. *Independent sets and eigenspaces*. PhD thesis, University of Waterloo, 2004.

- [65] E A Nordhaus and Jerry W Gaddum. On complementary graphs. *The American Mathematical Monthly*, 63(3):175–177, 1956.
- [66] Włodzimierz Ogryczak and Arie Tamir. Minimizing the sum of the k largest functions in linear time. *Information Processing Letters*, 85(3):117–122, 2003.
- [67] Michael L Overton and Robert S Womersley. Optimality conditions and duality theory for minimizing sums of the largest eigenvalues of symmetric matrices. *Mathematical Programming*, 62(1-3):321–357, 1993.
- [68] GuanFeng Ren and YueHua Bu. k -fold coloring of planar graphs. *Science China Mathematics*, 53(10):2791–2800, 2010.
- [69] Gert Sabidussi. Graphs with given group and given graph-theoretical properties. *Canadian Journal of Mathematics*, 9:515–525, 1957.
- [70] Hiroki Sayama. Estimation of Laplacian spectra of direct and strong product graphs. *Discrete Applied Mathematics*, 205:160–170, 2016.
- [71] C. Shannon. The zero error capacity of a noisy channel. *IRE Trans. Inf. Theory*, 2:8–19, 1956.
- [72] Yaroslav Shitov. Counterexamples to Hedetniemi’s conjecture. *Annals of Mathematics*, 190(2):663–667, 2019.
- [73] Richard Singleton. Maximum distance q -nary codes. *IEEE Transactions on Information Theory*, 10(2):116–118, 1964.
- [74] Lennart Sinjorgo. *Multicoloring of highly symmetric graphs*. Master’s thesis, Tilburg University, The Netherlands, 2021.
- [75] Saul Stahl. n -Tuple colorings and associated graphs. *Journal of Combinatorial Theory, Series B*, 20(2):185–203, 1976.
- [76] A. P. Subramanian, H. Gupta, S. R. Das, and M. M. Buddhikot. Fast spectrum allocation in coordinated dynamic spectrum access based cellular networks. In *2007 2nd IEEE International Symposium on New Frontiers in Dynamic Spectrum Access Networks*, pages 320–330, 2007.
- [77] Garry J Tee. Eigenvectors of block circulant and alternating circulant matrices. *New Zealand Journal of Mathematics*, 36(8):195–211, 2007.
- [78] Vadim G Vizing. On an estimate of the chromatic class of a p -graph. *Discret. Analiz.*, 3:25–30, 1964.
- [79] Mihalis Yannakakis and Fanica Gavril. The maximum k -colorable subgraph problem for chordal graphs. *Information Processing Letters*, 24(2):133 – 137, 1987.