

Efficient Use of Quantum Linear System Algorithms in Interior Point Methods for Linear Optimization

Mohammadhossein Mohammadisiahroudi, Ramin Fakhimi,
and Tamás Terlaky

Received: date / Accepted: date

Abstract Quantum computing has attracted significant interest in the optimization community because it potentially can solve classes of optimization problems faster than conventional supercomputers. Several researchers proposed quantum computing methods, especially Quantum Interior Point Methods (QIPMs), to solve convex optimization problems, such as Linear Optimization, Semidefinite Optimization, and Second-order Cone Optimization problems. Most of them have applied a Quantum Linear System Algorithm at each iteration to compute a Newton step. However, using quantum linear solvers in QIPMs comes with many challenges, such as having ill-conditioned systems and the considerable error of quantum solvers. This paper investigates how one can efficiently use quantum linear solvers in QIPMs. Accordingly, an Inexact Infeasible Quantum Interior Point Method is developed to solve linear optimization problems. We also discuss how we can get an exact solution by Iterative Refinement without excessive time of quantum solvers. Finally, computational results with a QISKIT implementation of our QIPM using quantum simulators are analyzed.

Keywords Quantum Interior Point Method · Linear Optimization · Quantum Linear System Algorithm · Iterative Refinement

Mathematics Subject Classification (2000) 90C51 · 90C05 · 81P68

1 Introduction

Linear Optimization (LO) is defined as optimizing a linear function over a set of linear equality and inequality constraints. Several algorithms were developed to solve LO problems [16, 29].

Mohammadhossein Mohammadisiahroudi, Corresponding author
Lehigh University
Bethlehem, PA 18015, USA
mom219@Lehigh.edu

Ramin Fakhimi
Lehigh University
Bethlehem, PA 18015, USA
fakhimi@Lehigh.edu

Tamás Terlaky
Lehigh University
Bethlehem, PA 18015, USA
terlaky@Lehigh.edu

Karmarkar [25] developed the foundations of polynomial time and practically efficient Interior Point Methods (IPMs) for solving LO problems. Since Karmarkar’s publication, a large class of theoretically and practically efficient IPMs were developed, see, e.g., [40, 36, 43]. Contrary to the Simplex method, a feasible IPM reaches an optimal solution by traversing through the interior of the feasible region [36].

Contemporary IPMs reach an optimal solution by starting from an interior point and following the central path [36]. The most efficient IPMs are primal-dual methods, meaning they strive to satisfy the optimality conditions while maintaining strict primal and dual feasibility. It should be noted that basic IPMs need an initial feasible interior point. Some current commercial solvers apply Feasible IPMs (F-IPMs) based on the self-dual embedding formulation of LO problems, e.g., MOSEK, while Infeasible Interior Point Methods (I-IPMs) can start with an infeasible but positive solution. Theoretical analysis shows the best iteration complexity of F-IPMs is $\mathcal{O}(\sqrt{n}L)$, where n is the number of variables, and L is the binary length of the input data. On the other hand, the best iteration complexity of I-IPMs is $\mathcal{O}(nL)$. In practice, the performance of both feasible and infeasible IPM are similar [43].

A linear equation system is solved at each iteration of IPMs to calculate a Newton direction. There are three choices for the linear equation system: (i) Full Newton System, (ii) Augmented System, and (iii) Normal Equation System (NES). In classical computers, a general approach is applying Cholesky Factorization to solve the NES, because it has a symmetric positive definite coefficient matrix. A partial update technique improved the complexity of solving the system in each iteration. This approach leads to the best total complexity of $\mathcal{O}(n^3L)$ arithmetic operations for solving LO problems. However, this is not always efficient in practice [36]. Since Cholesky Factorization requires $\mathcal{O}(n^3)$ arithmetic operations for large dense matrices, several researchers studied inexact solution methodologies for solving Newton systems. Bellavia [9] proved the convergence of an Inexact I-IPM (II-IPM) for general convex optimization problems. Mizuno and his colleagues studied the convergence of II-IPMs [33, 20]. Korzak [31] and Baryamureeba and Steihaug [8] proved the convergence of II-IPM proposed by Kojima et al. [30]. Korzak [31] also showed that the total time complexity of his algorithm is polynomial. Al-Jeiroudi and Gondzio [24], and Monteiro and O’Neal [35] investigated the use of Preconditioned Conjugate Gradient (PCG) methods in II-IPMs. Zhou and Toh [46] also proved the convergence of an II-IPM for SDO problems. The best iteration complexity of II-IPMs is $\mathcal{O}(n^2L)$ which is $\Theta(n^{1.5})$ weaker than the best iteration complexity of exact F-IPMs.

Quantum computers have recently emerged as a powerful alternative to classic computers [15]. Starting from Deutsch’s Problem, a series of problems and algorithms have demonstrated theoretically exponential speedup compared to their classical counterparts [18, 19, 39]. One of the promising quantum algorithms is the HHL method [23] to solve linear system problems. The HHL method showed quantum advantage with respect to the dimension compared to classical linear systems solutions algorithm. However, this method has unfavorable dependence on the condition number, the sparsity of the matrix, and the inverse precision of the solution. Several researchers attempted to improve the performance of Quantum Linear System Algorithms (QLSAs), [42, 15]. This paper explores an efficient use of the QLSAs for solving the Newton system at each iteration of IPMs.

To investigate quantum speedup for continuous optimization, Brandão and Svore [11] proposed a non-IPM quantum algorithm based on the Multiplicative Weight Update Method (MWUM) of [6] to solve Semidefinite Optimization (SDO) problems. After this papers, many improved versions of Quantum MWUMs (QMWUMs) are proposed for SDO [12, 4], and LO [5]. The advantage of QMWUMs is that their complexity has linear dependence on the dimension for SDO and sublinear dependence for LO. The major issue with QMWUMs is that they are highly de-

pendent on inverse precision and an upper bound for the norm of the optimal solution, which are exponentially large for LO.

In another direction, a few studies were proposing Quantum IPMs (QIPMs). Kerenidis and his colleagues presented a series of papers on QIPMs for solving LO, SDO [26], and Second-order Cone Optimization (SOCO) problems [28]. Note that for both SDO and SOCO, they missed some important parts, such as the need for symmetrization/scaling. So, their QIPMs are invalid for their conic optimization problems. Kerenidis and Prakash [26] claimed that their algorithm has $\mathcal{O}(\frac{n^2}{\epsilon^2} \kappa^3 \log(\frac{1}{\zeta}))$ complexity for LO problems where ζ is the final optimality gap, ϵ is the precision of the QLSA. Observe that, to reach final precision ζ , the Newton system needs to be solved with precision $\epsilon < \zeta$. Further, κ is an upper bound for the condition number of the Newton systems at each iteration. They used block encoding to construct the Newton system and Quantum Tomography Algorithm (QTA) to extract the classical solution. Casares and Martin-Delgado [13] also provided a hybrid predictor-corrector QIPM scheme to solve LO problems using the well-known predictor-corrector method proposed by Ye et al. [45]. In each step, it uses the QLSA proposed by Chakraborty et al. [14] to solve the Newton systems. They claimed that the complexity of their method is $\mathcal{O}(L\sqrt{n}(n+m)\|M\|_F \kappa^2 \epsilon^{-2})$, where m is the number of constraints, and $\|M\|_F$ is an upper bound to the Frobenius norm of the coefficient matrix in the Newton system at each iteration. The authors claimed a quantum speedup with respect to the dimension n compared to classical algorithms. However, they overlooked some elements in their time complexity. Thus, the actual complexity has higher dependence on n and the required precision.

In both mentioned QIPMs, an exact F-IPM framework was used, regardless of the inherent inexactness of QLSAs. So, the proposed time complexities are not attainable, since Inexact IPMs have higher iteration complexity. Augustino et al. [7] addressed this issue and proposed two convergent QIPMs for SDO. Their complexity shows polynomial speed-up with respect to dimension, but similar to former QIPMs, their complexity suffers from linear dependence on condition number of the Newton system and precision. A major reason is that QLSAs' time complexities depend on the condition number of the linear system, and it is linear in inverse precision and dimension to extract an accurate classical solution. In IPMs, the condition number of the Newton system typically goes to infinity as the algorithm approaches an optimal solution [36]. It is also worth mentioning that ϵ , the required precision for solution of the Newton system, needs to be significantly smaller than the final precision ζ . Consequently, these complexity bounds are not polynomial in the classical sense. This paper explores an efficient use of the QLSAs for solving the Newton system at each iteration of QIPMs. We propose an II-QIPM to find an exact solution. We also employ an iterative refinement scheme to avoid exponential complexity in finding an exact optimal solution.

This paper is structured as follows. Section 2 discusses the performance of existing QLSAs. Section 3 introduces the LO problem and its characteristics. In Section 4, we present an II-QIPM to solve LO problems, in which a QLSA is used for solving the NES. We also analyze the complexity of the proposed II-QIPM. In Section 5, we employ an Iterative Refinement scheme to find an exact optimal solution of a LO problem without excessive time of QLSAs. Section 6 presents the first implementation of the proposed II-QIPM with Iterative Refinement (IR-II-QIPM) and evaluates the algorithms through computational experiments. Finally, conclusions are presented in Section 7.

2 Quantum Linear Algebra

This section reviews the use of Quantum algorithms to solve Linear System Problems (LSP).

Definition 2.1 (LSP) The Linear System Problem: Find a vector $z \in \mathbb{R}^p$ such that it satisfies equation $Mz = \sigma$ with coefficient matrix $M \in \mathbb{R}^{p \times p}$ and right-hand side (RHS) vector $\sigma \in \mathbb{R}^p$.

In the rest of the paper, $\|M\| = \|M\|_2$ is the 2-norm of matrix M , and $\|M\|_F$ is the Frobenius norm of M . We sometimes use $\tilde{\mathcal{O}}$ which suppresses the polylogarithmic factors in the "Big-O" notation. The quantities of the polylogarithmic factors are indicated as subscripts of $\tilde{\mathcal{O}}$. We use \mathbb{R}^n for the set of n -dimensional vectors of real numbers and \mathbb{C}^n for the set of n -dimensional vectors of complex numbers. It should be mentioned that the complexity of a classical algorithm when solving LSP means the number of arithmetic operations and the complexity of a quantum algorithm is the number of quantum gates.

The LSP can have either one, many, or no solutions. A basic approach for solving an LSP is Gaussian elimination, or LU factorization, with $\mathcal{O}(p^3)$ arithmetic operations. If M is a square symmetric positive semidefinite (PSD) matrix, we can also apply Cholesky factorization with $\mathcal{O}(p^3)$ arithmetic operations. The best complexity for an iterative algorithm with respect to p is $\mathcal{O}(pd\sqrt{\kappa}\log(1/\epsilon))$ arithmetic operations for the Conjugate Gradient method solving systems with symmetric PSD matrices, where d is the maximum number of non-zero elements in any row or column of M , κ is the condition number of M , and ϵ is the error allowed. If matrix M is just symmetric, one can use Lanczos algorithm with higher complexity. For an LSP with general square matrix M , the best iterative method is the GMRES algorithm, which has $\mathcal{O}(n^3)$ worst-case complexity [37]. To sum up, LSPs in general form are solvable in polynomial time in classical computing setting, but the worst-case complexity is $\mathcal{O}(n^3)$ for either matrix decomposition methods or iterative methods.

Before discussing QLSAs, we should mention that the $|z\rangle$ notation represent the quantum state corresponding to the unit classical vector z . We denote the basis state $|i\rangle$, which is a column vector with dimension p , one in coordinate i and zero in other coordinates [15]. QLSAs have different approaches, while all of them are solving Quantum Linear System Problems (QLSPs) defined as follows.

Definition 2.2 (QLSP) Let $M \in \mathbb{C}^{p \times p}$ be a Hermitian matrix with $\|M\| = 1$, $\sigma \in \mathbb{C}^p$, and $z := M^{-1}\sigma$. We define quantum states

$$|\sigma\rangle = \frac{\sum_{i=1}^p \sigma_i |i\rangle}{\|\sum_{i=1}^p \sigma_i |i\rangle\|} \quad \text{and} \quad |z\rangle = \frac{\sum_{i=1}^p z_i |i\rangle}{\|\sum_{i=1}^p z_i |i\rangle\|}.$$

For target precision $0 < \epsilon_{QLSP}$, the goal is to find $|\tilde{z}\rangle$ such that $\| |\tilde{z}\rangle - |z\rangle \| \leq \epsilon_{QLSP}$, succeeding with probability $\Omega(1)$.

Based on Definition 2.1 and Definition 2.2, QLSP is a different form of the LSP. It should be noted that at each iteration of QIPMs, instead of an LSP, we need to use a QLSA to solve a QLSP. Thus, we need to translate LSP to QLSP, solve the QLSP by QLSA and extract the solution by a QTA. Here, we analyze the details and costs of the process of translating LSPs to QLSPs, encoding in the quantum setting, solving them with a QLSA, and extracting classical solutions with a QTA as follows.

- (i) Model of Computation: the first important step is determining how to encode the input data in the quantum setting. There are two major input models. One is the sparse-access model which is used in the HHL algorithm [23] and then in other QLSAs [15, 41]. This is a quantum version of classical sparse matrix computation, and we assume access to unitaries that calculate the index of the l th non-zero element of the k th row of a matrix M when given (k, l) as input. A different input model, now known as the quantum operator input model, is

proposed in Low and Chuang [32], which is based on the idea of block-encoded matrices. In this input model, one has access to unitaries that store the coefficient matrix:

$$U = \begin{pmatrix} M/\alpha & \cdot \\ \cdot & \cdot \end{pmatrix},$$

where $\alpha \geq \|M\|$ is a normalization factor chosen to ensure that U has norm at most 1. Chakraborty et al. [14] showed that this quantum operator input model is more efficient than the sparse-access model and oracles to encode input data using block-encoding has favorable complexity compared to the sparse-access model. On the other hand, most of the block-encoding approaches use Quantum Random Access Memory (QRAM). However, the sparse-access model can be implemented in the standard gate-based quantum circuit model. Despite efficient encoding procedures, the quantum operator input model can not be implemented with current quantum computers since there is no physical implementation of QRAM. In our analysis, we assume that the data is stored in QRAM, and we use the quantum operator model by Chakraborty et al. [14] for the QLSA, which enjoys the best complexity to date. Using the QRAM structure, Kerenidis and Prakash [27] showed that one can implement ϵ -approximate block-encoding of M with $\mathcal{O}(\text{polylog}(\frac{p}{\epsilon}))$ complexity. Further, given M in the sparse-access input model, there is an ϵ -approximate block-encoding of M that can be implemented in complexity $\mathcal{O}(\text{polylog}(\frac{p\alpha}{\epsilon}))$ [14]. Thus, our results also apply to the sparse-access input model if we have the data in that form. Using these results and assuming access to QRAM, Chakraborty et al. [14] proposed a QLSA, in which they construct state $|\sigma\rangle$, build and implement a block-encoding of matrix M with $\mathcal{O}(\text{polylog}(\frac{p}{\epsilon}))$ complexity. Since we are interested in using QLSAs in a hybrid approach, we need to consider the cost of storing data in a classical form to QRAM which is $\mathcal{O}(p^2)$ for a fully dense matrix [14]. This cost should be paid once, and in the next section, it is shown that this cost will be dominated by the classical operations in each iteration of QIPMs.

- (ii) Translating LSP to QLSP: Based on the definition of QLSP, the coefficient matrix of the system must be Hermitian. If M is not Hermitian, one can construct $\bar{M}\bar{z} = \bar{\sigma}$, where

$$\bar{M} = \begin{bmatrix} 0 & M \\ M^\dagger & 0 \end{bmatrix}, \quad \bar{\sigma} = \begin{pmatrix} \sigma \\ 0 \end{pmatrix},$$

and find the vector $\bar{z} = \begin{pmatrix} 0 \\ z \end{pmatrix}$, where M^\dagger denotes the conjugate transpose of M . The size of the problem increases from p to $2p$. QLSP assumes $\|M\| = 1$. In the structure of block encoding, we address this normalization, but for sparse encoding, we need to normalize the system $\bar{M}\bar{z} = \bar{\sigma}$ where

$$\bar{M} = \frac{M}{\|M\|}, \quad \bar{\sigma} = \frac{\sigma}{\|M\|}, \quad \text{and} \quad \bar{z} = z.$$

Although we are using block-encoding, we scale the matrix in advance and let $\alpha = 1$ in the block encoding. This scaling will affect the precision and complexity, but in this way, we can have better complexity analysis since, in QIPMs, we need to bound the residual. We also have a similar scaling in the definitions of states $|z\rangle$ and $|\sigma\rangle$. The corresponding QLSP problem is $\bar{M}|z\rangle = |\bar{\sigma}\rangle$, and we can find an inexact solution $|\tilde{z}\rangle$ with $\| |\tilde{z}\rangle - |z\rangle \| \leq \epsilon_{QLSP}$. Since $\| |\tilde{z}\rangle \| = 1$, we need to scale back the solution with $\tilde{z} = \|\bar{\sigma}\| |\tilde{z}\rangle$. This scaling affects the target precision. In IPMs, we are interested in finding a solution where the residual is bounded as

$\|\sigma - M\tilde{z}\| \leq \epsilon_{LSP}$. Thus, we have

$$\begin{aligned} \|\sigma - M\tilde{z}\| &\leq \|M\| \|\bar{\sigma} - \overline{M}\tilde{z}\| \\ &\leq \|M\| \|\bar{\sigma}\| \|\bar{\sigma} - \overline{M}\tilde{z}\| \\ &\leq \|M\| \frac{\|\sigma\|}{\|M\|} \|\overline{M}|z\rangle - \overline{M}|\tilde{z}\rangle\| \\ &\leq \|\sigma\| \|\ |z\rangle - |\tilde{z}\rangle \| \\ &\leq \|\sigma\| \epsilon_{QLSP}. \end{aligned}$$

Thus, we must set the target error of the QLSP as $\epsilon_{QLSP} = \epsilon_{LSP}/\|\sigma\|$.

- (iii) **QLSA**: After preprocessing and encoding, we can apply the QLSA to solve the QLSP. The HHL algorithm [23] solves QLSPs with $\tilde{\mathcal{O}}_p(\frac{d^2\kappa^2}{\epsilon_{QLSP}})$ complexity. Several researchers attempted to improve the performance of the HHL algorithm. As the first attempt, Amplitude Amplification decreases the dependence on κ^2 to κ [1]. Wossnig et al. [42] proposed a QLSA with $\tilde{\mathcal{O}}_p(\|M\|_F \frac{\kappa}{\epsilon_{QLSP}})$ complexity by using the Quantum Singular Value Estimation. In another direction, Childs et al. [15] developed two QLSAs with exponentially better dependence on error with $\tilde{\mathcal{O}}_{p,\kappa,\frac{1}{\epsilon_{QLSP}}}(d\kappa)$ complexity. They proposed two approaches using Fourier and Chebyshev series representations. The best QLSA with respect to complexity uses block encoding and QRAM with $\tilde{\mathcal{O}}_{p,\frac{1}{\epsilon_{QLSP}}}(\alpha\kappa)$ complexity [14]. The normalization factor α is equal to 1 for our approach since we apply the QLSA to QLSP, which is normalized in advance. The details of these methods are out of the scope of this paper. For further details, see [17].
- (iv) **QTA**: QLSAs provide a quantum state proportional to the solution. We cannot extract the classical solution by a single measurement. We need Quantum Tomography Algorithms (QTAs) to extract the classical solution. There are several papers improving QTAs, see, e.g., [28]. We used the best QTA by [3], with complexity $\mathcal{O}(\frac{p\varrho}{\epsilon_{QTA}})$, where ϱ is bound for the norm of the solution. In our approach, ϱ is equal to 1 since we applied QTA to QLSP and scaled back the solution after QTA. Since the error is additive, we may choose

$$\epsilon_{QTA} = \epsilon_{QLSA} = \frac{\epsilon_{LSP}}{2\|\sigma\|}.$$

Table 1 presents the complexity of different classical and quantum algorithms for solving an LSP. Here, the complexity of a QTA is considered in the complexity of QLSAs. As shown in Table 1, although the complexity of solving LSP using QLSA+QTA will have a similar dependence on p compared to Conjugate Gradient (CG), the complexity of CG method is valid for solving LSPs with PSD matrices and the complexity of QLSA+QTA in Table 1 are valid for solving LSPs in a general form. QLSAs have better dependence on dimension p compared to factorization and elimination techniques. Generally, QLSA has worse dependence on κ , $\frac{1}{\epsilon_{LSP}}$, $\|M\|$, and $\|\sigma\|$. In our QIPM, we solve a modified NES which has symmetric PSD matrix and both CG and QLSA+QTA are applicable. Although CG has better complexity than QLSA+QTA based on Table 1, we use QLSA+QTA to solve the NES since it enables us to also build the Newton system in quantum computer which has polylogarithmic dependence on n . This approach enable us to avoid matrix-matrix product, which is inevitable in classical IPM using CG. Thus, the cost per iteration of IPM can leverage the quadratic dependence on dimension but the linear dependence on condition number and inverse precision can lead to exponential complexity. The following sections will discuss how we can deal with errors and condition numbers when we use QLSA in IPMs.

Table 1: Complexity of solving an LSP.

Algorithm	Complexity	Comment
Factorization methods (e.g. LU)	$\mathcal{O}(p^3)$	
Conjugate Gradient	$\mathcal{O}(pd\sqrt{\kappa}\log(\frac{1}{\epsilon}))$	For LSPs with PSD matrices.
HHL [23] + QTA [3]	$\mathcal{O}(pd + p\frac{d^2\kappa^2\ \sigma\ ^2}{\ M\ ^2\epsilon^2}\text{polylog}(\frac{p\kappa}{\epsilon}))$	Using sparse encoding.
QLSA [15] + QTA [3]	$\mathcal{O}(pd + p\frac{d\kappa\ \sigma\ }{\ M\ \epsilon}\text{polylog}(\frac{p\kappa}{\epsilon}))$	Using sparse encoding.
QLSA [14] + QTA [3]	$\mathcal{O}(p^2 + p\frac{\kappa\ \sigma\ }{\epsilon}\text{polylog}(\frac{p}{\epsilon}))$	Using block encoding.

3 The Linear Optimization Problem

Here we consider the standard form of Linear Optimization (LO) problem as follows:

$$\begin{aligned}
& \min c^T x & \max b^T y \\
\text{(P)} \quad & \text{s.t. } Ax = b, & \text{(D)} \quad \text{s.t. } A^T y + s = c, \\
& x \geq 0, & s \geq 0,
\end{aligned} \tag{1}$$

where $A : m \times n$ matrix with $\text{rank}(A) = m$, vectors $y, b \in \mathbb{R}^m$, and $x, s, c \in \mathbb{R}^n$. Problem (P) is called the primal problem and (D) is called the dual problem. Due to the Strong Duality Theorem [36], all optimal solutions, if exist, belong to the set \mathcal{PD}^* , which is defined as

$$\mathcal{PD}^* = \left\{ (x, y, s) \in \mathbb{R}^{n+m+n} : Ax = b, A^T y + s = c, x^T s = 0, (x, s) \geq 0 \right\}.$$

Now, we can define the optimal partition of the LO problem as

$$\begin{aligned}
B &= \{j \in \{1, \dots, n\} : x_j^* > 0 \text{ for some } (x^*, y^*, s^*) \in \mathcal{PD}^*\}, \\
N &= \{j \in \{1, \dots, n\} : s_j^* > 0 \text{ for some } (x^*, y^*, s^*) \in \mathcal{PD}^*\}.
\end{aligned}$$

It is shown that $B \cup N = \{1, \dots, n\}$, and $B \cap N = \emptyset$ [36].

Assumption 1. *From now on, without loss of generality [36], we assume that the Interior Point Condition (IPC) holds, i.e., there exists a solution (x, y, s) such that*

$$Ax = b, x > 0, A^T y + s = c, \text{ and } s > 0.$$

The IPC warranties [36] that the optimal set \mathcal{PD}^* is bounded, so there exists $\omega \geq 1$ such that

$$\omega \geq \max\{\|(x^*, s^*)\|_\infty : (x^*, s^*) \in \mathcal{PD}^*\}.$$

The central path is the curve defined by

$$\mathcal{CP} = \left\{ (x, y, s) \in \mathbb{R}^{n+m+n} : Ax = b, A^T y + s = c, x_i s_i = \mu \text{ for } i \in \{1, \dots, n\}, x, s, \mu > 0 \right\}.$$

By the IPC, the central path is well-defined, and an interior feasible solution (x, y, s) exists for all $\mu > 0$. Now, for any $0 < \gamma_1 < 1$ and $1 \leq \gamma_2$, we define an infeasible neighborhood of the central path for $\mu = \frac{x^T s}{n}$ as the following definition in [43].

$$\mathcal{N}(\gamma_1, \gamma_2) = \{(x, y, s) \in \mathbb{R}^{n+m+n} : (x, s) > 0, x_i s_i \geq \gamma_1 \mu \text{ for } i \in \{1, \dots, n\}, \|(R_P, R_D)\| \leq \gamma_2 \mu\},$$

where $R_P = b - Ax$, and $R_D = c - A^T y - s$. Assuming that the input data is integral, we denote the binary length of the input data by

$$L = mn + m + n + \sum_{i,j} \lceil \log(|a_{ij}| + 1) \rceil + \sum_i \lceil \log(|c_i| + 1) \rceil + \sum_j \lceil \log(|b_j| + 1) \rceil.$$

The following lemma is a classical result first proved by Khachiyan [29].

Lemma 3.1 *Let $(x^*, y^*, s^*) \in \mathcal{PD}^*$ be a basic solution. If $x_i^* > 0$, then we have $x_i^* \geq 2^{-L}$. If $s_i^* > 0$, then we have $s_i^* \geq 2^{-L}$.*

Theorem 3.1 (Chapter 3 in [43]) *An exact optimal solution can be obtained by a strongly polynomial rounding procedure when $(x, y, s) \in \mathcal{N}(\gamma_1, \gamma_2)$ and $\mu \leq 2^{-2L}$.*

If the IPC holds, then the optimal set \mathcal{PD}^* is bounded, and we can find the upper bound for all the coordinates of all optimal solutions as described in the following lemma.

Lemma 3.2 (Chapter 5 in [44]) *Assuming the IPC, then for any $(x^*, y^*, s^*) \in \mathcal{PD}^*$, $\max_i\{x_i^*\} \leq 2^L$, and $\max_i\{s_i^*\} \leq 2^L$.*

For the theoretical purpose, we can use $\omega = 2^L$ from Lemma 3.2, but in practice, for concrete LO problems, we may find a smaller bound. We define the set of ζ -optimal solutions as

$$\mathcal{PD}(\zeta) = \{(x, y, s) \in \mathbb{R}^{n+m+n} : (x, s) \geq 0, \frac{x^T s}{n} \leq \zeta, \|(R_P, R_D)\| \leq \zeta\}$$

4 An Inexact Infeasible Quantum IPM

To speed up IPMs, we use QLSAs to solve the Newton system at each iteration of IPMs. As discussed in Section 2, QLSAs inherently produce inexact solutions. Thus, one approach to use QLSA efficiently is to develop an Inexact Infeasible QIPM (II-QIPM). In this paper, we utilize the KMM method proposed by Kojima et al. [30] with the inexact Newton steps calculated by a QLSA. Given $(x^k, y^k, s^k) \in \mathcal{N}(\gamma_1, \gamma_2)$, let $\mu^k = \frac{(x^k)^T s^k}{n}$ and $0 < \beta_1 < 1$ be the centering parameter, then the Newton system is defined as

$$\begin{aligned} A\Delta x^k &= b - Ax^k, \\ A^T \Delta y^k + \Delta s^k &= c - A^T y^k - s^k, \\ X^k \Delta s^k + S^k \Delta x^k &= \beta_1 \mu^k e - X^k s^k, \end{aligned} \tag{2}$$

where e is all one vector with appropriate dimension, $X^k = \text{diag}(x^k)$, and $S^k = \text{diag}(s^k)$. Instead of solving the full Newton system, we may solve the Augmented system or the Normal Equation System (NES). From the Newton system (2), the NES is formulated as

$$M^k \Delta y^k = \sigma^k, \tag{NES}$$

where

$$\begin{aligned} D^k &= (X^k)^{1/2} (S^k)^{-1/2}, \\ M^k &= A(D^k)^2 A^T, \\ \sigma^k &= A(D^k)^2 c - A(D^k)^2 A^T y^k - \beta_1 \mu^k A(S^k)^{-1} e + b - Ax^k \\ &= b - \beta_1 \mu^k A(S^k)^{-1} e + A(D^k)^2 (c - A^T y^k - s^k). \end{aligned}$$

As we can see, the NES has a smaller size, m , than the full Newton system. Further, the coefficient matrix of the NES is symmetric and positive definite, thus Hermitian. Consequently, QLSAs can solve the NES efficiently. By its nature, a QLSA generates an inexact solution $\widetilde{\Delta y}^k$ with error bound $\|\Delta y^k - \widetilde{\Delta y}^k\| \leq \epsilon^k$. This error leads to residual r^k as

$$M^k \widetilde{\Delta y}^k = \sigma^k + r^k,$$

where $r^k = M^k(\widetilde{\Delta}y^k - \Delta y^k)$. After finding $\widetilde{\Delta}y^k$ inexactly by solving the NES using QLSA, we compute the inexact $\widetilde{\Delta}x^k$ and $\widetilde{\Delta}s^k$ classically as

$$\begin{aligned}\widetilde{\Delta}s^k &= c - A^T y^k - s^k - A^T \widetilde{\Delta}y^k, \\ \widetilde{\Delta}x^k &= \beta_1 \mu^k (S^k)^{-1} e - x^k - (D^k)^2 \widetilde{\Delta}s^k.\end{aligned}\tag{3}$$

As $\widetilde{\Delta}s^k$ and $\widetilde{\Delta}x^k$ are directly calculated by equations (3), one can verify that $(\widetilde{\Delta}x^k, \widetilde{\Delta}s^k, \widetilde{\Delta}y^k)$ satisfies

$$\begin{aligned}A\widetilde{\Delta}x^k &= b - Ax^k + r^k, \\ A^T \widetilde{\Delta}y^k + \widetilde{\Delta}s^k &= c - A^T y^k - s^k, \\ X^k \widetilde{\Delta}s^k + S^k \widetilde{\Delta}x^k &= \beta_1 \mu^k e - X^k s^k.\end{aligned}\tag{4}$$

To have an II-IPM using (NES) with iteration complexity $\mathcal{O}(n^2L)$, the residual norm must decrease at least $\mathcal{O}(\lambda_{\min}(A)\sqrt{n}\log n)$ time faster than $(x^k)^T s^k$ where $\lambda_{\min}(A)$ is the smallest singular value of A [46]. We can have wider residual bound but with higher iteration complexity of II-IPM [10]. In the literature of preconditioning the NES, some papers modified the equations (3) and the (NES) to transfer the residual from the first equation of (4) to its last equation. By these changes, we can get much better bounds [24, 35]. Since tight residual bound leads to the high complexity of QLSA+QTA, in this paper, we use a modification of the NES, which leads to $\mathcal{O}(n^2L)$ iteration complexity of II-QIPM, where the residual is decreasing with the rate of $\mathcal{O}(\sqrt{\mu^k})$.

Since A has full row rank, one can choose an arbitrary basis \hat{B} , and calculate $A_{\hat{B}}^{-1}$, $\hat{A} = A_{\hat{B}}^{-1}A$, and $\hat{b} = A_{\hat{B}}^{-1}b$. This calculation needs $\mathcal{O}(m^2n)$ arithmetic operations and happens just one time before the iterations of IPM. The cost of this preprocessing is dominated by the cost of II-QIPM, but it can be reduced by using the structure of A . For example, if the problem is in the canonical form, there is no need for this preprocessing. In the rest of this paper, all methodology is applied to the preprocessed problem with input data (\hat{A}, \hat{b}, c) . Now, we can modify the (NES) to

$$\hat{M}^k z^k = \hat{\sigma}^k \tag{MNES}$$

where

$$\begin{aligned}\hat{M}^k &= (D_{\hat{B}}^k)^{-1} A_{\hat{B}}^{-1} M^k ((D_{\hat{B}}^k)^{-1} A_{\hat{B}}^{-1})^T = (D_{\hat{B}}^k)^{-1} \hat{A} (D^k)^2 ((D_{\hat{B}}^k)^{-1} \hat{A})^T, \\ \hat{\sigma}^k &= (D_{\hat{B}}^k)^{-1} A_{\hat{B}}^{-1} \sigma^k = (D_{\hat{B}}^k)^{-1} \hat{b} - \beta_1 \mu^k (D_{\hat{B}}^k)^{-1} \hat{A} (S^k)^{-1} e + (D_{\hat{B}}^k)^{-1} \hat{A} (D^k)^2 (c - A^T y^k - s^k),\end{aligned}$$

where $D_{\hat{B}}^k = (X_{\hat{B}}^k)^{1/2} (S_{\hat{B}}^k)^{-1/2}$. We use the following procedure to find the Newton direction by solving (MNES) inexactly with QLSA+QTA.

- Step 1.** Find \tilde{z}^k such that $\hat{M}^k \tilde{z}^k = \hat{\sigma}^k + \hat{r}^k$ and $\|\hat{r}^k\| \leq \eta \frac{\sqrt{\mu^k}}{\sqrt{n}}$.
- Step 2.** Calculate $\widetilde{\Delta}y^k = ((D_{\hat{B}}^k)^{-1} A_{\hat{B}}^{-1})^T \tilde{z}^k$.
- Step 3.** Calculate $v^k = (v_{\hat{B}}^k, v_{\hat{N}}^k) = (D_{\hat{B}}^k \hat{r}^k, 0)$.
- Step 4.** Calculate $\widetilde{\Delta}s^k = c - A^T y^k - s^k - A^T \widetilde{\Delta}y^k$.
- Step 5.** Calculate $\widetilde{\Delta}x^k = \beta_1 \mu^k (S^k)^{-1} e - x^k - (D^k)^2 \widetilde{\Delta}s^k - v^k$.

The following Lemma shows how the inexact solution of (MNES) leads to residual only in the last equation of the Newton system.

Lemma 4.1 For the Newton direction $(\widetilde{\Delta x}^k, \widetilde{\Delta y}^k, \widetilde{\Delta s}^k)$, we have

$$\begin{aligned} A\widetilde{\Delta x}^k &= b - Ax^k, \\ A^T\widetilde{\Delta y}^k + \widetilde{\Delta s}^k &= c - A^T y^k - s^k, \\ X^k\widetilde{\Delta s}^k + S^k\widetilde{\Delta x}^k &= \beta_1\mu^k e - X^k s^k - S^k v^k. \end{aligned} \quad (5)$$

Proof. For the Newton direction $(\widetilde{\Delta x}^k, \widetilde{\Delta y}^k, \widetilde{\Delta s}^k)$, one can verify that

$$\begin{aligned} \widehat{M}^k \widehat{z}^k &= \widehat{\sigma}^k + \widehat{r}^k \\ M^k \widetilde{\Delta y}^k &= \sigma^k + A_{\widehat{B}} D_{\widehat{B}}^k \widehat{r}^k. \end{aligned}$$

For the first equation of (5), we can write

$$\begin{aligned} A\widetilde{\Delta x}^k &= A(\beta_1\mu^k(S^k)^{-1}e - x^k - (S^k)^{-1}X^k\widetilde{\Delta s}^k - v^k) \\ &= A(\beta_1\mu^k(S^k)^{-1}e - x^k - (S^k)^{-1}X^k(c - s^k - A^T y^k - A^T\widetilde{\Delta y}^k) - v^k) \\ &= \beta_1\mu^k A(S^k)^{-1}e - Ax^k - A(S^k)^{-1}X^k c + A(S^k)^{-1}X^k s^k + A(S^k)^{-1}X^k A^T y^k \\ &\quad + A(S^k)^{-1}X^k A^T \widetilde{\Delta y}^k - Av^k \\ &= \beta_1\mu^k A(S^k)^{-1}e - A(S^k)^{-1}X^k c + A(S^k)^{-1}X^k A^T y^k + \sigma^k + A_{\widehat{B}} D_{\widehat{B}}^k \widehat{r}^k - A_{\widehat{B}} D_{\widehat{B}}^k \widehat{r}^k \\ &= b - Ax^k. \end{aligned}$$

The second and third equations of (5) are obtained by Steps 4 and 5. \square

To have a convergent IPM, we need $\|S^k v^k\|_\infty \leq \eta\mu^k$, where $0 \leq \eta < 1$ is an enforcing parameter.

Lemma 4.2 For the Newton direction $(\widetilde{\Delta x}^k, \widetilde{\Delta y}^k, \widetilde{\Delta s}^k)$, if the residual $\|\widehat{r}^k\| \leq \eta\frac{\sqrt{\mu^k}}{\sqrt{n}}$, then $\|S^k v^k\|_\infty \leq \eta\mu^k$.

Proof. We have

$$\|S^k v^k\|_\infty = \|S_{\widehat{B}}^k v_{\widehat{B}}^k\|_\infty = \|S_{\widehat{B}}^k D_{\widehat{B}}^k \widehat{r}^k\|_\infty = \|(S_{\widehat{B}}^k)^{1/2} (X_{\widehat{B}}^k)^{1/2}\|_\infty \|\widehat{r}^k\|_\infty \leq \sqrt{n\mu^k} \|\widehat{r}^k\| \leq \eta\mu^k. \quad \square$$

In the following, we show that by satisfying $\|\widehat{r}^k\| \leq \eta\frac{\sqrt{\mu^k}}{\sqrt{n}}$, then the iterations of the II-QIPM remain in the $\mathcal{N}(\gamma_1, \gamma_2)$ neighborhood of the central path. The following theorem presents the complexity of solving the (MNES) system by utilizing the QLSA of Chakraborty et al. [14]. We can also use other QLSAs discussed in Section 2, leading to different complexity bounds.

Lemma 4.3 The QLSA by [14] and the QTA by [3] can build the (MNES) system, and produce a solution \widehat{z}^k for the (MNES) system satisfying $\|\widehat{r}^k\| \leq \eta\frac{\sqrt{\mu^k}}{\sqrt{n}}$ with $\tilde{\mathcal{O}}_{n, \frac{1}{\mu^k}, \|\widehat{\sigma}^k\|} (mn + m\sqrt{n} \frac{\kappa_E^k \|\widehat{\sigma}^k\|}{\sqrt{\mu^k}})$ complexity, where $E^k = (D_{\widehat{B}}^k)^{-1} \widehat{A} D^k$, and κ_E^k is the condition number of E^k .

Proof. Building the (MNES) system in classical computer needs some matrix multiplications, which costs $\mathcal{O}(m^2 n)$ arithmetic operations. We can write (MNES) as $E^k (E^k)^T \widehat{z}^k = \widehat{\sigma}^k$. As we can see, calculating E^k and $\widehat{\sigma}^k$ needs just $\mathcal{O}(mn)$ arithmetic operations. Chakraborty et al. [14] proposed an efficient way to build and solve a linear system in the form $E^k (E^k)^T \widehat{z}^k = \widehat{\sigma}^k$, with $\tilde{\mathcal{O}}(\text{polylog}(\frac{n}{\epsilon_{QLSA}^k}) \kappa_E^k)$ complexity. Also, we need to find the target precision for QLSA and

QTA such that $\|\hat{r}^k\| \leq \eta \frac{\sqrt{\mu^k}}{\sqrt{n}}$ is satisfied. Thus, to have $\|\hat{r}^k\| \leq \eta \frac{\sqrt{\mu^k}}{\sqrt{n}}$, it is sufficient to require $\epsilon_{LSP}^k \leq \eta \frac{\sqrt{\mu^k}}{\sqrt{n} \|\hat{M}^k\|}$. Based on the discussion of Section 2, we need to have

$$\epsilon_{QLSP}^k \leq \eta \frac{\sqrt{\mu^k}}{\sqrt{n} \|\hat{\sigma}^k\|}, \quad \epsilon_{QLSA}^k \leq \eta \frac{\sqrt{\mu^k}}{2\sqrt{n} \|\hat{\sigma}^k\|}, \quad \text{and} \quad \epsilon_{QTA}^k \leq \eta \frac{\sqrt{\mu^k}}{2\sqrt{n} \|\hat{\sigma}^k\|}.$$

With this target precision, the QLSA by [14] has $\mathcal{O}(\text{polylog}(\frac{n^{1.5} \|\hat{\sigma}^k\|}{\sqrt{\mu^k}}) \kappa_E^k)$ complexity, and the QTA by [3] has $\mathcal{O}(\frac{m\sqrt{n} \|\hat{\sigma}^k\|}{\sqrt{\mu^k}})$ complexity. Since calculating E^k and $\hat{\sigma}^k$ classically needs $\mathcal{O}(mn)$ arithmetic operations and the cost of storing them in QRAM is also $\mathcal{O}(mn)$, the total cost of building and solving the (NES) system is

$$\tilde{\mathcal{O}}_{n, \frac{1}{\mu^k}, \|\hat{\sigma}^k\|} (mn + m\sqrt{n} \frac{\kappa_E^k \|\hat{\sigma}^k\|}{\sqrt{\mu^k}}).$$

The proof is complete. \square

We present the II-QIPM as Algorithm 1 for solving LO problems. In this algorithm, we use QLSA and QTA to solve the NES.

Algorithm 1 II-QIPM

- 1: Choose $\zeta > 0$, $\gamma_1 \in (0, 1)$, $\gamma_2 > 0$, $0 < \eta < \beta_1 < \beta_2 < 1$,
 - 2: Choose $\omega \geq \max\{1, \|x^*, s^*\|_\infty\}$.
 - 3: $k \leftarrow 0$, $(x^0, y^0, s^0) \leftarrow (\omega e, 0e, \omega e)$, and $\gamma_2 \leftarrow \max\left\{1, \frac{\|(R_P^0, R_D^0)\|}{\mu^0}\right\}$
 - 4: **while** $(x^k, y^k, s^k) \notin \mathcal{PD}_\zeta$ **do**
 - 5: $\mu^k \leftarrow \frac{(x^k)^T s^k}{n}$
 - 6: $E^k \leftarrow (D_B^k)^{-1} \hat{A} D^k$ and $\hat{\sigma}^k \leftarrow (D_B^k)^{-1} \hat{b} - \beta_1 \mu^k (D_B^k)^{-1} \hat{A} (S^k)^{-1} e + (D_B^k)^{-1} \hat{A} (D^k)^2 (c - A^T y^k - s^k)$
 - 7: $\epsilon_{QLSA}^k \leftarrow \eta \frac{\sqrt{\mu^k}}{2\sqrt{n} \|\hat{\sigma}^k\|}$ and $\epsilon_{QTA}^k \leftarrow \eta \frac{\sqrt{\mu^k}}{2\sqrt{n} \|\hat{\sigma}^k\|}$
 - 8: $(\Delta x^k, \Delta y^k, \Delta s^k) \leftarrow \text{solve MNES}(\beta_1)$ by QLSA+QTA with precision ϵ_{QLSA}^k and ϵ_{QTA}^k
 $\hat{\alpha}^k \leftarrow \max\{\bar{\alpha} \in [0, 1] \mid \text{for all } \alpha \in [0, \bar{\alpha}] \text{ we have}$
 - 9: $((x^k, y^k, s^k) + \alpha(\Delta x^k, \Delta y^k, \Delta s^k)) \in \mathcal{N}(\gamma_1, \gamma_2)$ and
 $(x^k + \alpha \Delta x^k)^T (s^k + \alpha \Delta s^k) \leq (1 - \alpha(1 - \beta_2))(x^k)^T s^k\}$
 - 10: $(x^{k+1}, y^{k+1}, s^{k+1}) \leftarrow (x^k, y^k, s^k) + \hat{\alpha}^k (\Delta x^k, \Delta y^k, \Delta s^k)$
 - 11: **if** $\|x_{k+1}, s_{k+1}\|_\infty > \omega$ **then**
 - 12: **return** Primal or dual is infeasible.
 - 13: $k \leftarrow k + 1$
 - 14: **return** (x^k, y^k, s^k)
-

It can be easily verified that e.g., $\beta_1 = 0.5$, $\beta_2 = 0.9995$, $\eta = 0.4$ and $\gamma_1 = 0.5$ yield a valid choice, i.e. satisfying the conditions in the first line of Algorithm 1. In the following, we prove the polynomial complexity of Algorithm 1.

4.1 Convergence of the II-QIPM

In this section, a convergence proof and iteration complexity bound are provided for the proposed II-QIPM. The analysis closest to ours can be found in [24] and [35] where they are preconditioning

the Newton systems which are then solved by CG. However, we modified the NES and solved it by QLSA+QTA under different conditions and assumptions. First, in Lemma 4.4, we study basic properties of the proposed II-QIPM as presented in Algorithm 1. Then, Lemma 4.7 shows that the sequence $\{\hat{\alpha}^k\}$ is strictly positive for all k . The iteration complexity of Algorithm 1 is proved in Theorem 4.1. In the proof, we need to analyze values of α^k such that at iteration k of Algorithm 1, the Newton step satisfies all the conditions in line 9. To ease notation, we use

$$x^k(\alpha) = x^k + \alpha \Delta x^k, \quad y^k(\alpha) = y^k + \alpha \Delta y^k, \quad s^k(\alpha) = s^k + \alpha \Delta s^k, \quad \mu^k(\alpha) = \frac{x^k(\alpha)^T s^k(\alpha)}{n},$$

$$R_P^k(\alpha) = b - Ax^k(\alpha), \quad \text{and} \quad R_D^k(\alpha) = c - A^T y^k(\alpha) - s^k(\alpha).$$

The following lemma shows some properties of the proposed II-QIPM.

Lemma 4.4 *At iteration k of Algorithm 1, for any $\alpha \in [0, 1]$, with $\mu^k = \frac{(x^k)^T s^k}{n}$, we have*

$$R_P^k(\alpha) = (1 - \alpha)R_P^k, \quad (6a)$$

$$R_D^k(\alpha) = (1 - \alpha)R_D^k, \quad (6b)$$

$$(x^k(\alpha))^T s^k(\alpha) \geq (1 + \alpha(\beta_1 - \eta - 1))n\mu^k + \alpha^2(\Delta x^k)^T \Delta s^k, \quad (6c)$$

$$x_i^k(\alpha)s_i^k(\alpha) \geq (1 - \alpha)x_i^k s_i^k + \alpha(\beta_1 - \eta)\mu^k + \alpha^2 \Delta x_i^k \Delta s_i^k \text{ for } i \in \{1, 2, \dots, n\}. \quad (6d)$$

Proof. To prove (6a) and (6b), for any $\alpha \in [0, 1]$, by (2) we have

$$R_P^k(\alpha) = b - A(x^k + \alpha \Delta x^k) = b - Ax^k - \alpha A \Delta x^k = b - Ax^k - \alpha(b - Ax^k) = (1 - \alpha)R_P^k,$$

$$R_D^k(\alpha) = c - A^T(y^k + \alpha \Delta y^k) - s^k - \alpha \Delta s^k = c - A^T y^k - s^k - \alpha(A^T \Delta y + \Delta s) = (1 - \alpha)R_D^k.$$

To prove (6c), using (5), we have

$$\begin{aligned} (x^k + \alpha \Delta x^k)^T (s^k + \alpha \Delta s^k) &= (x^k)^T s^k + \alpha[(x^k)^T \Delta s^k + (s^k)^T \Delta x^k] + \alpha^2(\Delta x^k)^T \Delta s^k \\ &\geq (x^k)^T s^k + \alpha[n\beta_1\mu^k - (x^k)^T s^k - n\eta\mu^k] + \alpha^2(\Delta x^k)^T \Delta s^k \\ &= [1 + \alpha(\beta_1 - \eta - 1)](x^k)^T s^k + \alpha^2(\Delta x^k)^T \Delta s^k. \end{aligned}$$

Using (5) again, we can similarly prove (6d) for all $i \in \{1, 2, \dots, n\}$ as follows:

$$\begin{aligned} (x_i^k + \alpha \Delta x_i^k)(s_i^k + \alpha \Delta s_i^k) &= x_i^k s_i^k + \alpha(x_i^k \Delta s_i^k + s_i^k \Delta x_i^k) + \alpha^2 \Delta x_i^k \Delta s_i^k \\ &\geq x_i^k s_i^k + \alpha(\beta_1 \mu^k - x_i^k s_i^k - \eta \mu^k) + \alpha^2 \Delta x_i^k \Delta s_i^k \\ &= (1 - \alpha)x_i^k s_i^k + \alpha(\beta_1 - \eta)\mu^k + \alpha^2 \Delta x_i^k \Delta s_i^k. \end{aligned}$$

Thus, the proof is complete. \square

Let us define the following functions

$$G_i^k(\alpha) = x_i^k(\alpha)s_i^k(\alpha) - \gamma_1\mu^k(\alpha) \text{ for } i \in \{1, \dots, n\},$$

$$g^k(\alpha) = x^k(\alpha)^T s^k(\alpha) - (1 - \alpha)(x^k)^T s^k,$$

$$h^k(\alpha) = (1 - \alpha(1 - \beta_2))(x^k)^T s^k - x^k(\alpha)^T s^k(\alpha).$$

We use the defined functions to check if a step with length α sufficiently reduces the complementarity gap and keeps the next iterate in the neighborhood $\mathcal{N}(\gamma_1, \gamma_2)$. It is obvious that $h^k(\alpha) \geq 0$ means the Armijo condition $(x^k(\alpha))^T s^k(\alpha) \leq (1 - \alpha(1 - \beta_2))(x^k)^T s^k$ holds, and the next lemma shows how we check that an iterate is in the neighborhood of the central path.

Lemma 4.5 For step length $0 < \alpha \leq 1$, if $G_i^k(\alpha) \geq 0$ and $g^k(\alpha) \geq 0$ then $(x^k(\alpha), y^k(\alpha), s^k(\alpha)) \in \mathcal{N}(\gamma_1, \gamma_2)$.

Proof. It is easy to verify that that conditions $G_i^k(\alpha) \geq 0$ and $g^k(\alpha) \geq 0$ lead to

$$\begin{aligned} x_i^k(\alpha) s_i^k(\alpha) &\geq \gamma_1 \mu^k(\alpha) \text{ for } i \in \{1, \dots, n\}, \\ (x^k + \alpha \Delta x^k)^T (s^k + \alpha \Delta s^k) &\leq (1 - \alpha(1 - \beta_2)) (x^k)^T s^k, \end{aligned}$$

respectively. Since $g^k(\alpha) \geq 0$, i.e. $x^k(\alpha)^T s^k(\alpha) \geq (1 - \alpha)(x^k)^T s^k$, we have

$$\frac{\|(R_P^k(\alpha), R_D^k(\alpha))\|}{\|(R_P^0, R_D^0)\|} = \frac{(1 - \alpha)\|(R_P^k, R_D^k)\|}{\|(R_P^0, R_D^0)\|} \leq \frac{(1 - \alpha)\mu^k}{\mu^0} \leq \frac{\mu^k(\alpha)}{\mu^0}.$$

Further, as $\gamma_2 = \frac{\|(R_P^0, R_D^0)\|}{\mu^0}$, we can conclude that $(x^k(\alpha), y^k(\alpha), s^k(\alpha)) \in \mathcal{N}(\gamma_1, \gamma_2)$. \square

In order to prove polynomial complexity of II-QIPM we need to find a positive lower bound for the step length $\hat{\alpha}^k$. The following lemma is bounding some remaining elements to get the step length bound.

Lemma 4.6 There exist $0 \leq \nu^k = \mathcal{O}(n^2 \mu^k)$ such that $\left| \Delta x_i^k \Delta s_i^k - \gamma_1 \frac{(\Delta x^k)^T \Delta s^k}{n} \right| \leq \nu^k$ for $i \in \{1, 2, \dots, n\}$ and $|(\Delta x^k)^T \Delta s^k| \leq \nu^k$.

Proof. To prove this lemma, we need to do the following steps

- Step 1. finding bound $\mathcal{C}_1^k = \mathcal{O}(n\mu^k)$ such that $\omega\theta^{k-1}\|x^k, s^k\|_1 \leq \mathcal{C}_1^k$;
- Step 2. finding bound $\mathcal{C}_2^k = \mathcal{O}(n\sqrt{\mu^k})$ such that $\|D^{-1}\Delta x^k\| \leq \mathcal{C}_2^k$ and $\|D\Delta s^k\| \leq \mathcal{C}_2^k$;
- Step 3. finding bound $0 \leq \nu^k = \mathcal{O}(n^2\mu^k)$ such that

$$|(\Delta x^k)^T \Delta s^k| \leq \nu^k \text{ and } \left| \Delta x_i^k \Delta s_i^k - \gamma_1 \frac{(\Delta x^k)^T \Delta s^k}{n} \right| \leq \nu^k \text{ for } i \in \{1, 2, \dots, n\}.$$

Step 1. Let us define $\theta^k = \prod_{i=0}^k (1 - \hat{\alpha}^i)$. One can verify that

$$R_P^k = \theta^{k-1} R_P^0, \text{ and } R_D^k = \theta^{k-1} R_D^0. \quad (7)$$

Based on the definition of $\mathcal{N}(\gamma_1, \gamma_2)$ and the choice of γ_2 , we have

$$\frac{\|(R_P^k, R_D^k)\|}{\mu^k} \leq \frac{\|(R_P^0, R_D^0)\|}{\mu^0}, \quad (8)$$

implying $\mu^k \geq \theta^{k-1} \mu^0$. We also define

$$(\bar{x}^k, \bar{y}^k, \bar{s}^k) = \theta^{k-1}(x^0, y^0, s^0) + (1 - \theta^{k-1})(x^*, y^*, s^*) - (x^k, y^k, s^k),$$

where, $(x^*, y^*, s^*) \in \mathcal{PD}^*$. One can verify that

$$\begin{aligned} A^T \bar{y}^k + \bar{s}^k &= 0, \\ A \bar{x}^k &= 0. \end{aligned}$$

Since \bar{s}^k is in the row space of A and \bar{x}^k is in the null space of A , we have $(\bar{x}^k)^T \bar{s}^k = 0$, or equivalently,

$$[\theta^{k-1}x^0 + (1 - \theta^{k-1})x^* - x^k]^T [\theta^{k-1}s^0 + (1 - \theta^{k-1})s^* - s^k] = 0.$$

Since $(x^*, s^*, x^k, s^k) \geq 0$, $x^* s^* = 0$, and $(x^0, s^0) = (\omega e, \omega e)$, we can write

$$[\theta^{k-1}x^0 + (1 - \theta^{k-1})x^* - x^k]^T [\theta^{k-1}s^0 + (1 - \theta^{k-1})s^* - s^k] = 0, \quad (9)$$

$$(\theta^{k-1})^2 (s^0)^T x^0 + \theta^{k-1} (1 - \theta^{k-1}) [(s^0)^T x^* + (s^*)^T x^0] + (s^k)^T x^k \geq \theta^{k-1} [(s^0)^T x^k + (x^0)^T s^k], \quad (10)$$

$$(\theta^{k-1})^2 n\mu^0 + 2\theta^{k-1} (1 - \theta^{k-1}) n\mu^0 + (s^k)^T x^k \geq \theta^{k-1} \omega [(e)^T x^k + (e)^T s^k], \quad (11)$$

$$\theta^{k-1} n\mu^k + 2(1 - \theta^{k-1})\mu^k + n\mu^k \geq \theta^{k-1} \omega \|x^k, s^k\|_1. \quad (12)$$

Inequality (11) is obtained by using $(x^0, s^0) = (\omega e, \omega e)$ and $\|(x^*, s^*)\|_\infty \leq \omega$, and the last inequality is obtained by using (7). Thus, let \mathcal{C}_1^k be defined as the left-hand side of (12), we have $\mathcal{C}_1^k = \mathcal{O}(n\mu^k)$.

Step 2. In this step we define

$$(\bar{x}^k, \bar{y}^k, \bar{s}^k) = (\Delta x^k, \Delta y^k, \Delta s^k) + \theta^{k-1}(x^0, y^0, s^0) - \theta^{k-1}(x^*, y^*, s^*),$$

where, $(x^*, y^*, s^*) \in \mathcal{PD}^*$. Similar to Step 1, one can verify that

$$A^T \bar{y}^k + \bar{s}^k = 0,$$

$$A \bar{x}^k = 0,$$

$$(\bar{x}^k)^T \bar{s}^k = 0.$$

Consequently, one can verify

$$\begin{aligned} & \|D^{-1}(\Delta x^k + \theta^{k-1}(x^0 - x^*)) + D(\Delta s^k + \theta^{k-1}(s^0 - s^*))\| \\ &= \|D^{-1}(\Delta x^k + \theta^{k-1}(x^0 - x^*))\| + \|D(\Delta s^k + \theta^{k-1}(s^0 - s^*))\| \\ &\leq \|(XS)^{-1/2}(\|XSe - \beta_1 \mu^k e\| + \eta \mu^k) + \theta^{k-1}\| \|D^{-1}(x^0 - x^*)\| + \theta^{k-1} \|D(s^0 - s^*)\|. \end{aligned}$$

Now, we have

$$\|D^{-1} \Delta x^k\| \leq \|(XS)^{-1/2}(\|XSe - \beta_1 \mu^k e\| + \eta \mu^k) + 2\theta^{k-1}\| \|D^{-1}(x^0 - x^*)\| + 2\theta^{k-1} \|D(s^0 - s^*)\|.$$

According to pages 116-118 of [43], we can derive the following inequalities:

$$\begin{aligned} \|(XS)^{-1/2}\| &\leq \frac{1}{\sqrt{\gamma_1} \sqrt{\mu^k}}, \\ \|XSe - \beta_1 \mu^k e\| &\leq n\mu^k, \\ \theta^{k-1} \|D^{-1}(x^0 - x^*)\| + \theta^{k-1} \|D(s^0 - s^*)\| &\leq \theta^{k-1} \|x^k, s^k\|_1 \|(XS)^{-1/2}\| \max(\|x^0 - x^*\|, \|s^0 - s^*\|) \\ &\leq \frac{\omega \theta^{k-1} \|x^k, s^k\|_1}{\sqrt{\gamma_1} \sqrt{\mu^k}} \leq \frac{C_1}{\sqrt{\gamma_1} \sqrt{\mu^k}}. \end{aligned}$$

Thus, $\|D^{-1} \Delta x^k\| \leq \frac{n\mu^k + \eta\mu^k + C_1^k}{\sqrt{\gamma_1} \sqrt{\mu^k}} = \mathcal{C}_2^k$, where $\mathcal{C}_2^k = \mathcal{O}(n\sqrt{\mu^k})$. We can similarly show that $\|D \Delta s^k\| \leq \mathcal{C}_2^k$.

Step 3. Based on Steps 1 and 2, we have

$$(\Delta x^k)^T \Delta s^k \leq \|D^{-1} \Delta x^k\| \|D \Delta s^k\| \leq (\mathcal{C}_2^k)^2,$$

$$|\Delta x_i^k \Delta s_i^k| \leq \|D^{-1} \Delta x^k\| \|D \Delta s^k\| \leq (\mathcal{C}_2^k)^2,$$

$$\left| \Delta x_i^k \Delta s_i^k - \gamma_1 \frac{(\Delta x^k)^T \Delta s^k}{n} \right| \leq (1 + \frac{\gamma_1}{n}) (\mathcal{C}_2^k)^2 \leq 2(\mathcal{C}_2^k)^2.$$

Thus, $\nu^k = 2(\mathcal{C}_2^k)^2 = \mathcal{O}(n^2 \mu^k)$. □

In the next lemma, we present a strictly positive lower bound for $\hat{\alpha}^k$. In the proof, we use parameters δ_1 , δ_2 , and δ_3 defined as follows:

$$\delta_1 = \frac{(1 - \gamma_1)(\beta_1 - \eta)}{n} > 0, \quad \delta_2 = \beta_1 - \eta > 0, \quad \delta_3 = \beta_2 - \beta_1 + \eta > 0. \quad (13)$$

Lemma 4.7 *At line 9 of Algorithm 1, at iteration k , we have*

$$\hat{\alpha}^k \geq \tilde{\alpha}^k := \min \left\{ 1, \min\{\delta_1, \delta_2, \delta_3\} \frac{(x^k)^T s^k}{\nu^k} \right\} > 0.$$

Proof. It is enough to show that the conditions of Lemma 4.5 hold for all $\alpha \in [0, \tilde{\alpha}^k]$. Based on Lemma 4.4, for any $\alpha \in [0, \tilde{\alpha}^k]$, we have

$$G_i^k(\alpha) = (x_i^k + \alpha \Delta x_i^k)(s_i^k + \alpha \Delta s_i^k) - \gamma_1 \frac{(x^k + \alpha \Delta x^k)^T (s^k + \alpha \Delta s^k)}{n} \quad (14a)$$

$$= (1 - \alpha)x_i^k s_i^k + \alpha(\beta_1 - \eta)\mu^k + \alpha^2 \Delta x_i^k \Delta s_i^k - \gamma_1 \frac{(1 + \alpha(\beta_1 - \eta - 1))(x^k)^T s^k + \alpha^2 (\Delta x^k)^T \Delta s^k}{n} \quad (14b)$$

$$\geq \alpha^2 (\Delta x_i^k \Delta s_i^k - \frac{\gamma_1}{n} (\Delta x^k)^T \Delta s^k) + (1 - \alpha)(x_i^k s_i^k - \frac{\gamma_1}{n} (x^k)^T s^k) + \alpha(\beta_1 - \eta)(1 - \gamma_1)\mu^k \quad (14c)$$

$$\geq -\alpha^2 \nu^k + \alpha \delta_1 (x^k)^T s^k \geq \alpha(\delta_1 (x^k)^T s^k - \nu^k \tilde{\alpha}^k) \geq 0. \quad (14d)$$

Equality (14b) follows from equation (6a) of Lemma 4.4, and inequality (14d) is due to the definition of $\tilde{\alpha}^k$ and the neighborhood $\mathcal{N}(\gamma_1, \gamma_2)$. Similarly, we show that $g_i^k(\alpha) \geq 0$ as

$$\begin{aligned} g_i^k(\alpha) &= x^k(\alpha)^T s^k(\alpha) - (1 - \alpha)(x^k)^T s^k \\ &\geq (1 + \alpha(\beta_1 - \eta - 1))(x^k)^T s^k + \alpha^2 (\Delta x^k)^T \Delta s^k - (1 - \alpha)(x^k)^T s^k \\ &\geq \alpha(\beta_1 - \eta)(x^k)^T s^k - \alpha^2 \nu^k \\ &\geq \alpha \delta_2 (x^k)^T s^k - \alpha^2 \nu^k \\ &\geq \alpha(\delta_2 (x^k)^T s^k - \nu^k \tilde{\alpha}^k) \geq 0. \end{aligned}$$

Again, by (6c) of Lemma 4.4, we have for all $\alpha \in [0, \tilde{\alpha}^k]$

$$\begin{aligned} h^k(\alpha) &= (1 - \alpha(1 - \beta_2))(x^k)^T s^k - (x^k + \alpha \Delta x^k)^T (s^k + \alpha \Delta s^k) \\ &= (1 - \alpha(1 - \beta_2))(x^k)^T s^k - (1 + \alpha(\beta_1 - \eta - 1))(x^k)^T s^k - \alpha^2 (\Delta x^k)^T \Delta s^k \\ &\geq \alpha(\beta_2 - \beta_1 + \eta)(x^k)^T s^k - \alpha^2 \nu^k \\ &\geq \alpha \delta_3 (x^k)^T s^k - \alpha^2 \nu^k \\ &\geq \alpha(\delta_3 (x^k)^T s^k - \nu^k \tilde{\alpha}^k) \geq 0. \end{aligned}$$

We showed that for all $\alpha \in [0, \tilde{\alpha}^k]$, all the conditions of Lemma 4.5 hold. Thus, we can conclude that $\tilde{\alpha}^k \leq \hat{\alpha}^k$ and the proof is complete. \square

By Lemma 4.7, we have a strictly positive lower bound for step length $\tilde{\alpha}^k$ to remain in the neighborhood of the central path while we decrease the optimality gap. In what follows, using the results of the previous lemmas, we establish the iteration complexity of Algorithm 1.

Theorem 4.1 *If Algorithm 1 does not terminate in line 11, then it reaches a ζ -optimal solution in at most $\mathcal{O}(n^2 \log \frac{1}{\zeta})$ iterations.*

Proof. Based on Lemma 4.7, we have

$$\hat{\alpha}^k \geq \alpha^k \geq \tilde{\alpha}^k := \min \left\{ 1, \min \{ \delta_1, \delta_2, \delta_3 \} \frac{n\mu^k}{\nu^k} \right\} \in (0, 1].$$

Hence, by the definition of the neighborhood, we have

$$(x^k)^T s^k \leq (1 - \tilde{\alpha}^k(1 - \beta_2))^k (x^0)^T s^0.$$

This implies that $\lim_{k \rightarrow \infty} (x^k)^T s^k = 0$. Thus, the algorithm terminates in finite number of steps. Since the iterates are in the $\mathcal{N}(\gamma_1, \gamma_2)$ neighborhood of the central path, a ζ -optimal solution is obtained when $\mu^k \leq \frac{\zeta}{\gamma_1}$. The algorithm stops when

$$\mu^k \leq (1 - \tilde{\alpha}^k(1 - \beta_2))^k \mu^0 \leq \frac{\zeta \mu^0}{\|R_p^0, R_D^0\|}.$$

By the definition of $\tilde{\alpha}^k$ and Lemma 4.6, we have $\frac{1}{\tilde{\alpha}^k} = \mathcal{O}(n^2)$. We can conclude that $k = \mathcal{O}(n^2 \log \frac{\omega}{\zeta})$. The proof is complete. \square

Remark 4.1 The sequences of $\{\mu^k\}$, primal infeasibility $\{\|Ax^k - b\|\}$, and dual infeasibility $\{\|A^T y^k + s^k - c\|\}$, generated by II-QIPM, converge linearly to zero.

Remark 4.2 Based on Theorem 3.1, we can calculate an exact solution by a rounding procedure if $\zeta \leq 2^{-4L}$. Consequently, the iteration complexity of II-QIPM for finding an exact solution is $\mathcal{O}(n^2 L)$.

In the next section, we provide the total time complexity of II-QIPM.

4.2 Total time complexity of the II-QIPM

As discussed in Lemma 4.3, we need to solve the NES at each iteration of the II-QIPM by subsequent application of QLSA and QTA, which requires $\tilde{\mathcal{O}}_{n, \frac{1}{\mu^k}, \|\hat{\sigma}^k\|} (mn + m\sqrt{n} \frac{\kappa_E^k \|\hat{\sigma}^k\|}{\sqrt{\mu^k}})$ computational cost. We can calculate the total time complexity of the II-QIPM as the product of the complexity of the QLSA and the number of iterations of the II-QIPM. The computational cost of the QLSA depends on κ_E^k , $\|\hat{\sigma}^k\|$, $\|E^k\|_F$, and μ^k which change through the algorithm. In the following theorem, we bound them properly and obtain the detailed total time complexity of the proposed II-QIPM algorithm.

Theorem 4.2 *The total time complexity of the proposed II-QIPM with QLSA by Chakraborty et al. [14] and QTA by Apeldoorn et al. [3] is*

$$\tilde{\mathcal{O}}_{n, \frac{1}{\zeta}, \omega, \|\hat{A}\|, \|\hat{b}\|} \left(n^2 \left[mn + \frac{m\sqrt{n}\kappa_{\hat{A}}(\|\hat{A}\| + \|\hat{b}\|)}{\zeta^3} \right] \right).$$

Proof. To establish the total time complexity of II-QIPM, we need to analyze how the matrices M^k and E^k evolve through the iterations. As in [36], considering the optimal partition B and N , we have

$$\frac{x_i^k}{s_i^k} = \mathcal{O}\left(\frac{1}{\mu^k}\right) \rightarrow \infty \text{ for } i \in B \quad \text{and} \quad \frac{x_i^k}{s_i^k} = \mathcal{O}(\mu^k) \rightarrow 0 \text{ for } i \in N. \quad (15)$$

Appropriate bounds are provided in the following results.

- (i) Based on Theorem 4.1, we have at the termination $\frac{1}{\mu^k} = \mathcal{O}(\frac{1}{\zeta})$ and $\mu^k \leq \mathcal{O}(\mu^0) = \mathcal{O}(\omega^2)$.
- (ii) Since $\|E^k\| \leq \|(D^k)^{-1}\|\|\hat{A}\|\|D^k\|$, and $\|D^k\| = \mathcal{O}(\frac{1}{\sqrt{\mu^k}}) = \mathcal{O}(\frac{1}{\sqrt{\zeta}})$ by (15). Similarly, $\|(D^k)^{-1}\|_F = \mathcal{O}(\frac{1}{\sqrt{\zeta}})$, and we have $\|E^k\| = \mathcal{O}(\frac{\|\hat{A}\|}{\zeta})$. Let $\kappa_{\hat{A}}$ be the condition number of \hat{A} . Thus, we have

$$\kappa_E^k = \mathcal{O}(\kappa_{\hat{A}}\zeta^{-2}).$$

- (iii) In the time complexity of II-QIPM, we also have $\frac{\|\sigma^k\|}{\sqrt{\mu^k}}$ coming from the precision of QLSA and QTA. We can easily verify that

$$\begin{aligned} \frac{\|\hat{\sigma}^k\|}{\sqrt{\mu^k}} &\leq \frac{\|(D^k)^{-1}\|}{\sqrt{\mu^k}} (\|\hat{b}\| + \|\hat{A}X^k(S^k)^{-1}\| \|c - A^T y^k - s^k\| + \beta_1 \|\mu^k \hat{A}(S^k)^{-1} e\|) \\ &\leq \frac{1}{\mu^k} (\|\hat{b}\| + \|\hat{A}X^k(S^k)^{-1}\| \|R_D^k\| + \beta_1 \|\hat{A}(S^k)^{-1} e\| \mu^k). \end{aligned}$$

Based on the definition of the neighborhood of the central path, $\|R_D^k\| \leq \gamma_2 \mu^k$. Thus, we can get bounds $\mathcal{O}(\frac{\|\sigma^k\|}{\sqrt{\mu^k}}) = \mathcal{O}(\frac{\|\hat{A}\| + \|\hat{b}\|}{\zeta})$.

- (iv) Based on Lemma 4.3, the complexity of QLSA by Chakraborty et al. [14] and QTA by Apeldoorn et al. [3] for building and solving the MNES is

$$\tilde{\mathcal{O}}_{n, \frac{1}{\zeta}, \|\hat{A}\|, \|\hat{b}\|} (mn + m\sqrt{n} \frac{\kappa_{\hat{A}}(\|\hat{A}\| + \|\hat{b}\|)}{\zeta^3}).$$

Thus, the detailed time complexity of the proposed II-QIPM with QLSA by Chakraborty et al. [14] and QTA by Apeldoorn et al. [3] is

$$\tilde{\mathcal{O}}_{n, \frac{1}{\zeta}, \omega, \|\hat{A}\|, \|\hat{b}\|} \left(n^2 \left[mn + \frac{m\sqrt{n}\kappa_{\hat{A}}(\|\hat{A}\| + \|\hat{b}\|)}{\zeta^3} \right] \right). \quad (16)$$

The time complexity is achieved by multiplying the number of iterations of II-QIPM and the total cost of each iteration, including building and solving it by QLSA+QTA. Thus, the proof is complete. \square

Let $\phi := \|\hat{A}\| + \|\hat{b}\|$, the complexity of II-QIPM can be simplified as

$$\tilde{\mathcal{O}}_{n, \frac{1}{\zeta}, \omega, \phi} \left(n^4 \frac{\phi \kappa_{\hat{A}}}{\zeta^3} \right). \quad (17)$$

In the complexity of II-QIPM, the $\frac{1}{\zeta}$ factors come from bounding the condition number and from QTA. Based on Theorem 3.1, $\frac{1}{\zeta} = 2^{\mathcal{O}(L)}$ leading to exponential complexity. We discuss how we can solve this and improve the complexity of the algorithm by stopping II-QIPM early, e.g., $\zeta = 10^{-2}$, and using the Iterative Refinement scheme discussed in the Section 5 to improve the precision.

Remark 4.3 Some QLSAs, such as Harrow et al. [23] and Childs et al. [15], take advantage of the sparsity of the NES. Consider (MNES), the sparsity of $\hat{M}^k = \hat{A}X^k(S^k)^{-1}\hat{A}^T$ is independent of X^k and S^k . So, let d be the maximum number of nonzero elements in any row or column of the matrix $\hat{A}\hat{A}^T$. Since matrix \hat{A} has two blocks $\begin{bmatrix} I & A_B^{-1}A_N \end{bmatrix}$, we have $d \leq \min\{m, n - m + 1\}$.

As matrix \hat{A} determines the sparsity of matrix M^k , so we can take advantage of the sparsity structure of matrix \hat{A} . In case \hat{A} is mostly sparse, e.g., $n - m \ll m$, but has a few dense columns, then this structure can be exploited. As described in [2], the sparse part can be separated to solve a sparse linear system by QLSA, and then the use of the Sherman-Morrison-Woodbury [38] formula allows calculating the solution of the original linear system efficiently.

5 Iterative Refinement Method

Based on Remark 4.2, we need $\zeta \leq 2^{-4L}$ to have an exact solution for the LO problem with integer data. The proposed II-QIPM has exponential complexity in finding an exact solution. An Iterative Refinement (IR) scheme can be employed to achieve polynomial complexity. By this scheme, a series of LO problems are solved by the II-QIPM with low precision $\hat{\zeta}$, e.g., $\hat{\zeta} = 10^{-2}$, and an IR method improves the precision to reach an exact solution. In the classical IPM literature, the IR method by Gleixner et al. [22] and the Rational Reconstruction method by Gleixner and Steffy [21] employed low-precision methods to generate a high-precision solution. Here, we adopt the IR method to generate a high precision solution that allows the identification of an exact optimal solution. While using only low-precision II-QIPM, Theorem 5.1 is the foundation of the IR method.

Theorem 5.1 (Gleixner et al. [22]) *Let the primal problem (P) be given as (1). For $\tilde{x} \in \mathbb{R}^n$ and $\tilde{y} \in \mathbb{R}^m$ and scaling factor $\nabla > 1$ consider the refining problem (\bar{P})*

$$\min \left\{ \nabla \bar{c}^T x \mid Ax = \nabla \bar{b} \text{ and } x \geq -\nabla \tilde{x} \right\},$$

where $\bar{c} = c - A^T \tilde{y}$ and $\bar{b} = b - A \tilde{x}$. Then \bar{x} and \bar{y} are $\hat{\zeta}$ -optimal solution for problem (\bar{P}) if and only if $\tilde{x} + \frac{1}{\nabla} \bar{x}$ and $\tilde{y} + \frac{1}{\nabla} \bar{y}$ are the $\hat{\zeta}$ -optimal solution for problem (P).

As presented in Algorithm 2, we use the II-QIPM of Algorithm 1 to solve the refining model and update the solution with an intelligent scaling procedure. Theorem 5.2 shows that a polynomial number of iterations are sufficient to reach an exact optimal solution.

Algorithm 2 IR-II-QIPM

Require: $(A \in \mathbb{Z}^{m \times n}, b \in \mathbb{Z}^m, c \in \mathbb{Z}^n)$

- 1: Choose scaling multiplier $\rho \in \mathbb{N}$ such that $\rho > 1$
 - 2: $\hat{\zeta} \leftarrow 10^{-2}, \zeta \leftarrow 2^{-4L}$
 - 3: $k \leftarrow 0$
 - 4: $(x^{*0}, y^{*0}, s^{*0}) \leftarrow$ Algorithm 1 (A, b, c) with $\hat{\zeta}$ precision
 - 5: **while** $r^k > \zeta$ **do**
 - 6: $\pi^k \leftarrow \max \left\{ r^k, \frac{1}{\rho \nabla^k} \right\}$
 - 7: $\nabla^k \leftarrow 2^{\lceil \log(1/\pi^k) \rceil}$
 - 8: $\bar{b}^k \leftarrow b - Ax^{*k}$ and $\bar{c}^k \leftarrow c - A^T y^{*k}$
 - 9: $(\hat{x}^k, \hat{y}^k, \hat{s}^k) \leftarrow$ Algorithm 1 $(A, \nabla^k \bar{b}^k, \nabla^k \bar{c}^k)$ with $\hat{\zeta}$ precision
 - 10: $x^{*k+1} \leftarrow x^{*k} + \frac{1}{\nabla^k} \hat{x}^k$ and $y^{*k+1} \leftarrow y^{*k} + \frac{1}{\nabla^k} \hat{y}^k$
 - 11: $r^{k+1} \leftarrow \max \left\{ \max_j |\bar{b}_j^{k+1}|, \max_i (-\bar{c}_i^{k+1}), \sum_i |\bar{c}_i^{k+1} x_i^{*k+1}| \right\}$
 - 12: $k \leftarrow k + 1$
-

Theorem 5.2 *Let (A, b, c) be integer. The number of iterations of Algorithm 2 to get 2^{-4L} -precise solution is at most $\mathcal{O}(L)$.*

Proof. Based on Corollary 3.6 in [22], we have

$$\left\lceil \frac{\log(\zeta)}{\log(\hat{\zeta})} \right\rceil = \left\lceil \frac{-4L \log(2)}{-2} \right\rceil = \mathcal{O}(L).$$

□

One can observe that, except ω , all parameters in the complexity of II-QIPM as specified in (17), are constant in all iterations of the IR method. Lemma 5.1 provides an upper bound for ω^k at iteration k of the IR method.

Lemma 5.1 *At the k^{th} iteration of the IR method, let $(\hat{x}^{*k}, \hat{y}^{*k}, \hat{s}^{*k})$ be an exact optimal solution of the refining problem $(A, \nabla^k \bar{b}_{k-1}, \nabla^k \bar{c}_{k-1})$, and $\omega^k \geq \max\{1, \|\hat{x}^{*k}, \hat{s}^{*k}\|_\infty\}$. Then, $\omega^k = \mathcal{O}((2\rho)^L)$.*

Proof. From Theorem 5.2 and last line of Algorithm 2, we have

$$x^* = x_0^* + \sum_{k=1}^{\mathcal{O}(L)} \frac{1}{\nabla^k} \hat{x}^k, \text{ and } s^* = s_0^* + \sum_{k=1}^{\mathcal{O}(L)} \frac{1}{\nabla^k} \hat{s}^k.$$

Based on Lemma 3.1, we know that $\|x^*, s^*\|_\infty \leq 2^L$, then we have $\|\hat{x}^{*k}, \hat{s}^{*k}\|_\infty \leq \nabla^k 2^L$.

Based on the procedure of updating the scaling factor in Algorithm 2, we can drive $\nabla^k = \mathcal{O}(\rho^k)$. We can conclude that $\omega^k = \mathcal{O}((2\rho)^L)$. \square

In Theorem 5.3, we have the total time complexity of the IR method using the proposed II-QIPM to find an exact optimal solution for LO problems.

Theorem 5.3 *Let $\hat{\zeta} = 10^{-2}$, then the total time complexity of finding an exact optimal solution using the IR-II-QIPM Algorithm 1 for solving the LO problem (1) is polynomial with*

$$\tilde{\mathcal{O}}_{n, \|\hat{A}\|, \|\hat{b}\|} \left(n^2 L \left[mn + m\sqrt{n}\kappa_{\hat{A}} (\|\hat{A}\| + \|\hat{b}\|) \right] \right),$$

the arithmetic operations, where \hat{A} and \hat{b} are preprocessed A and b .

Proof. The proof follows from combining the result of Theorem 5.2, the total time complexity of the proposed II-QIPM in (16), and Lemma 5.1. \square

Corollary 5.1 *The simplified complexity of the proposed IR-II-QIPM using the (MNES) is*

$$\tilde{\mathcal{O}}_{n, \phi} (n^4 L \phi \kappa_{\hat{A}}).$$

For finding ζ -optimal solution the complexity of IR-II-QIPM is

$$\tilde{\mathcal{O}}_{n, \phi, \omega, \frac{1}{\zeta}} (n^4 \phi \kappa_{\hat{A}}).$$

Thus, the iterative refinement procedure speeds QIPM up exponentially with respect to precision. It also addresses the condition number since the growing condition number of the Newton system is replaced by the condition number of constant matrix \hat{A} .

The following section presents the results of our numerical results.

6 Numerical Experiments

This section provides numerical results for the proposed II-QIPM using the QISKIT AQUA quantum simulator. Due to the limited number of qubits available in quantum computers and simulators, we use the NES, which has a smaller dimension. The numerical results are run on a workstation with Dual Intel Xeon® CPU E5-2630 @ 2.20 GHz (20 cores) and 64 GB of RAM. For the computational experiments, we have developed a Python `qipm` package available for public use¹.

IBM has implemented a QLSA, which is similar to the HHL method, without block-encoding and QRAM. With the current technology, the number of available qubits in gate-based quantum computers is limited to about one hundred. One of the main issues with quantum computers is that they are not scalable compared to classical computers. Currently, larger NISQ² devices suffer more from the lack of precision. On the other hand, quantum simulator algorithms are computationally expensive. The maximum number of qubits in a quantum simulator is roughly similar to that in an actual quantum computer. The main advantage of using a quantum simulator is that we do not need to handle the noise of NISQ devices. Despite this, we still need to deal with a high error level due to insufficient qubits and the high cost of QLSA+QTA in order to find high-quality solutions.

We used two post-processing procedures to improve the performance of the QLSA.

- (i) We first scale the linear system solution such that it satisfies the equality $\|Mz\| = \|\sigma\|$.
- (ii) We also check the sign of the linear system solution by comparing it with its negate.

The condition number increases the solution time. Notably, the dimension of the linear system increases the solution time as a step function for building the quantum circuit. The dimension of the system must be a power of two. The simulator expands the system size to the smallest possible power of two. Table 2 presents the number of qubits in quantum circuits for achieving the same precision. The QISKIT simulator error oscillates between zero and the norm of an actual solution. In some cases, the IBM QISKIT simulator fails, and it reports a zero vector as a solution. While the simulator has a parameter for tuning precision, it violates the predefined precision. We could not find any meaningful relationship between the precision of the simulator and the condition number of the coefficient matrix, and the dimension of the system. In the following sections, we discuss the implementation of the proposed II-QIPM and IR-II-QIPM using the QISKIT simulator of QLSA and evaluate their performance.

κ	2^1	2^2	2^3	2^4	2^5	2^6	2^7	2^8	2^9	2^{10}	2^{11}
#qubits	6	6	7	8	9	10	11	12	12	13	15

Table 2: Size of the circuit for linear systems with different condition numbers.

6.1 Evaluation of the II-QIPM

As discussed in Section 2, QLSAs have better dependence on the size of the linear system than classical algorithms. However, unreliable qubits cause a significant error in the solution of a linear system. Here, we use the IBM QISKIT simulator to solve linear systems arising in the

¹ <https://github.com/qcol-lu/qipm>

² NISQ: Noisy Intermediate Scale Quantum

proposed II-QIPM. For a fair comparison, we only consider those experiments that reach the desired precision. The running time of the quantum computers and quantum simulators is not comparable. Thus, instead of running time, we use the number of iterations as a performance measure.

We use the random instance generator of [34], where the norm of primal and dual solutions is set to two. The norms of the coefficient matrix and the RHS vector are set to one and two, respectively. The condition number of the coefficient matrix is set to two. The desired precision of the II-QIPM is equal to 0.1. It is worth noting that we set ω to 10 and all the instances are feasible.

Figure 1a illustrates II-QIPM performance on instances solved to the desired precision. As illustrated in Figure 1a, the number of iterations fluctuates with increasing the number of variables. The noisy behavior of the QLSA and the relatively small-scaled instances justifies this observation. Figure 1b shows the number of iterations increases by increasing the desired precision. The number of iterations also increases by the norm of the RHS vector, while ω has less impact on the number of iterations (see Figures 1c and 1d). As discussed in Section 4.2, the number of iterations is affected by the error of QLSAs. Scaling the RHS vector norm increases the error of QLSA and, as a result, increases the number of iterations. Furthermore, the larger ω is, the further away the starting point is from the central optimal solution, i.e., the limit point of the central path. It implies that increasing ω should increase the number of iterations. However, the fast convergence of IPMs and the noisy behavior of the QLSA make it difficult to observe a direct relationship between ω and the number of iterations.




Fig. 1: Effect of different characteristics of a LO problem on the number of iterations.

6.2 Evaluation of the IR-II-QIPM

Even with smart parameter tuning and perfect implementation, the HHL simulator has limited precision. Here, we analyze the performance of the IR method when combined with the II-QIPM. One important parameter that highly affects the norm of the RHS vector and, consequently, the HHL simulator's error is ω . As the IR progresses, the norm of a solution to the LO subproblem increases. Thus, we need to set ω based on the desired precision. Here, we set ω to 1000 to avoid infeasibility reported by the II-QIPM.

Figure 2a shows the logarithmic error of IR-II-QIPM for different desired precisions. We set the LO precision to 0.01 to evaluate the effect of the IR on the proposed II-QIPM. Figure 2b illustrates that by using IR, we can reach higher precision. The final precision in both the pure II-QIPM and the IR-II-QIPM is set to 10^{-4} . However, we can still not reach the desired precision in half of the instances because of the QLSA's error. By reducing the condition number, the iterative




Fig. 2: Obtained precision versus the desired precision on the IR-II-QIPM.

refinement method speeds up the solution time of Newton systems. The condition number of the Newton system is bounded by $\mathcal{O}(\kappa_A)$. Figure 3 shows how the condition number of the solved linear systems in IR-QIPM is bounded, while the condition number of the solved linear systems in QIPM without IR goes to infinity.




Fig. 3: The condition number of linear systems in QIPM and IR-QIPM to get 10^{-6} -precision solution for a primal-degenerate LO with 10 variables and 5 constraints

7 Conclusion

This paper analyzes in detail the benefits and challenges of novel proposed Quantum Interior Point Methods. Specifically, we analyze the use of QLSAs within IPMs and present a convergent II-QIPM. Previous papers overlooked that when one uses QLSA with QTA, the solution of the Newton system is inexact, and the Newton system's condition number goes to infinity as IPMs approach the optimal set. Here, we also adopt the IR method to find an exact solution in polynomial time. After addressing issues in earlier QIPMs, we proved the correctness and convergence of the proposed II-QIPM and analyzed its performance, both theoretically and empirically.

Algorithm	Simplified Complexity	Comment
II-IPM with Cholesky	$\mathcal{O}(n^5 L)$	
IR-II-IPM with CG	$\mathcal{O}(n^5 L \kappa_{\hat{A}})$	Anticipated complexity and needing complete analysis.
QIPM of [26]	$\tilde{\mathcal{O}}_n(n^2 L \kappa^3 2^{4L})$	Unattainable due to using exact IPM complexity.
QIPM of [13]	$\tilde{\mathcal{O}}_n(n^2 L \kappa^2 2^{2L})$	Unattainable due to using exact IPM complexity.
Proposed IR-II-QIPM	$\tilde{\mathcal{O}}_{n,\phi}(n^4 L \phi \kappa_{\hat{A}})$	

Table 3: Time complexity of finding an exact solution

Table 3 compares the complexity result of the proposed IR-II-QIPM with an analogous classical II-IPM and two recent QIPMs. The proposed IR-II-QIPM has polynomial complexity, while the other QIPMs cannot find an exact optimal solution in polynomial time. The exponential complexity of those QIPMs is caused by QLSA's error and the increasing condition number of the Newton system. At first glance, one can get the impression that the other two QIPMs have better complexity with respect to dimension. Still, these time complexities cannot be attained since they only contain the iteration complexity of exact IPMs, while these QIPMs solve the Newton system inexactly. They also need appropriate bounds for condition number κ and precision ϵ based on their setting of QIPMs. To correct the complexity of the QIPMs, at least $\mathcal{O}(n^{1.5})$ must be added for inexact Newton steps and an appropriate upper bound for QLSAs' error.

The complexity of the proposed IR-II-QIPM has better dependence on n than both its classical counterparts and the realistic complexity of its quantum counterparts. Still, the complexity of the proposed method depends on constants $\kappa_{\hat{A}}$ and $\|\hat{A}\|$. One may apply scaling and preconditioning techniques for LO problems with large $\kappa_{\hat{A}}$ to decrease $\kappa_{\hat{A}}$. Monteiro and O'Neal [35] used the speculated optimal partition instead of predefined basis (\hat{B}, \hat{N}) as a preconditioned NES. The major problem of this approach is that the cost of calculating the precondition $A_{\hat{B}}^{-1}$ in each iteration will destroy the quantum speed up in QIPMs. A viable research direction is to explore how to mitigate the effects of condition number and norm of Newton systems with a quantum-friendly approach. In addition, the proposed iterative refinement approach can also be used to mitigate the impact of the condition number in classical IPMs with CG in solving the NES. The complexity analysis of Inexact IPM augmented with an iterative refinement method that uses classical iterative solvers for solving the Newton system is part of ongoing work. However, the anticipated complexity of such algorithms, as reported in Table 3, will have unfavorable dependence on dimension than the proposed IR-II-QIPM. Because at each iteration, there are some matrix-matrix products to build the Newton system. By using QLSA+QTA, we can avoid the costs of classical matrix-matrix products and achieve polynomial quantum speedups.

We analyzed how the state-of-the-art QLSA+QTA can solve a classical linear system problem with quadratic dependence on dimension and linear dependence on the condition number and precision. Although there are classical polynomial-time algorithms to solve LSPs in gen-

eral form, QLSA-QTA is more scalable with regard to dimension than classical iterative and direct approaches. The performance of QIPMs will be improved if faster QTAs and QLSAs are proposed, and it is worth investigating polynomial-time quantum algorithms to solve LSPs.

We can also investigate Inexact-Feasible IPMs that are more adaptable with QLSAs since Feasible IPMs have better complexity than infeasible IPMs. Another direction can be developing pure QIPMs where all calculations happen in the quantum setting. However, current NISQ devices have some limitations that prevent having a pure QIPM. Such a method will circumvent the use of QTA within and enjoy the fast QLSAs.

Our computational experiments show that the proposed II-QIPM embedded in the IR scheme with the QLSA simulator of QISKIT AQUA can solve problems with hundreds of variables to a user-defined precision. However, a classical computer can simulate only a limited number of qubits, limiting the number of constraints. Although there are Feasible IPMs with better complexity than the proposed IR-II-QIPM, this paper is a significant step toward using quantum solvers in classical methods correctly and efficiently. In addition, the IR-II-QIPM leverage the wide neighborhood of the central path and starts with an infeasible interior solution compared to F-IPMs. We also demonstrated for the first time that LO problems could practically be solved using quantum solvers.

Acknowledgements This work is supported by Defense Advanced Research Projects Agency as part of the project W911NF2010022: *The Quantum Computing Revolution and Optimization: Challenges and Opportunities*.

References

- [1] Andris Ambainis. “Variable time amplitude amplification and quantum algorithms for linear algebra problems”. *STACS’12 (29th Symposium on Theoretical Aspects of Computer Science)*. Vol. 14. LIPIcs. 2012, pp. 636–647. URL: <https://bit.ly/3I1dDGR>.
- [2] Erling D. Andersen, Cornelis Roos, Tamás Terlaky, Theodore Trafalis, and Joost P. Warners. “The use of low-rank updates in interior-point methods”. *Numerical Linear Algebra and Optimization*. Science Press, Beijing, China, 2004, pp. 3–14. URL: <https://bit.ly/3WM0Ly2>.
- [3] Joran van Apeldoorn, Arjan Cornelissen, András Gilyén, and Giacomo Nannicini. “Quantum tomography using state-preparation unitaries”. *arXiv preprint* (2022). DOI: [10.48550/arXiv.2207.08800](https://doi.org/10.48550/arXiv.2207.08800).
- [4] Joran van Apeldoorn and András Gilyén. “Improvements in quantum SDP-solving with applications”. *arXiv preprint* (2018). DOI: [10.48550/arXiv.1804.05058](https://doi.org/10.48550/arXiv.1804.05058).
- [5] Joran van Apeldoorn and András Gilyén. “Quantum algorithms for zero-sum games”. *arXiv preprint* (2019). DOI: [10.48550/arXiv.1904.03180](https://doi.org/10.48550/arXiv.1904.03180).
- [6] Sanjeev Arora, Elad Hazan, and Satyen Kale. “The multiplicative weights update method: a meta-algorithm and applications”. *Theory of Computing* 8.1 (2012), pp. 121–164. DOI: [10.4086/toc.2012.v008a006](https://doi.org/10.4086/toc.2012.v008a006).
- [7] Brandon Augustino, Giacomo Nannicini, Tamás Terlaky, and Luis F Zuluaga. “Quantum interior point methods for semidefinite optimization”. *arXiv preprint* (2021). DOI: [10.48550/arXiv.2112.06025](https://doi.org/10.48550/arXiv.2112.06025).
- [8] Venansius Baryamureeba and Trond Steihaug. “On the convergence of an inexact primal-dual interior point method for linear programming”. *The Large-Scale Scientific Computing*. Ed. by Ivan Lirkov, Svetozar Margenov, and Jerzy Waśniewski. Berlin, Heidelberg: Springer Berlin Heidelberg, 2006, pp. 629–637. DOI: [10.1007/11666806_72](https://doi.org/10.1007/11666806_72).

- [9] Stefania Bellavia. “Inexact interior-point method”. *Journal of Optimization Theory and Applications* 96.1 (1998), pp. 109–121. DOI: [10.1023/A:1022663100715](https://doi.org/10.1023/A:1022663100715).
- [10] Stefania Bellavia and Sandra Pieraccini. “Convergence analysis of an inexact infeasible interior point method for semidefinite programming”. *Computational Optimization and Applications* 29.3 (2004), pp. 289–313. DOI: [10.1023/B:COAP.0000044183.43473.34](https://doi.org/10.1023/B:COAP.0000044183.43473.34).
- [11] Fernando GSL Brandao and Krysta M Svore. “Quantum speed-ups for solving semidefinite programs”. *2017 IEEE 58th Annual Symposium on Foundations of Computer Science (FOCS)*. IEEE. 2017, pp. 415–426. DOI: [10.1109/FOCS.2017.45](https://doi.org/10.1109/FOCS.2017.45).
- [12] Fernando G.S.L. Brandão, Amir Kalev, Tongyang Li, Cedric Yen-Yu Lin, Krysta M Svore, and Xiaodi Wu. “Quantum SDP solvers: Large speed-ups, optimality, and applications to quantum learning”. *arXiv preprint* (2017). DOI: [10.48550/arXiv.1710.02581](https://doi.org/10.48550/arXiv.1710.02581).
- [13] Pablo AM Casares and Miguel Angel Martin-Delgado. “A quantum interior-point predictor-corrector algorithm for linear programming”. *Journal of Physics A: Mathematical and Theoretical* 53.44 (2020), p. 445305. DOI: [10.1088/1751-8121/abb439/meta](https://doi.org/10.1088/1751-8121/abb439/meta).
- [14] Shantanav Chakraborty, András Gilyén, and Stacey Jeffery. “The power of block-encoded matrix powers: Improved regression techniques via faster Hamiltonian simulation”. *arXiv preprint* (2018). DOI: [10.48550/arXiv.1804.01973](https://doi.org/10.48550/arXiv.1804.01973).
- [15] Andrew M. Childs, Robin Kothari, and Rolando D. Somma. “Quantum algorithm for systems of linear equations with exponentially improved dependence on precision”. *SIAM Journal on Computing* 46.6 (2017), pp. 1920–1950. DOI: [10.1137/16m1087072](https://doi.org/10.1137/16m1087072).
- [16] George B. Dantzig. *Linear Programming and Extensions*. Princeton, NJ, USA: Princeton University Press, 1963. DOI: [10.1515/9781400884179](https://doi.org/10.1515/9781400884179).
- [17] Danial Dervovic, Mark Herbster, Peter Mountney, Simone Severini, Nairi Usher, and Leonard Wossnig. “Quantum linear systems algorithms: a primer”. *arXiv preprint* (2018). DOI: [10.48550/arXiv.1802.08227](https://doi.org/10.48550/arXiv.1802.08227).
- [18] David Deutsch. “Quantum theory, the Church-Turing principle and the universal quantum computer”. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences* 400.1818 (1985), pp. 97–117. DOI: [10.1098/rspa.1985.0070](https://doi.org/10.1098/rspa.1985.0070).
- [19] David Deutsch and Richard Jozsa. “Rapid solution of problems by quantum computation”. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences* 439.1907 (1992), pp. 553–558. DOI: [10.1098/rspa.1992.0167](https://doi.org/10.1098/rspa.1992.0167).
- [20] Roland W. Freund, Florian Jarre, and Shinji Mizuno. “Convergence of a class of inexact interior-point algorithms for linear programs”. *Mathematics of Operations Research* 24.1 (1999), pp. 50–71. DOI: [10.1287/moor.24.1.50](https://doi.org/10.1287/moor.24.1.50).
- [21] Ambros Gleixner and Daniel E. Steffy. “Linear programming using limited-precision oracles”. *Mathematical Programming* 183.1 (2020), pp. 525–554. DOI: [10.1007/s10107-019-01444-6](https://doi.org/10.1007/s10107-019-01444-6).
- [22] Ambros M. Gleixner, Daniel E. Steffy, and Kati Wolter. “Iterative refinement for linear programming”. *INFORMS Journal on Computing* 28.3 (2016), pp. 449–464. DOI: [10.1287/ijoc.2016.0692](https://doi.org/10.1287/ijoc.2016.0692).
- [23] Aram W. Harrow, Avinatan Hassidim, and Seth Lloyd. “Quantum algorithm for linear systems of equations”. *Physical Review Letters* 103.15 (2009). DOI: [10.1103/physrevlett.103.150502](https://doi.org/10.1103/physrevlett.103.150502).
- [24] Ghussoun Al-Jeiroudi and Jacek Gondzio. “Convergence analysis of the inexact infeasible interior-point method for linear optimization”. *Journal of Optimization Theory and Applications* 141.2 (2009), pp. 231–247. DOI: [10.1007/s10957-008-9500-5](https://doi.org/10.1007/s10957-008-9500-5).
- [25] Narendra Karmarkar. “A new polynomial-time algorithm for linear programming”. *Proceedings of the Sixteenth Annual ACM Symposium on Theory of Computing*. STOC ’84.

- New York, NY, USA: Association for Computing Machinery, 1984, pp. 302–311. DOI: [10.1145/800057.808695](https://doi.org/10.1145/800057.808695).
- [26] Iordanis Kerenidis and Anupam Prakash. “A quantum interior point method for LPs and SDPs”. *ACM Transactions on Quantum Computing* 1.1 (2020), pp. 1–32. DOI: [10.1145/3406306](https://doi.org/10.1145/3406306).
- [27] Iordanis Kerenidis and Anupam Prakash. “Quantum recommendation systems”. *arXiv preprint* (2016). DOI: [10.48550/arXiv.1603.08675](https://doi.org/10.48550/arXiv.1603.08675).
- [28] Iordanis Kerenidis, Anupam Prakash, and Dániel Szilágyi. “Quantum algorithms for second-order cone programming and support vector machines”. *Quantum* 5 (2021), p. 427. DOI: [10.22331/q-2021-04-08-427](https://doi.org/10.22331/q-2021-04-08-427).
- [29] Leonid Genrikhovich Khachiyan. “A polynomial algorithm in linear programming”. *Doklady Akademii Nauk*. Vol. 244. 5. Reports of the Russian Academy of Sciences. 1979, pp. 1093–1096. URL: <http://mi.mathnet.ru/eng/dan42319>.
- [30] Masakazu Kojima, Nimrod Megiddo, and Shinji Mizuno. “A primal—dual infeasible-interior-point algorithm for linear programming”. *Mathematical Programming* 61.1 (1993), pp. 263–280. DOI: [10.1007/BF01582151](https://doi.org/10.1007/BF01582151).
- [31] János Korzák. “Convergence analysis of inexact infeasible-interior-point algorithms for solving linear programming problems”. *SIAM Journal on Optimization* 11.1 (2000), pp. 133–148. DOI: [10.1137/S1052623497329993](https://doi.org/10.1137/S1052623497329993).
- [32] Guang Hao Low and Isaac L Chuang. “Hamiltonian simulation by qubitization”. *Quantum* 3 (2019), p. 163. DOI: [10.22331/q-2019-07-12-163](https://doi.org/10.22331/q-2019-07-12-163).
- [33] Shinji Mizuno and Florian Jarre. “Global and polynomial-time convergence of an infeasible-interior-point algorithm using inexact computation”. *Mathematical Programming* 84.1 (1999), pp. 105–122. DOI: [10.1007/s10107980020a](https://doi.org/10.1007/s10107980020a).
- [34] Mohammadhossein Mohammadisiahroudi, Ramin Fakhimi, Brandon Augustino, and Tamás Terlaky. “Generating Linear, Semidefinite, and Second-order Cone Optimization Problems for Numerical Experiments”. *arXiv preprint arXiv:2302.00711* (2023).
- [35] Renato Monteiro and Jerome W. O’Neal. “Convergence analysis of a long-step primal-dual infeasible interior-point LP algorithm based on iterative linear solvers”. *Georgia Institute of Technology* (2003). URL: <https://bit.ly/3ss0kg1>.
- [36] Cornelis Roos, Tamás Terlaky, and Jean-Philippe Vial. *Interior Point Methods for Linear Optimization*. Springer Science & Business Media, 2005. DOI: [10.1007/b100325](https://doi.org/10.1007/b100325).
- [37] Yousef Saad. *Iterative Methods for Sparse Linear Systems*. SIAM, 2003. DOI: [10.1137/1.9780898718003](https://doi.org/10.1137/1.9780898718003).
- [38] Jack Sherman and Winifred J Morrison. “Adjustment of an inverse matrix corresponding to a change in one element of a given matrix”. *The Annals of Mathematical Statistics* 21.1 (1950), pp. 124–127. URL: <https://bit.ly/3jJFwy6>.
- [39] Daniel R. Simon. “On the power of quantum computation”. *SIAM Journal on Computing* 26.5 (1997), pp. 1474–1483. DOI: [10.1137/s0097539796298637](https://doi.org/10.1137/s0097539796298637).
- [40] Tamás Terlaky. *Interior Point Methods of Mathematical Programming*. Vol. 5. Applied Optimization Kluwer Academic Publisher, 1996. DOI: [10.1007/978-1-4613-3449-1](https://doi.org/10.1007/978-1-4613-3449-1).
- [41] Almudena Carrera Vazquez, Ralf Hiptmair, and Stefan Woerner. “Enhancing the quantum linear systems algorithm using Richardson extrapolation”. *ACM Transactions on Quantum Computing* 3.1 (2022), pp. 1–37. DOI: [10.1145/3490631](https://doi.org/10.1145/3490631).
- [42] Leonard Wossnig, Zhikuan Zhao, and Anupam Prakash. “Quantum linear system algorithm for dense matrices”. *Physical Review Letters* 120.5 (2018). DOI: [10.1103/physrevlett.120.050502](https://doi.org/10.1103/physrevlett.120.050502).
- [43] Stephen J. Wright. *Primal-Dual Interior-Point Methods*. SIAM, 1997. DOI: [10.1137/1.9781611971453](https://doi.org/10.1137/1.9781611971453).

-
- [44] Yinyu Ye. *Interior Point Algorithms: Theory and Analysis*. Vol. 44. John Wiley & Sons, 2011. URL: <https://bit.ly/3i9zt1U>.
- [45] Yinyu Ye, Michael J. Todd, and Shinji Mizuno. “An $O(\sqrt{n}L)$ -iteration homogeneous and self-dual linear programming algorithm”. *Mathematics of Operations Research* 19.1 (1994), pp. 53–67. DOI: [10.1287/moor.19.1.53](https://doi.org/10.1287/moor.19.1.53).
- [46] Guanglu Zhou and Kim-Chuan Toh. “Polynomiality of an inexact infeasible interior point algorithm for semidefinite programming”. *Mathematical Programming* 99.2 (2004), pp. 261–282. DOI: [10.1007/s10107-003-0431-5](https://doi.org/10.1007/s10107-003-0431-5).