

# Faces of homogeneous cones and applications to homogeneous chordality

João Gouveia\*

Masaru Ito<sup>†</sup>

Bruno F. Lourenço<sup>‡</sup>

December 8, 2024

## Abstract

A convex cone  $\mathcal{K}$  is said to be homogeneous if its group of automorphisms acts transitively on its relative interior. Important examples of homogeneous cones include symmetric cones and cones of positive semidefinite (PSD) matrices that follow a sparsity pattern given by a homogeneous chordal graph. Our goal in this paper is to elucidate the facial structure of homogeneous cones and make it as transparent as the faces of the PSD matrices. We prove that each face of a homogeneous cone  $\mathcal{K}$  is mapped by an automorphism of  $\mathcal{K}$  to one of its finitely many so-called *principal faces*. Furthermore, constructing such an automorphism can be done algorithmically by making use of a generalized Cholesky decomposition. Among other consequences, we give a proof that homogeneous cones are projectionally exposed, which strengthens the previous best result that they are amenable. Using our results, we will carefully analyze the facial structure of cones of PSD matrices satisfying homogeneous chordality and discuss consequences for the corresponding family of PSD completion problems.

**Keywords:** Homogeneous cones, facial structure, projectional exposedness, homogeneous chordal graphs, PSD completion

## 1 Introduction

A convex cone is said to be *homogeneous* if its group of automorphisms acts transitively on its relative interior. Conic programming over homogeneous cones is a corner stone of the modern optimization landscape as it contains as special cases semidefinite programming (i.e., optimization over positive semidefinite matrices) and linear programming (i.e., optimization over the nonnegative orthant). However, the positive semidefinite matrices and nonnegative orthants are only merely two very special examples of homogeneous cones, as they are also self-dual under an appropriate choice of inner product and, therefore, are *symmetric cones*.

An important class of homogeneous but not necessarily symmetric cones include the positive semidefinite matrices with certain sparsity patterns. Let  $G$  be a graph on  $n$  vertices and consider

---

\*CMUC, Department of Mathematics, University of Coimbra, 3001-454 Coimbra, Portugal. This author was supported by the Centre for Mathematics of the University of Coimbra (UIDB/00324/2020, funded by the Portuguese Government through FCT/MCTES). During part of this work, the author also held a visiting professorship at the Institute of Statistical Mathematics, Japan. Email: [jgouveia@mat.uc.pt](mailto:jgouveia@mat.uc.pt).

<sup>†</sup>Department of Mathematics, College of Science and Technology, Nihon University, 1-8-14 Kanda-Surugadai, Chiyoda-Ku, Tokyo 101-8308, Japan. This author was supported partly by the JSPS Grant-in-Aid for Early-Career Scientists 21K17711. Email: [ito.masaru@nihon-u.ac.jp](mailto:ito.masaru@nihon-u.ac.jp).

<sup>‡</sup>Department of Fundamental Statistical Mathematics, Institute of Statistical Mathematics, Japan. This author was supported partly by the JSPS Grant-in-Aid for Early-Career Scientists 23K16844. Email: [bruno@ism.ac.jp](mailto:bruno@ism.ac.jp).

the cone  $\mathcal{S}_+(G)$  of  $n \times n$  symmetric positive semidefinite (PSD) matrices whose sparsity pattern follows  $G$ , i.e.,  $x \in \mathcal{S}_+(G)$  if and only if  $x$  is an  $n \times n$  PSD matrix with  $x_{ij} = 0$  for all  $i \neq j$  such that  $(i, j)$  is not an edge of  $G$ . The case where  $G$  is a chordal graph has been extensively studied in the literature as it is typically more efficient to solve problems exploiting chordal sparsity instead of seeing it simply as a SDP instance, e.g., [FKMN01, NFF<sup>+</sup>03]. The dual of  $\mathcal{S}_+(G)$  is also an important object, since it corresponds to the PSD completable matrices with pattern given by  $G$ .

If  $G$  is not only chordal but has no induced subgraph that is a path on four vertices, then  $\mathcal{S}_+(G)$  becomes a homogeneous cone [Ish13] and such graphs were called *homogeneous chordal* in [TV23]. PSD matrices that follow a homogeneous chordal sparsity pattern have remarkable properties that may fail in general for chordal graphs. A notable example is that, under an appropriate ordering of the vertices, inverses of Cholesky factors still respect the sparsity pattern defined by  $G$ , see [TV23, Theorem 3.1] for more details. Therefore, for homogeneous chordal graphs,  $\mathcal{S}_+(G)$  and its dual form important classes of homogeneous cones that have practical relevance and potential for interesting applications.

Motivated by the particular case of homogeneous chordality, our goal in this paper is to elucidate the facial structure of general homogeneous cones and make it as clear as the facial structure of PSD matrices, which we will now briefly review. Let  $\mathcal{S}_+^n$  denote the set of real  $n \times n$  symmetric positive semidefinite matrices. If  $\mathcal{F}$  is a face of  $\mathcal{S}_+^n$ , then there exists a  $n \times n$  orthogonal matrix  $q$  and  $r \leq n$  such that

$$q\mathcal{F}q^* = \left\{ \begin{pmatrix} a & 0 \\ 0 & 0 \end{pmatrix} \mid a \in \mathcal{S}_+^r \right\}, \quad (1.1)$$

where  $q^*$  denotes the transpose of  $q$ , see [BC75, Section 6]. That is, each face of  $\mathcal{S}_+^n$  is linearly isomorphic to a smaller PSD cone and this isomorphism can be realized as an automorphism of  $\mathcal{S}_+^n$  since the map  $Q$  that takes  $x$  to  $qxq^*$  satisfies  $Q(\mathcal{S}_+^n) = \mathcal{S}_+^n$ . The matrix  $q$  does not come from thin air: the kernel of matrices in the relative interior of  $\mathcal{F}$  is unique and correspond to some fixed subspace  $V \subseteq \mathbb{R}^n$ . With that,  $\mathcal{F} = \{x \in \mathcal{S}_+^n \mid \ker x \supseteq V\}$  holds. Therefore, computing  $q$  is an entirely constructive endeavour, as it is enough to pick any  $x \in \text{ri } \mathcal{F}$  and let  $q$  be such that its rows are orthogonal and the last  $n - r$  rows form an orthonormal basis for  $\ker x$ .

Another interesting property is that if we let  $v := q^* \begin{pmatrix} e_r & 0 \\ 0 & 0 \end{pmatrix} q$ , where  $e_r$  is the  $r \times r$  identity matrix and define the map  $\mathcal{P}(x) := vxv$ , we have that  $\mathcal{P}$  is a projection (i.e.,  $\mathcal{P}^2 = \mathcal{P}$ ) satisfying  $\mathcal{P}(\mathcal{S}_+^n) = \mathcal{F}$ . In this way, not only the faces of  $\mathcal{S}_+^n$  are isomorphic to smaller PSD cones, they are also projected versions of it. Cones for which each face arises as a projection of the original cone are known as *projectionally exposed* and were first described by Borwein and Wolkowicz in the context of their facial reduction approach in [BW81]. One of the main goals of this paper is to prove analogous results to (1.1) for homogeneous cones and discuss their consequences.

Another motivation for this work comes from the study of facial exposedness in general cones. We recall that a closed convex cone  $\mathcal{K}$  is said to be *facially exposed* if every face arises as the intersection of  $\mathcal{K}$  with one of its supporting hyperplanes. Although facial exposedness is useful, certain applications require stronger facial exposedness properties such as *niceness* (also known as facial dual completeness) [Pat07, Pat13a, Pat13b], *amenability* [Lou21, LRS22] or the aforementioned projectionally exposedness [BW81, ST90].

Every projectionally exposed (p-exposed) cone is amenable, every amenable cone is nice and every nice cone is facially exposed. In dimension at most three, facial exposedness implies p-exposedness, see [PL88, Theorem 3.2] or [ST90, Theorem 4.6]. In dimension four, there exists a facially exposed cone that is not amenable [Ros14] and a nice cone that is not amenable [LRS22].

Regarding homogeneous cones, Truong and Tunçel showed that they are facially exposed [TT03,

Theorem 6]. Later, Chua and Tunçel showed that homogeneous cones are nice [CT06, Proposition 4 and Section 4.2]. Then, in [LRS22] it was shown that homogeneous cones are amenable. In this work, we take another step and show that homogeneous cones are, in fact, p-exposed. In this way, homogeneous cones form a large family of p-exposed cones and it is not currently known if there are any other interesting classes of p-exposed cones strictly containing homogeneous cones. Notably, homogeneous cones are also hyperbolicity cones [Gül97, Section 8], which is another important class of cones [Går59, Ren06]. However, the strongest result so far is that hyperbolicity cones are amenable [LRS23] and it is not known whether they are p-exposed in general.

Our main results are as follows.

- (i) We will show that results analogous to (1.1) hold for general homogeneous cones, see Theorem 3.5. In particular, given a homogeneous cone  $\mathcal{K}$  of rank  $r$  (see Section 2.1), there are  $2^r$  distinguished faces called *principal faces* that play a role similar to the right-hand-side of (1.1) in the sense that every face  $\mathcal{F}$  of  $\mathcal{K}$  can be mapped to a principal face via an automorphism of  $\mathcal{K}$ . Similarly to the case of PSD matrices, the automorphism can be explicitly constructed given any  $x \in \text{ri } \mathcal{F}$  by making use of a generalized Cholesky factorization (Algorithm 1).  
This will allow us to easily obtain certain results that are intuitive but are not trivial to extract from the existing literature. For example, Theorem 3.5 contains as a particular consequence the fact that proper faces of homogeneous cones are also homogeneous cone themselves and must have strictly smaller rank, which, as far as we know, has never been formally stated or verified previously. One application of this is the computation of the length of a longest chain of faces of homogeneous cones, see Corollary 3.7. We will also show that conjugate faces have complementary ranks, see Proposition 3.10.
- (ii) Among the consequences of our discussion, of particular note is the fact that homogeneous cones are projectionally exposed, which strengthens previous results in [TT03, CT06, LRS22].
- (iii) We discuss in the detail the facial structure of PSD matrices with sparsity pattern given by a homogeneous chordal graph  $G$ , see Section 4. In particular we show that every face of  $\mathcal{S}_+(G)$  is isomorphic to some face that arises by considering induced subgraphs of  $G$  and this isomorphism can be realized by an automorphism of  $\mathcal{S}_+(G)$ , see Theorem 4.2. We will also prove analogous results to the dual cone of  $\mathcal{S}_+(G)$  which corresponds to the PSD completable matrices determined by  $G$ . Then, we will check that when Algorithm 1 is specialized to the dual of  $\mathcal{S}_+(G)$ , it will allow us to compute PSD completions having certain desirable properties, see Theorem 4.3.

This work is divided as follows. In Section 2, we recall some properties of convex sets and its faces and then we discuss the T-algebraic framework of Vinberg for handling homogeneous cones. Section 3 contains the main results of this paper regarding the facial structure of homogeneous cones. In Section 4, we discuss several applications of the results in Section 3 to the facial structure of  $\mathcal{S}_+(G)$  and its dual. We conclude this work in Section 5 with a list of open questions and some remarks on how our discussion can be useful in facial reduction approaches.

## 2 Preliminaries

Let  $\mathcal{E}$  be a finite dimensional Euclidean space equipped with an inner product  $\langle \cdot, \cdot \rangle$  and corresponding induced norm  $\|\cdot\|$ . Let  $C \subseteq \mathcal{E}$  be a convex set. We denote its relative interior and closure by  $\text{ri } C$  and  $\text{cl } C$ , respectively. The smallest subspace of  $\mathcal{E}$  containing  $C$  is denoted by  $\text{span } C$ .

Let  $\mathcal{F} \subseteq C$  be convex. Then,  $\mathcal{F}$  is said to be a *face* of  $C$  if  $x, y \in C$  and  $\alpha x + (1 - \alpha)y \in \mathcal{F}$  for some  $\alpha \in (0, 1)$  implies that  $x, y \in \mathcal{F}$ . In this case, we write  $\mathcal{F} \trianglelefteq C$  and, by convention, we will only consider non-empty faces. In the particular case where  $C$  is a convex cone, we have that a nonempty convex cone  $\mathcal{F} \subseteq C$  is a face if and only if  $x, y \in C$  and  $x + y \in \mathcal{F}$  implies  $x, y \in \mathcal{F}$ . A face  $\mathcal{F}$  is said to be *proper* if  $\mathcal{F} \neq C$  and it is *maximally proper* if there is no proper face  $\hat{\mathcal{F}} \trianglelefteq C$  such that  $\mathcal{F} \subseteq \hat{\mathcal{F}}$  and  $\mathcal{F} \neq \hat{\mathcal{F}}$ .

Let  $\mathcal{K} \subseteq \mathcal{E}$  be a closed convex cone. We say that  $\mathcal{K}$  is *pointed* if  $\mathcal{K} \cap -\mathcal{K} = \{0\}$  holds. Two convex cones  $\mathcal{K}_1, \mathcal{K}_2$  are said to be *linearly isomorphic* if there exists a linear bijection  $A$  such that  $A(\mathcal{K}_1) = \mathcal{K}_2$  holds.

Next, let  $x \in \mathcal{K}$ . We denote by  $\text{Face}(x, \mathcal{K})$  the minimal face of  $\mathcal{K}$  containing  $x$ , i.e.,  $\text{Face}(x, \mathcal{K}) = \bigcap_{\mathcal{F} \trianglelefteq \mathcal{K}, x \in \mathcal{F}} \mathcal{F}$ . Let  $\mathcal{F} \trianglelefteq \mathcal{K}$  be a face and  $x \in \mathcal{K}$ , then

$$\mathcal{F} = \text{Face}(x, \mathcal{K}) \Leftrightarrow x \in \text{ri } \mathcal{F}, \quad (2.1)$$

e.g., see [Pat00, Proposition 3.2.2] or (2.1) can also be inferred by the discussion in [Bar73].

An *extreme ray* of  $\mathcal{K}$  is a face of dimension 1, i.e., it is of the form  $\mathcal{F} = \{\alpha x \mid \alpha \geq 0\}$  for some nonzero  $x \in \mathcal{K}$ . In this case, we say that  $x$  *generates an extreme ray* of  $\mathcal{K}$ .

The dual cone of  $\mathcal{K}$  is denoted by  $\mathcal{K}^* := \{y \in \mathcal{E} \mid \langle x, y \rangle \geq 0, \forall x \in \mathcal{K}\}$ . A face  $\mathcal{F} \trianglelefteq \mathcal{K}$  is said to be *exposed* if there exists  $y \in \mathcal{K}^*$  such that  $\mathcal{F} = \mathcal{K} \cap \{y\}^\perp$  holds. In this case, we say that  $y$  *exposes the face*  $\mathcal{F}$ . The conjugate face of  $\mathcal{F}$  is defined as  $\mathcal{F}^\Delta := \mathcal{K}^* \cap \mathcal{F}^\perp$  and satisfies

$$\mathcal{F}^\Delta = \mathcal{K}^* \cap \{x\}^\perp, \quad \forall x \in \text{ri } \mathcal{F}, \quad (2.2)$$

which follows from [Roc97, Theorem 6.4]. In view of (2.1) and (2.2), if  $\mathcal{F}$  is exposed by  $y$ , then  $\mathcal{F}$  is precisely the face that is conjugated to  $\text{Face}(y, \mathcal{K}^*)$ . We will also need a well-known lemma that connects extreme rays and maximal proper faces via conjugacy.

**Lemma 2.1.** *If  $\mathcal{F} \trianglelefteq \mathcal{K}$  is an exposed extreme ray, then  $\mathcal{F}^\Delta$  is a maximal proper face of  $\mathcal{K}^*$ .*

*Proof.* In this proof we need the well-known fact that a face is facially exposed if and only if  $\mathcal{F}^{\Delta\Delta} = \mathcal{F}$  holds. In addition, all maximal proper faces are facially exposed, e.g., see [Tam85, Corollary 2.2]. With these facts in mind, suppose that  $\hat{\mathcal{F}} \trianglelefteq \mathcal{K}^*$  is a maximally proper face containing  $\mathcal{F}^\Delta$ . As conjugacy inverts inclusion, we have  $\hat{\mathcal{F}}^\Delta \trianglelefteq \mathcal{F}^{\Delta\Delta} = \mathcal{F}$ , so  $\hat{\mathcal{F}}^\Delta = \{0\}$  or  $\hat{\mathcal{F}}^\Delta = \mathcal{F}$ . In the former case we have  $\hat{\mathcal{F}} = \hat{\mathcal{F}}^{\Delta\Delta} = \mathcal{K}^*$ , which contradicts the properness of  $\hat{\mathcal{F}}$ . So, we must be in the latter case and  $\hat{\mathcal{F}} = \hat{\mathcal{F}}^{\Delta\Delta} = \mathcal{F}^\Delta$  holds.  $\square$

A face  $\mathcal{F} \trianglelefteq \mathcal{K}$  is said to be *projectionally exposed* (or p-exposed) if there exists a linear map  $\mathcal{P} : \mathcal{E} \rightarrow \mathcal{E}$  such that  $\mathcal{P}(\mathcal{K}) = \mathcal{F}$  and  $\mathcal{P}^2 = \mathcal{P}$ . If all faces of  $\mathcal{K}$  are p-exposed, then  $\mathcal{K}$  is said to be p-exposed. P-exposedness was proposed in [BW81] and a comprehensive discussion on p-exposedness is given in [ST90].

If the projection  $\mathcal{P}$  can be taken to be self-adjoint with respect to  $\langle \cdot, \cdot \rangle$ , then  $\mathcal{F}$  is said to be *orthogonally projectionally exposed* (or o.p.-exposed). Clearly, this is a notion that depends on the choice of inner product, so to emphasize this choice, we will sometimes say that  $\mathcal{F}$  is *o.p.-exposed under*  $\langle \cdot, \cdot \rangle$ . If all faces of  $\mathcal{K}$  are o.p.-exposed with respect the same inner product, then  $\mathcal{K}$  is said to be o.p.-exposed. Being o.p.-exposed is a more restrictive property and, for example, an o.p.-exposed polyhedral cone must be linearly isomorphic to a nonnegative orthant, see [ST90, Theorem 3.7 and pg. 233] and [BLP87]. That said, all symmetric cones are o.p.-exposed under an appropriate inner product [Lou21, Proposition 33].

## 2.1 Homogeneous cones and T-algebras

Let  $\mathcal{K} \subseteq \mathcal{E}$  be a convex cone. The automorphisms of  $\mathcal{K}$ , denoted by  $\text{Aut}(\mathcal{K})$ , are the group of linear bijections  $Q : \mathcal{E} \rightarrow \mathcal{E}$  satisfying  $Q(\mathcal{K}) = \mathcal{K}$ , where the group operation is the function composition. We say that  $\mathcal{K}$  is *homogeneous* if  $\text{Aut}(\mathcal{K})$  acts transitively on the relative interior of  $\mathcal{K}$ , i.e., for every  $x, y \in \text{ri } \mathcal{K}$ , there must exist  $Q \in \text{Aut}(\mathcal{K})$  satisfying  $Q(x) = y$ .

*Symmetric cones* are homogeneous cones that are self-dual under some inner product and the theory of Euclidean Jordan algebras [Koe99, FK94] seems to be the standard algebraic framework to deal with them. For homogeneous cones, there are different, albeit closely related, algebraic frameworks described in many foundational works [Vin63, Rot66, Gin92]. Brief overviews of this state of affairs can be seen in [YN15, Section 1] and [TV23, Section 6].

Our goal in this paper is to describe the facial structure of homogeneous cones, which as far as we know, has not been described explicitly before. Nevertheless, there are previous works describing polynomials that are closely related to the boundary structure of homogeneous cones [Ish01, Nak14, Nak18, GIL24, Nak24]. A previous work by Ishi also contains important information on the action of the automorphism group on the boundary of the cone, see [Ish00]. We will revisit this point later in Section 3. In optimization, the study of homogeneous cones seems to have been initially motivated by self-concordant barriers and interior-point methods [G96, GT98, Chu09], with other works focusing on geometric and representational aspects [TX01, Chu03, TT03, CT06].

In this work, we will use the theory of T-algebras of Vinberg [Vin63] as it seems to be the most natural for our purpose. Unfortunately, as of this writing, although the original russian text of [Vin63] is freely available on the internet, it is not completely trivial to get a copy of the English translation. Because of that, as we review T-algebras, we refer extensively to more accessible references in English such as [Chu03, CT06, Chu09, KTX12].

To start, we say that a *matrix algebra of rank  $r$*  is an algebra  $\mathcal{A}$  over  $\mathbb{R}$  equipped with a bigradation, i.e., a decomposition as a direct sum  $\mathcal{A} = \bigoplus_{i,j=1}^r \mathcal{A}_{ij}$  where the  $\mathcal{A}_{ij}$  are subspaces satisfying the following properties:

$$\begin{aligned} \mathcal{A}_{ij}\mathcal{A}_{jk} &\subseteq \mathcal{A}_{ik} \\ \mathcal{A}_{ij}\mathcal{A}_{kl} &= \{0\} \quad \text{if } j \neq k. \end{aligned} \tag{2.3}$$

Given  $a \in \mathcal{A}$ , we can write  $a = \sum_{i,j=1}^r a_{ij}$  in a unique way with  $a_{ij} \in \mathcal{A}_{ij}$ . We will refer to  $a_{ij}$  as the  $(i, j)$  component of  $a$ . Multiplication in  $\mathcal{A}$  is analogous to the usual matrix multiplication since (2.3) implies that

$$(ab)_{ij} = \sum_{k=1}^r a_{ik}b_{kj} \tag{2.4}$$

holds for  $a, b \in \mathcal{A}$ . A *matrix algebra with involution* is a matrix algebra equipped with a linear bijection  $*$  :  $\mathcal{A} \rightarrow \mathcal{A}$  such that for

$$a^{**} = a, (ab)^* = b^*a^* \text{ and } \mathcal{A}_{ij}^* = \mathcal{A}_{ji}, \quad \text{for all } i, j \in \{1, \dots, r\}. \tag{2.5}$$

With that, we have  $(a^*)_{ij} = a_{ji}^*$ , where “ $a_{ji}^*$ ” should be read as  $(a_{ji})^*$ .

Then, a *T-algebra of rank  $r$*  is a matrix algebra of rank  $r$  with involution such that the following additional axioms are satisfied.

(a1) For each  $i$ ,  $\mathcal{A}_{ii}$  is a subalgebra isomorphic to  $\mathbb{R}$ .

Let  $\rho_i : \mathcal{A}_{ii} \rightarrow \mathbb{R}$  denote the algebra isomorphism and let  $e_i$  denote the unit element in  $\mathcal{A}_{ii}$ , i.e., the element satisfying  $\rho_i(e_i) = 1$ . Furthermore, define the function  $\text{tr} : \mathcal{A} \rightarrow \mathbb{R}$  by  $\text{tr}(a) := \sum_{i=1}^r \rho_i(a_{ii})$ .

(a2) For all  $a \in \mathcal{A}$  and all  $i, j \in \{1, \dots, r\}$  we have  $e_i a_{ij} = a_{ij}$  and  $a_{ji} e_i = a_{ji}$ .

(a3) For all  $a, b \in \mathcal{A}$ ,  $\text{tr}(ab) = \text{tr}(ba)$ .

(a4) For all  $a, b, c \in \mathcal{A}$  and we have  $\text{tr}((ab)c) = \text{tr}(a(bc))$ .

(a5) For all  $a \in \mathcal{A}$ , we have  $\text{tr}(aa^*) \geq 0$  with equality if and only if  $a = 0$ .

(a6) For all  $a, b, c \in \mathcal{A}$  and  $1 \leq i \leq j \leq k \leq l \leq r$ , we have  $a_{ij}(b_{jk}c_{kl}) = (a_{ij}b_{jk})c_{kl}$ .

(a7) For all  $a, b \in \mathcal{A}$ ,  $1 \leq i \leq j \leq k \leq r$  and  $1 \leq l \leq k \leq r$ , we have  $a_{ij}(b_{jk}b_{lk}^*) = (a_{ij}b_{jk})b_{lk}^*$ .

We note that these axioms are sometimes stated in different but equivalent forms, see [Chu09, Remarks 2-7] for some equivalences.

Defining  $e := e_1 + \dots + e_r$ , Axiom (a2) implies that  $ea = a$  and  $ae = a$  holds for all  $a \in \mathcal{A}$ , so  $e$  plays the role of identity element.

Given a T-algebra  $\mathcal{A}$ , the bilinear function

$$\langle a, b \rangle := \text{tr}(ab^*) = \sum_{1 \leq i, j \leq r} \rho_i(a_{ij}b_{ij}^*) \quad (2.6)$$

is an inner product on  $\mathcal{A}$  due to Axiom (a5), which induces the norm  $\|a\| := \sqrt{\langle a, a \rangle}$  over  $\mathcal{A}$ . The subspaces  $\mathcal{A}_{ij}$  are orthogonal to each other under this inner product by (2.3) and Axiom (a3).

**Remark 2.2** (Analogy with the usual matrices). *The space  $M^{n \times n}$  of real  $n \times n$  matrices can be seen as a T-algebra in a natural way, however, an important difference is that for  $a \in M^{n \times n}$ ,  $a_{ij}$  is not a scalar, but a matrix as well. For example, for  $a := \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$ , we have the following decomposition*

$$a = a_{11} + a_{12} + a_{21} + a_{22} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 2 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ 3 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ 0 & 4 \end{pmatrix}.$$

Now, we define certain subsets of  $\mathcal{A}$ . The subspace of ‘‘Hermitian’’ matrices of  $\mathcal{A}$  is given by

$$\mathcal{H} := \{a \in \mathcal{A} \mid a^* = a\}.$$

We also define sets of upper triangular submatrices, so that

$$\begin{aligned} \mathcal{T} &:= \bigoplus_{1 \leq i \leq j \leq r} \mathcal{A}_{ij}, \\ \mathcal{T}_+ &:= \{a \in \mathcal{T} \mid \rho_i(a_{ii}) \geq 0 \text{ if } 1 \leq i \leq r\}, \\ \mathcal{T}_{++} &:= \{a \in \mathcal{T} \mid \rho_i(a_{ii}) > 0 \text{ if } 1 \leq i \leq r\}. \end{aligned}$$

When it is necessary to emphasize the underlying algebra, we will alternatively write  $\mathcal{H}(\mathcal{A})$ ,  $\mathcal{T}(\mathcal{A})$ ,  $\mathcal{T}_+(\mathcal{A})$  or  $\mathcal{T}_{++}(\mathcal{A})$ . Finally, the homogeneous cone in  $\mathcal{H}$  associated to the T-algebra  $\mathcal{A}$  is defined as

$$\mathcal{K}(\mathcal{A}) := \{tt^* \mid t \in \mathcal{T}_{++}\}.$$

and its closure is given by

$$\text{cl } \mathcal{K}(\mathcal{A}) = \{tt^* \mid t \in \mathcal{T}_+\}.$$

Each element in  $\mathcal{K}(\mathcal{A})$  has a unique representation as  $tt^*$  for  $t \in \mathcal{T}_{++}$ , see [Chu09, Proposition 4] or [Vin63, Chapter III, Proposition 2]. This is analogous to the fact that a positive definite symmetric



matrix has a unique Cholesky factorization in terms of upper triangular matrices. This uniqueness breaks down for elements of  $\text{cl}\mathcal{K}(\mathcal{A})$  and, in general,  $tt^* = uu^*$  for  $t, u \in \mathcal{T}_+$  does not imply  $u = t$ . Nevertheless, we will discuss how to recover uniqueness in Section 3.1.

An important result by Vinberg establishes the correspondence between homogeneous cones and T-algebras as follows.

**Theorem 2.3** ([Vin63, Chapter III, §2 and Theorem 4]). *Let  $\mathcal{K}$  be a pointed open homogeneous convex cone contained in a real finite dimensional space. Then, there is a T-algebra  $\mathcal{A}$  for which  $\mathcal{K}(\mathcal{A}) = \mathcal{K}$ . Conversely, if  $\mathcal{K}(\mathcal{A}) = \mathcal{K}$  for some T-algebra  $\mathcal{A}$ , then  $\mathcal{K}$  is an open homogeneous convex cone.*

**Dual cone** Let  $\mathcal{B}$  be the matrix algebra

$$\mathcal{B} := \bigoplus_{i,j=1}^r \mathcal{B}_{ij}, \quad \text{where } \mathcal{B}_{ij} := \mathcal{A}_{r+1-i, r+1-j}, \quad (2.7)$$

endowed with the same involution as  $\mathcal{A}$ , which forms a T-algebra of rank  $r$ . Then the following identity holds [Vin63, p. 390].

$$\mathcal{K}(\mathcal{A})^* = \text{cl}\mathcal{K}(\mathcal{B}) = \{t^*t \mid t \in \mathcal{T}_+(\mathcal{A})\}. \quad (2.8)$$

**Automorphisms** We now describe how  $\mathcal{T}_{++}$  can be seen as a group acting over  $\mathcal{K}(\mathcal{A})$ , see [Chu09, Section 2.2] or [Vin63, Chapter III, §2] for more details. Let  $u \in \mathcal{T}_{++}$  and for  $x \in \mathcal{K}(\mathcal{A})$  denote by  $t_x$  the unique element in  $\mathcal{T}_{++}$  such that  $x = t_x t_x^*$  holds. We define the map  $\tilde{Q}_u : \mathcal{K}(\mathcal{A}) \rightarrow \mathcal{K}(\mathcal{A})$  satisfying

$$\tilde{Q}_u(x) := (ut_x)(ut_x)^*, \quad \forall x \in \mathcal{K}(\mathcal{A}). \quad (2.9)$$

Then,  $\tilde{Q}_u$  is a bijection over  $\mathcal{K}(\mathcal{A})$  and it turns out that

$$\tilde{Q}_u(x) = u((t_x t_x^*)u^*) + u(u(t_x t_x^*)) - (uu)(t_x t_x^*) = u(xu^*) + u(ux) - (uu)(x) \quad (2.10)$$

holds, see [Chu09, Proposition 2]. In particular, the map  $\tilde{Q}_u$  is linear over  $\mathcal{K}(\mathcal{A})$ .

Next, for  $a \in \mathcal{A}$ , define  $a_H := a + a^*$ . Taking (2.10) as a starting point, we define the *quadratic map*  $Q_a : \mathcal{H}(\mathcal{A}) \rightarrow \mathcal{H}(\mathcal{A})$  for  $a \in \mathcal{A}$  as the linear map satisfying

$$Q_a(b) := \frac{1}{2}(a(ba^*) + a(ab) - (aa)b)_H, \quad (2.11)$$

e.g., see [Chu09, Definition 8]. In view of (2.10),  $\tilde{Q}_u(x) = Q_u(x)$  holds for  $x \in \mathcal{K}(\mathcal{A})$  and  $Q_u$  can be seen as a linear extension of  $\tilde{Q}_u$ .

A useful property is that for  $a \in \mathcal{A}$ , the adjoint map  $Q_a^*$  satisfies

$$Q_a^* = Q_{a^*}, \quad (2.12)$$

e.g., see [Chu09, Section 2.3].

The linear maps  $Q_u$  for  $u \in \mathcal{T}_{++}$  form a subgroup of the automorphism group of  $\mathcal{K}(\mathcal{A})$  acting transitively on  $\mathcal{K}(\mathcal{A})$ . In fact, for any  $u \in \mathcal{T}_{++}$ , there exists a unique  $u^{-1} \in \mathcal{T}_{++}$  such that that  $uu^{-1} = u^{-1}u = e$  holds, see [Chu09, Proposition 1]. More generally, we have

$$Q_u Q_t = Q_{ut}, \quad \forall u, t \in \mathcal{T}_+, \quad (2.13)$$

$$Q_u^{-1} = Q_{u^{-1}}, \quad \forall u \in \mathcal{T}_{++}, \quad (2.14)$$

$$Q_u(tt^*) = (ut)(ut)^*, \quad \forall u, t \in \mathcal{T}_+, \quad (2.15)$$

where (2.13) follows from [Chu09, Corollary 1] by taking limits, (2.14) follows from [Chu09, Proposition 1 and Corollary 1] and (2.15) can be obtained by taking limits in (2.9) or by invoking [Chu09, Proposition 2] and (2.11).

In what follows, we will say that an automorphism  $Q$  of  $\mathcal{K}(\mathcal{A})$  is *triangular* if there exists  $u \in \mathcal{T}_{++}$  such that  $Q = Q_u$  holds.

**Principal subalgebras** Let  $I \subseteq \{1, \dots, r\}$  be a subset of indices. We will identify certain subalgebras of  $\mathcal{A}$  that correspond to “principal submatrices”. We define  $\mathcal{A}_I$  to be the following subalgebra

$$\mathcal{A}_I := \{a \in \mathcal{A} \mid a_{ik} = a_{ki} = 0, \forall i \in I, \forall k \in \{1, \dots, r\}\}.$$

For example,  $a \in \mathcal{A}_{\{r\}}$  if and only if the  $r$ -th “column” and the  $r$ -th “row” of  $a$  vanishes. We will say that  $\mathcal{A}_I$  is a *principal subalgebra* of  $\mathcal{A}$ .

$\mathcal{A}_I$  can be seen as a T-algebra of rank  $s := r - |I|$  in a natural way using the structure inherited from  $\mathcal{A}$ , where  $|I|$  is the number of elements of  $I$ . Let  $J := \{1, \dots, r\} \setminus I$  be the remaining indices ordered as

$$n_1 < n_2 < \dots < n_s.$$

This implies that  $i < j \Leftrightarrow n_i < n_j$ , for all  $i, j \in \{1, \dots, s\}$ . In essence, we are “renaming” the remaining indices in  $J$  to indices in the set  $\{1, \dots, s\}$ , in such a way that their ordering in  $J$  is preserved. This will be important to ensure compatibility with the T-algebra structure of  $\mathcal{A}$ .

The bigradation of  $\mathcal{A}_I$  inherited from  $\mathcal{A}$  is given by  $\mathcal{A}_I = \bigoplus_{i,j=1}^s \bar{\mathcal{A}}_{ij}$ , where  $\bar{\mathcal{A}}_{ij} := \mathcal{A}_{n_i n_j}$ . For  $i, j, k, l \in \{1, \dots, s\}$ , we have

$$\bar{\mathcal{A}}_{ij} \bar{\mathcal{A}}_{lk} = \mathcal{A}_{n_i n_j} \mathcal{A}_{n_k n_l}.$$

Since  $n_j = n_k$  holds if and only if  $j = k$ , we have

$$\begin{aligned} \bar{\mathcal{A}}_{ij} \bar{\mathcal{A}}_{jk} &\subseteq \bar{\mathcal{A}}_{ik} \\ \bar{\mathcal{A}}_{ij} \bar{\mathcal{A}}_{kl} &= \{0\} \quad \text{if } j \neq k. \end{aligned}$$

Defining  $\bar{\rho}_i := \rho_{n_i}, \bar{e}_i := e_{n_i}$  and restricting the trace and involution functions to  $\mathcal{A}_I$ , it is straightforward (albeit tedious) to check that Axioms (a1) through (a7) still hold. For example, the restriction of the involution  $*$  is still an idempotent bijection from  $\mathcal{A}_I$  to  $\mathcal{A}_I$ , because if  $a_{ik} = a_{ki} = 0$ , then  $a_{ik}^* = a_{ki} = 0$  must also hold. Furthermore, we have

$$\bar{\mathcal{A}}_{ij}^* = \mathcal{A}_{n_i n_j}^* = \mathcal{A}_{n_j n_i} = \bar{\mathcal{A}}_{ji}$$

For the rest of the axioms the idea is that since they are true over  $\mathcal{A}$  for all the possible indices  $\{1, \dots, r\}$  they must be true for any subset  $J \subset \{1, \dots, r\}$  of those indices. This takes care of the validity of axioms (a1) to (a5) over  $\mathcal{A}_I$ . Next, if  $1 \leq i \leq j \leq k \leq l \leq s$ , then we must have  $1 \leq n_i \leq n_j \leq n_k \leq n_l \leq s$ , so Axiom (a6) is also valid for  $\mathcal{A}_I$ . A similar argument holds for Axiom (a7).

Finally, we take a look at the automorphisms of  $\mathcal{K}(\mathcal{A}_I)$ . As before, the triangular matrices  $\mathcal{T}_{++}(\mathcal{A}_I)$  induce a group that acts transitively on  $\mathcal{K}(\mathcal{A}_I)$  and in the next result we will see that they can be seen as restrictions of automorphisms of  $\mathcal{K}(\mathcal{A})$ .

**Proposition 2.4.** *Given a triangular automorphism  $Q_u$  of  $\mathcal{K}(\mathcal{A}_I)$ , there exists a triangular automorphism  $Q_{\tilde{u}}$  of  $\mathcal{K}(\mathcal{A})$  such that the restriction of  $Q_{\tilde{u}}$  to  $\mathcal{H}(\mathcal{A}_I)$  coincides with  $Q_u$ .*



*Proof.* We “complete”  $u$  by adding 1’s to the diagonal by letting  $\tilde{u} = u + \sum_{i \in I} e_i$ . In this way,  $\tilde{u} \in \mathcal{T}_{++}(\mathcal{A})$ . Next, let  $t \in \mathcal{T}_{++}(\mathcal{A}_I)$ . By (2.15) we have  $Q_{\tilde{u}}(tt^*) = (\tilde{u}t)(\tilde{u}t)^*$ . As a consequence of the multiplication rules in (2.4) and Axiom (a2) we have

$$\tilde{u}t = \left( u + \sum_{i \in I} e_i \right) t = ut + \sum_{i \in I} \sum_{j=1}^r t_{ij}.$$

However, since  $t \in \mathcal{T}_{++}(\mathcal{A}_I)$ , we have  $t_{ij} = 0$  whenever  $i \in I$ , so the second summation is zero and we have  $\tilde{u}t = ut$ . Therefore

$$Q_{\tilde{u}}(tt^*) = (ut)(ut)^* = Q_u(tt^*)$$

holds which implies that  $Q_{\tilde{u}}$  and  $Q_u$  coincide over  $\mathcal{K}(\mathcal{A}_I) = \{tt^* \mid t \in \mathcal{T}_{++}(\mathcal{A}_I)\}$  and must coincide over  $\mathcal{H}(\mathcal{A}_I) = \text{span } \mathcal{K}(\mathcal{A}_I)$ .  $\square$

### 3 Facial structure of homogeneous cones

In this section, we prove our main results on the facial structure of general homogeneous cones. Before that, we must discuss the generalized Cholesky decomposition in homogeneous cones.

#### 3.1 Unique Cholesky factorization in homogeneous cones

Every element in the interior of a homogeneous cone has a unique representation as  $tt^*$ , with  $t \in \mathcal{T}_{++}$ . Then, a limiting argument tells us that elements in the boundary can be represented as  $tt^*$  with  $t \in \mathcal{T}_+$ , however this factorization is not unique in general. We will address this issue in this subsection.

We say that  $t \in \mathcal{T}_+$  is *proper* if  $t_{ii} = 0$  implies that the  $i$ -th “column” of  $t$  vanishes, i.e.,  $t_{ki} = 0$  for all  $k < i$ . It turns out that each  $x \in \text{cl } \mathcal{K}(\mathcal{A})$  has a unique decomposition in terms of *proper* triangular matrices. In order to prove that, first we need the following proposition.

**Proposition 3.1.** *Let  $\mathcal{A}$  be a  $T$ -algebra of rank  $r$ . Each  $e_i$  generates extreme rays of both  $\text{cl } \mathcal{K}(\mathcal{A})$  and  $\mathcal{K}(\mathcal{A})^*$ . The proper face  $(\text{cl } \mathcal{K}(\mathcal{A})) \cap \{e_i\}^\perp$  of  $\text{cl } \mathcal{K}(\mathcal{A})$  is equal to  $\text{cl}(\mathcal{K}(\mathcal{A}_{\{i\}}))$ .*

*Proof.* We first prove that  $(\text{cl } \mathcal{K}(\mathcal{A})) \cap \{e_i\}^\perp = \text{cl}(\mathcal{K}(\mathcal{A}_{\{i\}}))$ . Let  $x \in (\text{cl } \mathcal{K}(\mathcal{A})) \cap \{e_i\}^\perp$ , then  $0 = \langle x, e_i \rangle = \rho_i(x_{ii})$ , so that  $x_{ii} = 0$ . Next we do some computations similar to the ones appearing in the proof of [KTX12, Proposition 2.4]. First, let  $t \in \mathcal{T}_+$  be such that  $x = tt^*$  holds. Then,

$$0 = x_{ii} = \sum_{j=1}^r t_{ij}t_{ij}^*.$$

Taking the trace, we get

$$0 = \text{tr}(x_{ii}) = \sum_{j=1}^r \text{tr}(t_{ij}t_{ij}^*).$$

By Axiom (a5), each  $\text{tr}(t_{ij}t_{ij}^*)$  is nonnegative and therefore they must be zero. Again by Axiom (a5), we conclude that all the  $t_{ij}$  are zero for  $j \in \{1, \dots, r\}$ . In particular,  $t$  belongs to the set of upper triangular matrices of the principal subalgebra  $\mathcal{A}_{\{i\}}$  and thus  $x \in \text{cl}(\mathcal{K}(\mathcal{A}_{\{i\}}))$  holds. The converse is clear. Overall, this also shows that  $\text{cl}(\mathcal{K}(\mathcal{A}_{\{i\}}))$  is an exposed face, since  $e_i = e_i^*$  also belongs to  $\mathcal{K}(\mathcal{A})^*$  by (2.8).

Next, we verify that  $e_i$  generates an extreme ray of  $\text{cl } \mathcal{K}(\mathcal{A})$ . Suppose that  $x + y = e_i$  holds for  $x, y \in \text{cl } \mathcal{K}(\mathcal{A})$ . Then,  $\rho_k(x_{kk}) + \rho_k(y_{kk}) = 0$  for  $k \neq i$ . Since  $\rho_k(x_{kk})$  and  $\rho_k(y_{kk})$  are nonnegative,

we have  $\rho_k(x_{kk}) = \rho_k(y_{kk}) = 0 = \langle x, e_k \rangle = \langle y, e_k \rangle$ . From [KTX12, Proposition 2.4], this implies that  $0 = x_{lk} = x_{kl} = y_{lk} = y_{kl}$ , for all  $k \neq i$  and all  $l \in \{1, \dots, r\}$ . In particular, only  $x_{ii}$  and  $y_{ii}$  can be nonzero and therefore, they must be in the half-line generated by  $e_i$ .

We recall that the dual cone is the cone corresponding to the T-algebra  $\mathcal{B}$  defined in (2.7) and each  $e_i$  is the identity element of  $\mathcal{B}_{r+1-i, r+1-i}$ . Applying what we proved so far to  $\text{cl } \mathcal{K}(\mathcal{B}) = \mathcal{K}(\mathcal{A})^*$  shows that the  $e_i$  are also extreme rays of  $\mathcal{K}(\mathcal{A})^*$ .  $\square$

The following proposition was essentially proven by Gindikin [Gin92, Chapter 2, §1, Lemma 7], but we provide a different proof here from which we will be able to extract an algorithm.

**Proposition 3.2** (Unique Cholesky factorization). *Let  $x \in \text{cl } \mathcal{K}(\mathcal{A})$ , then  $x = tt^*$  for an unique proper  $t \in \mathcal{T}_+$ .*

*Proof.* We proceed by induction on the rank  $r$  of the T-algebra. The proposition is true if  $r = 1$ . So let us assume that it holds for all T-algebras of rank  $(r - 1) \geq 1$  and let us show that it holds for T-algebras of rank  $r$ , for  $r \geq 2$ .

Let  $\mathcal{A}$  be a T-algebra of rank  $r$  and let  $x \in \text{cl } \mathcal{K}(\mathcal{A})$ . Let  $t \in \mathcal{T}_+$  be such that  $x = tt^*$ .

Let  $\tilde{t} \in \mathcal{T}_+$  be the element that corresponds to the first  $r - 1$  columns of  $t$ . That is,  $\tilde{t}_{ij} := t_{ij}$ , for all  $i, j \in \{1, \dots, r - 1\}$  and  $\tilde{t}_{ir} := 0$  for  $i \in \{1, \dots, r\}$ . Then  $\tilde{t} \in \mathcal{A}_{\{r\}}$ . By the induction hypothesis, over the subalgebra  $\mathcal{A}_{\{r\}}$  (which has rank  $r - 1$ ) there is a unique proper  $\hat{t} \in \mathcal{T}_+(\mathcal{A}_{\{r\}})$  such that  $\tilde{t}\tilde{t}^* = \hat{t}\hat{t}^*$  holds.

For the last column of  $x$  we have

$$x_{ir} = \sum_{k=1}^r t_{ik}(t^*)_{kr} = t_{ir}t_{rr}^* = \rho_r(t_{rr})t_{ir}, \quad \forall i \in \{1, \dots, r\}. \quad (3.1)$$

In particular,  $x_{rr} = \rho_r(t_{rr})t_{rr} = t_{rr}^2$  holds. We then consider two cases.

If  $\rho_r(x_{rr}) = 0$ , i.e.,  $\langle x, e_r \rangle = 0$ , then the last row and the last column of  $x$  must vanish by Proposition 3.1, so  $x$  belongs to the closure of the homogeneous cone associated to the principal subalgebra  $\mathcal{A}_{\{r\}}$ , i.e.,  $x \in \text{cl } (\mathcal{K}(\mathcal{A}_{\{r\}}))$ . Since  $\hat{t}$  is also proper with respect to  $\mathcal{A}$  (after all, the last column of  $\hat{t}$  vanishes),  $\hat{t}$  is the desired proper element satisfying  $x = \hat{t}\hat{t}^*$ . We note that  $\hat{t}$  is unique over  $\mathcal{T}_+$  because (3.1) implies that all proper  $t$  satisfying  $x = tt^*$  must have  $t_{rr} = 0$ , which leads to  $t_{ir} = 0$  for all  $i$ .

Next, suppose that  $\rho_r(x_{rr}) > 0$ . Then, (3.1) implies that the last column of any  $t \in \mathcal{T}_+$  satisfying  $x = tt^*$  is unique and satisfies:

$$t_{ir} = x_{ir} / (\sqrt{\rho_r(x_{rr})}). \quad (3.2)$$

Let  $\bar{t} \in \mathcal{T}_+$  be such that the first  $r - 1$  columns of  $\bar{t}$  correspond to the columns of  $\hat{t}$  and the last column is given by (3.2). With that  $\bar{t}$  must be proper, since  $\hat{t}$  is proper and  $\rho_r(\bar{t}_{rr}) > 0$ . For convenience, denote the last column of  $\bar{t}$  by  $u$ . We have

$$\bar{t}\bar{t}^* = (\hat{t} + u)(\hat{t}^* + u^*) = \hat{t}\hat{t}^* + uu^* = \tilde{t}\tilde{t}^* + uu^* = tt^* = x. \quad (3.3)$$

We recall that  $t$  is an arbitrary element of  $\mathcal{T}_+$  satisfying  $x = tt^*$ , so (3.3) and (3.2) imply that the last column of  $t$  is uniquely determined and the remaining  $r - 1$  columns must be such that  $\hat{t}\hat{t}^* = \tilde{t}\tilde{t}^*$  holds. In particular, if  $t$  is proper, the uniqueness of  $\hat{t}$  over  $\mathcal{A}_{\{r\}}$  implies that  $\tilde{t}$  (i.e., the first  $r - 1$  columns of  $t$ ) coincides with  $\hat{t}$ . This concludes the proof.  $\square$

The proof of Proposition 3.2 shows that if  $x \in \text{cl}\mathcal{K}(\mathcal{A})$  and  $x = tt^*$  holds for a proper  $t \in \mathcal{T}_+$  then, the last column of  $t$  is either zero (if  $x_{rr} = 0$ ) or is uniquely determined by (3.2). Denoting such a column by  $u$  and the remaining  $r - 1$  columns of  $t$  by  $\hat{t}$ , (3.3) implies that  $\hat{t}$  satisfies

$$\hat{t}\hat{t}^* = x - uu^*.$$

Now,  $\hat{t}$  is an element of  $t \in \mathcal{T}_+(\mathcal{A}_{\{r\}})$ , so again, the last column of  $\hat{t}$  in  $\mathcal{A}_{\{r\}}$  (i.e., the  $(r-1)$ -th column of  $\hat{t}$  seen as an element of  $\mathcal{A}$ ), is uniquely determined and is either zero (if  $(x - uu^*)_{r-1,r-1} = 0$ ) or is given by (3.2) with  $x - uu^*$  in place of  $x$ . This leads to an algorithm for computing a generalized Cholesky decomposition, see Algorithm 1. Note that Algorithm 1 can also be used to test membership in  $\mathcal{K}(\mathcal{A})$ : given some  $x \in \mathcal{H}(\mathcal{A})$ , we invoke Algorithm 1 with  $x$  as input. Denoting by  $t$  the output of Algorithm 1, we have  $x \in \mathcal{K}(\mathcal{A})$  if and only if  $x = tt^*$ .

---

**Algorithm 1** Generalized Cholesky decomposition in a homogeneous cone

---

**Require:**  $x \in \text{cl}\mathcal{K}(\mathcal{A})$

- 1:  $u \leftarrow 0, t \leftarrow 0, y \leftarrow x$
  - 2: **for**  $i = r, \dots, 1$  **do**
  - 3:    $y \leftarrow y - uu^*$
  - 4:   **if**  $\rho_i(y_{ii}) > 0$  **then**
  - 5:      $t_{ji} \leftarrow y_{ji}/(\sqrt{\rho_i(y_{ii})}), \quad \forall j \in \{1, \dots, i\}$ .
  - 6:   **else**
  - 7:      $t_{ji} \leftarrow 0, \quad \forall j \in \{1, \dots, i\}$
  - 8:   **end if**
  - 9:    $u \leftarrow i$ -th column of  $t$
  - 10: **end for**
  - 11: **return**  $t$
- 

### 3.2 Faces and their orbits under the action of triangular matrices

In this subsection, we present our main results on the facial structure of homogeneous cones. We start by observing that the cones induced by principal subalgebras are faces that are orthogonally projectively exposed.

**Lemma 3.3.** *Let  $\mathcal{A}$  be a  $T$ -algebra of rank  $r$  and let  $I \subseteq \{1, \dots, r\}$  be nonempty. Then,  $\text{cl}(\mathcal{K}(\mathcal{A}_I))$  is an orthogonally projectively exposed face of  $\text{cl}\mathcal{K}(\mathcal{A})$ . In particular,  $Q_{e_I}(\text{cl}\mathcal{K}(\mathcal{A})) = \text{cl}(\mathcal{K}(\mathcal{A}_I))$  holds, where  $e_I := \sum_{i \notin I} e_i$ .*

*Proof.* From Proposition 3.1 we have that each  $\text{cl}\mathcal{K}(\mathcal{A}_{\{i\}})$  for  $i \in \{1, \dots, r\}$  is a face and we have

$$\text{cl}(\mathcal{K}(\mathcal{A}_I)) = \bigcap_{i \in I} \text{cl}(\mathcal{K}(\mathcal{A}_{\{i\}})) = \bigcap_{i \in I} \text{cl}\mathcal{K}(\mathcal{A}) \cap \{e_i\}^\perp.$$

Since an intersection of faces is a face, this shows that  $\text{cl}(\mathcal{K}(\mathcal{A}_I))$  is a face of  $\text{cl}\mathcal{K}(\mathcal{A})$ .

Now, let  $e_I$  be the identity element of  $\mathcal{A}_I$ , i.e.,  $e_I = \sum_{i \notin I} e_i$ . We will consider the quadratic map  $Q_{e_I}$  and show that  $Q_{e_I}$  is the desired projection.

First, we observe that  $Q_{e_I}$  is idempotent, since  $Q_{e_I}Q_{e_I} = Q_{e_I e_I} = Q_{e_I}$  holds, where the first equality follows from (2.13) and the second from (2.3) and Axiom (a2). The map  $Q_{e_I}$  is also self-adjoint by (2.12) since  $e_I^* = e_I$ . This shows that  $Q_{e_I}$  is an orthogonal projection.

Next, let  $x \in \mathcal{K}(\mathcal{A})$  and write  $x = tt^*$  for some  $t \in \mathcal{T}_{++}$ , then, in view of (2.15) we have

$$Q_{e_I}(x) = (e_I t)(e_I t)^*.$$

Denoting the “rows” of  $t$  by  $t^i$ , so that  $t = t^1 + \cdots + t^r$  holds and letting  $t_I := \sum_{i \notin I} t^i$  we have  $e_I t = t_I$  and

$$Q_{e_I}(x) = (t_I)(t_I)^*.$$

Computing the diagonal elements of  $Q_{e_I}(x)$  for  $i \in I$  we obtain

$$(Q_{e_I}(x))_{ii} = \sum_{j=i}^r (t_I)_{ij} ((t_I)_{ij})^* = 0,$$

since  $(t_I)_{ij} = 0$  for  $i \in I$ . Then, Proposition 3.1 and  $Q_{e_I}(x) \in \text{cl } \mathcal{K}(\mathcal{A})$  implies that  $Q_{e_I}(x) \in \text{cl}(\mathcal{K}(\mathcal{A}_I))$ . Therefore, the projection  $Q_{e_I}$  maps elements of  $\text{cl } \mathcal{K}(\mathcal{A})$  to elements of  $\text{cl}(\mathcal{K}(\mathcal{A}_I))$ . Finally, since  $e_I$  is the identity element in  $\mathcal{A}_I$ , we have  $Q_{e_I}(x) = x$  for all  $x \in \mathcal{H}(\mathcal{A}_I)$  by (2.11).  $\square$

Motivated by Lemma 3.3, in what follows we will call a face of the format  $\text{cl}(\mathcal{K}(\mathcal{A}_I))$  a *principal face*, since they are induced by principal subalgebras of  $\mathcal{A}$ .

Our next task is showing that every element of  $\text{cl } \mathcal{K}(\mathcal{A})$  can be transformed to the identity element of some principal subalgebra via a triangular automorphism.

**Lemma 3.4.** *Let  $\mathcal{A}$  be a  $T$ -algebra of rank  $r$  and  $x \in \text{cl } \mathcal{K}(\mathcal{A})$ . Let  $x = tt^*$  for a proper  $t \in \mathcal{T}_+$  as in Proposition 3.2. There exists a triangular automorphism  $Q_u$  of  $\mathcal{K}(\mathcal{A})$  such that*

$$Q_u(x) = e_I,$$

where  $I := \{i \mid \rho_i(t_{ii}) = 0\}$ ,  $e_I := \sum_{i \notin I} e_i$ .

*Proof.* Let  $t_1, \dots, t_r$  denote the “columns” of  $t$ , so that

$$t = \sum_{i=1}^r t_i = \sum_{i \notin I} t_i,$$

where the second equality holds because  $t$  is assumed to be proper.

For  $i, j \in \{1, \dots, r\}$  we have

$$t_i e_j = \sum_{1 \leq k \leq i \leq r} t_{ki} e_j$$

By the multiplication rules in (2.4) and Axiom (a2), we have

$$t_i e_j = \begin{cases} t_i & \text{if } i = j \\ 0 & \text{otherwise.} \end{cases}$$

With this in mind, we define  $\tilde{u} \in \mathcal{T}_{++}$  as follows: for  $i \in I$ , we set the  $i$ -th column of  $\tilde{u}$  to be equal to  $e_i$  and for  $i \notin I$ , we set the  $i$ -th column of  $\tilde{u}$  to be  $t_i$ . We have

$$\tilde{u} e_I = \left( \sum_{j \in I} e_j + \sum_{i \notin I} t_i \right) \left( \sum_{k \notin I} e_k \right) = \sum_{i \notin I} t_i = t.$$

With that,  $Q_{\tilde{u}}$  maps  $e_I = e_I e_I^*$  to  $(\tilde{u} e_I)(\tilde{u} e_I)^* = tt^* = x$ , see (2.15). By (2.14), letting  $u := \tilde{u}^{-1}$ ,  $Q_u = Q_{\tilde{u}}^{-1}$  is a triangular automorphism mapping  $x$  to  $e_I$ .  $\square$

Gathering all the pieces collected so far, we have the following result, which contains, in particular, an analogue of (1.1) and the subsequent discussion.

**Theorem 3.5.** *Let  $\mathcal{F}$  be a proper face of a homogeneous cone  $\text{cl}\mathcal{K}(\mathcal{A})$  of rank  $r$ . The following items hold.*

- (i)  $\mathcal{F}$  is projectionally exposed.
- (ii)  $\mathcal{F}$  is a homogeneous cone of rank  $s < r$  and there is a triangular automorphism  $Q_u$  of  $\mathcal{K}(\mathcal{A})$  such that  $Q_u(\mathcal{F}) = \text{cl}\mathcal{K}(\mathcal{A}_I)$  holds for some  $I \subsetneq \{1, \dots, r\}$ . Furthermore, letting  $x \in \text{ri}\mathcal{F}$  and writing  $x = tt^*$  for a proper  $t \in \mathcal{T}_+(\mathcal{A})$ , we have  $I = \{i \mid \rho_i(t_{ii}) = 0\}$  and  $s = r - |I|$ .
- (iii) There is a subgroup  $\mathcal{G}$  of automorphisms of  $\mathcal{F}$  acting simply transitively on the relative interior of  $\mathcal{F}$  such that each  $g \in \mathcal{G}$  is obtained through the restriction of a triangular automorphism of  $\mathcal{K}(\mathcal{A})$  to  $\mathcal{F}$ .

*Proof.* Let  $x \in \text{ri}\mathcal{F}$  be arbitrary and consider a decomposition of  $x = tt^*$  with a proper  $t$  as in Proposition 3.2. By Lemma 3.4, there exists a triangular automorphism  $Q_u$  that maps  $x$  to  $e_I$ , where  $I = \{i \mid \rho_i(t_{ii}) = 0\}$ . We have

$$Q_u(\mathcal{F}) = Q_u(\text{Face}(x, \text{cl}\mathcal{K}(\mathcal{A}))) = \text{Face}(Q_u(x), \text{cl}\mathcal{K}(\mathcal{A})) = \text{Face}(e_I, \text{cl}\mathcal{K}(\mathcal{A})) = \text{cl}\mathcal{K}(\mathcal{A}_I), \quad (3.4)$$

where the first equality follows from (2.1). The second equality holds because  $Q_u$  is an automorphism, so it maps a face to another face and is also a bijection between the relative interiors. Finally, the last equality holds because  $e_I \in \text{ri}(\text{cl}\mathcal{K}(\mathcal{A}_I)) = \mathcal{K}(\mathcal{A}_I)$ . We note that if  $|I| = 0$  we have  $e_I = e$  and  $\mathcal{K}(\mathcal{A}_I) = \mathcal{K}(\mathcal{A})$ , which would contradict the fact that  $\mathcal{F}$  is proper. We have thus obtained item (ii) since homogeneity is preserved by linear isomorphisms and  $\mathcal{K}(\mathcal{A}_I)$  is a homogeneous cone of rank  $r - |I| < r$ .

Next, let  $Q_{e_I}$  be the orthogonal projection mapping  $\text{cl}\mathcal{K}(\mathcal{A})$  to  $\text{cl}\mathcal{K}(\mathcal{A}_I)$  as in Lemma 3.3. Then, (3.4) implies that  $Q_{u^{-1}}Q_{e_I}Q_u$  is a projection mapping  $\text{cl}\mathcal{K}(\mathcal{A})$  to  $\mathcal{F}$ , which proves item (i).

Let  $\mathcal{G}$  be the group of triangular automorphisms of  $\mathcal{K}(\mathcal{A}_I)$  and we recall that  $\mathcal{G}$  acts simply transitively on  $\mathcal{K}(\mathcal{A}_I)$ . In view of (3.4),  $Q_u^{-1}\mathcal{G}Q_u$  acts simply transitively on  $\mathcal{F}$ . Also, Proposition 2.4 tells us that for every  $Q \in \mathcal{G}$ , there exists some triangular automorphism  $Q_v$  that coincides with  $Q$  over  $\mathcal{H}(\mathcal{A}_I)$  which implies that  $Q_u^{-1}QQ_u$  coincides with  $Q_{u^{-1}}Q_vQ_u$  over  $\mathcal{F}$ . We note that  $Q_{u^{-1}}Q_vQ_u$  is a triangular automorphism of  $\mathcal{K}(\mathcal{A})$  since  $Q_{u^{-1}}Q_vQ_u = Q_{u^{-1}vu}$  holds by (2.13) and there is no ambiguity regarding  $u^{-1}vu \in \mathcal{T}_{++}$  since  $(u^{-1}v)u = u^{-1}(vu)$  which is implied by Axiom (a6). Overall, we conclude that, indeed, there exists a subgroup acting simply transitively over  $\mathcal{F}$  such that each element arises as a restriction of a triangular automorphism of  $\mathcal{K}(\mathcal{A})$ . This, together with (3.4), proves items (ii) and (iii).  $\square$

Theorem 3.5 provides a relatively complete picture of the facial structure of a homogeneous cone. It tells us that not only the faces are projectionally exposed and homogeneous cones themselves, but their homogeneous structure can be obtained from the original cone.

Furthermore, each face is linearly isomorphic to a principal face through an automorphism of the cone. Finding the automorphism is an entirely algorithmic process. Given some  $x \in \text{ri}\mathcal{F}$ , we first decompose  $x$  as  $tt^*$  for  $t$  a proper matrix as in Algorithm 1, then we compute the index set  $I = \{i \mid \rho_i(t_{ii}) = 0\}$ , which reveals the principal face that is isomorphic to  $\mathcal{F}$ . The actual isomorphism can then be computed by finding  $u \in \mathcal{T}_{++}$  such that

$$ut = e_I \quad (3.5)$$

holds and  $u$  always exists by Lemma 3.4. Solving this system does not pose difficulties given the upper triangular structure of  $u$  and  $t \in \mathcal{T}_+$ . Then,  $Q_u$  is an automorphism of  $\text{cl}\mathcal{K}(\mathcal{A})$  mapping  $\mathcal{F}$  to a principal face. Letting  $\bar{e} := e - e_I$  and noticing that Proposition 3.1 implies that  $\bar{e}$  exposes the principal face induced by  $I$ , this discussion also leads to the following expression for  $\mathcal{F}$

$$\mathcal{F} = \{x \in \text{cl}\mathcal{K}(\mathcal{A}) \mid \langle Q_u(x), \bar{e} \rangle = 0\} = \{x \in \text{cl}\mathcal{K}(\mathcal{A}) \mid \langle x, Q_{u^*}(\bar{e}) \rangle = 0\}, \quad (3.6)$$

which also reveals that  $Q_{u^*}(\bar{e})$  is an exposing vector for  $\mathcal{F}$ . Following the proof of Theorem 3.5, a projection mapping  $\text{cl}\mathcal{K}(\mathcal{A})$  to  $\mathcal{F}$  is  $Q_v$ , for  $v := u^{-1}e_I u$ . We will see a complete example of this process in Section 4.

Recalling the discussion of the faces of PSD matrices in (1.1), we can see several similarities but two important difference are that, in general,  $Q_v$  may not be self-adjoint and  $Q_u$  is not necessarily a linear isometry.

**Some consequences and discussion** The results so far together with Algorithm 1 provide an algorithmic way of identifying elements of  $\text{cl}\mathcal{K}(\mathcal{A})$  that generate extreme rays.

**Corollary 3.6** (Identifying extreme rays). *Let  $x \in \text{cl}\mathcal{K}(\mathcal{A})$  and write  $x = tt^*$  for some proper  $t \in \mathcal{T}_+$ . Then  $x$  generates an extreme ray if and only if  $\rho_i(t_{ii}) > 0$  for one and exactly one  $i \in \{1, \dots, r\}$ .*

*Proof.* Let  $\mathcal{F} = \text{Face}(x, \text{cl}\mathcal{K}(\mathcal{A}))$ . We observe that  $x$  generates an extreme ray if and only if the dimension of  $\mathcal{F}$  is one. By Theorem 3.5,  $\mathcal{F}$  is linearly isomorphic to  $\text{cl}\mathcal{K}(\mathcal{A}_I)$ , where  $I = \{i \mid \rho_i(t_{ii}) = 0\}$ . Since  $e_i \in \text{cl}\mathcal{K}(\mathcal{A}_I)$  for  $i \notin I$ , the dimension of  $\text{cl}\mathcal{K}(\mathcal{A}_I)$  is at least  $r - |I|$ . So the only way that  $\mathcal{F}$  can be an extreme ray is if  $r - |I| = 1$ , which happens if and only if  $\rho_i(t_{ii}) > 0$  for one and exactly one  $i \in \{1, \dots, r\}$ .  $\square$

Given a closed convex cone  $\mathcal{K}$ , a *chain of faces of length  $\ell$*  is a sequence of faces of  $\mathcal{K}$  such that  $\mathcal{F}_1 \subsetneq \dots \subsetneq \mathcal{F}_\ell$ . The length of a longest chain of faces of  $\mathcal{K}$ , denoted by  $\ell_{\mathcal{K}}$ , is an important quantity in the study of the regularization technique *facial reduction* [BW81, WM13, Pat13b]. In particular, it can be used to upper bound the number of steps required by facial reduction algorithms. Through this connection,  $\ell_{\mathcal{K}}$  appears in different contexts, e.g., as a way to upper bound the length of infeasibility certificates [LP18, Theorem 4]. The quantity  $\ell_{\mathcal{K}}$  also appears in the study of the expressive power of cones [Sau20]. Next, we compute  $\ell_{\mathcal{K}}$  for a homogeneous cone.

**Corollary 3.7.** *Let  $\text{cl}\mathcal{K}(\mathcal{A})$  be a homogeneous cone of rank  $r$ , then  $\ell_{\mathcal{K}} = r + 1$ .*

*Proof.* Take any chain of faces  $\text{cl}\mathcal{K}(\mathcal{A}) = \mathcal{F}_0 \supsetneq \dots \supsetneq \mathcal{F}_\ell$  of length  $\ell + 1$ . By item (ii) of Theorem 3.5, each  $\mathcal{F}_i$  is a homogeneous cone of rank smaller than that of  $\mathcal{F}_{i-1}$ . As  $\mathcal{K}$  has rank  $r$ , the length  $\ell + 1$  is at most  $r + 1$ . Using Proposition 3.1 inductively, we can construct a chain of faces of length  $r + 1$  as follows.

$$\{0\} = \text{cl}\mathcal{K}(\mathcal{A}_{\{1,2,\dots,r\}}) \subsetneq \text{cl}\mathcal{K}(\mathcal{A}_{\{1,2,\dots,r-1\}}) \subsetneq \dots \subsetneq \text{cl}\mathcal{K}(\mathcal{A}_{\{1\}}) \subsetneq \text{cl}\mathcal{K}(\mathcal{A}).$$

$\square$

The automorphism group of  $\text{cl}\mathcal{K}(\mathcal{A})$  acts on the set of (non-empty) faces in a natural way. It turns out that this action has finitely many orbits and each orbit contains a unique principal face.

**Corollary 3.8** (Orbits of faces). *Let  $\mathcal{F}(\text{cl}\mathcal{K}(\mathcal{A}))$  denote the set of faces of a rank  $r$  homogeneous cone  $\text{cl}\mathcal{K}(\mathcal{A})$ . The action of the triangular automorphisms on  $\mathcal{F}(\text{cl}\mathcal{K}(\mathcal{A}))$  has  $2^r$  orbits and each orbit has a unique representative among principal faces.*

*Proof.* By Theorem 3.5 each  $\mathcal{F} \trianglelefteq \text{cl}\mathcal{K}(\mathcal{A})$  is isomorphic to some principal face under a triangular automorphism. Since there are at most  $2^r$  principal faces, this is also an upper bound on the number of orbits. It only remains to show that there are indeed  $2^r$  orbits which can be done by arguing that each orbit contains at most one principal face.

Suppose that  $Q_t$  is a triangular automorphism and  $Q_t(\text{cl}\mathcal{K}(\mathcal{A}_I)) = \text{cl}\mathcal{K}(\mathcal{A}_J)$  holds. Then,  $|I| = |J|$  must also hold since rank is preserved by automorphisms. Let  $e_I$  and  $e_J$  be the identity elements of  $\mathcal{A}_I$  and  $\mathcal{A}_J$  respectively. We have  $e_I \in \text{ri}\text{cl}\mathcal{K}(\mathcal{A}_I) = \mathcal{K}(\mathcal{A}_I)$ , so  $Q_t e_I$  is an element of  $\mathcal{K}(\mathcal{A}_J)$  since linear transformations map relative interiors into relative interiors, e.g., see [Roc97, Theorem 6.6].

As  $\mathcal{K}(\mathcal{A}_J)$  is a homogeneous cone, there exists a triangular automorphism  $Q_{\bar{t}}$  of  $\mathcal{K}(\mathcal{A}_J)$  that maps  $Q_t e_I$  to  $e_J$ . Then, Proposition 2.4 implies that there exists a triangular automorphism  $Q_{\bar{t}}$  of  $\mathcal{K}(\mathcal{A})$  that maps  $Q_t e_I$  to  $e_J$ . Therefore,  $u := \bar{t}t$  is such that  $Q_u$  is a triangular automorphism of  $\mathcal{K}(\mathcal{A})$  satisfying  $Q_u(e_I) = e_J$ , by (2.13). Which in view of (2.15) implies that

$$Q_u(e_I) = (ue_I)(ue_I)^* = e_J = e_J e_J^*.$$

Following similar computations as in the proof of Lemma 3.4, we see that  $ue_I$  is a proper matrix. Since the decomposition in proper matrices is unique (Proposition 3.2), we conclude that  $ue_I = e_J$  must hold. Next, we verify that this implies  $I = J$ .

Recalling that  $e_I = \sum_{i \notin I} e_i$ , we have that  $(ue_I)_{ij} = \sum_{k=1}^r u_{ik}(e_I)_{kj}$  is 0 if  $j \in I$  and  $u_{ij}$  if  $j \notin I$ . Since  $ue_I = e_J$  must hold, we have  $u_{ij} = 0$  for  $i \neq j$ , i.e.,  $u$  only has nonzero components on its “diagonal”. Finally, since  $u \in \mathcal{T}_{++}$ ,  $ue_I = e_J$  forces that  $I = J$ .  $\square$

Corollary 3.8 is about the orbits under triangular automorphisms. If we consider the orbits under the *full* automorphism group, the number of orbits never increases, which implies that, in particular, it is still finite and principal faces can still be taken to be representatives of orbits. However, it may happen that each orbit may contain several principal faces<sup>1</sup>.

A finite number of orbits of faces is an interesting property that is not shared by all finite dimensional convex cones. For example, that is not the case for the  $p$ -cones for  $p \in (1, \infty)$ ,  $p \neq 2$  in dimension at least 3, as is implied by [IL19, Corollary 12]<sup>2</sup>.

Next, we take a look at projectional exposedness. It was shown in [Lou21, Proposition 33] that all symmetric cones are orthogonally projectationally exposed. In view of Theorem 3.5, the class of cones known to be projectationally exposed includes the homogeneous cones but it seems that the price to pay for losing self-duality is that we can no longer ensure that the projections are orthogonal (i.e., self-adjoint) under the inner product that comes from the T-algebra structure. We will revisit this issue in Proposition 4.5.

More generally, one may wonder how large the class of projectationally exposed cones is. Notably, for hyperbolicity cones and spectrahedral cones, two classes of cones larger than homogeneous cones, it is not currently known whether they are projectationally exposed in general. The strongest result so far is that they are amenable, see [LRS22, Corollary 3.5] and [LRS23].

**Connections to Ishi’s results** Here we will describe the connections of our discussion so far with a general result by Ishi [Ish00]. Let  $\mathcal{K}$  be an open homogeneous cone and  $H$  a group that acts linearly and simply transitively on  $\mathcal{K}$ , see [Ish00] for more details on the assumptions. Then, Ishi shows that there are  $2^r$  distinct orbits of the action of  $H$  onto  $\text{cl}\mathcal{K}$ , see [Ish00, Theorems A and

<sup>1</sup>For example, in the case of the positive semidefinite matrices  $S_+^n$ , all the extreme rays (which correspond to the faces generated by rank-1 PSD matrices) are in the same orbit under the action of the full automorphism group.

<sup>2</sup>In more details, Corollary 12 therein implies that the each orbit of the action of the automorphism group on the set of faces is finite. As there infinitely many faces, there must be an infinite number of orbits.



3.5]. This is done by invoking the theory of normal  $\mathfrak{j}$ -algebras, some Lie algebra theory and showing that the action of  $H$  onto  $\text{cl}\mathcal{K}$  can be expressed via generalized triangular matrices. Although the algebraic framework is different, the development in [Ish00] is somewhat close in spirit to what we have done here. In particular, translating to our language, Ishi proved that

$$\text{cl}\mathcal{K}(\mathcal{A}) = \bigcup_{I \subseteq \{1, \dots, r\}} \mathcal{O}_I,$$

where  $\mathcal{O}_I$  is the orbit of  $e_I = \sum_{i \notin I} e_i$  under the action of the triangular automorphisms and the union is disjoint. This is essentially equivalent to Corollary 3.8. Although there is no discussion of facial structure in [Ish00], using Ishi’s result as a starting point, it seems feasible to give another proof of Theorem 3.5. But in this case, some arguments regarding projectional exposedness and the realization of faces as homogeneous cones would still be necessary. While the discussion here has some overlap with [Ish00], our point of view is different and we emphasize certain concrete aspects such as Algorithm 1, how to explicitly identify faces, exposing vectors and the automorphisms needed to reveal them as in (3.5) and (3.6). Our development is also entirely done in the T-algebraic language, which is important from an algorithmic point of view, as it gives easy access to a self-concordant barrier of optimal parameter, e.g., see [Chu09, Section 3.1], [GT98, Theorem 4.1].

### 3.3 The dual cone

A T-Algebra for the dual cone  $\mathcal{K}(\mathcal{A})^*$  can be obtained as in (2.7). Under the bigradation induced by  $\mathcal{B}$ , upper triangular elements in  $\mathcal{B}$  correspond to lower triangular elements in  $\mathcal{A}$ . All the development done so far can thus be translated to the dual side by appealing to (2.7). Still, we believe it is useful to explicitly state certain dual results and translate the terminology, since there are certain subtleties.

For convenience, let  $\mathcal{L}_{++}(\mathcal{A}) := \mathcal{T}_{++}(\mathcal{A})^*$ ,  $\mathcal{L}_+(\mathcal{A}) := \mathcal{T}_+(\mathcal{A})^*$  and  $\mathcal{L}(\mathcal{A}) := \mathcal{T}(\mathcal{A})^*$ . A *triangular automorphism* of  $\mathcal{K}(\mathcal{A})^*$  is a linear map of the form  $Q_l$  for  $l \in \mathcal{L}_{++}(\mathcal{A})$ . Analogous properties to (2.13), (2.14), (2.15) hold for  $l, t \in \mathcal{L}_+$ . For example,  $Q_l(tt^*) = (lt)(lt)^*$  holds for  $l, t \in \mathcal{L}_+(\mathcal{A})$ , see also [Chu09, Proposition 7].

A potential pitfall is the definition of proper (lower) triangular matrices. Applying the definition of proper (upper) triangular matrices to  $\mathcal{B}$  and translating the indices back to  $\mathcal{A}$ , we see the “correct way” of defining properness. Indeed,  $t \in \mathcal{L}_+(\mathcal{A})$  is said to be *proper* if  $\rho_i(t_{ii}) = 0$  implies that the  $i$ -th “column” (not the row!) of  $t$  vanishes<sup>3</sup>. Analogous to Algorithm 1, we also have a Cholesky factorization in proper lower triangular matrices. Essentially, we apply Algorithm 1 to  $\mathcal{B}$  and translate back the indices to  $\mathcal{A}$  in two steps. First, the indices in  $t_{ij}$  and  $y_{ij}$  are replaced with  $r + 1 - i$  and  $r + 1 - j$ , respectively and, then we simplify the resulting *for loop*. The result is Algorithm 2.

The only remaining technical detail is that principal faces need to be replaced with their duals taken *inside the corresponding subalgebra*. More precisely, for a principal face  $\text{cl}\mathcal{K}(\mathcal{A}_I)$ , let

$$\mathcal{K}^D(\mathcal{A}_I) := \{tt^* \mid t \in \mathcal{L}_{++}(\mathcal{A}_I)\},$$

so that

$$\text{cl}(\mathcal{K}^D(\mathcal{A}_I)) = \{tt^* \mid t \in \mathcal{L}_+(\mathcal{A}_I)\} \tag{3.7}$$

<sup>3</sup>This is somewhat confusing, so here is a quick example. For  $r = 3$ , let  $t$  be an upper triangular element in  $\mathcal{B}$  and let us look at the third column, which corresponds to  $t_{13}, t_{23}, t_{33}$ . In view of (2.7), if we decompose  $t$  in the bigradation of  $\mathcal{A}$  (i.e., express  $t$  as sum of elements in the  $\mathcal{A}_{ij}$ ’s),  $t$  corresponds to a lower triangular  $\hat{t} \in \mathcal{L}(\mathcal{A})$  and the third column of  $t$  gets mapped to the first column of  $\hat{t}$ , so that  $t_{33} = \hat{t}_{11}$ ,  $t_{23} = \hat{t}_{21}$  and  $t_{13} = \hat{t}_{31}$ .

---

**Algorithm 2** Dual Generalized Cholesky decomposition in a homogeneous cone
 

---

**Require:**  $x \in \mathcal{K}(\mathcal{A})^*$

```

1:  $l \leftarrow 0, t \leftarrow 0, y \leftarrow x$ 
2: for  $i = 1, \dots, r$  do
3:    $y \leftarrow y - ll^*$ 
4:   if  $\rho_i(y_{ii}) > 0$  then
5:      $t_{ji} \leftarrow y_{ji} / (\sqrt{\rho_i(y_{ii})}), \quad \forall j \in \{i, \dots, r\}$ .
6:   else
7:      $t_{ji} \leftarrow 0, \quad \forall j \in \{i, \dots, r\}$ .
8:   end if
9:    $l \leftarrow i$ -th column of  $t$ 
10: end for
11: return  $t$ 

```

---

holds. A face of  $\mathcal{K}(\mathcal{A})^*$  as in (3.7) is said to be a *principal dual face*. We note that  $\text{cl}(\mathcal{K}^D(\mathcal{A}_I))$  is the dual of  $\text{cl}\mathcal{K}(\mathcal{A}_I)$  inside  $\mathcal{H}(\mathcal{A}_I)$ , i.e., we have  $\text{cl}(\mathcal{K}^D(\mathcal{A}_I)) = \text{cl}\mathcal{K}(\mathcal{A}_I)^* \cap \mathcal{H}(\mathcal{A}_I)$ .

**Theorem 3.9.** *Let  $\mathcal{K}(\mathcal{A})$  be a homogeneous cone of rank  $r$  and  $\mathcal{F}$  be a proper face of  $\mathcal{K}(\mathcal{A})^*$ .*

- (i)  $\mathcal{F}$  is projectionally exposed.
- (ii)  $\mathcal{F}$  is a homogeneous cone of rank  $s < r$  and there is a triangular automorphism  $Q_I$  of  $\mathcal{K}(\mathcal{A})^*$  such that  $Q_I(\mathcal{F}) = \text{cl}(\mathcal{K}^D(\mathcal{A}_I))$  holds for some  $I \subsetneq \{1, \dots, r\}$ . Furthermore, letting  $x \in \text{ri}\mathcal{F}$  and writing  $x = tt^*$  for a proper  $t \in \mathcal{L}_+(\mathcal{A})$ , we have  $I = \{i \mid \rho_i(t_{ii}) = 0\}$  and  $s = r - |I|$ .
- (iii) There is a subgroup  $\mathcal{G}$  of automorphisms of  $\mathcal{F}$  acting simply transitively on the relative interior of  $\mathcal{F}$  such that each  $g \in \mathcal{G}$  is obtained through the restriction of a triangular automorphism of  $\mathcal{K}(\mathcal{A})^*$  to  $\mathcal{F}$ .

*Proof.* Following the discussion so far, we apply Theorem 3.5 to  $\mathcal{B}$  as in (2.7) and translate the indices back to the bigradation of  $\mathcal{A}$ . □

Analogous to (3.5), the automorphism that maps  $\mathcal{F}$  to a principal dual face can be obtained by finding  $l \in \mathcal{L}_{++}(\mathcal{A})$  satisfying  $lt = e_I$ .

We conclude this subsection by observing that conjugate faces in a homogeneous cone have complementary ranks.

**Proposition 3.10.** *Let  $\mathcal{K}(\mathcal{A})$  be a homogeneous cone of rank  $r$  and let  $\mathcal{F} \trianglelefteq \text{cl}\mathcal{K}(\mathcal{A})$  be a face of rank  $s$ . Then,  $\mathcal{F}^\Delta = \mathcal{K}(\mathcal{A})^* \cap \mathcal{F}^\perp$  has rank  $r - s$ .*

*Proof.* Let  $x \in \text{ri}\mathcal{F}$ . By Theorem 3.5, there exists a triangular automorphism  $Q_{\hat{u}}$  of  $\mathcal{K}(\mathcal{A})$  that maps  $\mathcal{F}$  into a principal face  $\text{cl}\mathcal{K}(\mathcal{A}_I)$ . We may assume that  $Q_{\hat{u}}$  maps  $x$  to  $e_I$ , see the discussion around (3.5). Letting  $u := \hat{u}^{-1}$ ,  $Q_u = Q_{\hat{u}}^{-1}$  does the opposite: it maps  $e_I$  to  $x$ . Recalling that  $\mathcal{K}(\mathcal{A})^* \cap \mathcal{F}^\perp = \mathcal{K}(\mathcal{A})^* \cap \{x\}^\perp$ , we have by (2.12)

$$\mathcal{K}(\mathcal{A})^* \cap \{x\}^\perp = \{y \in \mathcal{K}(\mathcal{A})^* \mid \langle Q_u(e_I), y \rangle = 0\} = \{y \in \mathcal{K}(\mathcal{A})^* \mid \langle e_I, Q_{u^*}(y) \rangle = 0\}.$$

We have that  $e_I = \sum_{i \notin I} e_i$  and  $|I| = r - s$ .  $Q_{u^*}$  is a triangular automorphism of  $\mathcal{K}(\mathcal{A})^*$ , so  $Q_{u^*}(y) \in \mathcal{K}(\mathcal{A})^*$  if and only if  $y \in \mathcal{K}(\mathcal{A})^*$ . In addition, as the  $e_i$ 's are in  $\text{cl}\mathcal{K}(\mathcal{A})$ , for  $y \in \mathcal{K}(\mathcal{A})^*$  we have

$$\langle e_I, Q_{u^*}(y) \rangle = 0 \Leftrightarrow \langle e_i, Q_{u^*}(y) \rangle = 0, \forall i \notin I.$$

Therefore, from Proposition 3.1, for  $y \in \mathcal{K}(\mathcal{A})^*$ , we have  $\langle e_I, Q_{u^*}(y) \rangle = 0$  if and only if every row and column of  $Q_{u^*}(y)$  that is indexed by  $i$  not in  $I$  vanishes. This happens if and only if  $Q_{u^*}(y)$  is in the principal *dual* face  $\text{cl}(\mathcal{K}^D(\mathcal{A}_J))$ , where  $J := \{1, \dots, r\} \setminus I$ . In particular,

$$\mathcal{K}(\mathcal{A})^* \cap \{x\}^\perp = Q_{u^*}^{-1}(\text{cl}(\mathcal{K}^D(\mathcal{A}_J))).$$

That is,  $\mathcal{K}(\mathcal{A})^* \cap \{x\}^\perp$  is linearly isomorphic to a principal *dual* face of rank  $r - |J| = |I| = r - s$ , which concludes the proof.  $\square$

We note that the proof of Proposition 3.10 gives a recipe for determining the conjugate face  $\mathcal{F}^\Delta$ . If  $x \in \text{ri}\mathcal{F}$  and  $x = tt^*$  for a proper  $t$ , we let  $u \in \mathcal{T}_{++}$  be such that  $ue_I = t$ , where  $I := \{i \mid \rho_i(t_{ii}) = 0\}$ . With that,  $Q_{u^*}$  is an automorphism of  $\mathcal{K}(\mathcal{A})^*$  that maps  $\mathcal{F}$  to  $\text{cl}(\mathcal{K}^D(\mathcal{A}_J))$ , where  $J := \{i, \dots, r\} \setminus I$ .

## 4 Applications to homogeneous chordality

For this section, in contrast to Remark 2.2 and the previous sections, given a real matrix  $a \in M^{n \times n}$ , we return to the usual matrix convention where  $a_{ij}$  indicates the  $(i, j)$ -entry of  $a$ . Then, given a graph  $G$  with vertices  $V = \{1, \dots, n\}$  and edges  $E$ , we associate to it a convex cone  $\mathcal{S}_+(G) \subseteq \mathcal{S}_+^n$  given by

$$\mathcal{S}_+(G) := \{x \in \mathcal{S}_+^n \mid x_{ij} = 0 \text{ for all } i \neq j \text{ such that } \{i, j\} \notin E\}$$

and we denote by  $\mathcal{S}(G)$  the space spanned by  $\mathcal{S}_+(G)$ . If  $G$  is chordal, the cone  $\mathcal{S}_+(G)$  is generated by its rank one matrices, e.g., [AHMR88, Theorem 2.3]. When  $G$  not only is chordal but also does not contain induced subgraphs isomorphic to a path on 4 vertices, by [Ish13, Theorem A] they are also homogeneous and we will refer to such a graph  $G$  as being a *homogeneous chordal graph*. This class of cones was also extensively studied in [TV23].

The T-algebra underlying  $\mathcal{S}_+(G)$  is relatively simple, but the ordering of the vertices matter. Chordal homogeneous graphs have a very interesting ordering, called a *trivially perfect elimination ordering*. This means that the following two properties are satisfied.

- (o1) If  $\{i, j\}, \{i, k\} \in E$  and  $i < j < k$  hold, then  $\{j, k\} \in E$ .
- (o2) If  $\{i, j\}, \{j, k\} \in E$  and  $i < j < k$  hold, then  $\{i, k\} \in E$ .

See more details in [TV23, Section 2.3]. An ordering that only satisfies (o1) is called a *perfect elimination ordering* and all chordal graphs possess one. An ordering satisfying both (o1) and (o2) is a privilege of homogeneous chordality. In what follows, we will assume that the vertices are ordered following a trivially perfect elimination ordering.

Let  $\mathcal{A}(G) := \{a \in M^{n \times n} \mid a_{ij} = a_{ji} = 0 \text{ for all } i \neq j \text{ such that } \{i, j\} \notin E\}$ . We take the straightforward decomposition  $\mathcal{A}(G) = \bigoplus_{i,j=1}^r \mathcal{A}_{ij}$  where for  $\{i, j\} \in E$  or  $i = j$ ,  $\mathcal{A}_{ij} \cong \mathbb{R}$  is the subspace of  $M^{n \times n}$  of matrices that are zero outside the  $(i, j)$ -entry, i.e.,  $a \in \mathcal{A}_{ij}$  if and only if  $a_{kl} = 0$  for  $(k, l) \neq (i, j)$ . Otherwise, we set  $\mathcal{A}_{ij}$  to the zero subspace  $\{0\}$ . Overall,  $\mathcal{A}(G)$  is the set of  $n \times n$  matrices with the sparsity pattern prescribed by  $G$ . The involution is defined as the usual matrix transposition and the multiplication is defined as matrix multiplication, followed by the projection onto the sparsity pattern defined by  $G$ . We denote such a projection by  $\pi_G$  so that  $\pi_G(a)$  is the result of “zeroing out” the entries of  $a_{ij}$  for which  $\{i, j\} \notin E$  and  $i \neq j$ . With that, denoting the multiplication in  $\mathcal{A}(G)$  by  $\star$ , we have for  $a, b \in \mathcal{A}(G)$

$$a \star b = \pi_G(ab)$$

and the  $\mathcal{A}_{ij}$  satisfy (2.3) so it is indeed a bigradation of  $\mathcal{A}(G)$ .

To see that this is indeed a T-algebra we have to check the seven axioms. For the sake of clarity and preciseness, we recall once more that for  $a \in \mathcal{A}(G)$ ,  $a_{ij}$  denotes the  $(i, j)$ -entry of  $a$  rather than the element of  $\mathcal{A}_{ij}$  corresponding to  $a$ . This is a subtle, but luckily, unimportant distinction<sup>4</sup>.

Axioms (a1), (a2), (a3) and (a5) follow directly from the basic definitions and properties of the usual matrix multiplication, since the projection onto the sparsity pattern plays no role in them.

Axiom (a4) needs a little more thought. We have to check that for matrices  $a, b, c$  with the right sparsity pattern  $\text{tr}((a \star b) \star c) = \text{tr}(a \star (b \star c))$  holds. As  $\pi_G$  never zeroes diagonal entries, this is equivalent to checking that  $\text{tr}(\pi_G(ab)c) = \text{tr}(a\pi_G(bc))$  holds. First we prove  $\text{tr}(\pi_G(ab)c) = \text{tr}(abc)$  by showing that  $\pi_G(ab)c$  and  $abc$  have the same diagonal entries. Note that if  $[\pi_G(ab)]_{ij}$  is different from  $[ab]_{ij}$ , this means  $\{i, j\}$  is not in  $E$ , so  $c_{ji} = 0$ . Therefore

$$[\pi_G(ab)c]_{ii} = \sum_{j=1}^n [\pi_G(ab)]_{ij} c_{ji} = \sum_{j=1}^n [ab]_{ij} c_{ji} = [abc]_{ii}.$$

The same works for  $a\pi_G(bc)$  so we have axiom (a4).

For axioms (a6) and (a7) the ordering of the vertices is quite essential. Let us check that (o2) leads to (a6). Translating to the usual linear algebra language, (a6) amounts to requiring that for  $a \in \mathcal{A}_{ij}, b \in \mathcal{A}_{jk}, c \in \mathcal{A}_{kl}$  such that  $1 \leq i \leq j \leq k \leq l \leq r$  we have

$$a \star (b \star c) = (a \star b) \star c. \quad (4.1)$$

We start by observing that if, say,  $i = j$ , since  $b \star c \in \mathcal{A}_{jl}$  and  $b \in \mathcal{A}_{jk}$ , we would have  $a \star (b \star c) = a_{ii}(b \star c) = (a_{ii}b) \star c = (a \star b) \star c$ , by (a2). By similar arguments, (4.1) is true if  $j = k$  or  $k = l$ . Therefore, in order to check (a6), we may then assume that  $i < j < k < l$ . In this case, if  $\{i, j\}, \{j, k\}$  or  $\{k, l\}$  is not in  $E$ , then  $a, b$  or  $c$  would be zero, so both sides of (4.1) would be zero. But if they are all in  $E$  then by (o2) all the possible edges between  $\{i, j, k, l\}$  are in  $E$ , so the projection  $\pi_G$  will not have any effect and we simply have the associativity of matrix multiplication.

For axiom (a7) we proceed similarly. Let  $a \in \mathcal{A}_{ij}, b \in \mathcal{A}_{jk}, c \in \mathcal{A}_{lk}$  such that  $1 \leq i \leq j \leq k \leq r, 1 \leq l \leq k \leq r$  and we will check that

$$a \star (b \star c^*) = (a \star b) \star c^*, \quad (4.2)$$

which implies (a7). We start by noting that, analogously, if  $i = j, j = k$  or  $l = k$  holds, then the validity of (4.2) follows from (a2). So we consider the case where  $i < j < k$  and  $l < k$ . If  $\{i, j\}, \{j, k\}, \{l, k\}$  or  $\{i, l\}$  is not in  $E$  then both sides of (4.2) are zero, so we may assume that these edges are all in  $E$ . Then, (o2) implies that  $\{i, k\} \in E$ . We have three possibilities. If  $j = l$ , we already have all edges between  $\{i, l, k\} = \{i, j, k\}$  in  $E$ , so (4.2) is true because  $\pi_G$  does not zero any entries. The other two possibilities are  $i < j < l < k$  or  $i < l < j < k$ . In either case, since  $\{i, j\}, \{i, l\} \in E$ , we have  $\{j, l\}$  by (o1). and we conclude that all possible six edges between  $\{i, j, k, l\}$  are in  $E$ . Therefore, the projection  $\pi_G$  does not zero any elements in all the products appearing in (4.2) and we have the usual associativity of matrix multiplication, which implies (4.2).

We also observe that the inner product that comes with this T-algebra structure (see (2.6)) coincides with the usual matrix trace inner product since  $\text{tr}(a \star b^*) = \text{tr}(\pi_G(ab^*)) = \text{tr}(ab^*)$ .

Next we will verify that  $\mathcal{S}_+(G)$  coincides with the dual cone  $\mathcal{K}(\mathcal{A}(G))^*$ . For simplicity we write  $\mathcal{T}_{++}(G) := \mathcal{T}_{++}(\mathcal{A}(G)), \mathcal{T}_+(G) := \mathcal{T}_+(\mathcal{A}(G)), \mathcal{T}(G) := \mathcal{T}(\mathcal{A}(G)), \mathcal{L}_{++}(G) := \mathcal{T}_{++}(G)^*, \mathcal{L}_+(G) :=$

---

<sup>4</sup>After all, the  $(i, j)$ -entry of  $a \star b$  corresponds exactly to the  $\mathcal{A}_{ij}$  component of  $a \star b$  in the bigradation of  $\mathcal{A}(G)$ .

$\mathcal{T}_+(G)^*$  and  $\mathcal{L}(G) := \mathcal{T}(G)^*$ . In particular,  $\mathcal{L}_+(G)$  and  $\mathcal{T}_+(G)$  are, respectively, lower and upper triangular matrices with nonnegative diagonal and sparsity pattern determined by  $G$ .

The dual cone  $\mathcal{K}(\mathcal{A}(G))^*$  corresponds to elements of the form  $l \star l^*$  for  $l \in \mathcal{L}_+(G)$ . An interesting consequence of the ordering of the graph is that the projection is not needed. This is a known fact, but for the sake of completeness, we verify this. Let  $l \in \mathcal{L}(G)$  and  $i > j$ , then

$$(ll^*)_{ij} = \sum_{i \geq k \text{ and } j \geq k} l_{ik}l_{jk} = \sum_{j \geq k} l_{ik}l_{jk} = l_{ij}l_{jj} + \sum_{k=1}^{j-1} l_{ik}l_{jk}. \quad (4.3)$$

If  $\{i, j\} \in E$ , then  $(\pi_G(ll^*))_{ij} = (ll^*)_{ij}$ , so suppose that  $\{i, j\} \notin E$ . If there exists  $k$  such that  $k < j < i$  and  $l_{ik}$  and  $l_{jk}$  are both nonzero, then  $\{k, j\}, \{k, i\} \in E$ , which implies that  $\{i, j\} \in E$  by (o1), a contradiction. Therefore, for every  $k$  such that  $k < j < i$ , the term  $l_{ik}l_{jk}$  is zero. This implies that  $(ll^*)_{ij} = l_{ij}l_{jj} = 0 = (\pi_G(ll^*))_{ij}$  by (4.3) and the assumption that  $\{i, j\} \notin E$ . We conclude that, in fact,  $l \star l^* = ll^*$  holds for  $l \in \mathcal{L}_+(G)$ .

This tells us that  $\mathcal{K}(\mathcal{A}(G))^* = \{ll^* \mid l \in \mathcal{L}_+(G)\} \subseteq \mathcal{S}_+(G)$ . The converse and a summary of the discussion so far are given in the next theorem.

**Theorem 4.1** (T-algebra structure of homogeneous chordal cones). *Let  $G = (V, E)$  be a homogeneous chordal cone where the vertices are ordered following a trivially perfect elimination ordering. Then  $\mathcal{A}(G)$  is a T-algebra satisfying*

$$\mathcal{K}(\mathcal{A}(G))^* = \{ll^* \mid l \in \mathcal{L}_+(G)\} = \mathcal{S}_+(G)$$

and the following properties

- (i)  $l \star l^* = ll^*$ , for all  $l \in \mathcal{L}(G)$ .
- (ii)  $u \star t = ut$ , for all  $u, t \in \mathcal{T}(G)$ .
- (iii)  $l \star t = lt$ , for all  $l, t \in \mathcal{L}(G)$ .

*Proof.* It remains to verify the inclusion  $\mathcal{S}_+(G) \subseteq \{ll^* \mid l \in \mathcal{L}_+(G)\}$  and items (ii) and (iii). We start with the former. It is well-known that the extreme rays of  $\mathcal{S}_+(G)$  are generated by rank one matrices, e.g., [AHMR88, Theorem 2.3]. So, let  $v \in M^{n \times 1}$  (i.e., a column vector) be such that  $vv^*$  generates an extreme ray of  $\mathcal{S}_+(G)$ . Letting  $J := \{i \mid v_i \neq 0\}$ , we observe that for every pair  $i \neq j$ ,  $\{i, j\} \subseteq J$ , we must have  $\{i, j\} \in E$ , since  $(vv^*)_{ij} \neq 0$ . Let  $i$  be the smallest element of  $J$ . Replacing  $v$  with  $-v$  if necessary, we may assume that  $v_i$  is positive. Then, if we let  $l$  be the lower triangular matrix that has  $v$  as its  $i$ -column and is zero elsewhere, we have  $l \in \mathcal{L}_+(G)$ , since  $l_{ji} = v_j$  and  $\{j, i\} \in E$  whenever  $v_j \neq 0$  and  $i \neq j$ . We also have  $ll^* = vv^*$  which shows that  $\{ll^* \mid l \in \mathcal{L}_+(G)\}$  contains all the extreme rays of  $\mathcal{S}_+(G)$  and leads to the required inclusion.

For item (ii), let  $u, t \in \mathcal{T}(G)$ . We have that  $ut$  is also upper triangular, so  $(ut)_{ij} = 0$  if  $i > j$ . For  $i < j$  with  $\{i, j\} \notin E$ , we have

$$(ut)_{ij} = u_{ii}t_{ij} + u_{ij}t_{jj} + \sum_{k=i+1}^{j-1} u_{ik}t_{kj} = \sum_{k=i+1}^{j-1} u_{ik}t_{kj},$$

since  $t_{ij} = u_{ij} = 0$ . The indices of terms in the summation are such that  $i < k < j$  holds, so if  $u_{ik}$  or  $t_{kj}$  is nonzero we would have  $\{i, j\} \in E$  by (o2), a contradiction. Therefore,  $(ut)_{ij} = 0$  for  $i \neq j$ ,  $\{i, j\} \notin E$ , i.e.,  $u \star t = \pi_G(ut) = ut$ . This proves item (ii). Item (iii) then follows from item (ii) by taking adjoints.  $\square$

We note that we can also realize  $\mathcal{S}_+(G)$  as a “primal” homogeneous cone by constructing the T-Algebra  $\mathcal{B}(G)$  that is dual to  $\mathcal{A}(G)$  as in (2.7). From the ordering induced by  $\mathcal{B}(G)$ , matrices in  $\mathcal{S}_+(G)$  have upper triangular Cholesky decompositions.

Principal subalgebras of  $\mathcal{A}(G)$  are obtained by zeroing out some rows and columns of  $\mathcal{A}(G)$ . This corresponds to removing vertices of the underlying homogeneous chordal graph, an operation that in fact preserves homogeneous chordality. For  $H = (\hat{V}, \hat{E})$  an induced subgraph of  $G$ , we define

$$\mathcal{F}(H) := \{x \in \mathcal{S}_+(G) \mid x_{ii} = 0, i \notin \hat{V}\}$$

and, recalling Section 3.3, we observe that this coincides with the principal dual face  $\text{cl}(\mathcal{K}^D(\mathcal{A}(G)_I))$  induced by  $I = \{i \mid i \notin \hat{V}\}$ . Also,  $\mathcal{F}(H)$  (which is a cone of  $n \times n$  matrices) is linearly isomorphic to  $\mathcal{S}_+(H)$  (which is a cone of  $|\hat{V}| \times |\hat{V}|$  matrices). Theorems 3.5 and 3.9 then imply that every face of an homogeneous chordal cone  $\mathcal{S}_+(G)$  is isomorphic to a homogeneous chordal cone  $\mathcal{S}_+(H)$ , where  $H$  is an induced subgraph of  $G$ . In more details, we have the following theorem.

**Theorem 4.2.** *Let  $G = (V, E)$  be a homogeneous chordal graph and let  $\mathcal{F} \trianglelefteq \mathcal{S}_+(G)$ . Then,  $\mathcal{F}$  is projectionally exposed and there exists an automorphism  $Q$  of  $\mathcal{S}_+(G)$  that maps  $\mathcal{F}$  to  $\mathcal{F}(H)$ , where  $H$  is an induced subgraph of  $G$ . Additionally, if the vertices of  $G$  are in a trivially perfect elimination ordering the following items hold.*


- (i)  *$Q$  can be assumed to be of the form  $Q = Q_l$  for some  $l \in \mathcal{L}_{++}(G)$ . In particular  $Q_l(y) = lyl^*$  holds for  $y \in \mathcal{S}_+(G)$ .*
- (ii) *Suppose that  $x \in \text{ri } \mathcal{F}$  and  $x = tt^*$  holds for a proper matrix  $t \in \mathcal{L}_+(G)$ . Then, letting  $I = \{i \mid t_{ii} = 0\}$ ,  $H$  is the subgraph obtained by removing the vertices belonging to  $I$ . Furthermore,  $l$  in the previous item can be taken to be any  $l \in \mathcal{T}_{++}(G)$  satisfying  $lt = e_I$ , where  $e_I$  is the diagonal matrix that has 1 in its  $(i, i)$  component if  $i \notin I$  and zero elsewhere.*

*Proof.* Suppose that  $G$  follows a trivially perfect elimination ordering. By Theorem 4.1,  $\mathcal{A}(G)$  is a T-algebra and Theorem 3.9 tells us that  $\mathcal{F}$  is projectionally exposed and there is a triangular automorphism  $Q_l$  of  $\mathcal{K}(\mathcal{A}(G))^*$  that maps  $\mathcal{F}$  to a principal dual face of  $\mathcal{S}_+(G) = \mathcal{K}(\mathcal{A}(G))^*$ . Principal dual faces are obtained by zeroing out certain rows and columns, so each principal dual face corresponds to an induced subgraph  $H$  of  $G$ . Finally, Theorem 4.1 together with the dual version of (2.15) imply that for  $y = tt^* \in \mathcal{S}_+(G)$ , where  $t \in \mathcal{L}_+(G)$ , we have  $Q_l(t \star t^*) = Q_l(tt^*) = (lt)(lt)^* = ltt^*l^* = lyl^*$ , which proves item (i).

Item (ii) follows from Theorem 3.9 and the observation that, as discussed previously,  $\mathcal{F}(H)$  coincides with the principal dual face obtained by removing the vertices in  $I$ . The fact that  $l$  can be taken to be as in the statement follows from the discussion around (3.5) applied to the dual algebra and the existence of  $l$  follows from Lemma 3.4.

Finally, if  $G$  is in any other ordering, then  $\mathcal{S}_+(G)$  is linearly isomorphic to some  $\mathcal{S}_+(G')$  where  $G'$  is just a relabelling of  $G$  and is in a trivially perfect elimination ordering. Such a linear isomorphism is obtained by permuting rows and columns, so  $\mathcal{S}_+(G)$  still has the intended properties described in the statement.  $\square$


**An example** Let  $G$  be the following graph.



We note that the vertices follow a trivially perfect elimination ordering and that would not be the case if, say, the labels of 3 and 2 were exchanged. With that, we have

$$\mathcal{S}_+(G) = \left\{ \begin{pmatrix} x_{11} & 0 & x_{13} \\ 0 & x_{22} & x_{23} \\ x_{13} & x_{23} & x_{33} \end{pmatrix} \in \mathcal{S}(G) \mid \begin{pmatrix} x_{11} & 0 & x_{13} \\ 0 & x_{22} & x_{23} \\ x_{13} & x_{23} & x_{33} \end{pmatrix} \succeq 0 \right\}.$$

The nonzero proper principal faces  $\mathcal{S}_+(G)$  are the ones obtained from the following 6 induced subgraphs.



However, since  $\mathcal{S}_+(G)$  is not polyhedral, it has infinitely many faces that are not subfaces of the cones obtained by those induced subgraphs. For example, let  $x \in \mathcal{S}_+(G)$  be such that

$$x = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 2 \end{pmatrix}$$

and let us consider the problem of determining  $\mathcal{F} = \text{Face}(x, \mathcal{S}_+(G))$ , i.e., the minimal face of  $\mathcal{S}_+(G)$  containing  $x$ . Following Algorithm 2 we let

$$t := \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 1 & 0 \end{pmatrix}$$

and we may verify that, indeed,  $x = tt^*$  holds. We also have  $I = \{i \mid t_{ii} = 0\} = \{3\}$ , so according to Theorem 4.2,  $\mathcal{F}$  is isomorphic to  $\mathcal{F}(H)$ , where  $H$  is the induced subgraph obtained by removing the vertex with label 3. This automorphism can be obtained by finding  $l \in \mathcal{L}_{++}(G)$  such that

$$lt = e_I, \text{ where } e_I = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

One possible solution is

$$l = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -1 & -1 & 1 \end{pmatrix}.$$

Then,  $Q_l$  is an automorphism of  $\mathcal{S}_+(G)$  mapping  $x$  to  $(lt)(lt)^* = e_I e_I^* = e_I$ . Therefore,  $Q_l(\mathcal{F}) = Q_l(\text{Face}(x, \mathcal{S}_+(G))) = \text{Face}(e_I, \mathcal{S}_+(G)) = \mathcal{F}(H)$  is the principal dual face obtained by considering the subgraph induced by  $\{1, 2\}$ , i.e., zeroing out the third row and column of elements of  $\mathcal{S}_+(G)$ .

This leads to the following explicit description of  $\mathcal{F}$ .

$$\begin{aligned} \mathcal{F} &= \{x \in \mathcal{S}_+(G) \mid (Q_l(x))_{33} = 0\} = \{x \in \mathcal{S}_+(G) \mid (lxl^*)_{33} = 0\} \\ &= \{x \in \mathcal{S}_+(G) \mid x_{11} - 2x_{31} + x_{22} - 2x_{32} + x_{33} = 0\}. \end{aligned} \tag{4.4}$$

The face  $\mathcal{F}$  is also projectionally exposed and following the proof of item (i) of Theorem 3.5,  $\hat{l} = l^{-1}e_I l$  is a lower triangular matrix such that  $Q_{\hat{l}}$  maps  $\mathcal{S}_+(G)$  onto  $\mathcal{F}$  and fixes  $\mathcal{F}$ . We also note that  $Q_{\hat{l}}$  is not self-adjoint.



## 4.1 The PSD completion side

Let  $G = (V, E)$  be a homogeneous chordal graph. The dual cone of  $\mathcal{S}_+(G)$  is the cone of positive semidefinite (PSD) completable matrices that follow the sparsity pattern defined by  $G$ . That is,  $x \in \mathcal{S}_+(G)^*$  if and only if  $x \in \pi_G(\mathcal{S}_+^n)$ . A PSD completion of  $x$  is any  $w \in \mathcal{S}_+^n$  satisfying  $x = \pi_G(w)$ . While there is an extensive literature on completion problems, there seems to be few works that address the actual facial structure of  $\pi_G(\mathcal{S}_+^n)$ .

Now, suppose that the vertices are ordered according to a trivially perfect elimination ordering. Since  $\mathcal{S}_+(G)$  is  $\mathcal{K}(\mathcal{A}(G))^*$ , the cone  $\pi_G(\mathcal{S}_+^n) = \mathcal{S}_+(G)^*$  coincides with  $\text{cl}\mathcal{K}(\mathcal{A}(G))$ . Overall, we have

$$\pi_G(\mathcal{S}_+^n) = \{u \star u^* \mid u \in \mathcal{T}_+(G)\} = \{\pi_G(uu^*) \mid u \in \mathcal{T}_+(G)\}.$$

An element may have multiple PSD completions and it is of interest to obtain completions that have certain desired properties. For example, if there exists a positive definite completion to some  $x \in \pi_G(\mathcal{S}_+^n)$ , we may want to obtain one that maximizes the determinant.

In the next theorem, we gather several facts about the facial structure of  $\pi_G(\mathcal{S}_+^n)$  together with a proof that Algorithm 1 leads to a maximum rank decomposition. As before, if  $H = (\hat{V}, \hat{E})$  is an induced subgraph of  $G$ , then it also corresponds to a face of  $\pi_G(\mathcal{S}_+^n)$  in a natural way. By an abuse of notation, we define  $\pi_H(\mathcal{S}_+^n) := \{x \in \pi_G(\mathcal{S}_+^n) \mid x_{ij} = x_{ji} = 0, \forall i, j \notin \hat{V}\}$ . Put otherwise,  $x \in \pi_H(\mathcal{S}_+^n)$  if it has a PSD completion  $w$  satisfying  $w_{ii} = 0$  for  $i \notin \hat{V}$ .

**Theorem 4.3** (Faces of homogeneous PSD completable cones). *Let  $G = (V, E)$  be a homogeneous chordal cone and let  $\mathcal{F} \trianglelefteq \pi_G(\mathcal{S}_+^n)$ . Then,  $\mathcal{F}$  is projectively exposed and there exists an automorphism  $Q$  of  $\pi_G(\mathcal{S}_+^n)$  that maps  $\mathcal{F}$  to  $\pi_H(\mathcal{S}_+^n)$ , where  $H$  is an induced subgraph of  $G$ . Additionally, if the vertices of  $G$  are in a trivially perfect elimination ordering the following items hold.*

- (i)  $Q$  can be assumed to be of the form  $Q = Q_u$  for some  $u \in \mathcal{T}_{++}(G)$ . In particular  $Q_u(y) = \pi_G(uyu^*)$  holds for  $y \in \pi_G(\mathcal{S}_+^n)$ .
- (ii) Suppose that  $x \in \text{ri}\mathcal{F}$ . Then,  $x = \pi_G(tt^*)$  holds for an unique proper matrix  $t \in \mathcal{T}_+(G)$ . Then, letting  $I = \{i \mid t_{ii} = 0\}$ ,  $H$  is the subgraph obtained by removing the vertices belonging to  $I$ . Furthermore,  $u$  in the previous item can be taken to be any  $u \in \mathcal{T}_{++}(G)$  satisfying  $ut = e_I$ , where  $e_I$  is the diagonal matrix that has 1 in its  $(i, i)$  component if  $i \notin I$  and zero elsewhere.

In what follows, let  $x, t$  be as in item (ii) and let  $r := |\{i \mid t_i \neq 0\}|$ .

- (iii) The maximum possible rank of a PSD completion of  $x$  is  $r$  and  $tt^*$  is a maximum rank completion.
- (iv) If  $r = n$ , then  $tt^*$  is the maximum determinant completion.

*Proof.* As in the proof of Theorem 4.2, changing the ordering of the vertices of  $G$  amounts to permuting rows and columns, thus leading to linearly isomorphic cones. So we might as well assume that  $G$  follows a trivially perfect elimination ordering.

Both the fact that  $\mathcal{F}$  is projectively exposed and that there exists an automorphism mapping  $Q$  to some principal faces follow from Theorem 3.5. Under a trivially perfect elimination ordering,  $Q = Q_u$  for some  $u \in \mathcal{T}_{++}(G)$ . Then, if  $y \in \pi_G(\mathcal{S}_+^n)$  and  $y = t \star t^* = \pi_G(tt^*)$  for some  $t \in \mathcal{T}_{++}(G)$ , we have  $Q_u(y) = (ut) \star (ut)^* = \pi_G(utt^*u^*) = \pi_G(uyu^*)$ , which proves item (i).

Item (ii) is a direct consequence of the uniqueness of the generalized Cholesky factorization in Proposition 3.2 and item (ii) of Theorem 3.5.

For item (iii), we start by observing that  $\mathcal{F}$  is a homogeneous cone of rank  $r$ , also by item (ii) of Theorem 3.5. Then,  $\mathcal{F}^\Delta$  is a homogeneous cone of rank  $n - r$ , by Proposition 3.10. Let  $y \in \text{ri } \mathcal{F}^\Delta$  and write  $y = ll^*$ , where  $l$  is a proper lower triangular matrix in  $\mathcal{L}_+(G)$ . By Theorem 3.9, since  $\mathcal{F}^\Delta$  has rank  $n - r$ , the diagonal of  $l$  has exactly  $n - r$  positive elements, so  $y$  has rank  $n - r$ . Let  $w$  be a PSD completion of  $x$ , then

$$0 = \langle y, x \rangle = \langle y, \pi_G(w) \rangle = \langle \pi_G(y), w \rangle = \langle y, w \rangle,$$

where the last equality follows from  $y \in \mathcal{S}_+(G)$ . This immediately implies that the rank of  $w$  is at most  $r$ . Analogously, the diagonal of  $t$  has exactly  $r$  positive elements, so  $tt^*$  has rank  $r$  and it is a maximum rank completion of  $x$ .

For item (iv), since  $r = n$ , letting  $l := t^*$ , we note that  $l \in \mathcal{L}_{++}(G)$ . Let  $\bar{y} := l^{-1}(l^{-1})^*$  and  $\bar{w} := tt^*$ . In our T-algebra context,  $l^{-1}$  is the element in  $\mathcal{L}_{++}(G)$  satisfying  $l^{-1} \star l = e$ , where  $e$  is the identity matrix, see Section 2.1 and the discussion after (2.12). However, in view of item (iii) of Theorem 4.1,  $l^{-1} \star l = l^{-1}l = e$  holds, so  $l^{-1}$  is, in fact, the usual matrix inverse. Overall, we have  $(tt^*)^{-1} = l^{-1}(l^{-1})^*$ , i.e.,  $\bar{w}^{-1} = y$  holds in the usual linear algebraic sense. Also,  $y \in \text{ri } \mathcal{K}(\mathcal{A}(G))^* = \text{ri } \mathcal{S}_+(G)$  holds, since the diagonal of  $l^{-1}$  is positive.

We consider the following primal dual pair of problems.

$$\min_{w \in \mathcal{S}_+^n, \pi_G(w) = x} -\log \det(w) \quad \max_{y \in \mathcal{S}_+(G)} \langle x, y \rangle + \log \det(y) + r$$

The optimality conditions are  $\pi_G(w) = x$ ,  $w \in \text{ri } \mathcal{S}_+^n$ ,  $w^{-1} = y$ ,  $y \in \text{ri } \mathcal{S}_+(G)$ . Therefore,  $\bar{y}$  and  $\bar{w}$  are optimal solutions and  $\bar{w}$  is the maximum determinant completion.  $\square$

We remark that an analogous result to item (iv) was proved before in [TV23, Appendix B.4]. Also, Theorem 4.3 implies that  $x$  generates an extreme ray of  $\pi_G(\mathcal{S}_+^n)$  if and only if the maximum possible rank of PSD completions of  $x$  is one. As it may be of independent interest, we will check that this property also holds when  $G$  is only chordal.

**Proposition 4.4.** *Let  $G = (V, E)$  be a chordal graph,  $x \in \pi_G(\mathcal{S}_+^n)$  be nonzero and denote by  $r$  the maximum possible rank of a PSD completion of  $x$ . Then,  $x$  generates an extreme ray if and only if  $r = 1$ .*

*Proof.* First, suppose that  $x$  generates an extreme ray. Let  $\mathcal{F} := \text{Face}(x, \pi_G(\mathcal{S}_+^n))$ , i.e., the minimal face of  $\pi_G(\mathcal{S}_+^n)$  containing  $x$ . Let  $\mathcal{F}^\Delta = \mathcal{S}_+(G) \cap \{x\}^\perp$  be the conjugate face and  $y \in \text{ri } \mathcal{F}^\Delta$ . Then, if  $w$  is a PSD completion of  $x$  we have

$$0 = \langle x, y \rangle = \langle \pi_G(w), y \rangle = \langle w, \pi_G(y) \rangle = \langle w, y \rangle = 0.$$

Because  $w$  and  $y$  are PSD matrices, we have  $wy = yw = 0$  and, in particular,  $\ker w \supseteq \text{im } y$ . Also  $w \neq 0$ , since  $x \neq 0$ .

Since  $x$  generates an extreme ray,  $\mathcal{F}^\Delta$  is a maximum proper face of  $\mathcal{S}_+(G)$  by Lemma 2.1. Let  $y' \in \mathcal{S}_+(G)$  be such that it generates an extreme ray that is not in  $\mathcal{F}^\Delta$ . Since  $\mathcal{F}^\Delta$  is maximal, we have  $\text{Face}(y + y', \mathcal{S}_+(G)) = \mathcal{S}_+(G)$ , which implies that  $y + y'$  is positive definite and, in particular, has rank  $n$ . Because  $G$  is chordal, extreme rays correspond to rank one matrices ([AHMR88, Theorem 2.3]), so  $y'$  has rank one, which implies that  $y$  has rank  $n - 1$ .

Finally, since  $\ker w \supseteq \text{im } y$  holds, we have  $\text{rank } w \leq 1$ , which in view of  $w \neq 0$ , leads to  $\text{rank } w = 1$ . This tells us that  $r = 1$ .

For the converse, suppose that  $r = 1$  holds and  $x_1 + x_2 = x$ , for  $x_1, x_2 \in \pi_G(\mathcal{S}_+^n)$ . Let  $w_i$  be a PSD completion of  $x_i$ . Then  $w_1 + w_2$  is a PSD completion of  $x$ , so  $w_1 + w_2$  has rank one. This can only happen if  $w_1 = \alpha w_2$  for some  $\alpha \geq 0$ , which implies that  $x_1 = \alpha x_2$  and  $x_1, x_2$  are in the half-line generated by  $x$ , so  $x$  is an extreme ray.  $\square$

Beyond Proposition 4.4, we do not know if items (i) and (ii) of Theorem 4.3 have analogues for the case where  $G$  is only chordal.

Moving on, as discussed previously, faces of homogeneous cones are projectionally exposed but it may happen that the projection cannot be taken to be self-adjoint. We will observe here that if  $G$  is chordal, then  $\pi_G(\mathcal{S}_+^n)$  is orthogonally projectionally exposed under the trace inner product if and only if  $G$  is a disjoint union of cliques.

**Proposition 4.5.** *Let  $G = (V, E)$  be a chordal graph, then the following are equivalent.*

- (i)  $\pi_G(\mathcal{S}_+^n)$  is orthogonally projectionally exposed under the trace inner product.
- (ii)  $\pi_G(\mathcal{S}_+^n) = \mathcal{S}_+(G)$  (i.e.,  $\mathcal{S}_+(G)$  is self-dual)
- (iii)  $G$  is a disjoint union of cliques.

*Proof.* We start by observing that a pointed cone that is orthogonally projectionally exposed must be contained in its dual, e.g., [BLP87, Proposition 2.2]. Therefore, if  $\pi_G(\mathcal{S}_+^n)$  is orthogonally projectionally exposed, then  $\pi_G(\mathcal{S}_+^n) \subseteq \pi_G(\mathcal{S}_+^n)^* = \mathcal{S}_+(G)$ . However,  $\mathcal{S}_+(G) \subseteq \pi_G(\mathcal{S}_+^n)$  always holds, so we have  $\pi_G(\mathcal{S}_+^n) = \mathcal{S}_+(G)$ . This shows (i)  $\Rightarrow$  (ii).

If  $\pi_G(\mathcal{S}_+^n) = \mathcal{S}_+(G)$ , then, since  $G$  is chordal,  $\pi_G(\mathcal{S}_+^n)$  is generated by PSD rank one matrices and  $G$  must be a disjoint union of cliques by [GIL24, Theorem 4.2]. This shows (ii)  $\Rightarrow$  (iii).

Finally, if  $G$  is a disjoint union of cliques then, up to permutations of rows and columns,  $\pi_G(\mathcal{S}_+^n)$  is a direct product of positive semidefinite cones, so it is orthogonally projectionally exposed under the trace inner product, which shows the implication (iii)  $\Rightarrow$  (i).  $\square$

Proposition 4.5 does not exclude the possibility that some homogeneous chordal  $G$  that is not a disjoint union of cliques be such that  $\pi_G(\mathcal{S}_+^n)$  is orthogonally projectionally exposed *under a different inner product*. Also, we mention in passing that the implication (ii)  $\Rightarrow$  (i) is a special case of a more general phenomenon, see [GL23, Proposition 4.18].

## 5 Conclusion and open questions

In this paper our main goal was to elucidate the facial structure of homogeneous cones under the T-algebra framework and discuss applications to homogeneous chordality.

Here we briefly describe a potential practical application. When solving a semidefinite program that fails Slater's condition, one of the default ways of restoring constraint qualifications and reducing the size of the problem is through facial reduction [BW81, Pat13b, WM13]. This is done by reformulating the problem over a smaller face of the cone, which leads to a smaller SDP. The reformulation typically use the projectional exposedness of  $\mathcal{S}_+^n$  and expressions such as (1.1). Unfortunately, this often destroys sparsity if  $q$  is as in (1.1), since  $qxq^*$  does not necessarily satisfy any sparsity pattern present in  $x$ . Therefore, one must face the inevitable choice of either keeping sparsity or regularizing the problem. From a theoretical point of view, the issue is that the map  $x \mapsto qxq^*$  is an automorphism of  $\mathcal{S}_+^n$  but not necessarily an automorphism of the PSD cone that corresponds to the sparsity pattern of the problem.

However, if the sparsity pattern is homogeneous chordal, we have the tantalizing option of both regularizing the problem *and* preserving sparsity. If the feasible region of some SDP is contained in a proper face of a  $\mathcal{S}_+(G)$  as in Theorem 4.2, we can use an automorphism of  $\mathcal{S}_+(G)$  to reformulate the problem over a principal face, which is an operation that indeed preserves sparsity. Furthermore, since principal faces correspond to zeroing rows and columns, we can further reformulate the problem as a smaller dimensional SDP. For example, if the feasible region is contained in the

face  $\mathcal{F}$  as in (4.4), we can reformulate the problem as a  $2 \times 2$  SDP that still respect the same sparsity pattern by making use of the map  $Q_l$  that takes  $\mathcal{F}$  to the corresponding rank 2 principal face. Similar considerations apply to problems over  $\pi_G(\mathcal{S}_+^n)$ , in view of Theorem 4.3. It would be interesting to explore these possibilities in future works.

We conclude with some open questions.

- (a) *Suppose that  $G = (V, E)$  is chordal but not necessarily homogeneous. Are  $\mathcal{S}_+(G)$  and  $\pi_G(\mathcal{S}_+^n)$  projectively exposed? Do analogues of Theorem 4.2 and Theorem 4.3 hold?* For example, it would be interesting to check whether, assuming only chordality, the faces of  $\mathcal{S}_+(G)$  are linearly isomorphic to the finitely many faces that arise by considering induced subgraphs of  $G$  and if such an isomorphism can be realized through an automorphism of  $\mathcal{S}_+(G)$ . The smallest example for which we do not know the answer to these questions is when  $G$  is a path on four vertices. Analogously to the previous discussion, a positive answer would have interesting implications for facial reduction approaches.
- (b) We proved homogeneous cones are projectively exposed but they may fail to be orthogonally projectively exposed *under the inner product that comes from the  $T$ -algebra structure*, see Proposition 4.5. We do not know whether an arbitrary homogeneous cone can become orthogonally projectively exposed by changing the inner product appropriately. Currently, the only homogeneous cones known to be orthogonally projectively exposed are symmetric cones [Lou21, Proposition 33], are those the only ones?
- (c) Homogeneous cones are spectrahedral and, therefore, are hyperbolicity cones as well. These are two classes of cones that strictly contain homogeneous cones. Are they projectively exposed?

Regarding (b), this problem is related to finding nontrivial conditions that ensure that a homogeneous cone is actually symmetric, e.g., [YN16]. Related to that, motivated by certain considerations in quantum physics that are out-of-scope here, it was recently shown that if the automorphism group of a homogeneous cone acts transitively on the set of extreme rays, then it must be a symmetric cone [BUvdW23, Theorem 2].

## Acknowledgements

The support of the Institute of Statistical Mathematics in the form of a visiting professorship for the first author is gratefully acknowledged. We also thank Hideyuki Ishi for several helpful comments.

## References

- [AHMR88] Jim Agler, William Helton, Scott McCullough, and Leiba Rodman. Positive semidefinite matrices with a given sparsity pattern. *Linear algebra and its Applications*, 107:101–149, 1988.
- [Bar73] George Phillip Barker. The lattice of faces of a finite dimensional cone. *Linear Algebra and its Applications*, 7(1):71–82, 1973.
- [BC75] George Phillip Barker and David Carlson. Cones of diagonally dominant matrices. *Pacific Journal of Mathematics*, 57(1):15–32, 1975.
- [BLP87] George Phillip Barker, Michael Laidacker, and George Poole. Projectively exposed cones. *SIAM Journal on Algebraic Discrete Methods*, 8(1):100–105, 1987.

- [BUvdW23] Howard Barnum, Cozmin Ududec, and John van de Wetering. Self-duality and Jordan structure of quantum theory follow from homogeneity and pure transitivity. *ArXiv e-prints*, 2023.
- [BW81] Jonathan M. Borwein and Henry Wolkowicz. Regularizing the abstract convex program. *Journal of Mathematical Analysis and Applications*, 83(2):495 – 530, 1981.
- [Chu03] Chek Beng Chua. Relating homogeneous cones and positive definite cones via  $T$ -algebras. *SIAM Journal on Optimization*, 14(2):500–506, 2003.
- [Chu09] Chek Beng Chua. A  $T$ -algebraic approach to primal-dual interior-point algorithms. *SIAM Journal on Optimization*, 20(1):503–523, 2009.
- [CT06] Chek Beng Chua and Levent Tunçel. Invariance and efficiency of convex representations. *Mathematical Programming*, 111(1-2):113–140, 2006.
- [FK94] Jacques Faraut and Adam Korányi. *Analysis on symmetric cones*. Oxford mathematical monographs. Clarendon Press, Oxford, 1994.
- [FKMN01] Mitsuhiro Fukuda, Masakazu Kojima, Kazuo Murota, and Kazuhide Nakata. Exploiting sparsity in semidefinite programming via matrix completion I: General framework. *SIAM Journal on Optimization*, 11(3):647–674, 2001.
- [G96] Osman Güler. Barrier functions in interior point methods. *Mathematics of Operations Research*, 21(4):860–885, 1996.
- [Går59] Lars Gårding. An inequality for hyperbolic polynomials. *Journal of Mathematics and Mechanics*, pages 957–965, 1959.
- [GIL24] João Gouveia, Masaru Ito, and Bruno F. Lourenço. Minimal hyperbolic polynomials and ranks of homogeneous cones. *Arxiv e-prints*, 2024.
- [Gin92] Simon Gindikin. *Tube domains and the Cauchy problem*. American Mathematical Society, 1992.
- [GL23] João Gouveia and Bruno F. Lourenço. Self-dual polyhedral cones and their slack matrices. *SIAM Journal on Matrix Analysis and Applications*, 44(3):1096–1121, 2023.
- [GT98] Osman Güler and Levent Tunçel. Characterization of the barrier parameter of homogeneous convex cones. *Mathematical Programming*, 81(1):55–76, 1998.
- [Gül97] Osman Güler. Hyperbolic polynomials and interior point methods for convex programming. *Mathematics of Operations Research*, 22(2):350–377, 1997.
- [IL19] Masaru Ito and Bruno F. Lourenço. The automorphism group and the non-self-duality of  $p$ -cones. *Journal of Mathematical Analysis and Applications*, 471(1–2):392–410, 2019.
- [Ish00] Hideyuki Ishi. Positive Riesz distributions on homogeneous cones. *Journal of the Mathematical Society of Japan*, 52(1), 2000.
- [Ish01] Hideyuki Ishi. Basic relative invariants associated to homogeneous cones and applications. *Journal of Lie Theory*, 11(1):155–171, 2001.

- [Ish13] Hideyuki Ishi. On a class of homogeneous cones consisting of real symmetric matrices. *Josai mathematical monographs*, 6:71–80, 2013.
- [Koe99] Max Koecher. *The Minnesota Notes on Jordan Algebras and Their Applications*. Number 1710 in Lecture Notes in Mathematics. Springer, Berlin, 1999.
- [KTX12] Lingchen Kong, Levent Tunçel, and Naihua Xiu. Existence and uniqueness of solutions for homogeneous cone complementarity problems. *Journal of Optimization Theory and Applications*, 153(2):357–376, 2012.
- [Lou21] Bruno F. Lourenço. Amenable cones: error bounds without constraint qualifications. *Mathematical Programming*, 186:1–48, 2021.
- [LP18] Minghui Liu and Gábor Pataki. Exact duals and short certificates of infeasibility and weak infeasibility in conic linear programming. *Mathematical Programming*, 167(2, Ser. A):435–480, 2018.
- [LRS22] Bruno F. Lourenço, Vera Roshchina, and James Saunderson. Amenable cones are particularly nice. *SIAM Journal on Optimization*, 32(3):2347–2375, 2022.
- [LRS23] Bruno F. Lourenço, Vera Roshchina, and James Saunderson. Hyperbolicity cones are amenable. *Mathematical Programming*, 204(1–2):753–764, 2023.
- [Nak14] Hideto Nakashima. Basic relative invariants of homogeneous cones. *Journal of Lie Theory*, 24(4):1013–1032, 2014.
- [Nak18] Hideto Nakashima. Basic relative invariants of homogeneous cones and their Laplace transforms. *Journal of the Mathematical Society of Japan*, 70(1), 2018.
- [Nak24] Hideto Nakashima. An example of homogeneous cones whose basic relative invariant has maximal degree. *arXiv e-prints*, 2024.
- [NFF<sup>+</sup>03] Kazuhide Nakata, Katsuki Fujisawa, Mitsuhiro Fukuda, Masakazu Kojima, and Kazuo Murota. Exploiting sparsity in semidefinite programming via matrix completion II: implementation and numerical results. *Mathematical Programming*, 95:303–327, 2003.
- [Pat00] Gábor Pataki. The geometry of semidefinite programming. In Henry Wolkowicz, Romesh Saigal, and Lieven Vandenbergh, editors, *Handbook of semidefinite programming: theory, algorithms, and applications*. Kluwer Academic Publishers, 2000.
- [Pat07] Gábor Pataki. On the closedness of the linear image of a closed convex cone. *Math. Oper. Res.*, 32(2):395–412, 2007.
- [Pat13a] Gábor Pataki. On the connection of facially exposed and nice cones. *Journal of Mathematical Analysis and Applications*, 400(1):211–221, 2013.
- [Pat13b] Gábor Pataki. Strong duality in conic linear programming: Facial reduction and extended duals. In *Computational and Analytical Mathematics*, volume 50 of *Springer Proc. in Math. Stat.*, pages 613–634. Springer New York, 2013.
- [PL88] George D. Poole and Michael Laidacker. Projectionally exposed cones in  $R^3$ . *Linear Algebra Appl.*, 111:183 – 190, 1988.

- [Ren06] James Renegar. Hyperbolic programs, and their derivative relaxations. *Foundations of Computational Mathematics*, 6(1):59–79, 2006.
- [Roc97] Ralph T. Rockafellar. *Convex Analysis*. Princeton University Press, 1997.
- [Ros14] Vera Roshchina. Facially exposed cones are not always nice. *SIAM Journal on Optimization*, 24(1):257–268, 2014.
- [Rot66] Oscar S. Rothaus. The construction of homogeneous convex cones. *Annals of Mathematics*, 83(2):358–376, 1966. Correction: *ibid* 87 (1968), 399.
- [Sau20] James Saunderson. Limitations on the expressive power of convex cones without long chains of faces. *SIAM Journal on Optimization*, 30(1):1033–1047, 2020.
- [ST90] Chen-Han Sung and Bit-Shun Tam. A study of projectionally exposed cones. *Linear Algebra and its Applications*, 139:225 – 252, 1990.
- [Tam85] Bit-Shun Tam. On the duality operator of a convex cone. *Linear Algebra and its Applications*, 64:33–56, 1985.
- [TT03] Van Anh Truong and Levent Tunçel. Geometry of homogeneous convex cones, duality mapping, and optimal self-concordant barriers. *Mathematical Programming*, 100(2):295–316, 2003.
- [TV23] Levent Tunçel and Lieven Vandenbergh. Linear optimization over homogeneous matrix cones. *Acta Numerica*, 32:675–747, 2023.
- [TX01] Levent Tunçel and Song Xu. On homogeneous convex cones, the Carathéodory number, and the duality mapping. *Math. Oper. Res.*, 26(2):234–247, 2001.
- [Vin63] Ernest B. Vinberg. The theory of homogeneous convex cones. *Transactions of the Moscow Mathematical Society*, 12:340–403, 1963. (English Translation).
- [WM13] Hayato Waki and Masakazu Muramatsu. Facial reduction algorithms for conic optimization problems. *Journal of Optimization Theory and Applications*, 158(1):188–215, 2013.
- [YN15] Takashi Yamasaki and Takaaki Nomura. Realization of homogeneous cones through oriented graphs. *Kyushu Journal of Mathematics*, 69(1):11–48, 2015.
- [YN16] Takashi Yamasaki and Takaaki Nomura. A characterization of symmetric cones by the degrees of basic relative invariants. *Kyushu Journal of Mathematics*, 70(2):237–257, 2016.