

# Objective-Function Free Multi-Objective Optimization: Rate of Convergence and Performance of an Adagrad-like algorithm

Marianna De Santis\*      Gabriele Eichfelder†      Margherita Porcelli‡

February 2, 2026

## Abstract

We propose an Adagrad-like algorithm for multi-objective unconstrained optimization that relies on the computation of a common descent direction only. Unlike classical local algorithms for multi-objective optimization, our approach does not rely on the dominance property to accept new iterates, which allows for a flexible and function-free optimization framework. New points are obtained using an adaptive stepsize that does not require neither knowledge of Lipschitz constants nor the use of line-search procedures. The rate of convergence is analyzed and is shown to be  $\mathcal{O}(1/\sqrt{k+1})$ . The method is extensively validated on a broad class of unconstrained multi-objective problems and simple multi-task learning instances, and compared against a first-order line-search algorithm. Additionally, we present a preliminary study of the behavior under noisy multi-objective settings, highlighting the robustness of the method.

1

## 1 Introduction

Recently, a class of algorithms for unconstrained nonconvex optimization has been proposed where the value of the (single) objective function is never computed. This class, referred to as Objective-Function-Free Optimization (OFFO) algorithms, has recently been very popular in the context of noisy problems, in particular in deep learning applications (see [9, 24] among many others), where they have shown remarkable insensitivity to the noise level. Also, in these applications evaluating the loss function often requires looking at millions of data points, making its repeated computation per iteration often prohibitive. This context is particularly suitable for the OFFO framework. At the same time, stochastic multi-objective optimization problems

---

\*Dipartimento di Ingegneria dell'Informazione, Università degli Studi di Firenze, Via di Santa Marta 3, 50139 Florence, Italy ([marianna.desantis@unifi.it](mailto:marianna.desantis@unifi.it)), ORCID 0000-0002-1189-5917.

†Institute of Mathematics, Technische Universität Ilmenau, Po 10 05 65, D-98684 Ilmenau, Germany ([gabriele.eichfelder@tu-ilmenau.de](mailto:gabriele.eichfelder@tu-ilmenau.de)), ORCID 0000-0002-1938-6316.

‡Dipartimento di Ingegneria Industriale, Università degli Studi di Firenze, Viale Morgagni 40/44, 50134 Firenze, Italy, ISTI-CNR, Italy and member of the INdAM Research Group GNCS ([margherita.porcelli@unifi.it](mailto:margherita.porcelli@unifi.it)) ORCID 0000-0003-0183-1204.

<sup>1</sup>During the submission of this manuscript, we learned of independent work by Gonçalves, Grapiglia and Melo [16] posted on Optimization Online on January 30th, 2026, which proposes a similar Adagrad-Norm algorithm for multi-objective optimization. Nevertheless, the theoretical analysis of the method and the numerical experience are different.

involving both noise and multiple objectives found several applications in many fields such as finance, energy, transportation, logistics, and supply chain management, see e.g. the survey [22] and references therein.<sup>2</sup>

OFFO contains several well-known algorithms such as Adagrad [9, 30], Adam [24], RM-Sprop [35], ADADELTA [39] which have been proposed and widely used in the machine learning community, emerging as state-of-the-art techniques to train neural networks. All these methods only rely on current and past gradient information to adaptively determine the step size at each iteration. The OFFO algorithms have been introduced to unify and to extend the complexity and convergence theory of many algorithms for non convex problems by using a trust-region framework [18, 17]. Indeed, the work [18] re-interpreted the deterministic Adagrad as a general trust-region method in infinity norm where, unlike standard methods, no objective is evaluated to measure the progress towards a minimizer and to enforce descent. These methods have been extended to several class of problems including constrained optimization [1, 2, 19], multilevel optimization [20] and neural network pruning applications [31], including both deterministic and stochastic approaches.

In this work, we introduce the further challenge of optimizing multiple objective functions without evaluating any element function. This framework is particularly motivated by multi-task learning in neural networks, where a single model must simultaneously optimize multiple task-specific losses, see e.g. [3, 28].

The majority of local algorithms for multi-objective optimization use the fact that if a point does not satisfy suitable necessary optimality conditions, then it can be easily improved, in the sense that a new point can be easily defined, with respect to a specific quality measure, which is usually the objective functions value. Most of algorithms proposed in this respect, both for unconstrained and constrained multi-objective problems, extend the classical iterative scalar optimization algorithms, such as the steepest descent [11], Newton [10, 15], external penalty [13], interior point [14], just to name a few. The mentioned approaches produce sequences of points able to converge to single Pareto critical points. In particular, at every iteration, these algorithms look for a new improved point, that is, a point that dominates the previous one. Designing an algorithm that does not rely on the dominance property to accept new points is then a particular challenge in the context of multi-objective optimization.

Our framework is inspired by the OFFO work [18] by leveraging on the Adagrad-Norm algorithm [38] for deterministic non-convex optimization that relies on  $\ell_2$ -norm trust-region subproblems. Our contribution can be summarized as follows:

1. We develop an adaptive strategy that uses a first-order common descent direction for which we provide a through theoretical characterization. The adaptive stepsize rule relies on the current and past common descent directions.
2. No knowledge of the Lipschitz constants neither the employment of a line-search procedure (cfr [28]) is needed to define the adaptive stepsize.
3. We prove the convergence of the method with global convergence rate of  $\mathcal{O}(1/\sqrt{k+1})$ .
4. We extensively validate our method on a large class of purely unconstrained multi-objective problems and on simple multitask problems, in comparison with a first-order line-search algorithm.
5. We provide a preliminary study on the behaviour of both our method and the line-search based one on noisy multi-objective problems.

---

<sup>2</sup>Although noisy problems are a relevant motivation for our proposal, we consider here deterministic (noiseless) algorithms which are crucial to understand the behavior of stochastic ones, and provide some numerical experience on noisy multi-objective problems.

The paper is organized as follows. In Section 2, we review basic results on multi-objective optimization and introduce the notation used throughout the manuscript. Section 3 presents the **MO-Adagrad** algorithm and its theoretical analysis, including results on the common descent direction and a global convergence rate of  $\mathcal{O}(1/\sqrt{k+1})$ . In Section 4, we report our numerical experience. Finally, Section 5 concludes the paper.

## 2 Notation and Basics of Multi-objective Optimization

We consider the unconstrained multi-objective optimization problem

$$\min_{x \in \mathbb{R}^n} f(x) \quad (\text{MOP})$$

with  $f: \mathbb{R}^n \rightarrow \mathbb{R}^m$ , and each  $f_j: \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $j \in \{1, \dots, m\}$  being continuously differentiable. We use for any  $m \in \mathbb{N}$  the set of indices  $[m] := \{1, \dots, m\}$ . We denote the gradient of one of the functions  $f_j$  in some point  $x \in \mathbb{R}^n$  by  $\nabla f_j(x)$ . Moreover, we assume that the multi-objective optimization problem (MOP) has at least one weakly efficient point. Recall that a point  $\bar{x} \in \mathbb{R}^n$  is called efficient for (MOP) if for any  $x \in \mathbb{R}^n$  the conditions  $f_j(x) \leq f_j(\bar{x})$  for all  $j \in [m]$  already imply  $f(x) = f(\bar{x})$ , and it is called weakly efficient in case there is no  $x \in \mathbb{R}^n$  with  $f_j(x) < f_j(\bar{x})$  for all  $j \in [m]$ . For vectors  $y^1, y^2 \in \mathbb{R}^m$  the inequalities  $y^1 \leq y^2$  and  $y^1 < y^2$  are understood componentwise.

Analogously to single-objective optimization, one can define local concepts, like locally weakly efficient and locally efficient points: a point  $\bar{x} \in \mathbb{R}^n$  is called locally efficient for (MOP) if there exists a neighborhood  $\mathcal{N}(\bar{x})$  of  $\bar{x}$  such that for any  $x \in \mathcal{N}(\bar{x})$  the conditions  $f_j(x) \leq f_j(\bar{x})$  for all  $j \in [m]$  already imply  $f(x) = f(\bar{x})$ , and it is called locally weakly efficient in case there exists a neighborhood  $\mathcal{N}(\bar{x})$  of  $\bar{x}$  such that there is no  $x \in \mathcal{N}(\bar{x})$  with  $f_j(x) < f_j(\bar{x})$  for all  $j \in [m]$ .

Additionally, we assume, as it is often done in the literature for descent-type algorithms for multi-objective optimization, that the gradients of the functions  $f_j$ ,  $j \in [m]$  are Lipschitz continuous, i.e., for each  $j \in [m]$  there exists a Lipschitz constant  $L_j > 0$  with

$$\|\nabla f_j(x) - \nabla f_j(x')\|_2 \leq L_j \|x - x'\|_2 \quad \text{for all } x, x' \in \mathbb{R}^n. \quad (1)$$

We will make use of

$$L_{\max} := \max_{j \in [m]} L_j, \quad (2)$$

which denotes the maximum among the Lipschitz constants of the gradients  $\nabla f_j$ ,  $j \in [m]$ .

In our proofs, the function  $\Phi: \mathbb{R}^n \rightarrow \mathbb{R}$  will play an important role, which we define by

$$\Phi(x) = \max_{j \in [m]} f_j(x). \quad (3)$$

This function is often used in the context of multi-objective trust region methods [37]. There, for accepting a candidate point as a new iteration point one requires that there is a sufficient decrease for this function  $\Phi$ . This is a weaker criterion compared to requiring that a sufficient decrease is reached in each objective function  $f_j$ . This function is bounded from below under our assumptions:

**Lemma 1.** *Under our assumption that (MOP) has a weakly efficient point  $\bar{x} \in \mathbb{R}^n$  there exists  $\Phi_{\text{low}} \in \mathbb{R}$  such that*

$$\inf_{x \in \mathbb{R}^n} \Phi(x) \geq \Phi_{\text{low}} > -\infty.$$

*Proof.* Define  $\Phi_{\text{low}} := \min_{j \in [m]} f_j(\bar{x})$  and assume there exists a sequence  $(x^k)_{k \in \mathbb{N}} \subseteq \mathbb{R}^n$  with  $\lim_{k \rightarrow \infty} \Phi(x^k) = -\infty$ , i.e., we have for all  $j \in [m]$  that  $\lim_{k \rightarrow \infty} f_j(x^k) = -\infty$ . Then there exists  $N \in \mathbb{N}$  such that for all  $j \in [m]$  we have  $f_j(x^N) < \Phi_{\text{low}}$  and thus  $f_j(x^N) < f_j(\bar{x})$ , in contradiction to  $\bar{x}$  weakly efficient for (MOP).  $\square$

In this work,  $\Phi$  will be used for deriving the convergence analysis, but will never be used in the proposed algorithm.

A necessary condition for a point  $x \in \mathbb{R}^n$  to be weakly efficient for (MOP) is that  $x$  is a Pareto critical point for (MOP) as defined next. The necessary and sufficient optimality conditions will be formulated in Lemma 6.

**Definition 2.** We call  $x \in \mathbb{R}^n$  a Pareto critical point for (MOP) if there are scalars  $\lambda_j \geq 0$ ,  $j \in [m]$  with  $\sum_{j=1}^m \lambda_j = 1$  and

$$\sum_{j=1}^m \lambda_j \nabla f_j(x) = 0.$$

Within numerical algorithms, the following function  $\omega$  is often used.

**Definition 3.** We define the criticality measure  $\omega: \mathbb{R}^n \rightarrow \mathbb{R}_+$  as the function which associates to each  $x \in \mathbb{R}^n$  the optimal value of

$$\begin{aligned} \min_{\lambda \in \mathbb{R}^m} \quad & \left\| \sum_{j=1}^m \lambda_j \nabla f_j(x) \right\|_2^2 \\ \text{s.t.} \quad & \sum_{j=1}^m \lambda_j = 1, \quad \lambda \geq 0. \end{aligned} \tag{\Omega(x)}$$

The following result is easy to see:

**Lemma 4.** A point  $x \in \mathbb{R}^n$  is Pareto critical for (MOP) if and only if  $\omega(x) = 0$ .

In the literature, one can often find the following characterization for Pareto criticality of a point  $x \in \mathbb{R}^n$ , see [11, Def. 2.2].

$$\forall d \in \mathbb{R}^n \exists j \in [m]: \nabla f_j(x)^\top d \geq 0. \tag{4}$$

This condition is equivalent to that the optimal value of the optimization problem

$$\min_{\|d\|_2^2 \leq 1} \max_{j \in [m]} \nabla f_j(x)^\top d \tag{mgrad(x)}$$

is zero with optimal solution  $d = 0$ , see [11, Lemma 3]. The problem (mgrad(x)) was, for instance, used in [33, 23]. We show that this characterization is in fact equivalent to our definition above. While this is somewhat widely known, we give a formal proof below.

**Lemma 5.** A point  $x \in \mathbb{R}^n$  is a Pareto critical point for (MOP) if and only if (4) holds.

*Proof.* As we could not find a proof in the easy accessible literature of this well-known result, we give it here for completeness. A proof using Gordon's Theorem can be found in the Master's thesis [32]. First, assume that (4) holds, i.e.,  $d = 0$  is an optimal solution of (mgrad(x)) and thus  $(0, 0)$  is an optimal solution of

$$\begin{aligned} \min_{(t,d) \in \mathbb{R}^{1+n}} \quad & t \\ \text{s.t.} \quad & \nabla f_j(x^k)^\top d - t \leq 0, \quad \forall j \in [m], \\ & \|d\|_2^2 \leq 1. \end{aligned} \tag{mgrad'(x)}$$

For  $(\text{mgrad}'(\mathbf{x}))$ , the Slater constraint qualification holds, and thus the optimal solution has to satisfy the KKT optimality conditions. Note that the constraint  $\|d\|_2^2 \leq 1$  is not active in the optimal solution while all the other constraints are active. Thus, there exist multipliers  $\lambda_j \geq 0$ ,  $j \in [m]$ , with

$$1 + \sum_{j=1}^m \lambda_j(-1) = 0 \quad \text{and} \quad 0 + \sum_{j=1}^m \lambda_j \nabla f_j(\mathbf{x}) = 0 \quad (5)$$

and, as a consequence,  $\mathbf{x}$  is a Pareto critical point for  $(\text{MOP})$ . On the other hand, if  $\mathbf{x}$  is Pareto critical for  $(\text{MOP})$ , then the conditions (5) are satisfied for some  $\lambda_j \geq 0$ ,  $j \in [m]$ . Moreover, for  $(0, 0)$  all constraints of the form  $\nabla f_j(\mathbf{x})^\top d - t \leq 0$  are active, and thus the complementarity conditions are satisfied. As  $(\text{mgrad}'(\mathbf{x}))$  is a convex optimization problem, this is sufficient for  $(0, 0)$  being an optimal solution of  $(\text{mgrad}'(\mathbf{x}))$ , and thus that (4) is satisfied.  $\square$

As a consequence, the optimal value of  $(\text{mgrad}(\mathbf{x}))$  is non-positive, and zero if and only if  $\mathbf{x} \in \mathbb{R}^n$  is Pareto critical for  $(\text{MOP})$ . Moreover, by Lemma 5 and, for instance, [10, Theorem 3.1], we have the following necessary and sufficient optimality condition:

**Lemma 6.** *If a point  $\mathbf{x} \in \mathbb{R}^n$  is locally weakly efficient for  $(\text{MOP})$ , then  $\mathbf{x}$  is Pareto critical. If the functions  $f_j$ ,  $j \in [m]$  are convex and  $\mathbf{x} \in \mathbb{R}^n$  is Pareto critical for  $(\text{MOP})$ , then  $\mathbf{x}$  is weakly efficient for  $(\text{MOP})$ . If the functions  $f_j$ ,  $j \in [m]$  are even strictly convex and  $\mathbf{x} \in \mathbb{R}^n$  is Pareto critical for  $(\text{MOP})$ , then  $\mathbf{x}$  is efficient for  $(\text{MOP})$ .*

**Further useful notations.** The notation  $v_{i,k}$  denotes the  $i$ -th component of a vector  $v_k \in \mathbb{R}^n$ . Given two sequences  $\{\alpha_k\}$  and  $\{\beta_k\}$  of non-negative reals, we also say that  $\alpha_k$  is  $\mathcal{O}(\beta_k)$  if there exists a finite constant  $\kappa$  such that  $\lim_{k \rightarrow \infty} (\alpha_k / \beta_k) \leq \kappa$ .

### 3 The Multi-Objective Adagrad Algorithm

Building on ideas originally developed for single objective objective-function-free optimization, we propose the algorithm outlined in Algorithm 1, named **M0-Adagrad**. At every iteration, after computing a common descent direction for the objective functions, denoted by  $-g_k^s \in \mathbb{R}^n$  and computed by solving  $(\Omega(\mathbf{x}))$  in  $\mathbf{x} = \mathbf{x}^k$  (see Step 2), the update step  $s^k$  is obtained by scaling this direction with suitable weights  $w_k$ . These weights (see Step 3) accumulate the  $\ell_2$ -norm of past gradients in an analogous fashion as in the Adagrad-norm [38] algorithm.

In the following, we formally define our direction  $g_k^s$  and report a number of theoretical results related to it, needed for the subsequent convergence rate analysis.

**Definition 7.** *Let  $\mathbf{x}^k \in \mathbb{R}^n$  be any iteration point and denote with  $\lambda^k \in \mathbb{R}^m$  an optimal solution of  $(\Omega(\mathbf{x}))$  for  $\mathbf{x} = \mathbf{x}^k$ . Then we define the vector  $g_k^s \in \mathbb{R}^n$  by*

$$g_k^s := \sum_{j=1}^m \lambda_j^k \nabla f_j(\mathbf{x}^k)$$

and, in case of  $\omega(\mathbf{x}^k) > 0$ , the negative of the normed vector of  $g_k^s$  by

$$n_k^s := -\frac{1}{\|g_k^s\|_2} g_k^s.$$

---

**Algorithm 1** M0-Adagrad

---

1: Initialization:  $x_0 \in \mathbb{R}^n$ , a constant  $\varsigma \in (0, 1)$ , and the initial weight vector  $w_{-1} = \sqrt{\varsigma}$  are given. Set  $k = 0$ .

2: Solve  $(\Omega(x))$  with  $x = x^k$  to obtain  $g_k^s \in \mathbb{R}^n$

3: Compute the weights

$$w_k = \sqrt{(w_{k-1})^2 + \|g_k^s\|_2^2} \quad (6)$$

4: Compute the step

$$s^k = -\frac{g_k^s}{w_k} \quad (7)$$

5: Define

$$x^{k+1} = x^k + s^k,$$

increment  $k$  by one and return to Step 2.

---

Note that  $\|g_k^s\|_2^2 = \omega(x^k)$  and thus  $g_k^s = 0$  if and only if  $\omega(x^k) = 0$ . For the following, we define a function  $h: \mathbb{R}^n \rightarrow \mathbb{R}$  for some fixed  $x \in \mathbb{R}^n$  by

$$h(d) := \left( \max_{j \in [m]} \nabla f_j(x)^\top d \right) + \frac{1}{2} \|d\|_2^2.$$

For our proof of the convergence of our algorithm it will be important to relate the negative of the objective function of (mgrad(x)) with  $x = x^k$  in  $d = -g_k^s$ , i.e. of  $\max_{j \in [m]} \nabla f_j(x^k)^\top (-g_k^s)$ , with  $\omega(x^k)$ . We will obtain this in the forthcoming Lemma 9. For the proof of this Lemma 9, we first relate the value of  $h$  as defined above and with  $x = x^k$  in  $d = -g_k^s$  to that of  $\omega(x^k)$ . We also clarify that the obtained directions  $-g_k^s$  and  $n_k^s$  are in fact descent directions in case  $x^k$  is not yet a Pareto critical point.

**Lemma 8.** *Let  $x^k \in \mathbb{R}^n$  be any iteration point which is not a Pareto critical point for (MOP), that is,  $\omega(x^k) > 0$ . Then the following statements hold.*

(a) *The direction  $n_k^s$  is an optimal solution of (mgrad(x)) for  $x = x^k$ .*

(b) *The direction  $-g_k^s$  is the unique optimal solution of*

$$\min_{d \in \mathbb{R}^n} h(d) \quad (\text{pgrad}(x))$$

*for  $x = x^k$ , and for the optimal value  $h(-g_k^s)$  we have*

$$\omega(x^k) = \|g_k^s\|_2^2 = -2h(-g_k^s). \quad (8)$$

(c) *The direction  $n_k^s$ , and thus also  $-g_k^s$ , is a joint descent direction, i.e. there exists  $t_0 > 0$  such that*

$$\forall j \in [m], \forall t' \in (0, t_0]: f_j(x^k + t' n_k^s) < f_j(x^k).$$

*Proof.* We start with proving (a). To this end, we set

$$t^k := \max_{j \in [m]} \nabla f_j(x^k)^\top n_k^s$$

and we show that  $(t^k, n_k^s)$  is feasible for the optimization problem (mgrad'(x)) and satisfies the KKT optimality conditions for this problem with Lagrange multipliers  $\lambda_j^k$ ,  $j \in [m]$  and

$\xi^k := \|g_k^s\|_2/2 \geq 0$ . As (mgrad'(x)) is a convex optimization problem, this is sufficient for  $(t^k, n_k^s)$  being optimal for (mgrad'(x)) and thus for  $n_k^s$  being optimal for (mgrad(x)) for  $x = x^k$ .

First, note that  $\|n_k^s\|_2^2 = 1$ . The Lagrange function reads as

$$L(t, d, \lambda^k, \xi^k) = t + \sum_{j=1}^m \lambda_j^k (\nabla f_j(x^k)^\top d - t) + \xi^k (d^\top d - 1).$$

As  $\lambda^k$  is feasible for  $(\Omega(x))$  for  $x = x^k$  we obtain that the partial derivative of  $L$  as defined above w.r.t.  $t$  equals zero. For the partial derivative w.r.t.  $d$  we obtain the condition

$$\sum_{j=1}^m \lambda_j^k \nabla f_j(x^k) + 2\xi^k d = 0,$$

which is satisfied for  $d = n_k^s$  by the definition of  $\xi^k$  and of  $n_k^s$ . It remains to show that the complementarity conditions are fulfilled, i.e., that the Lagrange-multipliers to the inactive constraints are zero. As  $\|n_k^s\|_2^2 = 1$  it remains to show that for all  $j \in [m]$  we have

$$\lambda_j^k (\nabla f_j(x^k)^\top n_k^s - t^k) = 0. \quad (9)$$

Let  $J^k := \{j \in [m] \mid t^k = \nabla f_j(x^k)^\top n_k^s\}$ . Note that  $J^k \neq \emptyset$ . Now assume there is an index  $\ell \in [m] \setminus J^k$  with  $\lambda_\ell^k > 0$  and let  $\ell' \in J^k$ . Then this implies  $\nabla f_{\ell'}(x^k)^\top n_k^s < \nabla f_\ell(x^k)^\top n_k^s$ . By the definition of  $n_k^s$  we get

$$\gamma := (\nabla f_{\ell'}(x^k) - \nabla f_\ell(x^k))^\top g_k^s < 0.$$

Next we define for any  $\varepsilon \in (0, \lambda_\ell^k)$  the vector  $\tilde{\lambda} \geq 0$  by  $\tilde{\lambda}_j := \lambda_j^k$  for all  $j \in [m] \setminus \{\ell, \ell'\}$  and

$$\tilde{\lambda}_\ell := \lambda_\ell^k - \varepsilon, \quad \tilde{\lambda}_{\ell'} := \lambda_{\ell'}^k + \varepsilon.$$

Then  $\tilde{\lambda}$  is feasible for  $(\Omega(x))$  for  $x = x^k$  as  $\lambda^k$  is feasible for  $(\Omega(x))$  for  $x = x^k$ . We get for the objective function value of the problem  $(\Omega(x))$  for  $x = x^k$  that

$$\begin{aligned} \left\| \sum_{j=1}^m \tilde{\lambda}_j \nabla f_j(x^k) \right\|_2^2 &= \left\| \left( \sum_{j=1}^m \lambda_j^k \nabla f_j(x^k) \right) + \varepsilon (\nabla f_{\ell'}(x^k) - \nabla f_\ell(x^k)) \right\|_2^2 \\ &= \|g_k^s + \varepsilon (\nabla f_{\ell'}(x^k) - \nabla f_\ell(x^k))\|_2^2 \\ &= \|g_k^s\|_2^2 + \varepsilon^2 \|\nabla f_{\ell'}(x^k) - \nabla f_\ell(x^k)\|_2^2 + 2\varepsilon (\nabla f_{\ell'}(x^k) - \nabla f_\ell(x^k))^\top g_k^s \\ &= \|g_k^s\|_2^2 + \varepsilon (\varepsilon \|\nabla f_{\ell'}(x^k) - \nabla f_\ell(x^k)\|_2^2 + 2\gamma). \end{aligned}$$

Recall that  $\gamma < 0$ . In case  $\|\nabla f_{\ell'}(x^k) - \nabla f_\ell(x^k)\|_2^2 = 0$  this is a contradiction to  $\|g_k^s\|_2^2$  the optimal value of  $(\Omega(x))$  for  $x = x^k$ . Otherwise, for  $\varepsilon$  small enough, we also get a contradiction to  $\|g_k^s\|_2^2$  the optimal value of  $(\Omega(x))$  for  $x = x^k$ . Thus the assertion is proven.

For proving part (b), instead of (pgrad(x)) we examine the following optimization problem

$$\begin{aligned} \min_{(t, d) \in \mathbb{R}^{1+n}} \quad & t + \frac{1}{2} \|d\|_2^2 \\ \text{s.t.} \quad & \nabla f_j(x)^\top d - t \leq 0, \quad \forall j \in [m] \end{aligned} \quad (\text{pgrad}'(x))$$

for  $x = x^k$ . As this problem has a strictly convex objective function, there is not more than one optimal solution  $(\bar{t}^k, \bar{d}^k)$  and we aim on showing that this is  $\bar{t}^k := \max_{j \in [m]} f_j(x^k)^\top (-g_k^s)$  and  $\bar{d}^k = -g_k^s$ . We do this by showing that the KKT conditions are satisfied, as those are sufficient

for this convex optimization problem. It is easy to verify that in  $(\bar{t}^k, \bar{d}^k)$  with the Lagrange multipliers  $\lambda^k \geq 0$  the derivative of the Lagrange function vanishes. To see this recall that  $\sum_{j=1}^m \lambda_j^k = 1$  as  $\lambda^k$  is an optimal solution of  $(\Omega(x))$  for  $x = x^k$ . Next, note that the condition (9) can be written as follows, after multiplying the equation with  $\|g_k^s\|_2$ ,

$$\lambda_j^k (\nabla f_j(x^k)^\top (-g_k^s) - \bar{t}^k) = 0$$

for all  $j \in [m]$ . With the same argumentation as in the proof of (a) we can argue that this condition has to be fulfilled, which is the complementarity condition of  $(\text{pgrad}'(x))$  in  $x = x^k$ . Thus  $(\bar{t}^k, \bar{d}^k)$  is a KKT point of  $(\text{pgrad}'(x))$  and hence the unique optimal solution. As a consequence,  $\bar{d}^k = -g_k^s$  is the unique optimal solution of  $(\text{pgrad}(x))$  for  $x = x^k$ .

It remains to show that (8) holds, and we do this by showing that the dual problem of  $(\text{pgrad}'(x))$  is (next to some scalar multiplications and sign changes) the problem  $(\Omega(x))$ . In [28] it was already claimed that these problems are dual to each other, but the factor  $-0.5$  has to be taken into account, see below. The formulation of the dual problem of  $(\text{pgrad}'(x))$ , including this factor, can be found in [5] on pages 413-414, and the calculation of  $\bar{d}(\lambda)$  below can also be found in [12]. We give a rigorous proof here for completeness.

We always fix, as before,  $x = x^k$ . For formulating the objective function of the dual problem to  $(\text{pgrad}'(x))$  we need to minimize the Langrange-function w.r.t.  $t \in \mathbb{R}$  and  $d \in \mathbb{R}^n$  for fixed  $\lambda \geq 0$ . As the Lagrange-function to  $(\text{pgrad}'(x))$  is defined by

$$\begin{aligned} L(t, d, \lambda) &= t + \frac{1}{2} d^\top d + \sum_{j=1}^m \lambda_j (\nabla f_j(x^k)^\top d - t) \\ &= t (1 - \sum_{j=1}^m \lambda_j) + \frac{1}{2} d^\top d + \sum_{j=1}^m \lambda_j \nabla f_j(x^k)^\top d, \end{aligned}$$

we get that the infimum is  $-\infty$  unless  $\sum_{j=1}^m \lambda_j = 1$ . In case of a  $\lambda \geq 0$  with  $\sum_{j=1}^m \lambda_j = 1$  the infimum is attained in any  $\bar{t}(\lambda) \in \mathbb{R}$  and

$$\bar{d}(\lambda) = - \sum_{j=1}^m \lambda_j \nabla f_j(x^k).$$

Thus the objective function of the dual problem reads, for any  $\lambda \geq 0$  with  $\sum_{j=1}^m \lambda_j = 1$ , as

$$q(\lambda) := \inf_{(t, d)} L(t, d, \lambda) = \frac{1}{2} \bar{d}(\lambda)^\top \bar{d}(\lambda) + \sum_{j=1}^m \lambda_j \nabla f_j(x^k)^\top \bar{d}(\lambda) = -\frac{1}{2} \bar{d}(\lambda)^\top \bar{d}(\lambda).$$

Here, the dual problem can be formulated as

$$\begin{aligned} \max_{\lambda \in \mathbb{R}^m} \quad & -\frac{1}{2} \|\bar{d}(\lambda)\|_2^2 \\ \text{s.t.} \quad & \sum_{j=1}^m \lambda_j = 1, \quad \lambda \geq 0, \end{aligned}$$

which corresponds to

$$\begin{aligned} -\frac{1}{2} \min_{\lambda \in \mathbb{R}^m} \quad & \|\sum_{j=1}^m \lambda_j \nabla f_j(x^k)\|_2^2 \\ \text{s.t.} \quad & \sum_{j=1}^m \lambda_j = 1, \quad \lambda \geq 0, \end{aligned}$$

with optimal value, cf.  $(\Omega(x))$  for  $x = x^k$ , equal to  $-0.5\omega(x^k) = -0.5\|g_k^s\|_2^2$ . As  $(\text{pgrad}'(x))$  is convex and the Slater CQ holds, we have strong duality. The optimal value of  $(\text{pgrad}'(x))$  is  $h(-g_k^s)$ , and thus the assertion is shown.

For part (c), as  $n_k^s$  is optimal for  $(\text{mgrad}(x))$  for  $x = x^k$  by (a) and as the optimal value of  $(\text{mgrad}(x))$  for  $x = x^k$  has to be negative by Lemma 5 and by the non-criticality of  $x^k$ , we have  $\max_{j \in [m]} \nabla f_j(x^k)^\top n_k^s < 0$ . Note that it suffices to have that for all  $j \in [m]$  it holds  $\nabla f_j(x^k)^\top n_k^s < 0$  to have that  $n_k^s$  is a joint descent direction, cf. [33, Lemma 2.5].  $\square$

Note that problem (pgrad(x)) as well as the reformulation (pgrad'(x)) have already been studied in [11] for calculating a steepest descent direction. As noted there in Lemma 1 this problem can also be used to characterize Pareto criticality: a point  $x$  is Pareto critical if and only if the optimal value of (pgrad(x)) is zero. Otherwise, the optimal value is negative. It has also been stated that another descent direction can be obtained by solving the problem (pgrad(x)), in [11] w.r.t. the norm  $\|\cdot\|_\infty$ . We have just shown that, next to scaling, the directions obtained from (pgrad(x)) and ( $\Omega(x)$ ) can be considered to be the same.

We have shown that the direction  $-g_k^s$  is a joint, also called common, descent direction. It has also been called negative multi-gradient in [28].

**Lemma 9.** *Let  $x^k \in \mathbb{R}^n$  be any iteration point. Then*

$$\|g_k^s\|_2^2 = -\max_{j \in [m]} \nabla f_j(x^k)^\top (-g_k^s) \quad (10)$$

holds.

*Proof.* First, assume that  $x^k$  is a Pareto critical point for (MOP). Then we have  $\omega(x^k) = 0$  and  $g_k^s = 0$ . Then (10) trivially holds. Next, we assume that  $x^k$  is not a Pareto critical point for (MOP) and thus  $\omega(x^k) > 0$ . Then by Lemma 8(b) we get  $\|g_k^s\|_2^2 = -2h(-g_k^s)$ . By a simple calculation we derive (10).  $\square$

### 3.1 Convergence rate analysis

Recall that by (3) we have  $\Phi(x) = \max_{j \in [m]} f_j(x)$  and that  $L_{\max}$  was defined in (2)

**Lemma 10.** *Suppose that Algorithm 1 is applied to (MOP). Then, for all  $k \geq 0$  it holds*

$$\Phi(x^{k+1}) \leq \Phi(x^k) - \frac{\|g_k^s\|_2^2}{w_k} + \frac{L_{\max}}{2} \|s^k\|_2^2, \quad (11)$$

and

$$\Phi(x^0) - \Phi(x^{k+1}) \geq \sum_{\ell=0}^k \frac{\|g_\ell^s\|_2^2}{w_\ell} - \frac{L_{\max}}{2} \sum_{\ell=0}^k \frac{\|g_\ell^s\|_2^2}{w_\ell^2}. \quad (12)$$

*Proof.* Recall that by Lemma 9 we have  $-\|g_k^s\|_2^2 = \max_{j \in [m]} \nabla f_j(x^k)^\top (-g_k^s)$ . Considering that

$$x^{k+1} - x^k = s^k = -\frac{g_k^s}{w_k},$$

by the mean value theorem applied to each function  $f_j$ ,  $j \in [m]$ , we have:

$$\begin{aligned} \Phi(x^{k+1}) &\leq \max_{j \in [m]} \left\{ f_j(x^k) + \nabla f_j(x^k)^\top s^k + \frac{L_j}{2} \|s^k\|_2^2 \right\} \\ &\leq \Phi(x^k) + \frac{1}{w_k} \max_{j \in [m]} \nabla f_j(x^k)^\top (-g_k^s) + \frac{L_{\max}}{2} \|s^k\|_2^2 \\ &= \Phi(x^k) - \frac{\|g_k^s\|_2^2}{w_k} + \frac{L_{\max}}{2} \|s^k\|_2^2, \end{aligned}$$

that is (11). Furthermore, summing for  $\ell = 0, \dots, k$  it gives

$$\begin{aligned} \Phi(x^{k+1}) &\leq \Phi(x^0) - \sum_{\ell=0}^k \frac{\|g_\ell^s\|_2^2}{w_\ell} + \frac{L_{\max}}{2} \sum_{\ell=0}^k \sum_{i=1}^n \left( \frac{g_{i,\ell}^s}{w_\ell} \right)^2 \\ &= \Phi(x^0) - \sum_{\ell=0}^k \sum_{i=1}^n \frac{(g_{i,\ell}^s)^2}{w_\ell} + \frac{L_{\max}}{2} \sum_{\ell=0}^k \frac{\|g_\ell^s\|_2^2}{w_\ell^2}, \end{aligned}$$

that is inequality (12).  $\square$

Using similar reasoning as in the proof of the Lemma 10, we can show that  $s^k$  is a common descent direction and a descent direction for  $\Phi$ :

**Lemma 11.** *There exists  $t_0 > 0$  such that for all  $t' \in (0, t_0]$ ,*

$$f_j(x^k + t' s^k) < f_j(x^k) \quad \forall j \in [m]$$

and

$$\Phi(x^k + t' s^k) < \Phi(x^k).$$

*Proof.* By the mean value theorem applied to each function  $f_j$ ,  $j \in [m]$  and with Lemma 9, we have, for all  $t > 0$ :

$$\begin{aligned} f_j(x^k + t s^k) &\leq f_j(x^k) + t \nabla f_j(x^k)^T s^k + \frac{L_j}{2} t^2 \|s^k\|_2^2 \\ &\leq f_j(x^k) + \frac{1}{w_k} t \nabla f_j(x^k)^\top (-g_k^s) + \frac{L_j}{2} t^2 \|s^k\|_2^2 \\ &= f_j(x^k) + t^2 \left( \frac{1}{t} \frac{\nabla f_j(x^k)^\top (-g_k^s)}{w_k} + \frac{L_j}{2} \|s^k\|_2^2 \right). \end{aligned}$$

As  $\nabla f_j(x^k)^\top (-g_k^s) < 0$ , as  $-g_k^s$  is a common descent direction, there exists  $t_0 > 0$  such that

$$\frac{1}{t'} \frac{\nabla f_j(x^k)^\top (-g_k^s)}{w_k} + \frac{L_j}{2} \|s^k\|_2^2 < 0$$

for all  $t' \in (0, t_0]$  so that

$$\Phi(x^k + t' s^k) < \Phi(x^k),$$

for all  $t' \in (0, t_0]$ .  $\square$

We next proceed by recalling the following lemmas, that will be used in our main result concerning the convergence rate. The proof of the first one is obtained via private communication with authors and is reported for the sake of completeness in Appendix A. The proof of Lemma 13 can be found in [18, Lemma 3.1].

**Lemma 12.** *Suppose that*

$$au \leq b + c \log(u), \quad (13)$$

for some  $a, c > 0$  and  $u \geq 1$ . Then

$$u \leq \max \left\{ e^{b/c}, \frac{4c^2}{a^2} \right\}.$$

**Lemma 13.** *Let  $\{c_k\}_{k \geq 0}$  be a non-negative sequence and  $\xi > 0$ . Then*

$$\sum_{j=0}^k \frac{c_j}{(\xi + \sum_{\ell=0}^j c_\ell)} \leq \log \left( \frac{\xi + \sum_{j=0}^k c_j}{\xi} \right). \quad (14)$$

Thanks to Lemma 1 we have that a value  $\Phi_{low}$  exists such that

$$\Phi(x) \geq \Phi_{low}, \text{ for all } x \in \mathbb{R}^n.$$

We are now ready to state a (first) result on the global convergence rate of Algorithm 1. For that, the average of a sequence of values  $\beta^\ell \in \mathbb{R}$  is defined as

$$\text{average } \beta^\ell = \frac{1}{k+1} \sum_{\ell=0}^k \beta^\ell.$$

**Theorem 14.** *Let  $\Gamma_0 := \Phi(x^0) - \Phi_{low}$ . Suppose that Algorithm 1 is applied to (MOP). Then the following bound holds:*

$$\text{average } \omega(x^\ell) = \text{average } \|g_\ell^s\|_2^2 \leq \frac{\theta}{k+1} \quad (15)$$

with

$$\theta = \max \left\{ \varsigma, \sqrt{\frac{\varsigma}{2}}, \frac{\varsigma}{2} e^{\frac{2\Gamma_0}{L_{\max}}}, \sqrt{\frac{\varsigma}{2}} \frac{24L_{\max}^2}{\varsigma} \right\}.$$

*Proof.* Recall that by (8) we have  $\omega(x^k) = \|g_k^s\|_2^2$  and we have further  $\Gamma_0 = \Phi(x^0) - \Phi_{low} \geq \Phi(x^0) - \Phi(x^{k+1})$  for all  $k$ . From (12), we derive

$$\sum_{\ell=0}^k \frac{\|g_\ell^s\|_2^2}{w_\ell} \leq \Gamma_0 + \frac{L_{\max}}{2} \sum_{\ell=0}^k \frac{\|g_\ell^s\|_2^2}{w_\ell^2}. \quad (16)$$

Using Lemma 13 with  $c_j = \|g_j^s\|_2^2$  and the fact that  $w_k = \sqrt{\varsigma + \sum_{j=0}^k \|g_j^s\|_2^2}$  we have

$$\sum_{\ell=0}^k \frac{\|g_\ell^s\|_2^2}{w_\ell^2} = \sum_{\ell=0}^k \frac{\|g_\ell^s\|_2^2}{\varsigma + \sum_{j=0}^k \|g_j^s\|_2^2} \leq \log \left( 1 + \frac{1}{\varsigma} \sum_{\ell=0}^k \|g_\ell^s\|_2^2 \right).$$

Therefore, including the above inequalities in (16) we get

$$\sum_{\ell=0}^k \frac{\|g_\ell^s\|_2^2}{w_\ell} \leq \Gamma_0 + \frac{L_{\max}}{2} \log \left( 1 + \frac{1}{\varsigma} \sum_{\ell=0}^k \|g_\ell^s\|_2^2 \right). \quad (17)$$

Assume now that

$$\sum_{\ell=0}^k \|g_\ell^s\|_2^2 \geq \max \left\{ \varsigma, \frac{\varsigma}{2} e^{\frac{2\Gamma_0}{L_{\max}}} \right\}, \quad (18)$$

which implies

$$\Gamma_0 \leq \frac{L_{\max}}{2} \log \left( \frac{2}{\varsigma} \sum_{\ell=0}^k \|g_\ell^s\|_2^2 \right), \quad (19)$$

and

$$\varsigma + \sum_{\ell=0}^k \|g_\ell^s\|_2^2 \leq 2 \sum_{\ell=0}^k \|g_\ell^s\|_2^2.$$

Furthermore, recalling the definition of  $w_k$  in (6), we have that

$$w_k = \sqrt{\varsigma + \sum_{\ell=0}^k \|g_\ell^s\|_2^2} \leq \sqrt{2 \sum_{\ell=0}^k \|g_\ell^s\|_2^2}. \quad (20)$$

Then, observing that  $w_\ell \leq w_k$  for all  $\ell \leq k$  and using inequalities (19) and (20) in (17) we obtain

$$\frac{\sum_{\ell=0}^k \|g_\ell^s\|_2^2}{\sqrt{2 \sum_{\ell=0}^k \|g_\ell^s\|_2^2}} \leq L_{\max} \log \left( \frac{2}{\varsigma} \sum_{\ell=0}^k \|g_\ell^s\|_2^2 \right)$$

that is

$$\frac{\sqrt{\varsigma}}{2} \sqrt{\frac{2}{\varsigma} \sum_{\ell=0}^k \|g_\ell^s\|_2^2} \leq 2L_{\max} \log \left( \sqrt{\frac{2}{\varsigma} \sum_{\ell=0}^k \|g_\ell^s\|_2^2} \right). \quad (21)$$

We can now apply Lemma 12 since (21) is identical to (13) with

$$a = \frac{\sqrt{\varsigma}}{2}, \quad b = 0, \quad c = 2L_{\max}, \quad u = \sqrt{\frac{2}{\varsigma} \sum_{\ell=0}^k \|g_\ell^s\|_2^2}, \quad (22)$$

and  $u \geq 1$  due to (18), and obtain the bound

$$\sum_{\ell=0}^k \|g_\ell^s\|_2^2 \leq \sqrt{\frac{\varsigma}{2}} \max \left\{ 1, \frac{24L_{\max}^2}{\varsigma} \right\}.$$

Hence taking the average gives that

$$\text{average}_{\ell \in \{0, \dots, k\}} \|g_\ell^s\|_2^2 \leq \sqrt{\frac{\varsigma}{2}} \max \left\{ 1, \frac{24L_{\max}^2}{\varsigma} \right\} \frac{1}{k+1}.$$

Suppose that (18) fails, then we would have

$$\text{average}_{\ell \in \{0, \dots, k\}} \|g_\ell^s\|_2^2 \leq \max \left\{ \varsigma, \frac{\varsigma}{2} e^{\frac{2\Gamma_0}{L_{\max}}} \right\} \frac{1}{k+1}.$$

□

## 4 Numerical results

In this section, we present the numerical tests conducted to evaluate the performance of **M0-Adagrad**. We implemented **M0-Adagrad** in MATLAB and the test have been run on an Intel® Core™ i5-14400F processor running at 2.80GHz under Linux.

For comparison, we implemented a standard descent method in which  $-g_k^s$  (obtained solving  $(\Omega(x))$ ) serves as common descent direction and the step length is determined by using an Armijo-like rule as proposed in [11]. The method is labeled for **M0-Descent** in the forthcoming sections. We report the scheme of **M0-Descent** in Algorithm 2 for completeness. In our implementation of **M0-Descent**, we set  $\beta = 0.1$ .

To solve  $(\Omega(x))$  we adopt `fmincon` within MATLAB. Note that  $(\Omega(x))$  is of dimension  $m$ , meaning that computing  $-g_k^s$  is computationally convenient when dealing with a small number of objective functions.

In our numerical experience, we first provide an overview on the performance of **M0-Adagrad** on bi-objective unconstrained problems built from the CUTEst collection from [21] as well as on standard unconstrained multi-objective problems from literature (see e.g. [34]). We also consider noisy problem instances to assess the behavior of **M0-Adagrad** under different noise levels. Then,

---

**Algorithm 2** MO-Descent

---

- 1: Initialization:  $x_0 \in \mathbb{R}^n$ ,  $\beta \in (0, 1)$ . Set  $k = 0$ .
- 2: Solve  $(\Omega(x))$  with  $x = x^k$  to obtain  $g_k^s \in \mathbb{R}^n$
- 3: Set  $t = 1$ .
- 4: **while**  $f_j(x^k - tg_k^s) > f_j(x^k) - \beta t \nabla f_j(x^k)^T g_k^s$  for some  $j \in [m]$  **do**
- 5:      $t = t/2$
- 6: **end while**
- 7: Set  $\alpha^k = t$ .
- 8: Define

$$x^{k+1} = x^k - \alpha^k g_k^s,$$

increment  $k$  by one and return to Step 2.

---

we report the results of preliminary numerical experiments on the use of MO-Adagrad for training multi-task problems. Indeed we considered two synthetic datasets: one represents a 4-classes classification task and a binary classification task, the other one consists of 2 binary classification tasks, both for a set of points in  $\mathbb{R}^2$ , see e.g. [3].

#### 4.1 Experiments on CUTEst instances.

In a first set of experiments, we constructed bi-objective unconstrained instances from the single-objective unconstrained problems in the CUTEst collection [21] by adding a squared  $\ell_2$ -norm regularizer as a second objective. We excluded all problems with optimal value equal to 0 in order to obtain proper bi-objective instances. Overall, we built 124 instances; the names of used CUTEst problems are reported in Table 1 along with their dimension. All runs were initialized from the standard starting points provided by [21]. We stopped the algorithms as soon as a Pareto critical point is reached or in case a limit of 100000 number of gradient (and function in case of MO-Descent) evaluations is reached. For a fair comparison, for MO-Descent we counted that number of function evaluations divided by the problem dimension and added this quantity to the number of gradient evaluations.


We report in Figure 1 the performance profiles [8] related to number of gradient (and weighted function) evaluations. Clearly, no function evaluations are required by MO-Adagrad. While MO-Descent requires a smaller number gradient and function evaluations in case a Pareto critical point is reached, MO-Adagrad turns to be more robust solving the 89% of instances (MO-Descent only solves the 77% of instances).

#### 4.2 Experiments on noisy problems

In a second set of experiments, we selected the unconstrained problems used in [34] as benchmark multi-objective optimization instances, while we constructed 11 bi-objective unconstrained instances combining single-objective unconstrained problems in the CUTEst collection [21] sharing the same dimension. In this case, the starting point is chosen as the average of the standard starting points for the two original problems. For the benchmark problems, we used 10 randomly generated starting points.

As before, we stopped the algorithms as soon as a Pareto critical point is reached or in case the limit of 100000 number of gradient (and function in case of MO-Descent) evaluation is reached. As the objective-function-free methods are mainly known for their ability of being robust to noise, we evaluated our approach under different levels of noise added both to the objectives and to the gradients. Specifically, we considered noise level of  $\rho \in \{5\%, 15\%, 25\%\}$

Figure 1: Performance profiles (log-scale) on the number of gradient (and function) evaluations on instances derived from CUTEst problems by including  $\ell_2$ -norm regularizer as a second objective.



by adding a relative Gaussian noise with unit variance (see e.g. [19]). We report in Table 2 the results obtained in terms of gradient and function evaluations. Consistent with our first set of experiments, in the noise-free setting, **MO-Adagrad** exhibits fewer failures than **MO-Descent**, which fails on 8 instances. When noise is added to both the gradients and the objective functions, **MO-Descent** shows improved performance, whereas **MO-Adagrad** remains largely unchanged. In particular, **MO-Descent** reduces its number of failures from 8 to just 1 for all the considered values of  $\rho$ .

At first glance, this suggests that the presence of noise helps **MO-Descent** in identifying Pareto critical points. To investigate this further, we examined the locations of the Pareto critical points in the criterion space. We report in Table 3 the distance (on average for the benchmark instances) between the Pareto critical points detected in the presence of noise and those obtained in the noise-free setting. The results show that the points found by **MO-Descent** exhibit significantly larger distances compared to those obtained by **MO-Adagrad**. While this indicates that noise can drive exploration into different regions of the criterion space, it may also represents an undesirable behavior: **MO-Descent** may fail to reliably converge to the same solutions as in the noiseless setting, whereas **MO-Adagrad** demonstrates more robust convergence, highlighting its reliability in noisy scenarios.

### 4.3 Experiments on multi-task geometric classification

We consider a data set  $\mathcal{D} = \{(\mathbf{x}_i, y_i^{(1)}, y_i^{(2)})\}_{i=1}^N$  consisting of  $N = 10000$  points uniformly sampled in the square  $[-2, 2]^2 \subseteq \mathbb{R}^2$ . In the first example, called *Quadrants-Circle*, points are labeled with respect to two different criteria: Task 1 to belong to one of the four quadrants of  $\mathbb{R}^2$  (4-classes classification task); Task 2 to be inside or outside the unitary circle centered in the origin (binary classification task).

Since Task 2 (Circle) is not linearly separable in the original 2D space, the input vector  $\mathbf{x}_i$  is

Table 1: CUTEst instances considered to build bi-objective problems by including squared  $\ell_2$ -norm regularizer as a second objective.

Problem	$n$	Problem	$n$	Problem	$n$	Problem	$n$
ALLINITU	4	EGGCRATE	2	LSC1LS	3	PENALTY2	10
ARGLINA	200	EIGENALS	6	LSC2LS	3	POWELLBSLS	2
ARGLINB	10	EIGENBLS	6	LUKSAN12LS	98	POWERSUM	10
ARWHEAD	10	ELATVIDU	2	LUKSAN17LS	100	QING	5
BARD	3	ENGVAL1	10	LUKSAN22LS	100	QUARTC	10
BDQRTIC	10	ENGVAL2	3	MANCINO	10	ROSENBR	2
BEALE	2	ERRINROS	10	MARATOSB	2	ROSSIMP1	2
BIGGS6	6	ERRINRSM	10	METHANL8LS	31	ROSSIMP2	2
BOXBODLS	2	EXPFIT	2	MEXHAT	2	ROSSIMP3	2
BRKMCC	2	EXTROSNB	10	MEYER3	3	ROSZMAN1LS	4
BROWNAL	10	FLETBV3M	10	MGH17LS	5	S308	2
BROWNDEN	4	FLETCBV2	10	MGH17SLS	5	SENSORS	5
BROYDN3DLS	5	FLETCBV3	10	MODBEALE	10	SINQUAD2	10
CHNROSNB	5	FREUROTH	4	MSQRTALS	25	SINQUAD	10
CHNRSNBM	5	GAUSSIAN	3	MSQRTBLS	25	SPMSRTLS	4999
CLUSTERLS	2	GENROSE	10	n10FOLDTRLs	4	STREG	4
COOLHANSLS	9	GROWTHLS	3	NONCVXU2	10	THURBERLS	7
CRAGGLVY	10	GULF	3	NONCVXUN	10	TOINTGOR	50
CUBE	2	HATFLDD	3	NONDIA	10	TOINTGSS	10
CURLY10	15	HATFLDE	3	OSBORNEA	5	TOINTQOR	50
CYCLIC3LS	12	HIMMELBCLS	2	OSBORNEB	11	TQUARTIC	10
CYCLOOCFLS	20	HIMMELBH	2	OSCIGRAD	10	TRIDIA	5
DANIWOODLS	2	HYDCAR6LS	29	PALMER1C	8	TRIGON2	10
DENSCHNB	2	INDEFM	10	PALMER1D	7	VARDIM	10
DENSCHNC	2	JENSMP	2	PALMER2C	8	WATSON	12
DENSCHND	3	JUDGE	2	PALMER3C	8	WAYSEA1	2
DENSCHNF	2	KSSLS	10	PALMER4C	8	WOODS	4000
DEVGLA2	5	LANCZOS1LS	6	PALMER5C	6	YATP2CLS	35
DIXON3DQ	10	LANCZOS2LS	6	PALMER6C	8	YATP2LS	35
EDENSCH	10	LANCZOS3LS	6	PALMER7C	8	YFITU	3
EG2	10	LIARWHD	10	PALMER8C	8	ZANGWIL2	2

obtained via a feature engineering [4] that includes quadratic terms:

$$\mathbf{x}_i = [1, x_{i,1}, x_{i,2}, x_{i,1}^2, x_{i,2}^2]^T \in \mathbb{R}^5$$

where the first element 1 represents the bias term. The labels are defined as follows:

- $y_i^{(1)} \in \{1, 2, 3, 4\}$ : Quadrant Label.
- $y_i^{(2)} \in \{0, 1\}$ : Circle Label (1 if inside, 0 if outside).

As predicted models, we considered the Softmax Regression for Task 1 and the Logistic Regression for Task 2. Finally, we employed categorical Cross-Entropy and the binary Cross-Entropy as loss functions for the two tasks.

Overall, the final bi-objective problem is:

$$\min_{\mathbf{W}_1, \mathbf{w}_2} (J_1(\mathbf{W}_1), J_2(\mathbf{w}_2))$$

where  $\mathbf{W}_1 \in \mathbb{R}^{5 \times 4}$  and  $\mathbf{w}_2 \in \mathbb{R}^5$  denote the weights of Task 1 and Task 2, respectively, the objectives are

$$J_1(\mathbf{W}_1) = -\frac{1}{N} \sum_{i=1}^N \sum_{k=1}^4 \mathbb{1}(y_i^{(1)} = k) \log \left( \frac{\exp(\mathbf{w}_{1,k}^T \mathbf{x}_i)}{\sum_{j=1}^4 \exp(\mathbf{w}_{1,j}^T \mathbf{x}_i)} \right)$$

Table 2: Results on the instances obtained combining CUTEst problems and on the benchmark instances. For the CUTEst problems the starting point is defined as the average of the starting points of the two original problems. For the benchmark instances we use 10 different randomly generated starting points.

Problem	$n$	no noise		$\rho = 5\%$	
		MO-Descent	MO-Adagrad	MO-Descent	MO-Adagrad
BROWNDEN-ALLINITU	4	-	-	-	-
BROWNAL-ARWHEAD	10	<b>8.8</b>	18	9681.3	<b>18</b>
BROWNAL-VARDIM	10	<b>159.1</b>	15241	<b>126</b>	13866
ARWHEAD-VARDIM	10	-	<b>13</b>	2077.8	<b>372</b>
ZANGWIL2-ROSENBR	2	-	<b>76</b>	809.5	<b>50</b>
ZANGWIL2-CUBE	2	-	<b>66</b>	771	<b>98</b>
ZANGWIL2-ROSENBR	2	-	<b>76</b>	809.5	<b>50</b>
ZANGWIL2-WAYSEA1	2	-	<b>37</b>	1883	<b>38</b>
ROSENBR-WAYSEA1	2	-	<b>7</b>	9709.5	<b>519</b>
ROSENBR-CUBE	2	2059.5	<b>581</b>	1355	<b>706</b>
WAYSEA1-CUBE	2	-	<b>96</b>	17605.5	<b>296</b>
Lovison 3	2	<b>24.2(10)</b>	1685.8(10)	1995.6(10)	<b>1115.2(10)</b>
Lovison 4	2	<b>15.6(10)</b>	3422.4(10)	<b>270.2(10)</b>	3356.2(10)
MOP1	2	<b>3.0(10)</b>	2536.7(10)	<b>17.1(10)</b>	2517.1(10)
T1	2	<b>31.8(10)</b>	3844.7(10)	<b>837.0(10)</b>	1944.8(10)
T2	2	12.2(10)	<b>7.1(10)</b>	58.1(10)	<b>7.2(10)</b>
Problem	$n$	$\rho = 15\%$		$\rho = 25\%$	
		MO-Descent	MO-Adagrad	MO-Descent	MO-Adagrad
BROWNDEN-ALLINITU	4	-	-	-	-
BROWNAL-ARWHEAD	10	1425.1	<b>18</b>	68.7	<b>19</b>
BROWNAL-VARDIM	10	<b>667</b>	9505	<b>491</b>	4487
ARWHEAD-VARDIM	10	<b>53.2</b>	138	224.7	<b>67</b>
ZANGWIL2-ROSENBR	2	3452.5	<b>182</b>	914	<b>116</b>
ZANGWIL2-CUBE	2	1713.5	<b>71</b>	637.5	<b>44</b>
ZANGWIL2-ROSENBR	2	3452.5	<b>182</b>	914	<b>116</b>
ZANGWIL2-WAYSEA1	2	603	<b>39</b>	462	<b>39</b>
ROSENBR-WAYSEA1	2	80479	<b>1149</b>	54966	<b>1705</b>
ROSENBR-CUBE	2	1303	<b>660</b>	700.5	<b>553</b>
WAYSEA1-CUBE	2	7162	<b>162</b>	13514	<b>368</b>
Lovison 3	2	9847.5(10)	<b>1265.9(10)</b>	23562.0(10)	<b>1506.5(10)</b>
Lovison 4	2	<b>789.0(10)</b>	3190.6(10)	<b>1778.3(10)</b>	2875.0(10)
MOP1	2	<b>4.5(10)</b>	2401.6(10)	<b>4.5(10)</b>	2214.6(10)
T1	2	2627.6(10)	<b>2464.1(10)</b>	<b>1443.5(10)</b>	2093.6(10)
T2	2	65.1(10)	<b>7.2(10)</b>	68.2(10)	<b>6.6(10)</b>

and

$$J_2(\mathbf{w}_2) = -\frac{1}{N} \sum_{i=1}^N \left[ y_i^{(2)} \log(\hat{y}_i^{(2)}) + (1 - y_i^{(2)}) \log(1 - \hat{y}_i^{(2)}) \right] \quad (23)$$

where  $\mathbf{1}$  denotes the indicator function,  $\hat{y}_i^{(2)} = \sigma(\mathbf{w}_2^T \mathbf{x}_i)$  with  $\sigma$  the sigmoid function and  $\mathbf{w}_{1,j}$  denotes the  $j$ th column of  $\mathbf{W}_1$ , for  $j = 1, \dots, 4$ .

In the second example, named *Diagonals-Circle*, Task 1 is a non-linear binary classification task based on the XOR logic of the Cartesian quadrants. The model must distinguish between points in the main diagonal (Quadrants 1 and 3) and the anti-diagonal (Quadrants 2 and 4). This requires the inclusion of an interaction feature ( $x \cdot y$ ) to achieve linear separability. In this case, we used the objective function in (23) for both objectives.


In both examples, the data set is randomly split into a training set and a test set made up

Table 3: Distance from the Pareto critical points detected with no noise.

Problem	MO-Descent	MO-Adagrad	MO-Descent	MO-Adagrad	MO-Descent	MO-Adagrad
	$\rho = 5\%$	$\rho = 15\%$	$\rho = 25\%$	$\rho = 5\%$	$\rho = 15\%$	$\rho = 25\%$
BROWNL-ARWHEAD	6.53	<b>0.43</b>	6.51	<b>1.29</b>	10.20	<b>2.22</b>
BROWNL-VARDIM	$< 10e-4$	0.01	$< 10e-4$	0.05	$< 10e-4$	0.14
ROSENBR-CUBE	$< 10e-4$	0.04	$< 10e-2$	0.16	$< 10e-2$	0.09
Lovison 3	100.37	<b>32.58</b>	142.87	<b>113.09</b>	412.79	<b>210.07</b>
Lovison 4	15.41	<b>1.45</b>	27.22	<b>4.09</b>	25.21	<b>8.81</b>
MOP1	0.25	<b>0.11</b>	3.77	<b>0.46</b>	3.91	<b>0.76</b>
T1	37.56	<b>8.03</b>	59.61	<b>22.38</b>	82.10	<b>35.01</b>
T2	<b>0.05</b>	0.06	<b>0.16</b>	0.17	0.28	0.28

of 8000 and 2000 samples, respectively. A representation of the predicted labels on the testing set for 2 multi-task problems obtained with **MO-Adagrad** is reported in Figures 2 and 3.

Figure 2: *Quadrants-Circle*: Task 1 on the left, Task 2 on the right. Different colors denote different labels for the points.



In Table 4 we report the results obtained with **MO-Descent** and **MO-Adagrad** on the two multi-task examples. In particular, we let the solvers run for 1000 iterations and we retrieved the number of gradient/function evaluations ( $g_{eval}/f_{eval}$ ) and elapsed time in seconds (time) employed in the training phase, to get the maximum accuracy  $\mathcal{A}$  on the testing set. The accuracy  $\mathcal{A}$  is measured considering the minimum of the testing accuracies in the two tasks. We observe that the two solvers get comparable accuracies (slightly in favor of **MO-Descent**) but **MO-Adagrad** is faster by a factor 3 than **MO-Descent**. This is due to its objective function free nature of **MO-Adagrad** that makes it particularly suitable when, as in this multi-task framework, evaluating the objective function is roughly as expensive as evaluating the gradient.

## 5 Conclusions

In this work, we propose a novel Adagrad-like algorithm, **MO-Adagrad**, for unconstrained multi-objective optimization. Our method does not need line search based techniques or dominance-based criteria for accepting new points, while still converging to Pareto critical points. Through extensive numerical tests on a broad set of bi-objective instances, we show the robustness of

Figure 3: *Diagonals-Circle*: Task 1 on the left, Task 2 on the right. Different colors denote different labels for the points.




Table 4: Results on the multi-task geometric classification instances.

Problem	MO-Descent				MO-Adagrad			
	$g_{eval}$	$f_{eval}$	time	$\mathcal{A}$	$g_{eval}$	time	$\mathcal{A}$	
Quadrants - Circle	306	3957	24.9	99.8%	388	<b>6.9</b>	99.6%	
Diagonals - Circle	577	6324	27.3	99.9%	970	<b>10.0</b>	99.1%	

MO-Adagrad in comparison with a line search based first order method. The method looks robust and promising to deal with noisy problems, as known for its single-objective counterpart. Further preliminary experiments have been devoted to training multi-task problems, suggesting the potential ability of MO-Adagrad to deal with multi-task learning problems.

Future work will be devoted to extend MO-Adagrad to compute sets of nondominated Pareto points, as opposed to the single Pareto point currently produced. Recent advances in multiobjective optimization have seen growing interest in local algorithms that generate sequences of sets, with the aim of approximating the Pareto set of a multi-objective problem [6, 7, 25, 26, 27, 29]. However, avoiding the use of dominance-based acceptance criteria presents a significant challenge when generating sequences of Pareto sets, rather than individual points. Consequently, new techniques need to be explored to address this complexity.

#### Acknowledgment

The authors would like to thank Prof. Stefania Bellavia and her coauthors for providing Lemma 12 and its proof, which greatly contributed to simplifying our rate of convergence analysis.

#### Ethics approval and consent to participate

Not applicable.

#### Consent for publication

Not applicable.

## Funding

The work of M.P. was partially supported by INdAM-GNCS under the project CUP E53C24001950001. The research of M.P. was partially granted by the Italian Ministry of University and Research (MUR) through the PRIN 2022 “MOLE: Manifold constrained Optimization and LEarning”, code: 2022ZK5ME7 MUR D.D. financing decree n. 20428 of November 6th, 2024 (CUP B53C24006410006).

## Availability of data and materials

The data that support the findings of this study are available from the corresponding author upon request.

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

All authors contributed equally to the writing of this article. All authors reviewed the manuscript.

## A Proof of Lemma 12 from [2]

*Proof.* Suppose that  $u > e^{b/c}$ . Then  $b \leq c \log(u)$  and thus (see [36, eq. (14)])

$$au \leq 2c \log(u) \leq 2c \log(1+u) \leq \frac{2cu}{\sqrt{1+u}}.$$

Hence  $a\sqrt{1+u} \leq 2c$ , which is to say that  $u \leq (2c/a)^2 - 1$ , yielding the desired bound.  $\square$

## References

- [1] Stefania Bellavia, Serge Gratton, Benedetta Morini, and Philippe L Toint. Fast stochastic second-order adagrad for nonconvex bound-constrained optimization. *arXiv preprint arXiv:2505.06374*, 2025.
- [2] Stefania Bellavia, Serge Gratton, Benedetta Morini, and Philippe L Toint. An objective-function-free algorithm for general smooth constrained optimization. *In preparation*, 2026.
- [3] Stefania Bellavia, Francesco Della Santa, and Alessandra Papini. ATE-SG: Alternate through the epochs stochastic gradient for multi-task neural networks. *arXiv preprint arXiv:2312.16340*, 2025.
- [4] Christopher M. Bishop. *Pattern Recognition and Machine Learning*. Springer, New York, 2006. Chapter 4: Linear Models for Classification.
- [5] Jian Chen, Gao-Xi Li, and Xin-Min Yang. Variable metric method for unconstrained multiobjective optimization problems. *Journal of the Operations Research Society of China*, 11:409–438, 2022.
- [6] Guido Cocchi, Giampaolo Liuzzi, Stefano Lucidi, and Marco Sciandrone. On the convergence of steepest descent methods for multiobjective optimization. *Comput. Optim. Appl.*, 77:1–27, 2020.
- [7] Ana Luísa Custódio, JF Aguilar Madeira, A Ismael F Vaz, and Luís Nunes Vicente. Direct multisearch for multiobjective optimization. *SIAM J. Optim.*, 21(3):1109–1140, 2011.
- [8] Elizabeth D Dolan and Jorge J Moré. Benchmarking optimization software with performance profiles. *Mathematical programming*, 91(2):201–213, 2002.
- [9] John Duchi, Elad Hazan, and Yoram Singer. Adaptive subgradient methods for online learning and stochastic optimization. *Journal of machine learning research*, 12(7), 2011.
- [10] Jörg Fliege, L.M. Graña Drummond, and Benar Fux S Svaiter. Newton’s method for multiobjective optimization. *SIAM Journal on Optimization*, 20(2):602–626, 2009.
- [11] Jörg Fliege and Benar Fux Svaiter. Steepest descent methods for multicriteria optimization. *Mathematical Methods of Operations Research*, 51(3):479–494, August 2000.
- [12] Jörg Fliege, Ismael Vaz, and Luis Nunes Vicente. Complexity of gradient descent for multiobjective optimization. *Optimization Methods and Software*, 34(5):949–959, 2019.
- [13] Ellen H Fukuda, LM Graña Drummond, and Fernanda MP Raupp. An external penalty-type method for multicriteria. *TOP*, 24:493–513, 2016.

- [14] Ellen H Fukuda, LM Graña Drummond, and Fernanda MP Raupp. A barrier-type method for multiobjective optimization. *Optimization*, 69(11):2471–2487, 2020.
- [15] MLN Gonçalves, FS Lima, and LF Prudente. Globally convergent Newton-type methods for multiobjective optimization. *Comput. Optim. Appl.*, 83(2):403–434, 2022.
- [16] Max L. N. Gonçalves, Geovani Nunes Grapiglia, and Jefferson Gonçalves Melo. An adaptive line-search-free multiobjective gradient method and its iteration-complexity analysis. *Optimization Online* <https://optimization-online.org/?p=33379>, 30/01/2026.
- [17] Geovani N Grapiglia and Gabriel FD Stella. An adaptive trust-region method without function evaluations. *Computational Optimization and Applications*, 82(1):31–60, 2022.
- [18] Serge Gratton, Sadok Jerad, and Philippe L Toint. Complexity of a class of first-order objective-function-free optimization algorithms. *Optimization Methods and Software*, pages 1–31, 2024.
- [19] Serge Gratton, Sadok Jerad, and Philippe L Toint. Complexity of adagrad and other first-order methods for nonconvex optimization problems with bounds constraints. *arXiv preprint arXiv:2406.15793*, 2024.
- [20] Serge Gratton, Alena Kopaničáková, and Philippe L Toint. Multilevel objective-function-free optimization with an application to neural networks training. *SIAM Journal on Optimization*, 33(4):2772–2800, 2023.
- [21] Serge Gratton and Philippe L Toint. S2MPJ and CUTEST optimization problems for Matlab, Python and Julia. *Optimization Methods and Software*, pages 1–33, 2025.
- [22] Walter J Gutjahr and Alois Pichler. Stochastic multi-objective optimization: a survey on non-scalarizing methods. *Annals of Operations Research*, 236(2):475–499, 2016.
- [23] Nataša Krklec Jerinkić and Luka Rutešić. ASMOP: Additional sampling stochastic trust region method for multi-objective problems.
- [24] Diederik P Kingma. Adam: a method for stochastic optimization. *arXiv preprint arXiv:1412.6980*, 2014.
- [25] Matteo Lapucci and Pierluigi Mansueto. Improved front steepest descent for multi-objective optimization. *Oper. Res. Lett.*, 51(3):242–247, 2023.
- [26] Matteo Lapucci and Pierluigi Mansueto. A limited memory Quasi-Newton approach for multi-objective optimization. *Comput. Optim. Appl.*, 85(1):33–73, 2023.
- [27] Matteo Lapucci, Pierluigi Mansueto, and Davide Pucci. Effective front-descent algorithms with convergence guarantees. *arXiv preprint arXiv:2405.08450*, 2024.
- [28] Suyun Liu and Luis Nunes Vicente. The stochastic multi-gradient algorithm for multi-objective optimization and its application to supervised machine learning. *Annals of Operations Research*, 339(3):1119–1148, 2024.
- [29] Giampaolo Liuzzi, Stefano Lucidi, and Francesco Rinaldi. A derivative-free approach to constrained multi-objective nonsmooth optimization. *SIAM J. Optim.*, 26(4):2744–2774, 2016.
- [30] H Brendan McMahan and Matthew Streeter. Adaptive bound optimization for online convex optimization. *arXiv preprint arXiv:1002.4908*, 2010.
- [31] Margherita Porcelli, Giovanni Seraghi, and Philippe L. Toint. prunAdag: an adaptive pruning-aware gradient method. *Computational Optimization and Applications*, 93(1):85–119, 2026.
- [32] Jana Thomann. Decomposed descent methods in multiobjective optimization. Master’s thesis, Technische Universität Ilmenau, 2015.
- [33] Jana Thomann. *A trust region approach for multi-objective heterogeneous optimization*. PhD thesis, 2019. Technische Universität Ilmenau, Dissertation, 2019.
- [34] Jana Thomann and Gabriele Eichfelder. A trust-region algorithm for heterogeneous multiobjective optimization. *SIAM J. on Optimization*, 29(2):1017–1047, 2019.
- [35] Tijmen Tieleman and Geoffrey Hinton. Lecture 6.5-rmsprop, coursera: Neural networks for machine learning. *University of Toronto, Technical Report*, 6, 2012.
- [36] Flemming Topsøe. Some bounds for the logarithmic function. *Inequality theory and applications*, 4:137, 2007.
- [37] Kely D. V. Villacorta, Paulo R. Oliveira, and Antoine Soubeyran. A trust-region method for unconstrained multiobjective problems with applications in satisficing processes. *Journal of Optimization Theory and Applications*, 160(3):865–889, 2014.
- [38] Rachel Ward, Xiaoxia Wu, and Leon Bottou. Adagrad stepsizes: Sharp convergence over nonconvex landscapes. *Journal of Machine Learning Research*, 21(219):1–30, 2020.
- [39] Matthew D Zeiler. ADADELTA: an adaptive learning rate method. *arXiv preprint arXiv:1212.5701*, 2012.